

Strain-space model for Sars-CoV-2

Peter C. Jentsch, PhD ^{1,4} Finlay Maguire, PhD ^{3,5}
Samira Mubareka, MD, FRCPC ^{1,2}

¹Sunnybrook Research Institute, Toronto, Canada

²University of Toronto, Toronto, Canada

³Dalhousie University, Halifax, Canada

⁴Simon Fraser University, Burnaby, Canada

⁵Shared Hospital Laboratory, Toronto, Canada

April 14, 2022

- Dynamic model of Sars-CoV-2 evolution, representing antigenic diversity on a lattice (as in e.g. [Gog and Grenfell, 2002, Kryazhimskiy et al., 2007, Marchi et al., 2021])
- Antigenically distinct variants of the virus are mapped to 2D grid, distance between variants corresponds to the proportional reduction in maximum serum viral titre [Wilks et al., 2022, van der Straten et al., 2022]




Figure: Antigenic cartography of Sars-CoV-2, reproduced from [Wilks et al., 2022], Fig. 2


Model parameters/variables

Symbol	Description
N	Size of variant grid
S_{ij}	Population susceptible to variant $(i, j) \in [0, N]^2$
I_{ij}	Population infected by variant $(i, j) \in [0, N]^2$
R_{ij}	Recovered/Immune to variant $(i, j) \in [0, N]^2$
σ_{ijkl}	Probability that exposure to variant (i, j) causes immunity to variant (k, l)
β_{ij}	Transmission rate of variant (i, j)
ξ	Recovery rate of all strains
γ	Rate of immunity loss of all strains

Table: Table of symbols for Model 2

σ matrix

In practice, we assume σ_{ijkl} is just a 2-D gaussian distribution parameterized by the distance between (i, j) and (k, l) .



Model Equations

$$\frac{S_{ij}}{dt} = - \sum_{kl} \beta_{kl} \sigma_{ijkl} S_{ij} I_{kl} + \gamma R_{ij} \quad (1)$$

$$\frac{I_{ij}(t)}{dt} = \beta_{ij} S_{ij} I_{ij} - \xi I_{ij} + M (-4I_{ij} + I_{i-1,j} + I_{i+1,j} + I_{i,j-1} + I_{i,j+1}) \quad (2)$$

$$\frac{R_{ij}(t)}{dt} = \xi I_{ij} - \gamma R_{ij} \quad (3)$$

Boundary conditions: $I_{0,j} = 0, I_{j,0} = 0, I_{N,j} = 0, I_{j,N} = 0$
Initial conditions computed from genomic data in GISAID

- To incorporate more realistic mutation rates, we can go to continuous strain-space and use nonlocal reaction-diffusion dynamics as in [Rouzine and Rozhnova, 2018, Bessonov et al., 2021]

$$\frac{S_{ij}}{dt} = - \sum_{kl} \beta_{kl} \sigma_{ijkl} S_{ij} I_{kl} + \gamma R_{ij} \quad (4)$$

$$\frac{I_{ij}(t)}{dt} = \beta_{ij} S_{ij} I_{ij} - \xi I_{ij} + M (-4I_{ij} + I_{i-1,j} + I_{i+1,j} + I_{i,j-1} + I_{i,j+1}) \quad (5)$$

$$\frac{R_{ij}(t)}{dt} = \xi I_{ij} - \gamma R_{ij} \quad (6)$$



Bessonov, N., Bocharov, G., Meyerhans, A., Popov, V., and Volpert, V. (2021).

Existence and dynamics of strains in a nonlocal reaction-diffusion model of viral evolution.

SIAM Journal on Applied Mathematics, 81(1):107–128.



Gog, J. R. and Grenfell, B. T. (2002).

Dynamics and selection of many-strain pathogens.

Proceedings of the National Academy of Sciences, 99(26):17209–17214.



Kryazhimskiy, S., Dieckmann, U., Levin, S. A., and Dushoff, J. (2007).

On State-Space Reduction in Multi-Strain Pathogen Models, with an Application to Antigenic Drift in Influenza A.

PLOS Computational Biology, 3(8):e159.



Marchi, J., Lässig, M., Walczak, A. M., and Mora, T. (2021).

Antigenic waves of virus-immune coevolution.

Proceedings of the National Academy of Sciences,
118(27):e2103398118.



Rouzine, I. M. and Rozhnova, G. (2018).

Antigenic evolution of viruses in host populations.

PLoS Pathogens, 14(9):e1007291.



van der Straten, K., Guerra, D., van Gils, M., Bontjer, I.,
Caniels, T. G., van Willigen, H. D., Wynberg, E., Poniman,
M., Burger, J. A., Bouhuijs, J. H., et al. (2022).

Mapping the antigenic diversification of sars-cov-2.

medRxiv.



Wilks, S. H., Mühlemann, B., Shen, X., Türel, S.,
LeGresley, E. B., Netzl, A., Caniza, M. A.,
Chacaltana-Huarcaya, J. N., Daniell, X., Datto, M. B.,
Denny, T. N., Drosten, C., Fouchier, R. A. M., Garcia,
P. J., Halfmann, P. J., Jassem, A., Jones, T. C., Kawaoka,
Y., Krammer, F., McDanal, C., Pajon, R., Simon, V.,
Stockwell, M., Tang, H., van Bakel, H., Webby, R.,
Montefiori, D. C., and Smith, D. J. (2022).
Mapping SARS-CoV-2 antigenic relationships and
serological responses.
Preprint, Immunology.