# Characterizing evolutionary dynamics on a broader scale: a strain-space model for SARS-CoV-2

Peter C. Jentsch, PhD <sup>1,4</sup> Finlay Maguire, PhD <sup>3,5</sup> Samira Mubareka, MD, FRCPC <sup>1,2</sup>

<sup>1</sup>Sunnybrook Research Institute, Toronto, Canada

<sup>2</sup>University of Toronto, Toronto, Canada


<sup>3</sup>Dalhousie University, Halifax, Canada

<sup>4</sup>Simon Fraser University, Burnaby, Canada


<sup>5</sup>Shared Hospital Laboratory, Toronto, Canada

July 11, 2022

#### Infection spread with compartmental models




#### Modeling multiple infections is challenging




[Alizon and van Baalen, 2008]


# Complexity only increases with vaccination dynamics etc.



#### How to incorporate more data?



#### A useful approximation




J.R. Gog · J. Swinton


#### A status-based approach to multiple strain dynamics

Received: 5 April 2000 / Revised version: 24 July 2001 / Published online: 8 February 2002 – © Springer-Verlag 2002


## Begin with simple SIR model




#### Organize strains on a lattice




#### Strains mutate into adjacent strains



#### Infections provide cross-immunity to nearby strains




Cross-immunity to nearby strains




Mutation to adjacent strains

[Gog and Grenfell, 2002]

#### Full strain lattice model in 1 dimension




#### Extending the strain lattice to 2 dimensions




[Gog and Grenfell, 2002]

#### Extending the strain lattice to 2 dimensions




[Gog and Grenfell, 2002]

#### Quantifying antigenicity between viruses




### Projecting to low dimensions while preserving distances



[Pedregosa et al., 2011]

#### Projecting to low dimensions while preserving distances




[Pedregosa et al., 2011]


### Approximating genomic data in 2 dimesions



#### Approximating genomic data in 2 dimesions




#### Mapping SARS-CoV-2 in 2D




[Wilks et al., 2022]

#### Mapping SARS-CoV-2 in 2D




[Wilks et al., 2022]


# Methods of adding more genomes: polyclonal antibody binding studies




[Greaney et al., 2022]

#### Antibody Binding map







#### Methods of adding more genomes: homoplasic sites




[Page and Holmes, 2009]

#### Homoplasic mutations map





#### Evaluating the MDS approximation



#### Model parameters/variables

_	Symbol	Description
	N	Size of variant grid
	$S_{ij}$	Population susceptible to variant $(i, j) \in [0, N]^2$
	$I_{ij}$	Population infected by variant $(i, j) \in [0, N]^2$
	$R_{ij}$	Recovered/Immune to variant $(i, j) \in [0, N]^2$
	$\sigma_{ijkl}$	Probability that exposure to variant $(i, j)$ causes
		immunity
		to variant $(k, l)$
	$eta_{ij}$	Transmission rate of variant $(i, j)$
	v(t)	vaccination rate at time $t$
	s(t)	stringency at t
	ξ	Recovery rate of all strains
	$\gamma$	Rate of immunity loss of all strains

Table of symbols for Model 2

#### Model Equations

$$\frac{S_{ij}}{dt} = -\sum_{kl} s(t)\beta_{kl}\sigma_{ijkl}S_{ij}I_{kl} + \gamma R_{ij} - V(t)S$$
 (1)

$$\frac{I_{ij}(t)}{dt} = s(t)\beta_{ij}S_{ij}I_{ij} - \xi I_{ij} + M\left(-4I_{ij} + I_{i-1,j} + I_{i+1,j} + I_{i,j-1} + I_{i,j+1}\right) \tag{2}$$

$$\frac{R_{ij}(t)}{dt} = \xi I_{ij} - \gamma R_{ij} + V(t)S \tag{3}$$

Boundary conditions:  $I_{0,j} = 0, I_{j,0} = 0, I_{N,j} = 0, I_{j,N} = 0$ Initial conditions computed from genomic data in GISAID

#### Model Dynamics

(gifs of model dynamics)

#### Further work

- Differentiate between antigenic and genomic space
  - $\rightarrow$  non-local diffusion
- Model fitting
- Applications
  - ► Simple inference on antigenic space
  - ▶ VoC-aware NPI usage and vaccination

Alizon, S. and van Baalen, M. (2008).

Multiple infections, immune dynamics, and the evolution of virulence.

The American Naturalist, 172(4):E150–E168.

Gog, J. R. and Grenfell, B. T. (2002).

Dynamics and selection of many-strain pathogens.

Proceedings of the National Academy of Sciences,
99(26):17209–17214.

Greaney, A. J., Starr, T. N., and Bloom, J. D. (2022). An antibody-escape estimator for mutations to the sars-cov-2 receptor-binding domain.

Virus evolution, 8(1):veac021.

Lapedes, A. and Farber, R. (2001). The Geometry of Shape Space: Application to Influenza. Journal of Theoretical Biology, 212(1):57–69.

- Page, R. D. and Holmes, E. C. (2009).

  Molecular evolution: a phylogenetic approach.

  John Wiley & Sons.
- Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D., Brucher, M., Perrot, M., and Duchesnay, E. (2011). Scikit-learn: Machine learning in Python.

  Journal of Machine Learning Research, 12:2825–2830.
- Smith, D. J., Lapedes, A. S., de Jong, J. C., Bestebroer, T. M., Rimmelzwaan, G. F., Osterhaus, A. D. M. E., and Fouchier, R. A. M. (2004).

Mapping the Antigenic and Genetic Evolution of Influenza Virus.

Science, 305(5682):371-376.

Wilks, S. H., Mühlemann, B., Shen, X., Türeli, S., LeGresley, E. B., Netzl, A., Caniza, M. A., Chacaltana-Huarcaya, J. N., Daniell, X., Datto, M. B., Denny, T. N., Drosten, C., Fouchier, R. A. M., Garcia, P. J., Halfmann, P. J., Jassem, A., Jones, T. C., Kawaoka, Y., Krammer, F., McDanal, C., Pajon, R., Simon, V., Stockwell, M., Tang, H., van Bakel, H., Webby, R., Montefiori, D. C., and Smith, D. J. (2022). Mapping SARS-CoV-2 antigenic relationships and serological responses. Preprint, Immunology.

#### Mutation homoplasy

