

# Coupled models of structured contagion processes in human-environment systems

Peter C. Jentsch

Department of Applied Mathematics, University Of Waterloo

August 6th, 2021

Supervised by Chris T. Bauch and Madhur Anand

Peter C.  
Jentsch

Introduc-  
tion

First  
Project

Second  
Project

Third  
Project

References

# Introduction

Introduction

First  
Project

Second  
Project

Third  
Project

References

- A contagious process is one that is able to propagate itself through a set of hosts
- Contagious processes shape our lives in many important ways
- Current situation has thrust the importance of modeling infections into popular consciousness

Introduction

First  
Project

Second  
Project

Third  
Project

References

- Coupling can often dramatically affect stability, e.g. coupled oscillators
- Human-environment systems focus on dynamics of humanity and the environment viewed as a whole
- This perspective provides useful answers to important problems

Peter C.  
Jentsch

Introduc-  
tion

First  
Project

Second  
Project

Third  
Project

References

# First Project

Introduction

First  
Project

Second  
Project

Third  
Project

References

- Population beliefs about the usage of non-pharmaceutical interventions (NPIs) can have a major effective on the epidemic landscape
- Population age distribution is a key factor in both NPI usage and COVID-19 mortality
- Vaccine rollout important to reducing mortality in the long term (Bubar et al., 2020; Buckner, Chowell, and Springborn, 2020)

## Outline of project

- ① Describe age-structured compartmental model of COVID-19 infection in a population, coupled to model of social distancing dynamics.
- ② Fit model to data from Ontario, Canada
- ③ Evaluate outcomes from different vaccination strategies, comparing vaccination of vulnerable populations vs vaccination of susceptible populations

## Compartmental model overview

Introduction  
First Project  
Second Project  
Third Project  
References

### Disease Compartments

$S_i(t)$  : Susceptible  
 $S_{2,i}(t)$  : Vaccinated but still susceptible  
 $V_i(t)$  : Vaccinated and immune  
 $E_i(t)$  : Exposed  
 $P_i(t)$  : Pre-symptomatic  
 $I_{a,i}(T)$  : Infectious and asymptomatic

$I_{s,i}(t)$  : Infectious and symptomatic

$R_i(t)$  : Recovered

where  $i = 1 \dots 16$  comprises age structure

### Social compartments

$x(t)$  : Uses NPIs

$1 - x(t)$  : Does not use NPIs




Figure: Compartments

## Game theory as a model of NPI adoption

Introduction

First Project

Second Project

Third Project

References

		P2	use NPI	don't use NPI
		P1	use NPI	med risk
		use NPI	low risk, NPIs unpleasant	med risk, NPIs unpleasant
		use NPI	low risk, NPIs unpleasant	med risk, NPIs unpleasant
		don't use NPI	med risk, NPIs unpleasant	high risk
		don't use NPI	med risk	high risk

Table: NPI adoption as a two-player game (between P1 and P2)


## Vaccination strategies

We compare four vaccination strategies

- > 60 first
- < 20 first
- Uniform
- Contact-based

with respect to reduction in cumulative mortality after 5 years.

Contact-based vaccination strategy



**Figure: Vaccine allocation by age in the contact-based vaccination strategy.**  
Computed as the normalized leading eigenvector of the sum of the contact matrices.

## Parameterization

Introduction  
First Project  
Second Project  
Third Project  
References




Figure: Model parameterization used, physical distancing matches reported cases.

## Results

Introduction

First  
Project

Second  
Project

Third  
Project

References





Figure: Percentage reduction in mortality for four strategies depends on vaccination start date and the vaccination rate.

Vaccination begins on Jan 1, 21, 1.5% of pop. vaccinated per week



Vaccination begins on Jan 1, 21, 0.5% of pop. vaccinated per week





Figure: (a) timely vaccination prevents third wave, (b) partial vaccination and indirect protection help during the third wave

Vaccination begins on Sep 1, 21, 1.5% of pop. vaccinated per week



Vaccination begins on Sep 1, 21, 0.5% of pop. vaccinated per week




Figure: c) with a later start date but higher vaccination rate, partial vaccination and indirect protection help during the third wave as in b), (d) slow and late vaccination fails to prevent third wave.

## Results


Introduction

First  
Project

Second  
Project


Third  
Project

References



**Figure:** Later start to vaccination favours transmission-interrupting strategies for moderate vaccination rates. Each parameter pair is colored according to the strategy that prevents most deaths on average, over all realizations of the model.

## Results



**Figure:** More pre-existing natural immunity makes transmission-interrupting strategies more effective. Histogram of no. recovered at vaccination begin date, according to best strategy for that realization, over all parameter values in sensitivity analysis. Vertical lines are the median.

Introduction

First  
Project

Second  
Project

Third  
Project

References

- We described an age structured compartmental model of Sars-CoV-2 infection and vaccination coupled to a social model
- Showed that sometimes transmission interrupting strategies can be more effective
- Depends on the pre-existing immunity in the population

Peter C.  
Jentsch

Introduc-  
tion

First  
Project

Second  
Project

Third  
Project

References

## Second Project

Introduction

First  
Project

Second  
Project

Third  
Project

References

- Invasive forest pests cause incredible damage to ecosystems and lumber resources
- Evidence shows that movement of firewood is a major long distance vector (Koch et al. (2014))
- Education and awareness is a major way we try to reduce this vector



Figure: an Emerald Ash Borer, which devastated Ash populations in North America

Introduction

First  
Project

Second  
Project

Third  
Project

References

- ① Adapt a model such as Barlow et al. (2014) to a larger, more realistic network
- ② Use model to compare three possible prevention measures
  - Education/awareness
  - Inspection of moved firewood
  - Quarantine of highly susceptible forest patches
- ③ Assess measures across a range of parameter values and time horizons

Introduction

First Project

Second Project

Third Project

References

$$\frac{dS_i}{dt} = \underbrace{rS_i \left(1 - \frac{(S_i + I_i)}{K}\right)}_{\text{Logistic Growth Of Forest}} - \underbrace{AS_i(I_i + B_i)\theta_k(I_i - I_a)}_{\text{Infestation term}} \quad (1)$$

$$\frac{dI_i}{dt} = \underbrace{-\gamma I_i}_{\text{Death of infested trees}} + \underbrace{AS_i(I_i + B_i)\theta_k(I_i - I_a)}_{\text{Susceptibles become infested}} - d \underbrace{\sum_{j=1, j \neq i}^N P_{j,i}(1 - C_e)(1 - L_j)I_j}_{\text{Total infested wood leaving due to transport}} \quad (2)$$

$$\frac{dB_i}{dt} = \underbrace{-\gamma B_i}_{\text{Decay of firewood}} + d \underbrace{\sum_{j=1, j \neq i}^N P_{i,j}(1 - C_e)(1 - L_j)I_j}_{\text{Import of fallen wood}} \quad (3)$$

$$\frac{dL_i}{dt} = \sigma L_i(1 - L_i) \left( \underbrace{U}_{\text{Net cost to transport firewood}} + \underbrace{s(2L_i - 1)}_{\text{Social influence term}} + \underbrace{fI_i}_{\text{Impact of infestation}} \right) \quad (4)$$

We use  $T_i(t)$ , computed from equation 5 to be the total number of infested trees in patch  $i$  up to time  $t$ .

$$\frac{dT_i}{dt} = AS_i(I_i + B_i)\theta_k(I_i - I_a) \quad (5)$$

Define  $T(t) = \frac{1}{N} \sum_{i=1}^N T_i(t)$  to be the average total number of infested trees up to  $t$ .

Introduction

First Project

Second Project

Third Project

References




Figure: Travel network used to weight edges in firewood transport network

## Results


Introduction

First  
Project

Second  
Project

Third  
Project


References



**Figure:** Neither increasing  $U$  nor  $C_e$  are effective at long time scales. Total number of infested trees per node over 5 (a), 10 (b), and 20 (c) years

## Results

Introduction  
First Project  
Second Project  
Third Project  
References



**Figure:** The influence of infestation on transport strategy,  $f$ , can hinder the intervention by public outreach, in the long-term (after approximately 20 years). Efficacy of social incentives on infestation after time  $T$ . Inset graph shows an example of cross-section along the line  $f = 0.11$

Introduction

First Project

Second Project

Third Project

References




Figure: Increasing  $U$  becomes ineffective over time if  $A$  is sufficiently large. Efficacy of social incentives on infestation after time period  $T$  with respect to  $A$ , the intra-patch infestation parameter.

## Results


Introduction

First  
Project

Second  
Project

Third  
Project

References



**Figure:** The social incentive to not transport firewood,  $U$ , is more effective with lower pest spread rates. Efficacy of social incentives on infestation after time  $T$  intra-patch spreading rate  $A$ , affects infestation outcomes.

## Results


Introduction

First  
Project

Second  
Project

Third  
Project

References



**Figure:** Quarantine needs to include many patches for long term effects. Average total infested trees ( $T(t)$ ) after 5, 10 and 15 years (panels a), b), and c) respectively), assuming the quarantine begins one year after the pest is introduced.

Introduction

First  
Project

Second  
Project

Third  
Project

References

- Firewood inspection not likely to be effective in implementation
- Education, represented as social incentives, are able to decrease infection in the short term, but dependent on pest-specific parameters
- Patch quarantine can be effective if sufficiently many patches are isolated, and the pest is detected early

Peter C.  
Jentsch

Introduc-  
tion

First  
Project

Second  
Project

Third  
Project

References

## Third Project

Introduction

First  
Project

Second  
Project

Third  
Project

References

- Wildfire and bark beetles are disturbances integral to coniferous forest ecosystems in the western cordillera of North America (Kaufmann et al., 2008)
- Bark beetle outbreaks have always been destructive, but seem to be worse in recent decades
- Literature on causal relationship between bark beetle outbreaks and wildfire is extensive but inconclusive (Axelson, Alfaro, and Hawkes (2009))
- Existing modelling of these two coupled disturbances is sparse

Introduction

First  
Project

Second  
Project

Third  
Project

References

- Extend existing model of Duncan et al. (2015) to include wildfires
- Explore parameter regime of extended model
- Introduce forest stand thinning procedures to reduce MPB outbreaks
- Show that these stand thinning procedures are able to work due to increased stand heterogeneity

Introduction

First  
ProjectSecond  
ProjectThird  
Project

References

$$j_{n+1,1} = dJ_n + I_{n-2} + F_n \quad (6a)$$

$$j_{n+1,k} = (1 - d)j_{n,k-1} - \frac{\alpha_1}{T} P_n j_{n,k-1}, \quad k = 2 \dots K-1, K \quad (6b)$$

$$S_{n+1} = S_n + (1 - d)j_{n,K} - \left( I_n + \frac{\alpha_2}{T} P_n I_n \right) - \frac{\alpha_2}{T} P_n (S_n + (1 - d)j_{n,K}) - \sigma_F \xi_n \quad (6c)$$

$$I_{n+1} = r_1 I_n e^{-\beta_1(T-S_{n+1})} - \frac{\alpha_2}{T} P_n I_n + \sigma_I \xi_n \quad (6d)$$

$$F_{n+1} = P_n \left[ \frac{\alpha_1}{T} \sum_{k=1}^{K-1} j_{n,k} + \frac{\alpha_2}{T} (S_n + (1 - d)j_{n,K}) + \frac{\alpha_2}{T} I_n \right] + \sigma_F \gamma_n \quad (6e)$$

$$P_n = T - \sum_{i=1}^n F_i e^{-\kappa(n-i)} \quad (6f)$$

## Results

Introduction

First Project

Second Project

Third Project

References




Figure: Increasing burning rates  $\alpha_1, \alpha_2$  reduce MPB outbreak size, cause stability in fire regimes. Panels: a) Average size of largest MPB population, b) Average frequency of MPB outbreaks, c) Average size of largest fire season, d) Average frequency of severe fire seasons. All measured at equilibrium.

## Results

### Introduction

### First Project

### Second Project

### Third Project

### References




Figure: Without FTP,  
 $\alpha_1 = 0.02$ ,  $\alpha_2 = 0.0025$ , large  
even aged stands visible in  
juvenile age distribution (top  
panel).




Figure: With FTP,  
 $\alpha_1 = 0.02$ ,  $\alpha_2 = 0.0025$ ,  
 $\tau = 0.15$ ,  $m = 8$ , with greater  
heterogeneity in juvenile age  
structure (top panel).

## Results


Introduction

First Project

Second Project

Third Project

References



**Figure:** FTP most effective with small  $\alpha_1$ , CBP most effective with small  $\alpha_1, \alpha_2$ . a)  $\tau = 0.15, m = 8$ , b) with  $\tau = 0.15, m = 8$  applied every 5 years, c) controlled burning with  $\tau = 0.15, m = 8$ , with respect to burning rates  $\alpha_1, \alpha_2$ .

Introduction

First  
Project

Second  
Project

Third  
Project

References

- We show that increasing fire prevalence is able to dampen MPB outbreaks by increasing stand heterogeneity
- Stand heterogeneity can also be increased through forest thinning or prescribed burning, which also dampens MPB outbreaks.
- These results are consistent with ecological evidence (Seidl et al. (2016) and Kaufmann et al. (2008))

Introduction

First  
Project

Second  
Project

Third  
Project

References

- Three models of coupled human-environment systems
- Compared methods of mitigation for respective coupled infectious processes
- Results obtained from these answer a gap identified in literature

Introduc-  
tion

First  
Project

Second  
Project

Third  
Project

References

Thank you!

-  Axelson, Jodi N, René I Alfaro, and Brad C Hawkes (2009). “Influence of fire and mountain pine beetle on the dynamics of lodgepole pine stands in British Columbia, Canada”. In: *Forest Ecology and Management* 257.9, pp. 1874–1882.
-  Barlow, Lee-Ann et al. (2014). “Modelling interactions between forest pest invasions and human decisions regarding firewood transport restrictions”. In: *PLoS one* 9.4, e90511.
-  Bubar, Kate M et al. (2020). “Model-informed COVID-19 vaccine prioritization strategies by age and serostatus”. In: *medRxiv*.
-  Buckner, Jack H, Gerardo H Chowell, and Michael R Springborn (2020). “Optimal Dynamic Prioritization of Scarce COVID-19 Vaccines”. In: *medRxiv*.

-  Duncan, Jacob P et al. (2015). “A model for mountain pine beetle outbreaks in an age-structured forest: Predicting severity and outbreak-recovery cycle period”. In: *Bulletin of mathematical biology* 77.7, pp. 1256–1284.
-  Kaufmann, Merrill R et al. (2008). “The status of our scientific understanding of lodgepole pine and mountain pine beetles: a focus on forest ecology and fire behavior”. In.
-  Koch, Frank H et al. (2014). “Using a network model to assess risk of forest pest spread via recreational travel”. In: *PLoS one* 9.7, e102105.
-  Seidl, Rupert et al. (2016). “Spatial variability in tree regeneration after wildfire delays and dampens future bark beetle outbreaks”. In: *Proceedings of the National Academy of Sciences*, p. 201615263.