

Detailed Oncology Consultation Note Regarding Progression

Metropolis Comprehensive Cancer Center – Thoracic Oncology Service

DATE OF CONSULTATION: March 11, 2022

CONSULTING PHYSICIAN: Kenji Tanaka, MD (Medical Oncology)

REFERRING PHYSICIAN: Self (Established Patient)

REASON FOR CONSULTATION: Re-evaluation following routine surveillance imaging demonstrating disease progression after a prolonged response (>22 months) to first-line Pembrolizumab monotherapy. Discussion of second-line treatment options.

HISTORY OF PRESENT ILLNESS:

Mr. Smith is a 75-year-old gentleman with a history of Stage IV NSCLC (Adenocarcinoma confirmed via adrenal biopsy), diagnosed on March 9, 2020. His initial presentation was marked by significant fatigue and unintentional weight loss. Staging PET/CT revealed a 3.8 cm LUL primary lesion with mild FDG uptake, significant bilateral adrenal metastases (Right 5.5 cm, Left 4.8 cm, both intensely FDG-avid), and multiple sub-centimeter hepatic lesions highly suspicious for metastases (largest 1.2 cm in segment VI). Brain MRI was negative. Molecular profiling (NGS via Tempus xT on adrenal biopsy) was notable for **Wild-Type** status (EGFR/ALK/ROS1/BRAF/KRAS/MET/RET/NTRK negative). PD-L1 expression (IHC 22C3) was high: **TPS 70%, CPS 75, IC Score 3/+**. TMB was 7 mut/Mb.

Given his high PD-L1 expression and WT status, he commenced first-line **Pembrolizumab 200 mg IV every 3 weeks starting April 1, 2020**. He experienced an excellent and durable response. Clinically, his fatigue resolved within 2-3 months. Radiographically, he achieved a deep partial response by 6 months: the LUL primary became sub-centimeter and non-avid; adrenal metastases significantly decreased (nadir R 1.8 cm, L 1.5 cm, low avidity); hepatic lesions resolved completely. He tolerated Pembrolizumab extremely well, with no significant immune-related adverse events over nearly two years.

He remained stable in deep partial response on Pembrolizumab maintenance. Routine surveillance **CT Chest/Abdomen/Pelvis w/ contrast performed March 1, 2022**, compared to scans from November 15, 2021, demonstrated the first evidence of disease progression:

- **Adrenals:** Definite increase in size of both adrenal metastases. Right adrenal now measures 3.1 x 2.9 cm (vs. 1.9 x 1.7 cm). Left adrenal now measures 2.8 x 2.5 cm (vs. 1.6 x 1.4 cm). Both show increased central heterogeneity and subtle increase in peripheral enhancement.
- **Liver:** Re-appearance of several previously resolved hepatic lesions, now measuring 0.8 – 1.5 cm in segments VI, VII, and VIII. Appearance of 2 definitively new small hepatic lesions (<1 cm).
- **Chest:** LUL primary lesion remains stable/minimal (<1 cm). No new pulmonary nodules or concerning lymphadenopathy.
- **Impression:** Disease progression involving original adrenal sites and recurrence/progression in the liver after approximately **23 months** of Pembrolizumab therapy.

SUBJECTIVE (Today): Patient presents for discussion of these findings. He reports feeling generally well, denying any new or worsening symptoms including fatigue, pain, nausea, or weight loss. He was surprised

PATIENT: Smith, John Robert

MRN: SYN160 **DOB:** December 14, 1947

by the scan results given his continued well-being. Performance Status remains excellent, ECOG 0. He is prepared to discuss next steps.

PAST MEDICAL HISTORY: Coronary Artery Disease s/p CABG x 3 (2008, stable), Hypertension (on Lisinopril, Metoprolol), Hyperlipidemia (on Atorvastatin), GERD (on Pantoprazole). Former smoker (40 pack-years, quit 2000).

CURRENT MEDICATIONS: Pembrolizumab (discontinued), Lisinopril 20mg daily, Metoprolol Succinate ER 50mg daily, Atorvastatin 40mg daily, Pantoprazole 40mg daily, Aspirin 81mg daily.

REVIEW OF SYSTEMS: Entirely negative.

OBJECTIVE:

- *Vitals:* T 36.9, BP 130/76, HR 62, SpO2 97% RA. Wt stable. ECOG PS 0.
- *Exam:* Alert, vigorous appearing male. Exam completely unremarkable.
- *Relevant Labs (Baseline Mar 2020 → Pre-Prog Nov 2021 → Current Mar 2022):*
 - ALT: 25 → 18 → 22 U/L
 - AST: 20 → 15 → 19 U/L
 - ALP: 90 → 75 → 80 U/L
 - Total Bili: 0.7 → 0.5 → 0.6 mg/dL
 - Albumin: 4.0 → 4.1 → 4.0 g/dL
 - Creatinine: 1.0 → 1.1 → 1.0 mg/dL
 - Hgb: 13.5 → 14.2 → 14.0 g/dL
 - TSH: 2.1 → 1.8 → 2.3 mIU/L (remained euthyroid)

ASSESSMENT:

1. **Stage IV Lung Adenocarcinoma (WT, PD-L1 High):** Acquired resistance and disease progression documented after an exceptionally long response (23 months) to first-line Pembrolizumab. Progression sites are liver and adrenals. Patient remains asymptomatic with excellent performance status (ECOG 0).
2. **Comorbidities:** Stable.

PLAN:

1. **Discontinue Pembrolizumab:** Acknowledged excellent duration of benefit.
2. **Second-Line Systemic Therapy:** Standard of care is platinum-based doublet chemotherapy.
 - **Recommended Regimen: Carboplatin (AUC 5) + Pemetrexed (500 mg/m²) IV q3 weeks** for 4-6 cycles, followed by potential Pemetrexed maintenance if disease controlled/responding. Rationale: Adenocarcinoma histology, good PS, favorable toxicity profile relative to alternatives.
 - **Discussion:** Reviewed expected side effects (myelosuppression, fatigue, nausea, renal monitoring, B12/Folate need), treatment schedule, and palliative intent thoroughly. Patient motivated to proceed.

PATIENT: Smith, John Robert

MRN: SYN160 **DOB:** December 14, 1947

3. **Initiation Plan:**

- Target C1D1 start within 1-2 weeks, pending insurance verification.
- **Start Folic Acid 1 mg PO Daily today.**
- Schedule B12 1000 mcg IM injection prior to C1D1.
- Provide anti-emetic prescriptions (Ondansetron, Prochlorperazine).
- Schedule formal chemo education visit.

4. **Continue all other chronic medications.**

5. **Monitoring:** Labs (CBC w diff, CMP) prior to each cycle. Restaging CT C/A/P after 2-4 cycles.
Continue Brain MRI surveillance q6 months (last was Nov 2021 - neg; next due May 2022).

6. **Follow-up:** Return for chemo education and C1D1 infusion as scheduled.

M.D.
Kenji Tanaka, MD (Electronically Signed)

PATIENT: Smith, John Robert

MRN: SYN160 **DOB:** December 14, 1947