

DENDRITIC SPINES: THE LOCUS OF STRUCTURAL AND FUNCTIONAL PLASTICITY

Carlo Sala and Menahem Segal

Consiglio Nazionale delle Ricerche Institute of Neuroscience and Department of Medical Biotechnology and Translational Medicine, University of Milan, Milan, Italy; Neuromuscular Diseases and Neuroimmunology, Neurological Institute Foundation Carlo Besta, Milan, Italy; and Department of Neurobiology, The Weizmann Institute, Rehovot, Israel



Sala C, Segal M. Dendritic Spines: The Locus of Structural and Functional Plasticity. *Physiol Rev* 94: 141–188, 2014; doi:10.1152/physrev.00012.2013.—The introduction of high-resolution time lapse imaging and molecular biological tools has changed dramatically the rate of progress towards the understanding of the complex structure-function relations in synapses of central spiny neurons. Standing issues, including the sequence of molecular and structural processes leading to formation, morphological change, and longevity of dendritic spines, as well as the functions of dendritic spines in neurological/psychiatric diseases are being addressed in a growing number of recent studies. There are still unsettled issues with respect to spine formation and plasticity: Are spines formed first, followed by synapse formation, or are synapses formed first, followed by emergence of a spine? What are the immediate and long-lasting changes in spine properties following exposure to plasticity-producing stimulation? Is spine volume/shape indicative of its function? These and other issues are addressed in this review, which highlights the complexity of molecular pathways involved in regulation of spine structure and function, and which contributes to the understanding of central synaptic interactions in health and disease.

I.	INTRODUCTION	141
II.	METHODOLOGICAL CONSIDERATIONS	143
III.	ELECTROPHYSIOLOGICAL PROPERTIES OF ...	144
IV.	MOLECULAR CONSTRUCTION OF ...	148
V.	DEVELOPMENT OF DENDRITIC SPINES	163
VI.	SHORT-TERM MORPHOLOGICAL AND ...	164
VII.	LONG-TERM CHANGES AND DENDRITIC...	165
VIII.	MOLECULES IN SYNAPTIC PLASTICITY	167
IX.	DISEASE-RELATED CHANGES IN...	168
X.	PERSPECTIVES	173

I. INTRODUCTION

Ever since their first detection by Ramon y Cajal (408), dendritic spines have been postulated to underlie the neuronal locus of plasticity, where short-term alterations in synaptic strength are assumed to be converted to long-lasting memories that are embedded in stable morphological changes. Initially, dendritic spines were visualized using Golgi impregnation. This was followed in the late 1950s by electron microscopic (EM) reconstruction of neuronal tissue at the nanometer resolution. The EM picture introduced us to the postsynaptic density, the presynaptic terminal and its vesicles, and other organelles found in (e.g., spine apparatus) or absent from (e.g., mitochondria) the dendritic spine. Still, the search for morphological changes in dendritic spine following plasticity-producing stimulation was tedious, and the results were not always consistent. In one of the earliest EM studies, Fifkova and Van Harreveld (128) found that within 2–6 min after a

long (30 s) tetanic stimulation of the perforant path, spines in the dentate gyrus increased their size by 15%, followed by a further increase by 38% at 10–60 min after the stimulation. On the other hand, Desmond and Levi (101) found that the postsynaptic density (PSD) surface area per synapse increased markedly, but they did not report on a change in spine head volume. Geinisman (141) reported that following long-term potentiation (LTP) there is an increase in number of perforated synapses, with the PSD breaking down to two independent components, thus increasing the size of the overall synaptic contact. In a similar series of studies, using advanced morphometric three-dimensional reconstructed EM images, Stewart and colleagues (484) found that following chemical LTP, which presumably activates all synapses in the sliced hippocampus, there is an increase in the number of perforated synapses without a change in volume of the spines. In other studies by the same group (395) with anesthetized rats, a significant increase in thin and mushroom spine volumes was found, but without a change in synapse density. In further studies, they (322) describe an increase in perforated synapses in the dentate gyrus some 24 h after the tetanic stimulation in awake rats. In sharp contrast, Sorra and Harris (474), measuring three-dimensional reconstructed spines from serial section EM pictures, could not find any significant effect of LTP on morphological properties of spines. The different reported effects of LTP on spine properties are partly due to methodological differences, including the nature of stimulation used for generating plasticity, the brain area chosen for study, the condition of the tissue (in vivo vs. in vitro), the time after the

stimulation, and the methods of analysis. These are not trivial considerations, as there are reports that handling of the sliced tissue (e.g., cooling, blocking activity with drugs) may have considerable effect on dendritic spine density (238), regardless of the response to a tetanic stimulation, and probably irrespective of the physiological state of the tissue.

A different approach to the study of the role of dendritic spines in synaptic plasticity examined the effects of enriched environment on morphological and functional attributes of neural tissue, including dendritic spines. In one of the earlier studies, Globus et al. (147) found that allowing rodents to develop in an enriched environment resulted in an increase in dendritic spine density. In these early studies it was not possible to separate between direct effects on dendritic spines and global effects on the neural tissue, including an increase in blood supply, glia proliferation, and an increase in volume of the cortex. The power of the novel time lapse *in vivo* imaging for addressing the same issue studied by Globus et al. is exemplified in a recent study of the effects of enriched environment on dendritic spine density (219). The authors demonstrate that not only spine density is increased following life in an enriched environment, but there is also an increase in transient spines, and in turnover rate of spines. These observations are highly relevant to the assumed role of spines in storage of memories (see sects. VI and VII).

While the morphometric studies that compared populations of fixed spines have yielded important information, in that they indicated that dendritic spines undergo some morphological changes following an intense experience, be it a tetanic stimulation, growth conditions, hibernation, and hormonal and emotional variations, they were not so informative as to the exact changes that a given spine can undergo following such an intense experience, and the molecular mechanisms leading to these changes.

The ability to follow individual dendritic spines over time, using *in vivo* time-lapse imaging, first introduced two decades ago (160, 343), allowed a rapid progress in the description and analysis of changes that may take place in dendritic spines in response to plasticity-producing stimulation. The advantages of the time lapse imaging over the more traditional counting and measuring of different populations of spines are obvious; repeated-measures statistics are stronger than comparisons among different samples, especially in cases of great heterogeneity of spine shapes. On the other hand, the time lapse imaging is inherently restricted to a smaller simultaneous samples of spines likely to be located on the same segments of the dendrite, which may not represent the population of spines at large, and is restricted to light microscopic resolution, which is blind to possible changes in subsynaptic organelles.

Regardless of the pitfalls of the time-lapse live imaging, there was a tremendous increase in the amount of information accumulated in the past decade about the mechanisms and roles

of dendritic spines in neuronal plasticity, as evidenced by the large number of review articles published in the year of 2012 (53, 121, 129, 258, 306, 364, 386, 416, 427, 517 to list a few).

Despite the rapid accumulation of information on the molecular and functional attributes of dendritic spine regulation, some standing issues are still unsettled: Is spine formation triggered by incoming afferents or by an intrinsic process independent of afferent activity; does it develop from a filopodium that shrinks or from a dendritic shaft synapse that elongates to become a spine? Indications for both of these possibilities have been reported, but it is not entirely clear what is the dominant mode of spine formation, if there is any. Obviously, young neurons are endowed with filopodia, so it is likely that the initial formation of spines is caused by conversion from filopodia, whereas in adult neurons, where there is also a significant generation of novel spines, they are not preceded by filopodia. Exceptions to this generalization can also be found; when neurons fail to receive input, spines tend to shoot out protrusions, probably in search of glutamate excitation (415).

Likewise, once a spine is formed, how long does it take to become functional, and to react to presynaptic stimulation? There are reports of different time scales, from minutes to days and weeks, but the verdict is not out yet. Along the same line, once a spine is formed, how stable it is, and what might cause it to disappear/shrink, and at what rate? Finally, given the large number of different families of molecules assumed to be necessary for the formation of spines, are all spines equipped with the same molecular cascades, or are different spines endowed with different molecules, which may cause different morphologies and perhaps functions?

A. What Are the Functional Demands on "Plastic" Spines?

The key issue in spine research is the functional correlates of the great heterogeneity in spine morphologies. Why are there short and long, stubby and mushroom spines? Do they serve different functions, or are only the "mature" ones functional, and the other types "reserve" contacts, to be used upon request? On the functional end, the attempt to narrow down the wealth of different memory mechanisms into a minimal number of simple electrophysiological and molecular "building blocks" within dendritic spines is at best only a crude approximation. At present, there are a variety of memory processes: working, procedural, declarative, semantic, and perceptual, and they are associated with different brain areas and different time scales and may utilize different molecular cascades (326). In fact, despite extensive studies utilizing pharmacology and genetic manipulation of specific genes related to plasticity, there is still no clear link between different plasticity processes and spine morphology. It is plausible that different types of memories obey different electrophysiological rules and may leave different morphological fingerprints in the brain.

At the cellular level, there are different plasticity-producing stimulation patterns that are likely to use different molecular cascades and morphological changes. For example, while the dominant mode of producing LTP in brain areas such as the hippocampus involves activation of the N-methyl-D-aspartate (NMDA) receptor (168, 186), there is a distinct non-NMDA mediated LTP, which is assumed to activate release of calcium from stores (156, 410). Is it possible that these two ways to generate LTP are funneled into the same morphological changes in the spine? Also, spike timing dependent plasticity (STDP) (121), metaplasticity, and chemical LTP (448) are all assumed to involve changes in distributions of glutamate receptor subtypes following a change in intracellular calcium concentration and affect synaptic responses for extended periods of time. Are all these plastic processes sharing the same locus of change? Finally, there are strong indications that plastic changes take place also at the presynaptic terminal, and involve changes in release probability (113, 116), as well as appearance and disappearance of synaptic boutons (34). Furthermore, some plastic changes involve nonsynaptic mechanisms, including changes in glia properties (352) and dendritic excitability (131). These may not necessarily involve direct changes in spine morphology.

B. Longevity of Memories

Different types of memories have different rules and time course; motor memory (speech, bicycle ride, piano playing) is in general persistent, while working memory is highly labile. Can we assume that both types are “stored” in dendritic spines? Recent studies on motor learning *in vivo* addressed these issues, have been reviewed extensively (193, 243), and are discussed below (see sects. VI and VII).

One interesting diversion from the traditional view of memories being stores in stable dendritic spines are recent indications that ongoing phosphorylation state of a molecule (e.g., PKMzeta, Ref. 429) controls apparently stable memories. If so, are stable spines necessary for stable memories, or are these morphological changes an epiphenomenon, and some specific molecules are the ones to carry the “memories”?

II. METHODOLOGICAL CONSIDERATIONS

A. Labeling of Neurons

One of the mysteries of the Golgi stain is its selectivity. Over a century after its discovery, it is still not known why so few neurons stain with the Golgi method, and if there is something unique about the ones that do stain, that may bias the evaluation of changes produced by behavioral/chemical manipulations in the brain. A similar issue is not addressed systematically; with the use of transfection/infection methodologies, a favorite method for imaging individual neurons *in vivo* as well as in cultured brain slices and dissociated neuronal cultures,

only a small fraction of neurons take up the plasmid and express the protein of interest, e.g., green fluorescent protein (GFP). Furthermore, once the neuron takes up the plasmid, and if the promoter is ubiquitous, it is likely to recruit the protein-generating cellular machinery for producing large amounts of the transfected molecules. Intuitively, this should suppress the neuronal responses to other demands for proteins. However, this issue has not been addressed systematically, except to show that adjacent, transfected and nontransfected neurons share the same electrical properties (458, 504).

B. Estimation of Spine Dimensions From Fluorescent Images

A large number of recent publications use time-lapse imaging to estimate changes in spine dimensions following an acute exposure to plasticity-producing stimulation, e.g., caged glutamate. Some of these studies demonstrate immediate and striking, up to threefold, increases in spine volume (311). However, these changes may result partially from changes in fluorescence intensity rather than size of the imaged spine. The earlier studies believe to have accounted for this change in fluorescence intensity, and a tentative correlation between fluorescence intensity and spine size has been documented [see Holtmaat et al. (194), 10 spines imaged and reconstructed at the EM level], but this issue is not addressed systematically, and there is a clear increase in size of the fluorescent spot upon an increase in fluorescence intensity. One way to deal with these two possible sources of change, without being committed to one or the other, is to express the change as a “normalized fluorescence intensity” (583). Obviously, even if the measured change is in fluorescence intensity rather than size of the spine, it is a dramatic change that should be analyzed.

GFP-tagged molecules are used routinely in the study of localization, dynamics, and mobility of molecules following induction of changes in response to electrical or chemical stimulation. For example, GFP-tagged PSD-95 (e.g., Ref. 110), marks the postsynaptic side of the synapse. The postsynaptic density, as seen in EM pictures, is far smaller than the corresponding fluorescent image of GFP-PSD-95 (427), indicating that the fluorescent image provides by far an overestimation of organelle size, and is most likely affected by the amount of light emitted by the fluorophore. This has been confirmed in recent imaging methods using super-resolution microscopy (see sect. X and Refs. 298, 345).

C. Dendritic Spine Density

What is the “normal” dendritic spine density? The numbers range from 0.2 to 3.5 spines per 1 μm of dendrite (the latter is a count in human postmortem cortex, Ref. 35). Obviously, this number is dependent on age, cell type, and position along the dendrite. However, it is also dependent on the method of measurement. The most accurate count is based on three-

dimensional EM reconstructed dendrite which yield a larger density than a light microscope, single plane Golgi stained neuron. Likewise, the spine shape is also dependent on all these parameters, so it is of utmost importance to consider the baseline spine density before dwelling into changes produced by one treatment or another.

1. Direct versus indirect effects on spine morphology

There are several ways to generate changes in spine morphology. In general, one can activate individual spines, either by electrical stimulation or by flash photolysis of caged glutamate. The latter stimulation also activates nonsynaptic glutamate receptors, which may or may not reside on spine heads, but also on dendritic shaft. On the other hand, the entire network can be activated by protocols such as “chemical” LTP or long-term depression (LTD), which will cause obvious changes in a large proportion of the imaged spines, but not be specific to a given spine, and in fact can be caused by changes in dendritic excitability (131).

2. Spatiotemporal resolution

The detection of dynamic changes in spine parameters, including volume changes, transient rise in $[Ca^{2+}]_i$, spine motility as well as reactivity to flash photolysis of caged molecules, is dependent on the spatiotemporal resolution of the imaging system and on the concentration of the calcium dye employed. By the same token, flash photolysis of caged molecules, a popular vehicle to study properties of individually activated dendritic spines, can vary from nanosecond flash to several milliseconds. Obviously, the sphere of diffusion of the activated compound will vary as a function of the duration of flash so that the action of the photoreleased molecule may not be as localized as one wishes.

These are several cautionary notes that a critical evaluation of the vast literature on dendritic spines need to consider. Obviously, the more standard the methods are, the more generalized and reliable are the results.

III. ELECTROPHYSIOLOGICAL PROPERTIES OF DENDRITIC SPINES

The early theoretical view of dendrites and spines considered them as being passive cylindrical compartments. Based on cable theory, research addressed issues such as the electrotonic distance of the synapse from the soma (405). Already in the 1980s, Perkel, Rall, and Shepherd considered dendritic spines as having excitable membrane (325, 387, 456), and subsequently the theoretical research on dendritic spines gained momentum and began to be relevant to the rapidly accumulating experimental evidence for the presence of voltage-gated currents in dendritic spines.

With the increased use of molecular and high-resolution imaging tools for localized receptors and ion channels in different

compartments of neurons, the knowledge about the electrical properties of dendritic spines is becoming more relevant than ever before. Furthermore, the ability to alter the morphology of dendritic spines using insertion and deletion of specific genes also calls for a better understanding of the electrophysiological consequences of these alterations.

A. Voltage-Gated Sodium Channels in Spines

While theoretical considerations assumed that the spine neck is not a barrier to the transfer of the synaptic charge to the parent dendrite (449), experimental evidence for this assumption is rather scarce, for the simple reason that recording of electrical activity from spine head has not yet been achieved due to the small size of this organelle. Still, the estimation of the axial resistance of the spine neck is crucial in the attempt to extract the relevance of the physical distance of the spine head from the parent dendrite. Estimations of the spine neck resistance range from 1.2 G Ω (158), which is critical for the ability to generate large currents at the spine head, to ~ 4 –50 M Ω (490). The latter estimate is based on the measurement of diffusion of fluorescent molecules from spine head to the parent dendrite. A more recent estimate of spine neck resistance is based on voltage measurements in the dendrite adjacent to the spine and equalization of the magnitude of calcium transient in the spine head and the electrical responses in the dendrite, and amounted to ~ 500 M Ω (167). This latter study indicates that the spine head can generate a large local voltage transient, sufficient to sensitize adjacent spines. The functional implications of this will be dealt with below (see sect. VII). Interestingly, this recent study did not address the functional relevance of large heterogeneity of spine neck parameters.

The presence of voltage-gated Na channels in dendritic spines has been demonstrated using live imaging methods. Using the Na dye SBFI, Rose et al. (418) measured Na transients in dendritic spines of CA1 neurons in a slice, suggested to amount to a modest change of 4 mM for a train of 20 action potentials. Obviously, these were early days of imaging and the resolution was not so high. Later studies using second harmonic generation (360) were able to image membrane potential changes in dendritic spines, and reported a 10% change in fluorescence of FM4–64 for 100 mV. More recently, using voltage-sensitive dyes injected into individual neurons in layer 5 of the neocortex, Palmer and Stuart (377) compared back-propagating action potentials (bAP) with synaptic inputs. They found that bAP can invade the spines without significant loss in magnitude, and they estimated the size of the response to vary from 80 to 20 mV, depending on the distance from the soma, as expected for apical dendrites. Interestingly, they found that hyperpolarization of the soma can increase the size of the synaptic input, indicating that there is no involvement of voltage-gated channels in the amplification of the synaptic input. Furthermore, they estimated spine neck resistance to be on the order of 500 M Ω , which will reduce the synaptic response at the soma by $<15\%$. More recently, Acker et al. (3)

imaged single voxels at high temporal resolution using voltage-sensitive dyes in cortical slices, to find a 16% change in fluorescence/100 mV in individual dendritic spines. They confirmed the presence of voltage-sensitive fast changes in fluorescence, correlated with somatic action potentials. Interestingly, they did not report any difference between spines of different lengths and sizes. This indicates that the length of the spine neck is not relevant to the transfer of voltage signals from the parent dendrite to the spine head, but it does not predict the transfer from the head to the dendrite. Another, rather indirect means for localizing voltage-gated sodium channels in spine heads was used by Yuste and colleagues (19) who measured the voltage response to flash photolysis of caged glutamate, to find that blockade of Na channels with tetrodotoxin (TTX) reduced markedly the responses at the soma, indicating that the spine Na channels act to boost the glutamate responses. This suggestion contradicts that of Palmer and Stuart (see above).

Finally, while there is accumulating evidence, using voltage-sensitive dyes and Na dye for voltage-gated Na channels in the spine head, the morphological evidence for this is rather scarce. Using quantitative EM immunocytochemistry, Lorincz and Nusser (288) were unable to detect the presence of Nav channels in dendritic spines. These channels were found in somata and proximal dendrites, still at much lower density than in the axon initial segment, but were conspicuously absent from dendritic spines. Thus it is still unclear which are the channels that are activated, at very short latencies, by the bAP.

B. Potassium Channels in Spines

Of the variety of voltage-gated potassium channels residing on central neurons (212), special attention is devoted to the Kv4.2 channel that carries the transient I_A current which is assumed to reside primarily on remote dendrites and spines (13). In a recent EM mapping study, Kerti et al. (228) found this channel to distribute evenly on dendrites and spines, but with no preference to remote dendrites. An interesting association between Kv4.2 and an actin binding protein filamin has been demonstrated (388), and it was suggested that filamin doubles the current density generated through the Kv4.2 channels in transfected cell lines. However, a recent mapping study could not localize filamin within dendritic spines of cultured hippocampal neurons (357).

Another, voltage-independent/calcium-dependent potassium channel that is found in dendritic spines is the small potassium (SK2) type. It is activated by a rise in intracellular calcium, is selectively blocked by apamin, and resides at high concentrations near postsynaptic density within dendritic spines (44). Blockade of SK channels or their genetic deletion increases excitability of dendrites and spines and enhances their ability to undergo LTP (7). This channel is subject to modulation by the M1 muscarinic receptor, which reduces the sensitivity of this channel to intracellular calcium concentration, causes an

increase in the size of the excitatory postsynaptic potential (EPSP), and may underlie the effects of muscarinic drugs on learning and memory (145). Interestingly, the channel is activated by an influx of calcium through the NMDA receptor. In the process of LTP, this channel is assumed to be removed from the plasma membrane, and thus increase excitability of the dendritic spine (145).

Another voltage-insensitive potassium channel assumed to be selectively associated with the postsynaptic density is the G protein-coupled K channel (GIRK, Ref. 294). It complexes with protein kinase A, protein kinase C and phospholipase C, as well as protein phosphatase 1, protein phosphatase 2A, and regulators of G protein signaling, all of which modulate its activity. It is assumed to be activated by the metabotropic GABA receptor (GABA-B) and by serotonin to hyperpolarize the spine membrane and cause depotentiation of the excitatory synapse. Interestingly, morphine enhances GIRK channel activity in the hippocampus (351). While GIRK channels are not assumed to selectively localize in dendritic spines, their interactions with excitatory neurotransmission has important implications for dendritic spine plasticity.

C. Calcium Handling in Spines

Compared with NaV channels, much more is known about the presence and functions of calcium channels in dendritic spines. Intracellular calcium concentration ($[Ca^{2+}]_i$) is maintained at a low (50–100 nM) concentration in resting conditions and can rise, following stimulation, by several orders of magnitude. This rise in $[Ca^{2+}]_i$ activates cascades of kinases and phosphatases, resulting changes in cellular functions (see sect. VI). The dendritic spine is endowed with several species of voltage-gated calcium channels. These include channels of the L [Ca(v)1.2 and Ca(v)1.3], T, and R [Ca(v)2.3] types (43). The former channel is distributed in the soma and dendrites as well, but is concentrated in dendritic spine postsynaptic density and distal dendrites (277), but the T and R channels are probably responsible for the rise of $[Ca^{2+}]_i$ following back-propagating action potentials (43). These channels are localized primarily on spines of pyramidal neurons and selectively activate the SK channel, which is instrumental in rapid restoration of membrane potential following voltage-gated depolarization of the spine head (44). Interestingly, this selective action of the Ca(v)2.3 channel indicates that there are calcium microdomains within the dendritic spine head, as small as they may be. Other types of calcium channels are present in other cell types; cerebellar Purkinje cells are endowed with P-type channels, while thalamic neurons contain an abundance of T-type channels (89). These channels differ in their kinetics, voltage dependence, and drug sensitivity (see Ref. 517 for review).

Another category of routes for calcium entry into spines are the ligand-gated channels. These include two subtypes: ionotropic and metabotropic receptors. The most abundant one is

the glutamate receptor. Spine heads contain ionotropic glutamate receptors of the AMPA and the NMDA types, as well as metabotropic glutamate receptors. The NMDA receptors, when activated, allow both Na^+ and Ca^{2+} to influx into the spine. The NMDAR is voltage/Mg sensitive and, upon activation, is likely to be responsible for most of the influx of calcium into the spine head, in as much as blockade of the NMDA receptors suppresses the calcium transient recorded upon glutamate activation of the synapse (44, 428). In fact, it has been proposed that some spines may be endowed only with NMDA receptors, making them “silent synapses” at resting potential, and, when activated, cause an influx of calcium, leading to the recruitment of AMPA receptors into the spine heads (301). Spine heads contain several other receptor species, including the metabotropic glutamate receptor (mGluR), likely to cause activation of phospholipase C followed by release of calcium from stores (123, 151), and the nicotinic cholinergic receptor (119).

Two other possible modes of calcium entry into spines are the transient receptor potential (TRP) channel and the store-operated calcium entry (SOCE) channel. The TRPs constitute a family of channels that is subdivided into several species (355). There is accumulating evidence that TRPC3, a nonselective cation channel, allows calcium into cells, and it is assumed to mediate the effects of brain-derived neurotrophic factor (BDNF) on spine formation and synaptic activity (11, 348). It is further suggested that the effects of BDNF to cause trafficking of AMPA receptors into the synapse, an essential element in synaptic plasticity, are mediated partially by the TRP channel (348). NMDA-mediated calcium influx may also involve TRP channels, at least in olfactory bulb neurons (485). Another member of the TRP channel family, the TRPC6 has also been associated with excitatory synapses, and assumed to regulate their formation (585) via a CaMKIV-CREB pathway. Thus overexpression of TRPC6 results in a higher density of spines, higher rates of miniature excitatory postsynaptic currents (mEPSCs), and a better performance in a water-maze task.

The role of calcium stores in dendritic spine function and plasticity is still debated. Neuronal calcium stores are of two types, the inositol 1,4,5-trisphosphate (IP_3) receptor and endoplasmic reticulum store, studied extensively in cerebellar Purkinje cells (151) but is also ubiquitous in other brain regions as well, including the hippocampus and cortex. The IP_3 receptor is assumed to mediate the action of acetylcholine (ACh) and other neuromodulators, to cause release of $[\text{Ca}^{2+}]_i$ from stores and to cause a subsequent change in AMPA and NMDA receptor functions (123). Interestingly, while IP_3 receptors have been localized in dendritic spines of GABAergic cerebellar Purkinje cells and spiny striatal cells where they have been studied extensively (196), they are probably present in dendritic shaft but not in spines of pyramidal neurons (453), even though their function in synaptic plasticity has been extensively documented (123, 410). One possible mode of involvement of

endoplasmic reticulum in spine calcium-induced calcium release from IP_3 stores has been proposed recently (192); synaptic activation of glutamate receptors could evoke a delayed calcium surge in large spines that was blocked by mGluR antagonists and heparin (indicating IP_3 stores). The role of these delayed calcium surges in synaptic plasticity is not entirely clear.

The more controversial store is the one associated with the ryanodine receptor. This store has been studied extensively in muscle fibers but is also found in neurons throughout the brain (529). There are three species of ryanodine receptors (RyR1–3) that are differentially localized in dendrites and spines of central neurons. Its main attribute is that it is activated by calcium influx into the cell, meaning that the influx of calcium ions through the plasma membrane is amplified by release of calcium from RyR-associated calcium stores. Caffeine and low concentrations of ryanodine (0.5–1 μM) are agonists for this store, and high concentrations of ryanodine (100 μM) or cyclopiazonic acid (CPA) are antagonists of the store. Initial studies (254, 299) were unable to detect an effect of CPA on subthreshold synaptically evoked rise of $[\text{Ca}^{2+}]_i$ in CA1 neurons of the hippocampus, a response that was shown to be mediated by activation of the NMDA receptor. In contrast, Emptage et al. (112) demonstrated that release from stores is responsible for most of the rise of $[\text{Ca}^{2+}]_i$ that is seen following synaptic activation in CA1 neurons of cultured hippocampal slices. Interestingly, they found that blockade of calcium-induced calcium release from stores does not affect back-propagating action potential-induced calcium rise. This observation contrasts with that of Sabatini and co-workers (299, 428) who found that the decay time constant of the elevated $[\text{Ca}^{2+}]_i$ caused by back-propagating action potential is prolonged by CPA. Admittedly, Sabatini and co-workers (299, 428) found that this is a heterogeneous effect, likely due to heterogeneity of the concentration of RyRs in dendritic spines.

The issue remained controversial when Brunig et al. (61) reported that they could not detect any effect of caffeine on spine motility, unlike the effects of NMDA or AMPA receptor activation. Furthermore, Harvey and Svoboda (172) could not confirm an involvement of RyR in tetanic stimulation-induced spine head expansion. These observations contrast with the effects of caffeine on $[\text{Ca}^{2+}]_i$ and spine morphology reported before (251), and the description of the presence of calcium stores in dendritic spines (170). A more recent study proposes that the RyR2 and -3 isoforms mediate the action of BDNF on dendritic spines and on cognitive tasks associated with the hippocampus (5).

A possible molecular and structural substrate associated with the RyR is the spine apparatus, which is enriched with synaptopodin, an actin binding peptide that was originally detected in the kidney, and later in the brain (338). It has been proposed that synaptopodin is colocalized with the presence of the RyR

in dendritic spines of rat hippocampus (534). The association of synaptopodin with RyR may resolve the disagreements about the proposed involvement of RyR in synaptic plasticity, as only some large spines contain the spine apparatus and synaptopodin, while the others may not be affected by activators of the RyR (253).

Strikingly, recent work by Stutzmann and Mattson (486) associates Alzheimer's disease (AD) neuropathology with an overexpression of RyRs in dendrites and particularly in spines. Furthermore, caffeine caused a large facilitation of reactivity to tetanic stimulation only in neurons from young 3xTg mouse hippocampus. This supersensitivity to activation of the RyR was noticed before the emergence of neuropathology in these mice, indicating that activation of RyR stores may lead rather than lag behind the pathology of AD. These results indicate that the association of RyR with dendritic spines, and their role in release of calcium from stores in relation to synaptic plasticity, may be more critical than originally suggested, especially in the development of AD.

Once $[Ca^{2+}]_i$ rises in the dendritic spine, it is removed rapidly from the cytoplasm. Several routes for removal of excess calcium are found in dendritic spines, including extrusion via Ca^{2+} -ATPase pump and Na^+ - Ca^{2+} exchangers (62). Calcium can also be pumped into intracellular calcium stores that are present in the endoplasmic reticulum by activation of endoplasmic reticulum Ca^{2+} -ATPase. Interestingly, one of the main sinks for accumulation of excess intracellular calcium is the mitochondria, which is conspicuously absent from the dendritic spine (473). The absence of mitochondria, the relative sparse endoplasmic reticulum, and the small volume of the spine led to the suggestion that the spine can raise $[Ca^{2+}]_i$ momentarily to very high levels, much above those of the parent dendrite, so as to effectively activate local calcium-dependent phosphorylation/dephosphorylation of key proteins in the signal transduction cascades (88).

D. The Spine Neck: Barrier or Bridge?

As mentioned above, earlier studies (490) measured the dissipation of a fluorescent dye from the spine head to the parent dendrite, to estimate the spine neck resistance to be $<50\text{ M}\Omega$. In a later study, using photoactivatable GFP, Bloodgood and Sabatini (43) found large variations in the diffusional coupling between spine head to the parent dendrite, which are not explained only by spine neck length. In fact, the diffusional coupling was found to be regulated by pre- and postsynaptic activity, such that active neurons restrict the coupling between their spines and the parent dendrites. Further studies by Grun-ditz et al. (158) demonstrated that postsynaptic depolarization is sufficient to restrict the diffusion between the spine and dendrite. In fact, spine neck regulates the local depolarization of the spine head, to allow variable synaptically evoked calcium transients in the spine head (149).

The functional significance of spine neck length has three facets: first is the question of what chemical, if any, is transmitted between the spine head and the parent dendrite, and what might it signal to the dendrite. Second is the question of whether there are functional differences between short and long spines with respect to magnitude/quality of synaptic currents. Third is the question of what regulates spine neck length and its function as a variable barrier.

In as much as there are no differences between short and long spines in voltage responses to bAP arriving from the soma, Araya et al. (18) reported that shorter spines produce larger somatic EPSCs in response to flash photolysis of caged glutamate. In support of this is the recent suggestion (22) that the spine neck morphology may regulate the diffusion of glutamate receptors from the dendrite into the synapse, which may underlie the neck-length dependent variations in synaptic responses, and the suggestion that a synapse on filopodia is not as effective, and certainly not as stable as a synapse on a spine. On the other hand, Harris and Stevens (171), in an extensive EM reconstruction study, found a correlation between spine head size and PSD dimension, but could not find a correlation between PSD size and spine neck length. PSD size is correlated with magnitude of the synaptic response as has been shown by several groups. What then is the function of the synapse on long spines if they do not impact the soma? Spine neck length can vary as a function of exposure to plasticity-producing stimulation (228), so their functional integration into the dendritic signaling is likely to result from their exposure to enhanced stimulation.

What is the signal that is transmitted between spine head and the parent dendrite? Because of the fast kinetics of decay of elevated calcium, and the fact that the spine neck is a rather narrow ($\sim 0.1\text{--}0.2\text{ }\mu\text{m}$) and long ($1\text{--}2\text{ }\mu\text{m}$) tube, it was assumed that the path between the spine head and the dendritic shaft is not used for dissipation of excess calcium from the spine head (428). However, flash photolysis of caged calcium in dendritic spines (249) indicates that spine neck length can be considered a variable resistor for the diffusion of calcium from the spine head to the parent dendrite, such that short spine necks allow elevated calcium to diffuse from the spine head into the parent dendrite.

Small molecules can carry the message sent from activated spines to neighboring ones. One of them is Ras. It is generated by uncaging of glutamate near a dendritic spine. It has been associated with LTP of reactivity of the spine to glutamate and can spread into the parent dendrite and adjacent spines up to $10\text{ }\mu\text{m}$ away from the activated one (173). This rise of active Ras is likely to underlie the enhanced sensitivity of spines adjacent to the activated one to subsequent activation. RhoA is another such molecule that becomes persistently active following intense stimulation and can spread to adjacent spines (341).

IV. MOLECULAR CONSTRUCTION OF SPINES

The identification of the molecules and molecular families involved in spine formation and morphology started in the late 1990s when it became possible to transfect neuronal cultures with cDNAs expressing proteins potentially important for spine morphogenesis. Since then, several hundreds of proteins/hormones/growth factors have been proposed to act as spine morphogenic factors (TABLE 1). For most of these proteins their activity was demonstrated by transfecting wild-type or mutant species in hippocampal/cortical neurons in dissociated cultures or organotypic brain slices. For some of them, the animal model (knockout, knock-in, or transgene) has been developed so as to reinforce and extend the *in vitro* findings. Interestingly, the presence of functional mitochondria in dendrites seems to be required for the correct formation, maturation, and plasticity of dendritic spines as suggested by recent experiments in neurons deleted or transfected with mutant proteins, such as PGC-1 α , Drp1, OPA1, Sig-1Rs, PINK1/PARK6, and Centaurin- α 1, that regulate formation, morphology, function, and dendritic localization of mitochondria (77, 280, 492, 515, 574). In this review we classify the spine morphogenic factors depending on the molecular composition/function, and we emphasize the molecules that have been demonstrated to have a spine morphogenic activity in both culture and brain tissue.

A. Actin Binding and Cytoskeletal Proteins

The cytoskeleton of dendritic spines is composed mainly of filamentous (F)-actin that forms parallel filaments in the neck of the spines and lattice and twisted filaments in the spine head. Thus it is not surprising that all the molecules that regulate growth and shrinkage of dendritic spines will eventually modify positively or negatively actin polymerization, and even the overexpression of actin alone in neurons could increase dendritic spine density (215).

A number of actin binding proteins have been implicated in spine morphogenesis (FIGURE 1). Among them, Drebrin A was the first to be proposed as a regulator of spine morphology when overexpressed in cultured neurons (42, 176, 330). In cultured cortical neurons, the overexpression of Drebrin cDNA (Drebrin A, the adult isoform) positively modifies the length of dendritic spines (176) while in hippocampal cultures mature spines became less stable. It is suggested that Drebrin activity is dependent mainly on its F-actin interaction but also on its binding to some actin-regulatory proteins, such as profilin, myosin, gelsolin, and Ras (42, 303). Although these are consistent *in vitro* observations, Drebrin knockout mice have morphologically normal dendritic spines but show altered homeostatic plasticity, suggesting that Drebrin regulates a subset of actin involved in synaptic plasticity (17).

Among the other actin-binding proteins shown to be morphogenic for spines are myosins II and VI, NESH,

spinophilin/neurabin II, synaptopodin, acidic calponin, cortactin, SPIN90, VASP, and WAVE. These proteins regulate actin polymerization directly in the dendritic spines where they accumulate. On the other hand, none of them is specific for dendritic spines because these proteins regulate actin polymerization in many other cellular functions and subcellular compartments.

Among the molecular motors that bind and contract actin filaments, myosin IIB and myosin VI (Myo6) have been found to regulate actin polymerization in spines. These motors are formed by a hexameric protein complexes with two heavy chains and four light chains. Myosin II (in particular the IIB or the nonmuscle *MyH10* isoform) is highly localized at synapses and regulates dendritic spine morphology and synaptic strength (426, 582). In culture, blocking myosin IIB with blebbistatin or with shRNA against the heavy chain inhibits mature spine formation and induces filopodia-like protrusions (426). Also, spine maturation into a mushroom shape requires the contractile activity of myosin IIB (189). Interestingly, the myosin II regulatory light chain (MLC) can be phosphorylated and activated by GIT1/PAK complex (see sect. IVB) to induce dendritic spines formation (582). Another myosin II expressed in neurons and in spines is the muscle sarcomeric *MyH7B* isoform, knocking down of which alters spine head morphology (422). The comparison of *MyH10* and *MyH7B* knock down indicates that the two myosin II might have a cooperative mechanism to control spine morphology and synapse function (422).

The role of myosin VI (Myo6) in dendritic spines morphogenesis has been demonstrated both *in vitro* and *in vivo* (371). Myo6 also regulates AMPA receptor internalization because Myo6-deficient mouse (called Snell's waltzer) exhibits reduced synapse number, abnormal short dendritic spines, profound astrogliosis, and deficit in the stimulation-induced internalization of AMPAR (371). These data suggest that a limited number of specific actin-based motors control the structure and function of spines in addition to the polymerization state of actin.

Abi-1, Abi-2, NESH/Abi-3, members of the c-Abl tyrosine interactor (Abi) protein family, are linked to PSD scaffold proteins and spine F-actin (378, 401). In the PSD Abi-1 binds to ProSAP2/Shank3, a scaffold protein of the PSD (see sect. IVB) and is translocated to nuclei from synapses where it can activate gene transcription of E-box-containing genes upon NMDA receptor stimulation. Functional studies in hippocampal cultured neurons have demonstrated that Abi-1 knocking down by RNAi changes both dendrite morphology, by increasing the branching, and spine structure that became more similar to an immature shape. Thus Abi-1 can regulate dendrite and synapse formation working as a synapto-nuclear messenger specifically controlling gene transcription (401). However, Abi-1 can also activate Rac1 and F-actin

Table 1. List of major proteins involved in molecular construction of spines

Name	Main Funtion	Experimental Model	Methods	Reference Nos.
<i>Actin binding and cytoskeletal proteins</i>				
Abi-1	Actin binding protein	C	RNAi	378, 401
Abi-2	Actin binding protein	M	KO	155
Abi-3/NESH	Actin binding protein	C	Over/RNAi	26
α N-Catenin	Actin binding protein	C	Over	1
Arp2/3	Actin binding protein	M	KO	231
Calponin	Actin binding protein	C	Over	407
Cortactin	Actin binding protein	C	Over/RNAi	185
Drebrin A	Actin binding protein	C	Over/KO	17, 42, 176, 303
MLC	Actin binding protein	C	RNAi	582
Myosin IIB	Actin binding protein	C	Blebbistatin/RNAi	189, 422, 426
Myosin VI	Actin binding protein	M	KO	371
Neurabin I	Actin binding protein	C OBS	Over	369, 590
Neurabin II Spinophilin	Actin binding protein	M	KO	122
Profilin I/II	Actin binding protein	C OBS	Over/RNAi	4, 323
SPIN90	Actin binding protein	C M	Over/RNAi/KO	81, 236, 275
Synaptopodin	Actin binding protein	C	RNAi	368, 534
VASP	Actin binding protein	C	RNAi	282
WAVE1	Actin binding protein	C M	RNAi KO	237
WAVE3	Actin binding protein	C	Over	391
CP	Actin capping protein	C	RNAi	120
Eps8	Actin capping protein	C M	Over/RNAi/KO	320, 480
EB3, p140Cap/SNIP	Microtubule binding protein	C	RNAi	159, 210
MAP1B	Microtubule binding protein	M	RNAi	510
Actin	Cytoskeleton	OBS	Over	215
<i>Small GTPase and associated proteins</i>				
ARF6	GTPase	C M	Over/RNAi/KO	83, 329, 404
Cdc42	GTPase	OBS M	KO	501, 522
Rac1	GTPase	C OBS M	DN/CA/PI/KO	52, 496, 501, 502, 522
Rap1	GTPase	C	CA	557
Rap2	GTPase	M	Transgene	425
Rem2	GTPase	C	RNAi	143
RhoA	GTPase	OBS	CA	501
Rif	GTPase	C	DN/CA	195
Rnd1	GTPase	C	Over/RNAi	206
ARHGEF6/ α PIX	RhoGEF	OBS M	RNAi/KO	359, 406
ARHGEF7/ β PIX	RhoGEF	C OBS	RNAi	441
Dock180	RhoGEF	C	Over/RNAi	232
GEFT	RhoGEF	C	Over	60
Kalirin-7	RhoGEF	C OBS M	Over/RNAi/KO	64, 217, 295, 296, 383, 385, 556, 558, 570, 571
Lfc/GEF-H1	RhoGEF	C	Over/RNAi	222, 424
Tiam1	RhoGEF	C	Over/RNAi	328, 507, 508, 581
Vav	RhoGEF	C OBS M	KO	161
α 1-Chimerin	RhoGAP	C	Over/RNAi	526
oligophrenin1	RhoGAP	OBS M	RNAi/KO	229, 230
p19ORhoGAP	RhoGAP	M	KO	450
p25OGAP	RhoGAP	C OBS	RNAi	166, 202, 349
RhoGAP2	RhoGAP	C	Over	524
SRGAP2	RhoGAP	C	Over/RNAi	71

Continued

Table I.—Continued

Name	Main Funtion	Experimental Model	Methods	Reference Nos.
EPAC1/2	RapGEF	C M	Over/RNAi/KO	477, 550, 565
PDZGEF1	RapGEF	C	Over/RNAi	274
Rap1GAP	RapGAP	C	Over	313
SPAR	RapGAP	C	Over	190, 376
RasGRF1	RasGEF	C	Over/RNAi	274
SynGAP	RasGAP	C M	KO	528
EFA6A	ArfGEF	C M	Over/RNAi/KO	83
Centaurin α 1	ArfGAP	C	Over/RNAi	492
SrGAP3	RacGAP	M	KO	114, 539
PAK	Kinase	M	Transgene	180
PAK1	Kinase	OBS C	DN/CA/RNAi	48, 175
PAK3	Kinase	C OBS M	Over/CA/DN/RNAi	40, 46, 48, 50, 103, 256, 317, 359, 506

Cell surface receptors and adhesion molecules

α 3 Integrin	Adhesion protein	M	KO	227
α 5 Integrin	Adhesion protein	C	Over/RNAi	543
Arcadlin	Adhesion protein	M C	KO	568
DSCAM	Adhesion protein	M	KO	312
IL1RAPL1	Adhesion protein	C M	Over/KO	382, 523
N-cadherin	Adhesion protein	C OBS	Over/RNAi	96, 106, 316, 367, 430, 568
Neuroigin1,2,3,4	Adhesion protein	C M	Over/RNAi/KO	21, 45, 78, 79, 87, 208, 493
SALM2	Adhesion protein	C	Over/RNAi	242
Syndecan-2	Adhesion protein	C	Over/RNAi	117, 118, 283
Telencephalin	Adhesion protein	C M	KO	404
APP	Membrane protein	M	Transgene	270
TSPAN7	Membrane protein	C	Over/RNAi	31
Vezatin	Membrane protein	C M	KO	96
beta2-nAChR	Receptor	C M	Over/KO	289
GABAAR	Receptor	M	KO	183
GluA2	Receptor	C	Over/RNAi	380, 430
GluN1	Receptor	OBS M	RNAi/KO	10, 521
GluN2B	Receptor	M	KO	59
Npn-2	Receptor	M	KO	512
NgR1	Receptor	C M	Over/KO	6, 272
PGC-1 α	Receptor	C	Over/RNAi	77

Receptor tyrosine kinases and other kinases

EphB1/2/3	TK receptor	C M	Over/RNAi/KO	118, 184, 203, 224, 225, 331, 383
EphA4/ephrin-A3	TK receptor	C OBS M	Over/RNAi/KO	55, 66, 339, 587
ErbB2/B4	TK receptor	M	KO	30
ErbB4	TK receptor	OBS	Over/RNAi	278
p75NTR	TK receptor	M OBS	KO	576
TrkB	TK receptor	M	KO	68, 535
CaMKII	Kinase	C OBS	Over/RNAi	218, 366
CaMKIIalpha	Kinase	M	KO	562
CDKL5	Kinase	C	Over/RNAi	414
DCLK1	Kinase	C M	RNAi/KO	460
DGKf	Kinase	C M	Over/RNAi	233
LIMK-1	Kinase	M	KO	318
MARK4	Kinase	C	Over	573

Continued

Table I.—Continued

Name	Main Function	Experimental Model	Methods	Reference Nos.
NDR1/2	Kinase	C OBS	RNAi	520
PAR1b	Kinase	C	RNAi	177
PI3K	Kinase	C	Activation	92
PKM α	Kinase	C	Over	417
Plk2	Kinase	C M	KO	274
Wnt7a; Dvl1	Kinase	C M	KO	84
<i>Postsynaptic scaffold proteins and adaptor proteins</i>				
CASK	Scaffold	C	Over/RNAi	69
CTTNBP2	Scaffold	C	RNAi	75, 76
cypin	Scaffold	C	Over/RNAi	516
DISC1	Scaffold	C	Over/RNAi	181
GIT1	Scaffold	C M	RNAi/KO	321, 582
Homer1a	Scaffold	C	Over	432
Homer1b	Scaffold	C OBS	Over	179, 433
intersectin-s	Scaffold	C	RNAi	505
IQGAP1	Scaffold	M	KO	137
N-WASP	Scaffold	C	Over/RNAi	544
PICK1	Scaffold	C	Over/RNAi	347
PSD-95	Scaffold	C OBS M	Over/KO	111, 354, 398, 483, 533
Preso	Scaffold	C	Over/RNAi	273
SAP102	Scaffold	C	Over/RNAi	72
Shank1	Scaffold	C M	Over/KO	200, 433
Shank2	Scaffold	M	KO	444, 549
Shank3	Scaffold	C M	Over/RNAi/KO	108, 421, 530, 541
TANC1/2	Scaffold	C M	Over/KO	164
WAVE1	Scaffold	M	KO	468
afadin	Adaptor	M	KO	33
IRSp53	Adaptor	C M	Over/RNAi/KO	82, 234, 443
Numb	Adaptor	C	Over/RNAi	356
PAR-3	Scaffold	C	RNAi	581
PAR-6	Scaffold	C	Over/RNAi	580
<i>Micro RNA (miRNA), mRNA binding protein, and transcription factors</i>				
miR-29a/b	miRNA	C	Over	284
miR-125a	miRNA	C	Antisense	335
miR-125b	miRNA	C	Over/sponge	109
miR-132	miRNA	M	Over/sponge/transgene	109, 166, 202, 315, 381
miR-134	miRNA	C	Over/antisense	445
miR-138	miRNA	C	Antisense	463
miR-185	miRNA	M	KO	559
Satb1	DNA binding protein	C from M	KO	28
hnRNPK	RNA binding protein	C	Over/RNAi	402
Staufen2 (Stau2)	RNA binding protein	C	Over/RNAi	148
TLS	RNA binding protein	M	KO	134
Cux1/Cux2	Transcription factors	C M	KO	91
FoxO6	Transcription factor	M	KO	436

C, primary neuronal cultures; OBS, organotypic brain slices; M, genetically modified mice; Over, overexpression; DN, dominant negative; CA, constitutively active; PI, pharmacological inhibition.

polymerization upon calcium/calmodulin-dependent protein kinase II (CaMKII)- α phosphorylation at serine-88 induced by glutamate stimulation (378). Abi-2 is also strongly expressed in the brain, and its knockout in

mice impairs spine density, by reducing the number of mushroom shaped spines. These morphological alterations correlate with negatively altered both short- and long-term memory formation (155).




FIGURE 1. Actin binding proteins. The scheme shows the principal actin binding proteins (in ovals) and the main associated proteins (in the ellipse) divided for the principal activity on actin filaments (one filament in green).

The manipulation of NESH/Abi-3 expression has multiple effects: overexpression of NESH/Abi-3 in cultured neurons reduces the amount of mushroom-type spines and synapse density but increases the thin filopodia, while siRNA knockdown also reduces mushroom spine numbers but increases spine density (26). Whether NESH/Abi-3 activity is regulated by synaptic stimulation was not determined as yet.

Neurabin I and neurabin II (or Spinophilin) are both F-actin-binding proteins with a common domain organization forming NH₂-terminal actin binding domain, PP1 binding domains, PDZ and a COOH-terminal coiled-coil domain. The two proteins are able to associate in homo- and heterodimers with the coiled-coil domain and to cross-link F-actin. Immature cultured neurons overexpressing neurabin I have more dendritic filopodia, while older neurons have enlarged spines (369). The overexpression of the NH₂-terminal actin-binding domain is sufficient to increase actin polymerization, spine motility, density, and length (590). Similarly, spinophilin is predominantly localized in dendritic spines (as the name suggests) of pyramidal neurons and is necessary for the development of dendritic spines because knockout mice show more filopodia and immature spines compared with wild-type mice

of the same age and express defects in glutamatergic transmission (122). Molecularly both proteins associate in synapses with a similar complex of molecules, including Lfc, a Rho GEF that regulates the Rho-dependent organization of F-actin in spines (424).

Cortactin is another synaptic actin binding protein. Its knockdown by siRNA eliminates dendritic spines in hippocampal neurons, while its overexpression causes elongation of spines. Because the localization of cortactin in synapses is regulated by NMDA receptor activation, cortactin can also be considered as a regulator of activity-dependent spine morphogenesis (185). The synaptic proteins cortactin-binding protein 2 (CTTNBP2) binds to and recruits cortactin to spines; in the absence of CTTNBP2, the number of dendritic spines is reduced, and cortactin and other synaptic proteins are less associated with spines heads (75, 76).

Acidic calponin is another actin binding protein the expression of which is upregulated during dendritic spine plasticity following pilocarpine-induced seizures and which consequently regulates spine morphology and den-

sity through the reorganization and stabilization of actin filaments (407).

SPIN90/WISH [SH3 protein interacting with Nck, 90 kDa/ Wiskott-Aldrich syndrome protein (WASP) interacting SH3 protein] is another scaffold regulator of actin polymerization in dendritic spines. Overexpression of SPIN90 in cultured neurons increases the number and length of dendritic filopodia/spines via an N-WASP-independent mechanism, while knock-down of its expression and in knockout mice dendritic spine density and maturation is reduced (81, 236, 275). SPIN90 binds to PSD-95 and Shank and might work as their downstream effector on spine morphogenesis (81, 236, 275).

The multifunctional actin-binding protein vasodilator-stimulated phosphoprotein (VASP) is a major regulator of actin polymerization. In cultures VASP knockdown leads to a significant and specific decrease in the density of spines and synapses. Not surprisingly, the actin binding domains of VASP, Ena/VASP homology 1 and 2 (EVH1, EVH2), are required for VASP accumulation near spine actin (282). While VASP is considered an actin anti-capping protein, the actin-capping protein CP, a regulator of actin filament growth, has also been described to have a similar effect to that of VASP; CP knock-down in cultured neurons reduces spine and synapse development and maturation (120). Similarly, Eps8, another actin capping protein, is important for spine morphogenesis (320, 480).

The WAVE (Wiskott-Aldrich syndrome protein family Verprolin homologous protein) families of proteins are key regulators of actin polymerization through their ability to activate the actin-related protein (Arp2/3) complex. Loss of WAVE1 in cultured neurons and in vivo induces a decrease in mature dendritic spines (237, 468). On the other hand, WAVE3 overexpression in cultured neurons decreases spine numbers, but the remaining are eventually more mature (391). WAVE1 activity on spine actin is dependent on Cdk5 phosphorylation that inhibits its ability to recruit mitochondria in dendrites (488) and to regulate the actin polymerizing Arp2/3 complex (237). The Arp2/3 complex is critical for maintaining actin polymerization in spines because the postnatal knockout of the Arp2/3 subunit ArpC3 leads to progressive loss of dendritic spines and abolishes the activity-dependent enlargement of spines heads (231).

Although polymerized actin is still the main component of the spine cytoskeleton (220), recent evidence suggests that microtubules actively participate in the shaping of dendritic spine morphology. EB3 is microtubule tip-tracking protein that enters spines with the microtubules and modulates spine morphology by interacting with p140Cap/SNIP, a regulator of Src tyrosine kinase. Thus EB3-labeled growing microtubule ends control p140Cap recruitment to spines as well as modulate cortactin function and dendritic spine actin dynamics (159, 210). Localization of p140Cap to spines is also dependent on

the presence of partitioning-defective 1b (PAR1b), a cell polarity-regulating serine/threonine protein kinase. Thus PAR1b knock-down neurons exhibit reduced mushroom-like dendritic spines and more filopodia-like protrusions, without changes in total number of protrusions (177).

Microtubule-associated protein 1B (MAP1B) is highly expressed during the beginning of neuronal development and is present in dendritic spines. In MAP1B knockout mice, the density of mature dendritic spines is reduced but is compensated by an increase of immature filopodia-like protrusions. In these mice, changes in spine morphology are associated with a decrease in Rac1 activity and an increase in RhoA activity and decrease in phosphorylated cofilin. Thus MAP1B directly or indirectly regulates the actin cytoskeleton (510).

B. Small GTPase and Associated Proteins

Actin polymerization is controlled by extracellular and intracellular signals that all come together on small GTPase of the Rho family. The switch between the active GTP-bound and inactive GDP-bound states can change the activity of specific actin binding proteins which will promote or suppress the polymerization of actin filament.

For example, the actin-based ruffling motion of lamellipodia, which is regulated by small GTPases in many cell types, is also present in the peripheral submembrane region of dendritic spines (252).

The three members of the Rho family of small GTPases mostly involved in spine morphogenesis are RhoA, Rac1, and Cdc42 (52, 353, 363, 496, 501, 502, 522), distinguished in two opposite activities; RhoA inhibits, whereas Rac and Cdc42 promote the growth and/or stability of dendritic spines (FIGURE 2A). However, this classification is too simplistic because the effects of each individual GTPase on the final spine morphology depend also on the relative level of expression and activity of the three GTPase in each spine.

Rho family GTPases are regulated by a number of upstream and downstream molecules, and several nucleotide exchange factors (GEFs) and GTPase activating proteins (GAPs) regulate Rho GTPases in spines.

Eight Rho-GEF proteins have thus far been described to play a role in regulating spine morphology. Among them, only one which seems to be specific for RhoA is GEF-H1/ARHGEF2/Lfc [Lbc (lymphoid blast crisis)'s first cousin] (222, 424). Lfc translocates upon synaptic stimulation in a calcium-dependent manner from microtubules to spines where it binds to neurabin II and is associated with AMPA receptor complex. However, the effect on spine morphology is contradictory also because the shRNA inhibition of Lfc in cultured neurons seems to reduce RhoA activity but increase Rac1 activity (222, 424).




FIGURE 2. Small GTPase. The summary diagram shows how small GTPase of the Rho family (A) and of the RAS family (B) are controlled in regulating actin in spines. The GEF and GAP proteins (in blue and green ovals) and main associated proteins (in ellipses) are indicated.

A major and most studied spine RhoGEF is Kalirin-7, which can activate all three RhoGTPase. Kalirin-7 is highly concentrated in the PSD where it binds to PSD-95 and other scaffold proteins. Several studies (including *in vivo* experiments) demonstrate that Kalirin-7 promotes spine formation especially in cortical neurons (64, 295, 296, 385, 558). The activity of Kalirin-7 is also regulated by the phosphorylation of CaMKII and Cdk5 (296, 558).

ARHGEF6/ α PIX and ARHGEF7/ β PIX [α/β -p21-activated kinase (PAK) interacting exchange factor] are both GEF for Rac1 and Cdc42 in dendritic spines. Mutations in the ARHGEF6 gene are found in patients afflicted by a nonsyndromic form of X-linked intellectual disability, and its role in spine maturation has been demonstrated also in the KO mice (261, 359, 406, 582). β PIX is recruited at synapses through its specific interaction with COOH-terminal unique sequence at the scaffold protein Shank (see below) (379), and its activity is regulated by CaM-kinase kinase/CaM-kinase complex and synaptic activity (441).

The Rac1/Cdc42 GEF Tiam1 is also involved in dendrite and spine development, and it has been demonstrated *in vitro* to be associated with NMDA receptors and to be a downstream signaling of the EphB receptor-dependent spine development (see below) (507, 508). The activity of Tiam1 is also regulated by the scaffold polarity protein PAR-3 that is able to spatially restrict Tiam1 activity to dendritic spines (581).

The last three RhoGEF described to play a role in spine are GEFT, Vav, and Dock180. GEFT is diffusely expressed in the brain and induces both neurite outgrowth and dendritic spine enlargement, acting on the Rac and Cdc42-PAK pathways (60). With the use of KO mice it has been demonstrated that Vav mediates BDNF-stimulated Rac-GTP production and spine head enlargement in CA3-CA1 synapses as well as theta burst-stimulated LTP (161). Dock180 has been identified as the only RhoGEF, among the 70 tested, to be able to promote spine morphogenesis in a complex formed by RhoG/ELMO1/Dock180 signaling module (232).

Activated GTP-bound Rho GTPases are inhibited by specific GTPase activating proteins (RhoGAP), five of which seem to be important for dendritic spine actin remodeling. An extensively studied synaptic RhoGAP is oligophrenin because when its gene is mutated, a form of syndromic X-linked intellectual disability emerges (41). Both *in vitro* and *in vivo* (in the KO mice), oligophrenin suppression reduces spine length and maturation supposedly by over activation of RhoA although Rac1 and Cdc42 are also substrates of oligophrenin (153, 229).

p250GAP is an NMDA receptor-associated RhoGAP, yet the knock-down of which increases spine maturation by modulating RhoA (202, 349) or Rac1 activity (202), probably depending on the type of neurons studied. The p190RhoGAP is another GAP specific for RhoA that is regulated by Arg kinase

and partition scaffold protein PAR-6 and is important for maintaining the number and morphology of mature spines in adult mice (450, 580).

A peculiar RhoGAP specific for Rac1 is the non-kinase phorbol ester receptor α 1-chimerin, a protein that when overexpressed in cultured hippocampal neurons inhibits the formation of new spines and removes the existing ones by binding to synaptic GluNA2 subunit of the NMDA receptors (526). Two opposite effects on spines have been observed for two other synaptic RhoGAPs called SRGAP2 and RhoGAP2/ARGEF22. When overexpressed, SRGAP2 promotes spine maturation and limits spine density (71), while RhoGAP2 increases spine density (524) and both specifically stimulate the GTPase activity of Rac1. These data indicate that the general conclusion that RhoA inhibits, while Rac and Cdc42 stimulate the growth and/or stability of dendritic spines is too simplistic, and a more complex analysis of the relative contribution of each RhoGTPase to spine actin polymerization is needed.

Indeed, a recent study examined the spine activation of RhoA and Cdc42 using a fluorescent methodology and two-photon glutamate uncaging. Activation of both RhoA and Cdc42 was similar; an initial fast phase lasting 5 min which was followed by persistent activation enduring more than half an hour. However, the two GTPase have a different mobility: RhoA diffused and spread outside the spine up to 5 μ m along the dendrite while active Cdc42 remained concentrated in the stimulated spine. Interestingly, inhibition of the RhoA-ROCK pathway blocked the initial spine growth, and the Cdc42-Pak pathway was essential for the maintenance of the change in spine plasticity (341).

The major synaptic downstream effectors of RhoGTPase are the p21-activated kinases PAK1 and PAK3 that when activated promote formation and/or growth of spines. RNAi-mediated knock-down of PAK3 or transfection of dominant-negative PAK3 mutant induces abnormally long dendritic spines that lack synapses and a loss of mature spine synapses (46, 359). Similarly, PAK1 activity promotes spine maturation (48, 175).

However, in PAK3 knockout mice, spines are morphologically normal, but long-term synaptic plasticity is reduced and memory consolidation is impaired (317). Alterations in synaptic plasticity and a reduction in spine density have been found in cortical but not hippocampal neurons of transgenic mice overexpressing dominant negative PAK (for both PAK1 and PAK3) (180).

More recently, Dubos et al. (103) showed that inhibition of PAK3 resulted in an activity-dependent growth of new and unstable spines and the impairment of plasticity-mediated spine stabilization and formation of persistent spines (103). Thus PAK3 activity seems to regulate negatively spine growth (103). The role of PAK3 in mediating activity-dependent rear-

range of synaptic connectivity associated with learning is clearly suggested by the finding that in human, PAK3 mutation causes intellectual disability (8, 256). Lim-kinase 1 (LIMK-1) is a downstream molecule of PAK and Rho-associated kinase ROCK. Knockout mice for LIMK-1 manifest altered spine morphology, synaptic plasticity, and reduced spatial learning (318).

A link between Cdc42 and actin remodeling is Wiskott-Aldrich syndrome protein (N-WASP), which binds and activates the Arp2/3 complex and promotes actin branching and polymerization. N-WASP is localized in spines, and the knock-down of endogenous N-WASP or its inhibition with wiskostatin or the overexpression of an N-WASP mutant, which is unable to bind Arp2/3, induces a significant decrease in the number of spines and excitatory synapses in cultured neurons (544).

Finally, a specific synaptic scaffold protein, called G protein-coupled receptor kinase-interacting protein 1 (GIT1), is involved in the functional assembly of α PIX and PAK3. The importance of GIT1, that also contains an ADP-ribosylation factor (Arf) GAP domain, is demonstrated by the finding that its knockout both in cultured neurons and in transgenic mice reduces dendritic spine number (321, 582).

Ras family of GTPases and the related Rap GTPases are also involved in regulating spine morphogenesis (FIGURE 2B). The involvement of Ras and the downstream mitogen-activated protein kinase signaling pathways in regulating spine morphogenesis has been demonstrated by the observation that spine density in cortical pyramidal neurons is increased in transgenic mice expressing active Ha-Ras (138). In contrast, transgenic mice overexpressing constitutively active Rap2 (Rap2V12) in forebrain displayed fewer and shorter dendritic spines in CA1 hippocampal neurons (425).

A major regulator of Ras activity in spines is the RasGAP protein SynGAP that plays a major inhibitory role by developmentally repressing excitability synapse and dendritic spine formation during the critical period (86, 528). On the other hand, neurons overexpressing the postsynaptic RapGAPs SPAR (190, 375, 376, 476) and Rap1GAP (313) have spines with altered morphology and larger head. Thus these results suggest that Ras and Rap, in the postsynaptic compartment, have an opposite effect in controlling synapse strength, spine morphology, and synaptic plasticity. As far as synaptic plasticity is concerned, Ras is a positive effector of LTP, while Rap1 and Rap2 regulate LTD and depotentiation (589). Interestingly, the activities of Ras and Rap are coordinated by polo-like kinase 2 (Plk2), a homeostatic suppressor of overexcitation, that controls the phosphorylation-dependent ubiquitin-proteasome degradation of the Ras activator RasGRF1 and Rap inhibitor SPAR and the phosphorylation-activation of Ras inhibitor SynGAP and Rap activator PDZGEF1 (274).

It is still not clear how Ras and Rap and mitogen-activated protein kinase downstream pathways regulate dendritic spine formation (553). One effector could be Afadin, although it regulates dendritic spines mostly via cadherin function (33). Ras also binds to and interferes with drebrin activity on actin (42). Rap1 effect on spine morphology is also mediated by the adaptor protein AF-6, a scaffold protein associated with actin and other signaling proteins (557). However, both SynGAP and SPAR can regulate actin polymerization via cofilin function or directly by binding to actin (65, 376).

Among the Ras-related family of small GTPases, the ADP-ribosylation factor 6 (ARF6) and the related GEF (EFA6A) and GAP (Centaurin α 1) for ARF6 seem to regulate spine development and maintenance by controlling the conversion of filopodia to spines, or vice versa (329), and the stability of both early and mature spines by inducing the internalization of telencephalin [TLN/intercellular adhesion molecule-5 (ICAM5)] specifically at the filopodia and activating Rac1 (83, 333, 404).

Finally, two other small GTPase have been involved in regulation of spine number and morphology, namely, Rem2 and Rnd1 (143, 206). Another GTPase, Rif, is regulating filopodia formation (195). However, little is known about how are these molecules regulated and their respective effectors. Also, thus far, the data were collected only with in vitro experiments.

C. Cell Surface Receptors, Extracellular Matrix, and Adhesion Molecules

Most of the recently discovered adhesion proteins important for the formation and development of brain synapses are also involved in dendritic spine formation. Among them Neuroligin, SALM, netrin-G ligand, and IL1RAPL1 have been found to induce synapses and spine formation both in vitro and in vivo. Interestingly, all of these proteins bind to PSD-95, a major excitatory scaffold protein (see below).

The overexpression of Neuroligin1 in cultured rat hippocampal neurons promotes spine morphogenesis, together with excitatory postsynaptic differentiation, while knock-down by shRNA of three members of the family, Neuroligin1–3, reduces dendritic spine number (78, 79).

Similar results have been described for members of other synaptic adhesion molecules. In cultured rat neurons, overexpression and the knock-down of netrin-G ligand 1 and 2 (NGL-1, NGL-2) and SALM2, three synaptic adhesion-like molecules (SALM) isoforms increase or decrease the number of PSD-95-positive dendritic protrusions, respectively (21, 235, 242) (FIGURE 3A).

The overexpression of interleukin-1 receptor accessory protein-like 1 (IL1RAPL1) and 2 (members of a novel family of brain specific Toll/IL-1 receptors) induce the formation of new




FIGURE 3. Synaptic scaffolds and membrane proteins. The cartoon shows the major PSD-95 associated proteins (**A**) and synaptic membrane proteins and associated complexes (**B**) involved in regulating dendritic spine formation.

spines, while in IL1RAP1 knockout mice, the number of dendritic spines and excitatory synapses in the CA1 region of the hippocampus is reduced (382).

The activity of these molecules in dendritic spines correlates with their ability to induce the maturation of excitatory synapses by binding and recruiting pre- and postsynaptic partners. With the use of mutants of these proteins and overexpression in cultured neurons, it has been demonstrated that either the extracellular or the intracellular domains are sufficient to increase the number of dendritic spines. For example, the inhibition of Neuroigin1-dependent shedding of the extracellular domain makes the protein more capable of inducing dendritic spines but also the intracellular PDZ binding domain is required for this (489). For IL1RAP1, on the other hand, the binding to RhoGAP2 (see above) at the intracellular domain is required for the ability of IL1RAP1 to increase spine density when overexpressed in cultured neurons (524).

The cadherin-based homophilic cell adhesion system is also involved in spine morphogenesis, and it regulates pre- and

postsynaptic apposition and synaptic plasticity (437). Neuronal cadherin (N-cadherin) seems to be required for depolarization-induced spine enlargement when neuronal activity induces N-cadherin conformational change and redistribution at synapses (367). N-cadherin can also regulate spine morphology via the recruitment to spines of β -catenin, which interacts with the actin cytoskeleton, when its tyrosine dephosphorylates by receptor tyrosine phosphatases that belong to the LAR (leukocyte common antigen-related) family, a protein that is also essential for the formation of spines and synapses (106). However, the main role of N-cadherin is in the long-term maintenance and stability of dendritic spines (316). The regulated endocytosis of N-cadherin via protocadherin arcadlin and TAO2b/MEK3/p38 activation links neuronal activity with changes in dendritic spine morphology (568).

Some cadherin-associated membrane proteins contribute to dendritic spine morphogenesis: Vezatin is an integral protein indirectly associated with E-cadherin-catenin complex (but in neurons preferentially with N-cadherin) and actin cytoskeleton, and its conditional knockout in mice leads to a reduced

proportion of mature dendritic spines that correlate with an increased proportion of stubby spines (96). Finally, N-cadherin mediates the ability of the GluA2 subunit of the AMPA receptor to induce dendritic spine formation. Indeed, the extracellular NH₂-terminal domain of subunit GluR2, named NTD domain, is sufficient to induce spine growth through the *cis*- or *trans*-interaction of synaptic N-cadherin (380, 430).

Among the integrin family of adhesion protein the $\alpha 5$ subunit regulates spine morphogenesis and synapse formation in hippocampal neurons because its knockdown or the overexpression of a dominant negative mutant reduces the number of spines and synapses; $\alpha 5$ downstream signaling is dependent on Src kinase, Rac, and the signaling adaptor GIT1 protein (543). Mice knocked out for the $\alpha 3$ subunit develop normal spines up to P42 when they gradually lose hippocampal synapses and spines because the missing activation of p190RhoGAP that negatively modulate RhoA function (227).

In one of the first published overexpression experiments in neurons, Ethell and Yamaguchi (117) demonstrated that cell surface heparan sulfate proteoglycan syndecan-2 plays a critical role in spine development because its transient transfection accelerates spine maturation. Two years later, the same group demonstrated that syndecan-2 induces spine formation and clustering at synapses after being phosphorylated by EphB2 receptor tyrosine kinase (118). More recently it has been demonstrated that syndecan-2 downstream signaling leading to spinogenesis involves neurofibromin-mediated activation of PKA and Enabled (Ena)/vasodilator-stimulated (VASP) (283).

The role of extracellular matrix proteins in spine formation, maturation, and plasticity has been first indicated by Oray et al. (370a) who suggested that tPA may affect dendritic spine maturity much like monocular deprivation. More recently, other matrix components, like chondroitin sulfate proteoglycans (97a) and reelin (356a), have been shown to play a significant role in maturation of dendritic spines of cortical and hippocampal neurons.

Because molecules involved in axon guidance cues also have key roles in synapse formation and function, it is not surprising that some of these proteins are also involved in dendritic spine formation. Among them a clear role has been demonstrated for Sema3F and its holoreceptor components Npn-2, which are negative regulators of spine development and synaptic structure. Genetic ablation of either Sema3F or Npn-2 induces an increase of aberrantly shaped spines in granule cell dentate gyrus and pyramidal neurons of cortical layer V, two areas where Sema3F or Npn-2 are highly expressed (512). Nogo Receptor 1 (NgR1), a member of the Nogo-66 receptor family that controls axonal growth cone collapse and myelin-dependent growth inhibition after injury, has been recently described to control activity-dependent spine morphogenesis and turnover. Mice knockout for NgR1 showed altered spine

morphology in apical dendrites of hippocampal CA1 neurons and higher spines turnover in somatosensory neurons (6, 272). NgR1 activity on spines seems to be mediated by a direct binding to FGF1 and FGF2 (272). Thus both Sema3F/Npn-2 and FGF1-FGF2/NgR1 signaling complexes are among the few molecular cascades that inhibit spine overgrowth and suppress dynamic changes during development.

D. Receptor Tyrosine Kinases and Other Kinases

Several receptor tyrosine kinases have been implicated in regulating synapse formation (FIGURE 3B). Among them, the erythropoietin-producing hepatocellular carcinoma (Eph) receptors form a peculiar family of receptor tyrosine kinases (RTKs) with unique activity in brain synapses. Indeed, Eph receptors and ephrin ligands mediate both cell-to-cell adhesion and signaling in both directions by forming multimeric clusters that link juxtaposed cell membrane. EphB receptors are directly associated with NMDA receptors at the postsynaptic membrane, mediating NMDA receptor clustering upon activation by ephrin ligand (95). Among the different EphB receptor subtypes, the EphB2 seems to be specifically involved in spine morphogenesis, and its role has been demonstrated both in vitro and in vivo (118, 184, 203, 224, 225, 331, 383). EphB2 phosphorylates Syndecan-2 (see above); interacts with the GEF proteins intersectin, Numb, kalirin, and focal adhesion kinase (FAK)/Src/Grb2/paxillin; and activates Cdc42/N-WASP-actin, Rac-Pak-actin, and RhoA-actin pathways, respectively (203, 225, 331, 356, 383, 459, 505). Interestingly, the interaction of presynaptic ephrins with postsynaptic Eph receptors might coordinate the reciprocal interaction between presynaptic and postsynaptic functional size during structural changes accompanying synaptic plasticity (464).

As opposed to EphB, the activation of EphA4 by ephrin-A3, expressed also by astrocytes, induces spine retraction and spine modifications as observed in pyramidal neurons of EphA4 knockout mice (66, 339). Two pathways have been identified downstream of EphA4 activation: one involves inhibition of the scaffold protein Crk associated substrate (Cas), of the focal adhesion kinase (FAK) and of the proline-rich tyrosine kinase 2 (Pyk2) with the impairment of $\beta 1$ -integrin activity (55). The second involves activation of phospholipase C $\gamma 1$ (PLC $\gamma 1$) that induces spine retraction by modulating the association of the actin depolymerizing/severing factor cofilin with the plasma membrane (587). Thus EphB2 and EphA4 activation have opposite effects on spine morphogenesis and synaptic activity, while some amounts of the ephrin specific ligands may contribute to balancing their activity.

Neuregulin-1 (NRG1) signaling through tyrosine kinase receptor ErbB4 participates in regulating dendritic spine maturation, since synaptic activity induces the recruitment of ErbB4 into the synapse that selectively enhances AMPA synaptic currents and increases dendritic spine size (278). This was con-

firmed *in vivo* by the observation that knockout of ErbB2/B4 impairs dendritic spine maturation and alters the postsynaptic scaffold proteins interaction with glutamate receptors. These morphological alterations in ErbB2/B4-deficient mice are associated with behavioral manifestation of increased aggression and reduced prepulse inhibition (30).

Other than the intracellular kinases that are directly downstream signaling of the small GTPase (see above), few other intracellular kinases have been demonstrated to be important for dendritic spine morphogenesis. Not surprisingly, the CaMKII, the most abundant synaptic kinase, is one of them. CaMKII regulates synaptic plasticity by phosphorylating several substrates. Confirming earlier *in vitro* experiments (218), the kinase activity appears to be required for the tetanic stimulation to induce dendritic spine enlargement because it was absent in homozygous CaMKII α (K42R) knock-in mice that express normal quantity of a kinase-dead CaMKII α (562). On the other hand, there are indications that CaMKII β , the other CaMKII isoform, can increase spine volume by bundling F-actin through a direct stoichiometric interaction in the absence of the kinase domain (366). CaMKII also mediates the activity of Wnt7a-Dvl1 on dendritic spines. Wnt7a-Dvl1-deficient mice showed defects in spine morphogenesis and mossy fiber-CA3 synaptic transmission in the hippocampus, and CaMKII inhibition abolishes the effects of Wnt7a on spine growth and excitatory synaptic strength (84).

Two lipid kinases have been recently demonstrated to be important for dendritic spine morphogenesis. Diacylglycerol kinases (DGKs) are the principal kinases that terminate diacylglycerol (DAG) by phosphorylating DAG into phosphatidic acid (PA). Among the several DGKs isoforms, DGK ζ interacts with PSD-95, localizes at excitatory synapses, and regulates spine maintenance but not formation. Although these data are strongly supported by the observation that mice knockout for DGK ζ show reduced spine density and excitatory synaptic transmission, the molecular mechanisms underlying these actions are not known (233).

The second lipid kinase is the phosphoinositide-3-kinase (PI3K). This kinase is mainly acting when a stimulated tyrosine kinase receptor induces the recruitment and activation of the PI3K complex (formed by the catalytic and inhibitory subunits) to the receptor. Activated PI3K leads to the production of phosphatidylinositol 3,4,5-trisphosphate (PIP₃) from phosphatidylinositol 4,5-bisphosphate. PIP₃ acts as a second messenger by activating the Akt kinase, of which downstream targets include glycogen synthase kinase 3 (GSK3) and the mammalian target of rapamycin (mTOR). This signaling cascade is involved in regulating dendritic development (211), synaptic plasticity (259), and perhaps spine morphogenesis (92, 259). Interestingly, a recent study demonstrated that a PI3K-activating transduction peptide is sufficient to induce synaptogenesis and spinogenesis in both cultures of rat hip-

pocampal neurons and in CA1 hippocampal neurons *in vivo* (92). PI3K activation was effective also in adult neurons, suggesting that functional synaptogenesis can be induced in mature mammalian brains by the specific activation of PI3K (92).

Among the kinases implicated in long-term maintenance of neuronal plasticity and memory, protein kinase M ζ (PKM ζ) is an atypical isoform of protein kinase C (PKC), which when overexpressed on primary cultures of cortical neurons modifies spine morphology by reducing their length without changing their density. The regulation of spine morphology by PKM ζ may explain its role in long-term synaptic plasticity (417).

Two families of kinases, NDR1/2 and DCLK1/2, involved in regulating cellular morphogenesis in eukaryotes and cellular migration, respectively, have been recently found to play a role in dendrites and dendritic spine morphogenesis. Genetic manipulations of NDR1/2 have demonstrated that it positively regulates spine formation by activating Rabin8, a GDP/GTP exchange factor (GEF) of Rab8 GTPase, while DCLK1/2 negatively regulates spine formation by disassembling postsynaptic density complex (460, 520).

E. Postsynaptic Scaffold Proteins and Adaptor Proteins

The PSD is an assembly of several scaffold proteins that link synaptic receptors and membrane proteins to spine cytoskeleton and signaling molecules. Among the PSD scaffold proteins, the PSD-95 and the Shank families of proteins play a critical role in spine morphogenesis.

The four members of the PSD-95 family (PSD-95/SAP90, PSD-93/chapsyn-110, SAP102, and SAP97) have a common structure formed by three PDZ domains, a SRC homology (SH3) domain, and a guanylate kinase-like (GK) domain (455).

SAP97 is longer because it has also a LIN2/LIN7 (L27) domain at the NH₂-terminal before the first PDZ domain. PSD-95 family is part of the protein superfamily called MAGUK (membrane-associated guanylate kinase), which is marked by the existence in all members of one GK and at least one PDZ domain.

As described before, PSD-95 (actually the most abundant PSD scaffold protein) is an important player in synapse and dendritic spine formation because it is the scaffold protein that interacts with neuroligin, NGL, SALM, and IL1RAPL family of proteins, which in the postsynaptic membrane mediate the trans-synaptic adhesion with specific membrane-associated presynaptic proteins to induce synapse formation. The classical example is the trans-synaptic neuroligin-neurexin interaction that regulates pre- and postsynaptic differentiation (79, 154, 399). Actually, at least in cultured neurons, the numerical

balance between inhibitory and excitatory synapses is proportional to the amount of PSD-95 expression (142, 399).

On the cytoplasmic side, PSD-95 interacts also with a broad range of signaling molecules linking together cytoplasmic signal-transducing enzymes to membrane receptors and assembling signal coupling in the PSD. Thus, at least in cultured hippocampal neurons, the overexpression of PSD-95 increases the number of dendritic spines and induces the maturation of excitatory synapses (111, 398). This is done by recruiting transmembrane proteins but also intracellular signaling, scaffold and adaptor proteins such as kalirin-7 (383), SPAR (376), IRSp53 (82), Preso (273), TANC1/2 (164), nNOS (354), Shank2 (483), and cypin (516) that are directly or indirectly involved in spine formation. In vivo, deletion of PSD-95 partially reduces spine number in the striatum and in the hippocampus (533).

SAP102 is the other member of family potentially involved in spine morphogenesis. A specific NH₂-terminal domain, the expression of which is regulated by alternative splicing, regulates SAP102 interaction with GluN2B and promotes the formation of synapses and long spines, suggesting that SAP102 mediates NMDA receptor ability to induce dendritic spine formation (72). Finally, like PSD-95, CASK, a member of the MAGUK family of proteins, also promotes spine formation by linking synaptic adhesion molecules to the actin cytoskeleton, and its activity is regulated by SUMOylation (69).

Several in vitro and in vivo studies indicate that Shank family of PSD scaffold proteins is a major regulator of dendritic spine morphology. The Shank family is composed of three genes (*Shank1–3*) that codify for a multi-domain proteins containing an NH₂-terminal ankyrin repeat, followed by a SH3, a PDZ domain, a long proline-rich region, and a sterile alpha motif domain (SAM) at the COOH terminus (454).

The protein-protein interaction domains present in Shank link at least two glutamate receptor subtypes: the NMDA receptors/PSD-95/GKAP complex, through the PDZ domain that binds to the COOH-terminal of SAPAP/GKAP, another scaffold protein of the PSD, and the type 1 mGluRs/Homer interaction at the proline-rich domain.

Shank1 was the first to be associated with spine morphogenesis by the demonstration that co-overexpression of Shank1 and Homer1b (the long isoform of Homer1 gene) in hippocampal neurons induces the maturation of spines during development and the volume enlargement of mature spines, without altering their number. Overexpressed Shank1 and Homer1b are also able to accumulate in the PSD other synaptic proteins like GKAP/SPAP and GluN1 and the F-actin (179, 433).

Homer are scaffold proteins, structurally formed of an NH₂-terminal Ena/VASP homology 1 (EVH1) domain and a

COOH-terminal coiled-coil domain that mediates tetramerization with other Homer proteins (179). The EVH1 domain specifically binds to a PPXXF sequence that has been found in Shank1–3 proteins, mGluR1/5, inositol-1,4,5-trisphosphate (IP₃) receptor, ryanodine receptor, and in some members of the TRPC family of ion channels (555, 575). Interestingly, the short-splice variant of Homer1 (Homer1a) that is deleted from the coiled-coil domain and cannot tetramerize is a natural dominant negative, and importantly, synaptic activity modulates the expression of Homer1a (14, 432, 555).

As opposed to Shank1, the overexpression of Shank3 is able to induce the formation of new spines at least in cerebellar granule cells, while Shank3 knock-down by RNAi expression in hippocampal neurons reduces the total number and the mature-like spines. Shank overexpression has a global effect on synapse maturation because it induces also the maturation of the presynaptic compartment (108, 420, 433).

The ability of Shank and Homer to regulate spine maturation can be explained by the association of Shank with cortactin (346), Abp1 (403), αPIX (379), and Cdc42-binding protein IRSp53 (472) while Homer binds to oligophrenin-1 (153). Interestingly, Shank1 and Homer1b in the PSD form a mesh-like matrix structure that serves as a structural platform to assemble other synaptic proteins that contribute to the enlargement of dendritic spines (179). Thus neurons overexpressing Homer1b mutants, unable to tetramerize, or the dominant negative Homer1a splice variant have a reduced number of smaller spines, and a reduced number of synaptic AMPA and NMDA receptors (178, 432). Finally, Shank1 and Homer1b cooperation on spine enlargement can be explained also by the demonstration that their overexpression in hippocampal neurons recruits the entire endoplasmic reticulum compartment to dendritic spines (435).

The role of Shank1 (see below) in regulating dendritic spine shape has been demonstrated also in vivo in the knockout mice that showed alteration in PSD protein scaffold composition, smaller dendritic spines, and weaker excitatory synaptic transmission (200).

In contrast to Shank, the overexpression of PICK1 in hippocampal neurons reduces spine size, and its knockdown increases spine size. PICK1 is a synaptic scaffold protein that binds AMPAR subunits GluA2/3 that contain a PDZ and BAR domain proteins (165). The PICK1 effect appears to be mediated by its direct binding and inhibition of Arp2/3 complex activity (347).

IQGAP1 is a member of a family of scaffold proteins structurally formed by a calponin homology domain that bind actin, a WW motif allowing the association with ERK1/2, the calmodulin binding IQ domain, and a RasGAP-related domain that binds to Cdc42 and Rac1. Mice knockout for IQGAP1 show a specific reduction of dendritic spine number in specific brain

regions associated with cognition and emotion, such as the hippocampus and lateral amygdala. In these brain areas IQGAP1 directly interacts with GluN2A and contributes to ERK signaling in response to GluN2A-dependent NMDAR stimulation, linking this pathway to spine morphogenesis (137).

Insulin receptor substrate 53 (IRSp53) is PSD adaptor protein that links Rac1 to WAVE2 and binds to Shank and PSD-95. Overexpression of IRSp53 in cultured neurons increases the number of dendritic spines while its knockdown reduces the number, length, and width of spine. IRSp53 might play a role in linking PSD-95 to activated Rac1/Cdc42 and spine actin because IRSp53 ability to bind WAVE2 with its SH3 domain is required for inducing spine morphogenesis (82). However, dendritic spine morphology is normal in mice knocked out for IRSp53 although several synaptic functions, such as NMDA receptor-mediated transmission, LTP, and learning and memory are altered in these mice (234, 443).

F. Micro RNA, mRNA Binding Protein, and Transcription Factors

The identification of microRNAs (miRNAs) as noncoding RNAs that mediate posttranscriptional gene silencing and their implication in neuronal development and neurodegeneration have stimulated the search for specific miRNA that play a role in dendritic spines morphogenesis and their respective molecular targets. Among the miRNAs selectively expressed in the brain, at least five have been demonstrated to possess a functional role in controlling synaptogenesis and spine morphology: miR-134, miR-138, miR-132, miR-125b, and miR-29a/b. Interestingly, several miRNA target genes codify for proteins previously demonstrated to be involved in spine morphogenesis.

The first identified miRNA, miR-134, negatively regulates the size of dendritic spines and excitatory synaptic transmission by inhibiting the translation of *Limk1* that controls spine development (see above). Interestingly, BDNF increases *Limk1* expression by relieving the miR-134 translational inhibition via the TrkB/mTOR signaling pathway, which inactivates a not well defined miR-134-associated silencing complex (445).

With the use of a functional screen, miR-138 was found to be highly expressed in the brain and specifically localized in dendrites. miR-138 negatively regulates the size of dendritic spines in rat hippocampal neurons by inhibiting the expression of the enzyme acyl protein thioesterase 1 (APT1), which controls the depalmitoylation of synaptic proteins such as the α_{13} subunits of G proteins ($G\alpha_{13}$). In addition to $G\alpha_{13}$, other synaptic proteins could be a substrate of APT1, although the spine size reduction can be caused by the elevated level of palmitoylated $G\alpha_{13}$, in the absence of APT1, which might increase the activity of RhoA (463).

Rho-family GTPase activating protein p250GAP is the target of miR132. The expression of miR132 is regulated by synaptic activity and CREB during periods of active synaptogenesis. By reducing the expression of p250GAP, miR132 increases spine formation. This is done by modulating synapse-specific Kalirin7-Rac1 signaling (166, 202) (see also below). The role of miR132 in spine morphogenesis has also been demonstrated in vivo in transgenic mice that express miR132 in forebrain neurons. Hippocampal neurons in these mice showed a marked increase in dendritic spine density, but these morphological changes are associated with significant deficits in novel object recognition and with the decreased expression of the Rett Syndrome protein MeCP2 (166). miR-132 controls synaptic plasticity and spine maturation in visual cortex (315), while in olfactory bulb (OB) neurons regulates not only dendritic complexity and spine density but also the survival of newborn neurons, suggesting that this miRNA regulates a set of genes involved in neurogenesis and structural plasticity (381). Finally, miR-132, together with miR-125b, is associated with the Fragile-X mental retardation protein (FMRP) that regulates their activity. Overexpression of miR-125b reduces dendritic spine number and maturation, an opposite effect to that of miR-132 overexpression, and inhibits the expression of GluN2A (109). miR-125a controls PSD-95 expression and spine maturation by forming an inhibitory complex with AGO2 that can be disassembled by mGluR-dependent dephosphorylation of FMRP (335).

Finally, miR-29a/b, like miR-125b, also reduces mushroom-shaped dendritic spines with an associated increase in dendritic filopodia when overexpressed in hippocampal neurons; in this case, miR-125b probably downregulates the expression of *Arpc3*, a component of the ARP2/3 actin nucleation complex (284).

Local translation of specific mRNAs in proximity to dendritic spines has been implicated in synapse remodeling and synaptic plasticity. Not surprisingly, proteins involved in controlling mRNA targeting and translation in dendrites are directly involved in regulating spine morphogenesis. In vitro and in vivo experiments have demonstrated that the absence of TLS (translocated in liposarcoma), *Staufen2*, and heterogeneous nuclear ribonucleoprotein K (hnRNPK) selectively impair mRNA targeting of proteins that regulates the number of mature dendritic spines (134, 135, 148, 402). Translation of TLS transported dendritic mRNA is controlled by metabotropic glutamate receptor (mGluR5) activation, and some mRNA for cytoskeleton proteins, such as β -actin, are specifically guided by *Staufen2*. hnRNPK interact with Abi-1 forming in a multiprotein complex reciprocal regulating spine formation and synaptic maturation (402). Another major dendritic protein involved in controlling dendritic mRNA translation is the Fragile-X syndrome protein FMRP (see below).

Neuronal development is controlled by the activity of specific subsets of transcription factors (TFs) that coordinate the ex-

pression of proteins involved in neuronal morphogenesis. Recently a subclass-specific TFs have been found to control dendritic spines and the maturation and strength of the synapses. Cux1 and Cux2 encode the vertebrate homologs of the *Drosophila* homeobox transcription factor Cut, which controls the dendrite morphology of postmitotic populations in the peripheral nervous system (PNS) (157, 214, 247). Both Cux1 and Cux2 are also expressed in postmitotic brain neurons where they coordinate the expression of Xlr3b and Xlr4b, chromatin remodeling genes previously implicated in cognitive defects. Indeed, knockout mice for Cux2 show a specific reduction of dendritic spine numbers in cortical layers II-IV neurons (91). Similarly, knockout of FoxO6, a neuron-specific member of the FoxO family of transcription factors, impairs the number of dendritic spines in hippocampal neurons by altering the expression of genes involved in synaptic function associated with learning and memory (436).

Satb1 is a genome organizer protein highly expressed in mature cerebral neurons. In postnatal cortical pyramidal neurons of Satb1 knockout mice, the density of dendritic spines is reduced probably because the transcription control of some immediate early genes (IEGs) (*Fos*, *Fosb*, *Egr1*, *Egr2*, *Arc*, and *Bdnf*) is altered; Satb1 binds to genomic loci of these IEGs (28).

G. Steroid Hormones

The first indication that dendritic spine density changes during the estrous cycle of female rats was demonstrated more than 20 years ago (551). The density of dendritic spines was correlated with the fluctuations of both estradiol and progesterone levels occurring during the estrous cycle of females, and the ovariectomy of female rats induced a decrease in dendritic spine density in the hippocampus, which could be rescued by estradiol therapy (152). Interestingly, this estradiol cyclic regulation of spine synapse density was found only in the hippocampus and not in the neocortex or in the cerebellum (400), although it has been reported that estradiol promotes in vitro and in vivo the formation of dendritic spines in Purkinje neuron (431).

The original suggestion that circulating hormones are directly controlling spine density is now questioned by the observations that synapse density was not influenced by application of estradiol to hippocampal slice cultures irrespective of the sex of the animals providing the tissue (125, 257). On the other hand, in hippocampal slice cultures, the inhibition of aromatase activity (the enzyme responsible for 17 β -estradiol synthesis) resulted in loss of spine synapses that could be rescued by the application of exogenous estradiol (257, 586). Thus these data suggest that locally produced estradiol rather than estradiol originating from the reproductive organs regulates dendritic spine synapse density.

The level of estradiol synthesis in the hippocampus is also controlled by gonadotropin releasing hormone (GnRH). Al-

though GnRH is the key regulator of reproduction, by controlling the secretion of follicle stimulating hormone (FSH) and luteinizing hormone (LH) in the pituitary and thus steroid hormone synthesis in the gonads, it can also influence estradiol synthesis directly (209). Hippocampal slice culture treatment with GnRH induced the release of estradiol into the medium and increased dendritic spine density in a dose-dependent manner (400). These data suggest that estradiol regulates dendritic spines and synapse function also via a paracrine mechanism.

On the other hand, several studies demonstrated recently that dendritic spine densities are clearly different between male and female mice in vivo. For example, the density of spines in males is similar to their density in ovariectomized females, while the hippocampus of intact cyclic females could contain double that of males (93, 125, 462). Parental estradiol treatment of ovariectomized mice caused an increase in the number of mature mushroom spines (279). This sex-specific difference in spine number correlates with the plasma levels of estradiol that is up to five times higher in females at the stage of proestrus than in males, while there is no difference between males and ovariectomized females (125). It should be added that the estradiol concentration in hippocampus of males and ovariectomized females is extremely low (125). These studies indicate that circulating estradiol rather than locally synthesized estradiol regulates spine density.

Whatever its origin, estradiol has two time constants of effects on spines: a rapid (hours) and slow (days, discussed above) one. A 2-h exposure is sufficient to significantly increase the total density of spines in stratum radiatum of the CA1 region of the hippocampus (see review in Ref. 370). This rapid effect cannot be explained by the classical transcriptional genomic pathway of the canonical intracellular estradiol receptors. Thus a putative membrane estrogen receptor or a different, more rapid pathway for the intracellular estrogen receptor should be considered for the nongenomic response.

The most probable candidates for extranuclear, dendritic, and synaptic membrane estrogen receptors (ER) should be ER α and ER β (191). Both are also classical nuclear receptors, but they may work with different mechanisms at synapses, driving either kinases, metabotropic glutamate receptor (mGluR), or G protein activation (54, 475).

ER α has been localized in spines of hippocampal pyramidal and granule neurons by immunoelectron microscopic and Western blots analysis in the PSD where it is accumulated upon NMDA stimulation (191, 336).

ER β is also abundantly expressed in cortical neurons (327) and has been shown to be required for correct memory and hippocampal synaptic plasticity because mice knockout for ER β display impaired learning (287). Finally, in cultures, the rapid activation of ER β with the specific agonist WAY-

200070 increased spine density and PSD-95 accumulation in membrane regions (478). At the mRNA and protein levels, no differences were found between sexes for these receptors (423). Thus both $ER\alpha$ and $ER\beta$ are functionally present in dendritic compartments and can be rapidly activated to modulate synapse function and spine morphogenesis.

Downstream pathways that are responsible for rapid spine cytoskeletal rearrangements have been partially identified. Blockade of the ERK pathway pharmacologically or by reducing intracellular $[Ca^{2+}]$ abolishes the effect of estradiol on the spine density (340).

In cultured cortical neurons, direct activation of $ER\beta$ by WAY-200070 induces the phosphorylation of PAK and ERK1/2 (478) while activation of $ER\beta$ induced the phosphorylation and activation of WAVE1 (439). Estradiol can also activate FAK and actin polymerization by triggering c-Src/PI3K dependent phosphorylation on Tyr397 (438). In developing hypothalamus, estradiol promotes spine formation by a rapid, nongenomic activation of PI3 kinase in presynaptic neurons that increase the release of glutamate. This triggers higher glutamate receptor number and mitogen-activated protein kinases activation in postsynaptic neurons and protein synthesis-dependent formation of dendritic spines (447).

Finally, another possible mechanism of action of estrogen is through the regulation of GABAergic transmission. Estradiol application in vitro, acting probably on both $ER\alpha$ and $ER\beta$, appears to suppress GABAergic transmission, increasing excitability in hippocampal cells and facilitating the formation of new dendritic spines (342). On the other hand, in mice treated with the specific $ER\beta$ ligand LY3201, GABAergic synapse became stronger through the higher expression of GAD in cortex and hippocampus, and this correlates with a reduction of number of spines in hippocampal neurons, suggesting that that $ER\alpha$ and $ER\beta$ have opposite effects on synapse plasticity (498).

Among the other steroid hormones, glucocorticoid activity is also important for dendritic spine plasticity. Chronic glucocorticoid treatment leads to loss of stable spines, suggesting that glucocorticoids play a role in the development and maintenance of dendritic spines in cortical neurons (286). More recently it has been demonstrated that corticosterone reduces the expression levels of caldesmon (CaD), a ubiquitous actin-linked regulatory protein that binds to and stabilizes actin filaments, causing dendritic spines to become less stable. Corticosterone-mediated suppression of CaD expression levels is caused by inhibition of serum response factor-dependent transcription. Because CaD enlarges the spine-head size by stabilizing F-actin dynamics, CaD is a critical target in the corticosterone-induced damaging effects on dendritic spine morphogenesis (500).

In contrast, it has been demonstrated that stress-responsive glucocorticoid receptor (GR) is present on dendritic spines and regulates local signaling of the actin cytoskeleton. Corticosterone rapidly (60 min) increases the number of large spines and total levels of phosphorylated ERK1/2 proteins. Thus acute corticosteroids treatment promotes spine formation via multiple kinase networks including PKA and mitogen-activated protein kinase (246). In summary, corticosteroids might mediate both acute stress induced spine morphogenesis and synapse formation and chronic stress mediated spines elimination.

V. DEVELOPMENT OF DENDRITIC SPINES

A. Is Spine Formation Affected by Active Synaptic Interaction?

The rules that govern the formation of dendritic spines in the developing brain, as well as the question of whether these mechanisms are different from those that regulate spine formation in the adult brain are still not resolved. Two observations are common: in the young brain there are few spines and many filopodia which disappear in the mature neuron (126, 473). Dendritic filopodia, unlike growth cone-associated filopodia, are extremely sensitive to ambient activity; they form and are more motile if action potentials are blocked with TTX, or if synaptic activity is blocked (397), indicating that they may play a role in the formation of connections among neurons, and in the synchronized activity typical of young networks. On the other hand, most of the synapses in the young brain reside on the dendritic shaft (412). Do spines form from filopodia, which are motile and transient by nature, or are spines formed from existing shaft synapses on dendrites? An intermediate possibility is that dendritic filopodia shoots out to sense a source of glutamate release in an axon (e.g., Ref. 415), whereas a shaft synapse is formed at the same location and develops into a spine (169, 174). Removing the synapse away from the shaft, by the production of a spine, may allow the cell to generate more synapses with en-passage axons, and be able to scale down the efficacy of the synapse, such that a spine synapse will have a lower impact on the parent neuron. Synaptic connections leading to spontaneous network bursts have been seen also in young cultured and in vivo neurons (3–4 days old) before the formation of spines, and these are likely to result from shaft synapses (271). Likewise, the conversion of filopodia to spines is not likely to occur in mature neuron, when filopodia are rare, and spines can form within minutes to hours, as illustrated in cycling female rats (552) and during the recovery from hibernation (394, 396) as well as in living adult mice (561, 563). Interestingly, exposure of neurons to TTX, which blocks action potential discharges, causes filopodia to grow off existing spines even in mature neurons (415), indicating that the neuron is extremely sensitive to changes in ambient activity, to the extent that even if it does not support the existence of filopodia, they are generated

within minutes after the demand emerges. On the other hand, Reilly et al. (412) imaged developing neurons in culture and were unable to observe a shaft synapse that converts to a spine synapse over 31 h of monitoring. Thus, even though both shaft synapses and filopodia are abundant in juvenile neurons, the eventual spine synapse does not seem to develop from a shaft synapse. Still, young neurons are far more responsive to glutamate challenge and develop dendritic spines than adult neurons (591; see also Ref. 264), and the reasons for this are not entirely clear. Likewise, the fate of the shaft synapse as well as the stimulus that triggers the formation and location of a spine synapse need further exploration.

VI. SHORT-TERM MORPHOLOGICAL AND FUNCTIONAL PLASTICITY OF SPINES

A. Formation of New Spines

Early time-lapse imaging of dendrites following LTP-inducing protocol described the formation of new spines in cultured hippocampal slices (115). The new spines were detected within 30 min after the induction of LTP in the slice and amounted to an addition of ~15% to the existing population of spines. Interestingly, the authors did not report on a change in volume of the existing spines on these dendrites following LTP induction. Similar observations were made by Shi et al. (458) who described NMDA receptor-dependent formation of new filopodia in response to tetanic stimulation in cultured hippocampal slices. In neither of these studies was there any indication that the growth of the new spine or filopodia is associated with formation of new synapses, although the electrophysiological responses recorded in the slice showed the typical rapid growth of synaptic responses associated with LTP generation. These issues were addressed in a study with dissociated hippocampal neurons (150, 373) where new spines were found to be attached to synaptophysin-containing terminals. Still, the issue of when do nascent dendritic spines become functional was subject to further explorations. Bonhoeffer and colleagues (344) combined confocal microscopy with EM studies of cultured hippocampal slices to suggest that physical contacts with presynaptic fibers are made within tens of minutes after enhanced activation, but mature spines with presence of both pre- and postsynaptic elements, indicating a functional synapse, are formed only 15–19 h after the plasticity-evoking stimulation. In a similar approach, Zito et al. (591), also using cultured hippocampal slices, were able to record responses to flash photolysed glutamate from young (<2 h) spines, responses that in many respects were indistinguishable from mature spines of the same volume. This indicates that new spine already contain AMPA receptors, confirming earlier work with dissociated cultures (58, 150, 365). More recently, Kwon and Sabatini (264), working in acute cortical slices of young (P8–12) mice, found that new spines could emerge within <2 min after glutamate uncaging near the dendritic shaft of layer 2–3 pyramidal neurons. These new spines are functional in the sense that they respond to presyn-

aptic stimulation, much the same way as old dendritic spines. Interestingly, in this experiment glutamate uncaging did not cause expansion of existing spine heads (see below).

On the other extreme is the study of Knott et al. (241) done with time-lapse imaging of in vivo neocortex, suggesting that only 4 days after their formation dendritic spines become innervated and mature.

Altogether, these results do not provide a clear universal time course for the relations between spine formation and its functions. The time ranges between minutes, or as soon as a spine is formed, to hours and days. The reason for this broad time spectrum is not entirely clear, and it may have to do with the age of the tissue, the stimulation, and detection protocols. A recent observation addressed this age possibility by showing that nascent spines that are exposed repeatedly to flash photolysis of caged glutamate (LTP protocol) have a higher propensity to survive in mature form, compared with unstimulated ones (188). This stabilization is dependent on activation of the NMDA receptor and its linked CaMKII, but it is not clear that these novel mature spines are at all innervated.

B. Spine Expansion

Individual dendritic spines that are activated by electrical stimulation or by photoconversion of caged glutamate can undergo a nearly immediate (<1 min) expansion of their head volume by over threefold (311). This was accompanied by a rapid (2 min) increase in synaptic responses to cage glutamate, likely to be mediated by lateral diffusion of glutamate receptors into PSD, which can take place within a fraction of a second (182a). Interestingly, the authors proposed that only small spines can expand, while the large ones are resistant to the effects of caged glutamate and do not expand. The initial expansion was dependent on activation of NMDA receptors in the spine head while a long-lasting expansion was dependent on activation of kinases (566). Later studies reported that the change in spine volume was smaller than in the original report [Yang et al., 40% (566); Lai et al., 40% (267)] and may develop over a longer time period [20 min (267, 583) or longer (442)]. Regardless of the magnitude of the immediate effects, the ability of spines to change their volume over a short time is impressive indeed. Furthermore, spine expansion is associated with an increase in the density of glutamate receptors in the spines, as seen following induction of LTP (e.g., Refs. 188, 310). The question is if these volume changes are a necessary condition for the electrophysiological change or are just a by-product of the functional change. A corollary issue is if all types of plasticity (e.g., LTP, STDP, LTD) involve a volume change in the affected spines. The differences between the different reports on the latency and magnitude of spine head expansion may be related to the insertion of glutamate receptors into the spine heads such that spines will inflate only when glutamate receptors are added into their heads (248, 250).

A clear dissociation between spine expansion and LTP has been demonstrated by Yang et al. (566), where blockade of protein kinase A did not block the initial spine expansion, but its maintenance, whereas it did block generation of LTP. Morphological factors, such as spine neck length and initial head dimension, may cause spine expansion to be fast (172, 276, 311), or rather slow (267). In a recent study by Lynch and co-workers (255), they suggest that a difference between young (2 wk) and adult (8 wk) neurons recorded in acute hippocampal slices may underlie this heterogeneity, in that the young mice spines can show an immediate, twofold expansion, whereas spines in the adult slices are less prone to expansion, and when they do, they lack the immediate surge in spine volume, and inflate to roughly 40% above their normal size.

The role of the expanded spine in synaptic plasticity has been alluded to also in the study by Matsuo et al. (310), who found that newly synthesized GFP-GluR1 molecules tend to accumulate in large mushroom spines, and not in thin spines or stubby ones, 1 day after fear conditioning. This effect is transient and dissipates 2 days later. This observation is similar to a previous EM study by Ostroff et al. (372), showing that polyribosomes accumulate in dendritic spines 2 h after a tetanic stimulation. These spines also have larger postsynaptic density, indicating that protein synthesis contributes to the maintenance of spine inflation, and to the enhanced synaptic transmission in these spines.

Several exceptions to the correlation between spine expansion and magnitude of synaptic responses have recently been described. One of the most studied plastic change in synaptic properties is the homeostatic scaling (518). In a recent study, conducted with cultured slices of the hippocampus exposed to TTX, Soares et al. (467) describe a significant increase in mEPSC amplitude and frequency, without a change in spine size or density. They ascribe the functional change in synaptic currents to a change in the balance between different glutamate receptor subtypes.

A similar diversion from the general correlation between spine size and synaptic current has been described in a phenomenon called distance-dependent scaling (DDS) (465). It has been known that synapses on remote dendrites generate much larger synaptic currents than synapses on proximal dendrites. Smith et al. (465) could not find a correlation between the large DDS of synaptic current and spine size. In a recent study, Shipman et al. (461) proposed that this scaling is associated with a change in GluA2 subunit density on the remote spines, illustrating that a change in synaptic current is not necessarily associated with spine volume.

Finally, the spine volume variations have important implications with respect to the involvement of intracellular calcium concentration in synaptic plasticity. This was alluded to in a recent theoretical study (362), suggesting that changes in spine volume may result in different mag-

nitudes of surges of spine $[Ca^{2+}]_i$ which will affect the stability of synaptic plasticity.

VII. LONG-TERM CHANGES AND DENDRITIC SPINES

The longevity of dendritic spines has been the subject of intense investigation since the introduction of chronic time-lapse imaging of spines in the intact neocortex. In one of the earliest studies, Trachtenberg et al. (511) found a large fraction (50%) of stable spines in mouse barrel cortex. They further found that plasticity increases the turnover of spines, in line with a later suggestion that sensory deprivation prevents spine loss, found during normal development of the neocortex (592). In contrast, small retinal damage caused a threefold increase in spine turnover in the affected visual cortex (226), leaving the issue of whether sensory deprivation facilitates or suppresses spine turnover rate still open. In a different study, Holtmaat et al. (194) found that thick spines persist for a month, whereas thin spines are transient, with a slower turnover of spines in the visual compared with the somatosensory cortex. This corroborates the suggestion that thick spines are maintaining “memory,” as suggested in the *in vitro* studies (above). Furthermore, a long-term change in spine generation and elimination dynamics resulting in a reorganization of synaptic connectivity has been reported to take place in the mouse neocortex following extensive motor learning. These changes lasted many days after the training (561, 563). Some possible differences in the proportion of stable spines found between different studies may result from differences in type of cranial window produced to image the spines, as results with cranial window where the skull was completely removed produced different results from those where the skull was thinned down, but not removed, maintaining a healthier cortex (560).

While the general finding is that spine density/size increase with long-term memory, and that large spine heads indicate “memory” (223), there are notable exceptions. In two recent studies, fear conditioning is actually associated with a reduction in spine density. In one of these studies (440), the authors analyzed hippocampal neurons where GFP-GluR1 was generated by a *c-fos* promoter, indicating that these neurons are activated by the learning experience. Compared with home-caged mice or mice exposed to the context alone, fear-conditioned animals had fewer dendritic spines on the activated neurons. In the other study (267), the authors used time-lapse microscopy in living mice, focusing on dendritic spines of layer 5 pyramidal neurons of frontal association cortex. In these mice, fear conditioning was associated with elimination of dendritic spines, whereas fear extinction involved reappearance of spines, in close proximity to the ones that were eliminated by fear conditioning.

A. Spine Pruning and LTD

LTD has been considered as a parallel and opposite process to the more common LTP. Both processes involve a rise in

[Ca²⁺]_i, probably to different magnitudes and duration. There are several protocols for induction of LTD, including a low-frequency (1 Hz) stimulation, as well as an exposure to metabotropic receptor agonist. The electrophysiological and biochemical rules that govern LTD have been studied extensively, but the morphological basis of LTD is much less explored. While there is a general consensus that LTP or enhanced synaptic activity is correlated with an increase in spine size/density, there is less information on the possible parallel reduction in dendritic spine density/shape with LTD, which causes a reduction in reactivity to afferent stimulation. In an early study, Nagerl et al. (361), using an organotypic hippocampal slice, found that spines retract, in an NMDA-dependent manner, with low-frequency stimulation, which causes LTD. Interestingly, the ability to retract spines is drastically reduced with age of the culture, as is spontaneous formation and retraction of spines. Altogether, the number of retracted spines is rather low (2–3 spines/100 μm of dendrite). In another study, Kamikubo et al. (221) found that regardless of the means for induction of LTD, it is associated with synapse loss. Interestingly, the synapse loss was measured weeks after the induction of LTD, and it required three exposures to the LTD generating stimulation. Thus it is not clear if the morphological change is in any way related casually to the electrophysiological LTD.

In another recent study using LTD induction protocol in cultured slices of the hippocampus, a significant retraction of presynaptic boutons was described, even more striking than the earlier report on spine shrinkage (34). Altogether, these studies indicate that LTD is associated with a reduction in functional synapses with LTD, even though the actual proportion of reduced synapses does not match the marked electrophysiological changes that are seen following induction of LTD.

Does a neuron lose its synapses when spines are pruned, or does it retain the synapses on the dendritic shaft? Since a mature synapse contains an array of adhesion molecules linking the pre- and postsynaptic membranes, and the synapse is engulfed by glial cells, it is unlikely that the presynaptic terminal detaches from the postsynaptic side, especially since the synapse is tough enough so as to resist mechanical dissociation (as in the process of preparation of synaptosomes). However, recent EM study suggests that cortical neurons do lose their connection with afferent inputs (241). In contrast, there is a marked decrease in spine density in hibernating animals that is counteracted by an increase in shaft synapses (396, 536). When the animal rises from hibernation, the spines reappear, and the animal remembers tasks learned before hibernation, indicating that persistence of spines is not a prerequisite for memories (85). In fact, when trained 24 h after arousal from hibernation, animals remember better than controls (545). Likewise, female rats in estrus cycle, when their hippocampal spine density is down by 30%, are not less capable of remembering items learned during earlier time in the cycle. It is there-

fore unlikely that neurons lose their afferents once their spines disappear. On the other hand, when the cell loses its afferents, as is the case in the cultured hippocampal slice, the denervated spines disappear, only to reappear when new axons innervate the vacated region on the dendritic shaft (132).

In a similar study, conducted with cultured slices, chronic blockade of AMPA receptors led to disappearance of spines, but this was compensated by the appearance of shaft synapses, and by an increase in efficacy of synaptic transmission (309). Thus reduced synaptic activity, as is the case with LTD (above), should not necessarily lead to disappearance of synapse, even though the spines may dissolve.

Another interesting disparity is the observation that long-term sensory deprivation prevents rather than enhances spine pruning (592). Still, input deprivation produced four synapses away from the monitored neuron in the barrel cortex is different from a local ongoing blockade of activity by TTX, especially since the extrinsic sensory afferents constitute only a fraction of the excitatory input to the cortical neuron. In another study, blockade of activity with TTX increased rather than decreased the number of functional synapses, as evidenced by the staining with presynaptic proteins, but these studies were conducted with 2- to 3-day-old hippocampal slices, at an age where few if any spines are functional and the excitatory synapses are made on the dendritic shaft, so the rules that govern the structure-function relations may be different at this young age (271).

B. Elementary Units of Plasticity

What is the elementary unit of plasticity? Is it the individual dendritic spine or a cluster of them? Are they all placed on the same dendritic segments, or can they synchronize on different dendritic branches? As indicated (173), activated Ras can diffuse away from a spine into adjacent ones. In fact, adjacent spines on a dendritic branch are frequently activated synchronously in a network, which indicates that they probably react to common inputs (495). This synchronization contributes to the strengthening of the network “memory” and confirms earlier rules suggesting a need for cooperativity to produce LTP (546). Clusters of spines are formed on adjacent locations on dendrites in motor learning tasks, and these are longer lasting than individual spines grown in association with the learned task (133). Thus spine clustering may be an efficient way to maintain spines that are formed during learning, and this may have functional significance. In similar studies on sensory cortex, Makino and Malinow (300) described functional clustering of adjacent spines in response to sensory stimulation. Interestingly, sensory deprivation causes a nonselective enhancement of reactivity of spines, not specific to clusters of spines/dendrites. This homeostatic upregulation is similar to silencing-induced upregulation of synaptic activity, seen originally in cultured neurons (518). In a more recent study, learning of motor skills caused an increase in parallel fiber multiple

synapse boutons, contacting adjacent spines on the same dendrite (274a). Interestingly, spines adjacent to those innervated by a large bouton appeared to shrink, indicating that concerted weakening of adjacent spines may constitute a homeostatic balancing of synaptic strength. Interestingly, in young neurons, prior to formation of dendritic spines, there are already clustered bouts of activity in adjacent synapses (238a) that are dependent on sodium spikes. This indicates that synchronized activity of adjacent synapses is important for the formation and strengthening of developing networks.

VIII. MOLECULES IN SYNAPTIC PLASTICITY

Neuronal activity profoundly influences spine morphology. Calcium influx through NMDA receptor activation has been shown to play a critical role in regulating spine size and density because several molecular events that control actin polymerization in spines are Ca^{2+} dependent (311, 588). Brain-derived neurotrophic factor (BDNF) has also been suggested to play both a major neurotrophic role in controlling spine development but also stimulation-dependent spine growth. Both NMDA receptor and TrkB, the receptor for BDNF, activate signaling molecules present in the spines or recruited upon stimulation. Some of these recruited molecules play an essential role in synapse and dendritic spine plasticity.

The importance of NMDA receptor is supported also by genetic experiments. Deletion of the GluN1 subunit in cortical layer 2/3 pyramidal neurons leads to a decrease in spine density, but the remaining spines have a larger head size as well as an increase in the miniature excitatory postsynaptic current amplitude and frequency (521). In brain slices, knock-down of GluN1 increases the motility of spines and over time leads to their loss. Interestingly the C-tail of the receptor is required for the long-term stabilization of synapses and spines and not only for the ionotropic properties of NMDA receptors (10). However, the NMDA receptor composition and the different signaling downstream of its activation are important for the effect on spine morphology. For example, mice lacking GluN2B in pyramidal neurons of cortex and CA1 region of the hippocampus show substantial alterations in the NMDAR-dependent LTD and also exhibit decreased dendritic spine density (59). In medium spiny striatal neurons, the blockade of GluN2A-containing receptor with antagonist NVP-AAM077 induces a significant increase of spine head width while treatment with D1 receptor agonist (SKF38393), which induces a decrease of GluN2A-containing NMDA receptors, also increase spine head width. On the other hand, the GluN2B antagonist (ifenprodil) abolishes the spine morphological effect of the D1 agonist. Thus NMDA receptor composition at the corticostriatal synapses plays an important role in determining the structure and function of dendritic spines (527).

The most important molecule responding to a rise in $[\text{Ca}^{2+}]$ upon NMDA receptor activation is the calcium-sensing cal-

modulin (CaM) that activates the serine/threonine kinases CaMKI, CaMKII, and CaMKIV. These kinases can phosphorylate several proteins implicated in spine morphology and plasticity including Kalirin-7 and other signaling and scaffolding proteins (470, 542, 558). Recently it has been demonstrated that the CaMKII phosphorylation of Rpt6 proteasomal subunit and proteasome activation is required for NMDA receptor-dependent, activity-induced spine outgrowth (163).

Other neurotransmitters might modulate morphology and plasticity of dendritic spine. Among them is serotonin. Activation of the 5-HT_{2A} receptors in pyramidal neurons increases spine size via a Kalirin-7-Rac1-PAK-dependent pathways. These data might have an important implication in neuropsychiatric disorders for the link between serotonergic signaling, altered in multiple neuropsychiatric disorders, and dendritic spine morphogenesis (217).

As mentioned before, dopamine is also a relevant neurotransmitter in its ability to modulate dendritic spines plasticity. The number of dendritic spines is reduced in striatal medium spiny neurons of rats treated for 3 wk with 6-hydroxydopamine, the neurotoxin that selectively kills dopaminergic and noradrenergic neurons (471, 540). Woolfrey et al. (550) showed that D1/D5 receptors activation with selective agonist SKF-38393 induced spine shrinkage by activation of the RapGEF protein Epac2.

The cholinergic system has also been implicated in dendritic spine morphology. Cortical pyramidal neurons of layer V have lower spine density after the specific elimination of the cholinergic system using saporine-conjugated antibodies against nicotinic receptors (192 IgG) (457). In vivo studies have demonstrated that $\beta 2$ subunit of the acetylcholine nicotine receptor knockout induces a decrease of spine density in the higher order association areas (29). Thus activation of $\beta 2$ -containing nAChRs by nicotine in vivo or in organotypic slice culture rapidly increases the number of spines by inducing intracellular calcium rise and activation of CaMKII (289).

Finally, some genetic evidence suggests that inhibitory synapses also control dendritic spine development; mice lacking the GABA_A receptor $\alpha 1$ subunit showed an increased density of dendritic filopodia during the second and third postnatal weeks and a parallel decrease in density of mature mushroom spines (169). These studies suggest that correct balancing between excitatory and inhibitory synaptic transmission is important for regulating outgrowth of filopodia and spine maturation (183). Thus these data underline the relevance of the main neurotransmitters signaling in modulating dendritic spine plasticity.

Among the neurotrophic factors, BDNF and the high-affinity receptor TrkB (tropomyosin-related kinase B) have been strongly implicated in synaptic and dendritic spine formation

and plasticity (291, 293). Several studies have demonstrated the ability of BDNF to change dendritic spine density and morphology in a variety of neuronal populations (12, 291, 293, 499, 519). Not surprisingly, TrkB knockout mice have significantly fewer dendritic spines on CA1 hippocampal neurons (292). In contrast, hippocampal pyramidal neurons of mice carrying loss-of-function mutation of the pan-neurotrophin receptor p75^{NTR} have a higher spine density, suggesting a possible antagonistic action of p75^{NTR} on TrkB receptor (576). The activation and phosphorylation of ERK1/2 is necessary for BDNF to increase dendritic spine functions, but TrkB receptor can also interact and phosphorylate Tiam1, the Rac1 GEF protein (9, 328). Interestingly, BDNF seems to have a specific local activity because its mRNA is transported and translated in dendrites in an activity-dependent manner (12, 531).

As mentioned above, synaptic stimulation induces changes in spine morphology probably by recruiting some spine-specific actin regulatory proteins. This is the case with the two actin-binding proteins profilin II and N-catenin that are accumulated in spines by either synaptic NMDA receptor activation or LTP induction. Profilin II translocates into dendritic spines following the raise of postsynaptic Ca²⁺ levels and by the association with Ena/VASP family proteins and remains concentrated in spines for hours. By locally reducing actin dynamics, profilin II can stabilize spine morphology and reduce motility upon stimulation (4). In the lateral amygdala, the enhancement of associatively induced synaptic responses to fear learning is also molecularly mediated by the recruitment of profilin II to spines and the associated enlargement of PSD (268). These results suggest that profilin II links the synaptic activation of glutamate receptors with the functional stabilization of spine morphology to a mature shape. A major role in regulating dendritic and spine morphology of both profilin I and II have been supported also by genetic studies (323).

α N-catenin is a cadherin-associated protein that accumulates in activated synapses and links the adhesion molecules to actin filaments in association with β -catenin. In cultures, transfected α N-catenin promotes the maturation of dendritic spines by inhibiting their motility and stabilizing their shape. Thus endogenous α N-catenin may mediate neural activity-dependent spine stabilization by suppressing their motility and turnover when recruited at synapses (1).

Two other actin binding proteins Drebrin and Synaptopodin are recruited to spines and promote actin assembly and spine stabilization upon LTP induction or NMDA stimulation (136, 368, 494). As mentioned before, Synaptopodin regulates also Ca²⁺ concentration in spine and spine structure because the genetic deletion of synaptopodin alters synaptic plasticity probably because dendritic spines became devoid of spine apparatus (100).

As mentioned before, LTD is accompanied by a marked shrinkage of spines. Spine shrinkage requires activation of NMDA receptors and calcineurin and is mediated by the actin-severing protein cofilin (588). Calcineurin and phosphatidylinositol 3-kinase (PI3K) mediate dephosphorylation and promote cofilin translocation to dendritic spines, which is partially mediated by β -arrestin-2 (392).

Finally, activity-dependent remodeling of synapses is also mediated by protein kinase Polo-like kinase 2 (Plk2, also known as SNK). Synaptic activity induces Plk2 expression and targeting to dendritic spines where it phosphorylates and induces degradation of RapGAP protein SPAR that causes loss of mature dendritic spines (375).

IX. DISEASE-RELATED CHANGES IN DENDRITIC SPINES

Most of the neurological disorders characterized by alteration in cognition, emotion, and memory loss are often caused by altered synaptic connectivity and plasticity (for reviews, see Refs. 223, 384). Not surprisingly, anatomo-pathological analysis of brains of patients affected by Alzheimer's disease, schizophrenia, intellectual disabilities, and autism spectrum disorders often shows alteration in dendritic spine morphology (see **FIGURE 4**).


A. Alzheimer's Disease

Several studies suggest that synapse and dendritic spine dysfunctions are preceding and contributing to the eventual neuronal death and neurodegeneration in Alzheimer's disease (AD) (307, 572). This is also supported by many postmortem tissue samples from patients with AD that consistently showed prominent synapse and dendritic spine loss in the hippocampus and throughout the cortex, the principal areas affected by AD-related pathology (99, 240, 537).

Indeed, the number of synapses and spines eliminated is often greater than the expected level for the amount of lost neurons and better correlates with the cognitive decline, suggesting that the primary synaptic failure is a prominent pathogenic cause of AD (99, 537).

The identification of specific genes associated with the pathogenesis of AD will lead to better understanding the molecular mechanisms underlying synapse and spine alterations. Mutations in three major genes implicated in beta amyloid (A β) metabolism (APP, PSEN1, and PSEN2) have been associated with familial AD with an autosomal dominant form of inheritance and early onset of the disease (37, 38). These mutations cause a well-documented increase in production of pathogenic A β oligomers that at the end are responsible for inducing spine alterations, and reducing spine density (265, 452).

The role of these mutations has been demonstrated using several transgenic mouse models of AD that have been generated



Many other genes have been identified as high risk factors for the development of AD in the elderly. Among these genes, apolipoprotein E (ApoE) is the most important and studied risk factor. ApoE has three major isoforms: apoE2, E3, and E4. ApoE4 is associated with a high risk of developing the

But which are the molecular mechanisms causing spine degeneration in AD? Recent results suggest that some signaling pathways regulating synaptic plasticity are involved. For ex-

ample, cofilin and drebrin, the actin-binding proteins with opposite effects on actin dynamics (see before), are both affected in AD. Cofilin phosphorylation and activity are upregulated by A β 1–42 peptide in a concentration-dependent manner, resulting in alteration in actin polymerization (240, 302). In striking contrast, drebrin, the postsynaptic protein that binds and stabilizes actin in spines, is severely reduced in the brains of patients with AD and in transgenic animal models of the disease (90, 147).

PAK is another important factor that regulates spine morphology, the activation of which is strikingly reduced in both AD patients and in mice models of AD (584). mRNA and protein levels of kalirin-7 were also found to be considerably reduced in the hippocampus of patients, indicating an alteration of the kalirin-7/Rac1/PAK pathway causes spine pathology in AD (570, 571).

More recently it has been shown that the partitioning defective-1 (PAR-1)/microtubule affinity-regulating kinase (MARK) family kinases is a potential mediator of A β toxicity in dendritic spines. MARK4 overexpression in cultured neurons mimics A β effect on synapses and dendritic spines, while a peptide that is specifically able to inhibit the activities of all PAR-1/MARK family members eliminates the toxic effects of A β oligomers on dendritic spines and synapses (573).

Most of the signaling molecules altered by A β described above are downstream of NMDA receptor, and overactivation of the NMDAR/calcineurin/GSK-3 β pathways is specifically induced by A β treatment. Calcineurin is a calcium-sensitive phosphatase that regulates synaptic plasticity and is required for AMPA receptor internalization and LTD. Indeed, A β oligomer-induced AMPA receptor internalization and spine loss is prevented by calcineurin inhibition (197, 554).

Finally, it is important to note that AD patients have altered expression of many pre- and postsynaptic proteins and that, at least in cultured neurons, A β oligomers destabilize some synaptic protein complexes, like Shank and Homer platform; reduce the surface expression of NMDARs and AMPARs; and induce synapse and spine collapse (20, 197, 419, 466).

Identifying the molecular mechanisms of the A β toxicity in dendritic spines may be the focus of targeted pharmacological intervention and may open new therapeutic avenues towards the cure of AD.

B. Schizophrenia

Schizophrenia is a psychiatric disorder involving disturbing thoughts, emotions, cognition, and perception of reality, with an incidence of 0.5–1% of the population. It emerges in late adolescence or early adulthood. Postmortem studies in human brains have reported grey matter loss and altered spine density in specific brain regions such as the dorsolateral prefrontal

cortex (DLPFC), an area that shows reduced activity during cognitive tasks in the affected individuals (146, 149, 497). Similarly, a reduction in grey matter volume in superior temporal gyrus as well as in spine density has been consistently reported (491, 569). Likewise, a reduction in hippocampal volume associated with reduced spine density on subicular and CA3 dendrites has been found (245, 482). Altogether these results propose a strong association between loss of grey matter, reduced spine density, and functional hypoactivity in specific brain regions of patients affected by schizophrenia (244).

The discovery of specific gene mutations associated with schizophrenia accelerated the search for the molecular basis of spine alteration observed in schizophrenic patients. Among the hundreds of gene variants that have been associated with schizophrenia, up to now few of them have shown constant associations with schizophrenia: DISC1, ERBB4, and NRG1 are the three most studied in relation of synapse and spine function. However, rare but very penetrant CNVs have been recently found in many genes encoding synaptic proteins, but these genomic studies need to be confirmed by functional experiments (538). Also, mutations in some synaptic genes could be associated either with schizophrenia or intellectual disabilities and autism, suggesting that some pathogenic mechanisms are shared by the two neurological diseases (139).

The disrupted in schizophrenia 1 (DISC1) gene was identified in a Scottish pedigree with a disruption in the DISC1 open reading frame (479). Subsequently, DISC1 polymorphisms, SNPs, and frame shift mutations affecting DISC1 protein function have been linked to schizophrenia in other families (285, 446). More recently, DISC1 functions have been associated with neurodevelopmental disorders that clinically manifest schizophrenia plus mood or schizoaffective disorder (57).

In cultured cortical neurons, DISC1 knockdown reduces spine area (181), and in mice that carry DISC1 mutations that mimic a well-defined schizophrenia risk allele, spine numbers in dentate gyrus granule cells are reduced (262). However, in another transgenic mouse expressing an inducible human DISC1 mutant, spine density in cortical pyramidal neurons was increased when the expression of mutant DISC1 was induced prenatally, while combined prenatal and postnatal expression of mutant DISC1 increased spine density specifically in dentate granule cells (24). Thus DISC1 controls spine morphology in a specific developmental and mutation-dependent manner.

Molecularly, DISC1 is a scaffold protein interacting with a regulator of spine morphogenesis such as kalirin-7, and regulates kalirin-7 binding to PSD-95 and access of to Rac1 in response to NMDA receptor activation (181). Strikingly, kalirin knockout mice express strongly reduced spine density in the frontal cortex and present schizophrenic-like behavior including impairments in working memory, prepulse inhibition, and altered sociability (64, 556). Even more remarkably, the expression of kalirin-7 mRNA was reduced in the DLPFC

(187), and several missense mutations of the kalirin-7 gene were identified in association with schizophrenia (260).

Polymorphisms in NRG1 and a rare CNV for the NRG1 receptor ERBB4 have been strongly associated with schizophrenia (314, 451, 538). As mentioned above, mice knocked out for these genes show alteration in spine density, and these morphological alteration are associated with several schizophrenia-relevant behavioral phenotypes that can be treated with clozapine (30, 278, 314).

The 22q11.2 microdeletion syndrome accounts for 1–2% of schizophrenia cases, although only 20–25% of individuals affected by this syndrome manifest schizophrenia beyond other symptoms including severe intellectual disability. Morphologically, all mice models for the syndrome [Df(16)A^{+/-} and the knockout for single genes present in this region ZDHHC8 and DGCR8] show reduced hippocampal spine density and sizes (337, 481). ZDHHC8 is a palmitoyl transferase that contributes to PSD-95 palmitoylation and localization to synapses. Dgcr8, instead, regulates miRNA processing and neuronal development. Interestingly, Gogos and co-workers (559) found that in the Df(16)A^{+/-} mouse the altered spine development is caused by a specific reduction of a micro RNA, miR-185, localized in the deleted chromosomal region, that mediate the repression of a newly discovered Golgi apparatus resident protein termed Mirta22. The modes by which this protein regulates dendritic spine formation remain to be elucidated.

C. Intellectual Disability and Autism Spectrum Disorders

Intellectual disability (ID) and autism spectrum disorders (ASD) are severe neurodevelopmental disorders characterized, respectively, by low intelligence quotient (IQ) and impairment of social interactions and adaptation. The two clinical manifestations are often present in the same individual, and indeed, ~75% of people with ASD also show a nonverbal IQ below 70, a classical sign of ID.

This group of neurodevelopmental diseases affects 1–2% of children, with diagnosis usually occurring below 3 years of age (409, 509). Numerous mutated genes for synaptic proteins have been associated with both ID and ASD, and functional studies have shown that these genes are regulating synapse and dendritic spine formation and plasticity (532).

Neuropathological data for spine morphological alteration in ID and ASD patients have been collected using Golgi-impregnated postmortem human brain tissue. Two major phenotypes have been described. The reduced number of spines but normal morphology is the phenotype often associated with ID, whereas an increase in immature elongated spine density accounts more for ASD and fragile X syndrome (127, 201, 205). These data support the suggestion that develop-

mental retardation of neuronal connectivity constitutes the hallmark of brain dysfunction in these patients.

The genetic search for mutations in ID and ASD patients is still under way, and several reports have identified rare mutations and CNVs in many synaptic genes. We will summarize the function of some of these genes in relation to dendritic spine alteration. Not surprisingly, most of these genes code for proteins that are involved in synapse formation like adhesion and scaffold molecules or in regulating actin polymerization through small GTPase pathways (47, 525).

We mentioned before the relevance of the Neuroligin-Neurexin protein complex for synapse and dendritic spine formation. A strong association of mutations in Nrnx1, Nlgn1, Nlgn3, and Nlgn4 genes were found in patients with ASD (reviewed in Refs. 94, 98, 487). However, in addition to the strong in vitro evidence for the ability of Neuroligin proteins to regulate spine number and formation, the knockout mice and transgenic mice mimicking the human mutations do not show alterations in spine morphology (or dendritic spine morphology analysis were not reported) except for the α -neurexin, double or triple knockout of the three isoforms, that showed a modest reduction of dendritic spine number in cortex (45, 87, 104, 208, 493). Thus in vivo mutation or deletion of one Neuroligin and Neurexin perturbs functionally of the synapses, but the overall structure remains normal.

Mutations of the interleukin-1 receptor accessory protein-like 1 (IL1RAPL1) gene are linked to autism and ID (67). In addition to the role of IL1RAPL1 in regulating dendritic spine formation through the interaction with PSD-95 and RhoGAP2 protein (382, 524) (see also above), the extracellular domain of IL1RAPL1 is able to induce presynaptic differentiation by binding to PTP δ , suggesting that IL1RAPL1 has a function similar to other synaptic adhesion molecules (524).

The tetraspanin 7 (TSPAN7) protein is codified by the *TM4SF2* gene and is a member of the membrane protein tetraspanin superfamily. It is formed by four transmembrane domains linked by a short extracellular loop (EC1), a very short intracellular loop (IL), a longer extracellular loop (EC2), and short NH₂- and COOH-terminal cytoplasmic tails. Various mutations in the *TM4SF2* gene (2, 579) have been directly associated with nonsyndromic ID. A possible function of TSPAN7 in promoting filopodia and dendritic spine formation by binding to PICK1 and regulating AMPA receptor trafficking to synapses has been proposed (31).

Several genes linked to ID and autism code for proteins implicated in the Rac1/Cdc42/PAK and Rho pathways, like the upstream regulators of Rac/Cdc42 such as α PIX/ARHGEF6 and srGAP3/MEGAP (Slit Robo Rho GTPase-activating protein 3), or downstream effectors such as PAK3 and LIMK and the RhoGAP protein oligophrenin-1. In this review we have

already illustrated the function of most of these genes in regulating actin cytoskeleton in spines (see above).

A translocation breakpoint on chromosome 3p25, which causes the functional inactivation of srGAP3, has been described in a patient with severe intellectual disabilities (114).

SrGAP3 inhibits Rac by forming a complex with the scaffold protein WAVE1 (469). Knockout mice for SrGAP3 show various neuroanatomical changes, including enlarged dendritic spines, but also schizophrenia-related behavioral phenotype, suggesting that mutations of SrGAP3 may contribute to various neurodevelopmental diseases (539).

Implication of the Ras/Rap pathways in ID is currently suggested by the identification of mutations in the SynGAP1 gene and RAPGEF4, encoding the synaptically localized RapGEF Epac2 (25, 162). Interestingly, in the SynGAP1 mouse model, dendritic spine synapses develop prematurely during the early postnatal period, and this early spine maturation enhances excitability in the developing hippocampus, associated with classical behavioral abnormalities. SynGAP1 function is critical during development because inducing SynGAP1 mutations past the critical developmental period does not alter spine morphology (86). Epac2 missense mutations (mimicking human mutations) in cultured neurons increase dendritic spine number (Epac2-T809S) and volume (Epac2-V646F) (550), but in two independent *in vivo* studies, Epac2-deficient mice showed reduced dendritic spine motility and density (or normal density) in cortical neurons (477, 565).

The *fmr1* gene encodes the fragile X mental retardation protein (FMRP). The absence of FMRP, caused by the abnormal expansion of the CGG sequence in the gene, causes the fragile X (FX) syndrome, a major cause of congenital ID which is more frequent in males than in females. Several studies have shown that FX patients have neurons with an increased number of longer and thinner dendritic spines (27). *Fmr1* knockout mice modeling FX syndrome have recapitulated some of the immature spine morphologies observed in the brains of FX humans (204).

FMRP is an RNA-binding protein that regulates, in an activity-dependent manner, both the transport and translation of a subset of neuronal mRNAs to synapses (15, 16, 27, 32, 97, 124). The deregulated translation of mRNAs associated with FMRP might explain why in FX syndrome there are alterations in dendritic spine and synapse morphology and functions (32). Indeed, the translation of several synaptic proteins, such as PSD-95, SAPAP1–3, Shank1, Shank3, IRSp53, MAP1B, mGluR5, GluN1, GluN2B, and GluA1, is altered in synaptosomes of *fmr1* KO mice (39, 109, 199, 266, 281, 290, 334, 350, 577, 578). Additionally, some data link FMRP function to GTPase signaling. In *Drosophila*, the homologous FMRP regulates dendritic development by controlling actin cytoskeleton via the translational suppression of both Rac1 and profilin (411).

Mice knockout for *Fmr1* have impairment in LTP that can be rescued by BDNF treatment or by the stimulation of a Ras/PI3K pathway, indicating a potential role of the Ras signaling in this process and alterations in the connection between GTPases and actin polymerization in absence of FMRP (198).

The Phelan-McDermid syndrome (PMS; also named 22q13.3 deletion syndrome) is characterized by severe intellectual disability and autistic-like behavior (51, 305, 390, 548). The small deletion of chromosome 22 involves almost always a region that encodes the PSD scaffold protein Shank3. Deletion of one of the two Shank3 genes is assumed to cause the neurological pathology of PMS. De novo mutations in SHANK3 (107, 140, 332) and also in SHANK2 have been identified in individuals with ID and autism (36). The importance of Shank3 and Shank2 haploinsufficiency has been demonstrated in recent studies using knockout mice (56, 389, 530, 541). Cultured neurons knocked down for Shank3 and heterozygous or homozygous knockout mice for Shank3 have alterations in synaptic function and dendritic spine morphology (108, 530, 541). Thus these studies demonstrate that Shank3 is a major regulator of both synaptic morphology and function.

Mutations in tuberous sclerosis proteins 1 and 2 (TSC1, TSC2) or the phosphatase and tensin homolog (PTEN) genes are frequently diagnosed with autism, and each of these proteins regulates synaptic structure (63, 130, 503). Indeed, PTEN deficiency shows dendritic hypertrophy and elevated spine density while TSC1 or TSC2 ablation induces an enlargement of dendritic spines (263, 503).

The Angelman syndrome gene UBE3A, located at the maternal duplications of chromosome 15q11-q13, encodes an E3 ubiquitin ligase whose maternal deficiency reduces dendritic spine density and length in cerebellar, hippocampal, and cortical pyramidal neurons (102, 567). The multiple and not completely clear function of UBE3A suggest a potential link between autism and altered synaptic structure (297).

Rett syndrome is caused, in the majority of the patients, by mutations on the MECP2 gene, which encodes methyl-CpG-binding protein-2 (MeCP2), a protein that regulates gene transcription by binding to methylated DNA. Straight confirmation of MeCP2 contribution in the control of synapses is suggested by postmortem analysis of brains of female patients that showed lower spine density in the hippocampal CA1 pyramidal neurons (70, 216). Similarly, in mice deficient in MeCP2, spine dynamics is severely impaired early during development (269). CDKL5 is another gene that can be mutated in Rett syndrome. Interestingly, CDKL5 positively regulates dendritic spine morphology and synapse activity by phosphorylating NGL-1 and stabilizing its association in synapse with PSD95 (414).

Finally, it is important to note that when pharmacological and genetic manipulations successfully rescue both functional and behavioral phenotypes in many mouse model of intellectual disability and autism, there was always a parallel recovery of spine alterations, indicating the importance of these microstructural perturbations in the pathogenesis of these neurodevelopment diseases (269, 393, 413, 513).

X. PERSPECTIVES

This review attempts to outline and organize the bewildering number of molecules associated with the construction, maintenance, plasticity, and elimination of dendritic spines. The striking heterogeneity of spine morphologies on different types of neurons, as well as among adjacent spines in the same dendritic branch, remained puzzling: could it be that different molecular pathways act in different neurons/spines? Why is it that there are so many parallel molecular pathways for the formation/maturation of spines? Recent studies indicate that the functions of dendritic spines may be regulated by additional receptors residing on the spine head and/or neck. Among them is the GABAergic innervation that targets not only dendritic shafts but also selective spines and can suppress the calcium elevation in spines in response to synaptic stimulation (80). Another recent modulator of synaptic function is the nicotinic cholinergic receptor, found on dendritic spines of cortical neurons, which may enhance the activation of the NMDA receptor, and facilitate excitatory sensory transmission (564). Finally, some dendritic spines express receptors for corticotropin releasing factor, which may also modulate their ability to respond to afferent stimulation (74). These studies indicate that the functions of individual dendritic spines may be modulated by a host of factors, which challenges even more the understanding the role of dendritic spines in neuronal plasticity.

While the apparently simple dendritic spine appears to be more complex than ever realized, the tools for its investigation are improving as well. In fact, the major breakthroughs in the study of dendritic spines paralleled the development of imaging and molecular tools. Such was the case with the introduction of the high sensitivity cooled CCD, the confocal microscope which allowed three-dimensional reconstruction and time-lapse imaging of individual live spines and the two-photon microscope which allowed the investigation of dendritic spines in the cortex in vivo. The new generation of super-resolution microscopy, which breaks the limits of optical spatial resolution by an order of magnitude, include the sptPALM (304), the uPAINT (144), the STED (547), the STORM (182), and the DH-QPM (308). These tools are beginning to be introduced in the study of the distribution and dynamics of receptor molecules in dendritic spines. Initial observations (345) describe clusters of glutamate receptors in relations to PSD-95. Further experiments promise to revolutionize the understanding of the interrelations among molecular families in the spine and their relevance to plasticity-induced changes in spine morphology and function.

Several other imaging and molecular tools are being explored in the study of spine plasticity. Among them is the introduction of highly sensitive genetic calcium sensors (GCaMP6; Ref. 73) that can be used for imaging activity over extended periods of time, without the need to introduce dyes into the imaged cells. Selective activation of individual spines can be achieved by optogenetic activation of presynaptic fibers terminating on individual dendritic spines (374).

In conclusion, the combined use of high-resolution live microscopy, with the ability to stimulate and monitor individual molecular changes in clusters of spines in vivo over long periods of time, and the introduction of System Biology ability to monitor the simultaneous expression of hundreds of genes, will undoubtedly contribute to the understanding of the universal rules that govern the heterogeneous construction and function of central dendritic spines. We are almost there.

ACKNOWLEDGMENTS

C. Sala and M. Segal are grateful to members of their laboratories for critical discussions.

Address for reprint requests and other correspondence: M. Segal, Dept. of Neurobiology, The Weizmann Institute, Rehovot 76100, Israel (e-mail: Menahem.segal@weizmann.ac.il).

GRANTS

This work was supported by Israel Science Foundation Grant #805/09 (to M. Segal) as well as Comitato Telethon Fondazione Onlus Grant GGP11095 and Ministry of Health in the frame of ERA-NET NEURON (to C. Sala).

DISCLOSURES

No conflicts of interest, financial or otherwise, are declared by the authors.

REFERENCES

1. Abe K, Chisaka O, Van Roy F, Takeichi M. Stability of dendritic spines and synaptic contacts is controlled by alpha N-catenin. *Nat Neurosci* 7: 357–363, 2004.
2. Abidi FE, Holinski-Feder E, Rittinger O, Kooy F, Lubs HA, Stevenson RE, Schwartz CE. A novel 2 bp deletion in the TM4SF2 gene is associated with MRX58. *J Med Genet* 39: 430–433, 2002.
3. Acker CD, Yan P, Loew LM. Single-voxel recording of voltage transients in dendritic spines. *Biophys J* 101: L11–13, 2011.
4. Ackermann M, Matus A. Activity-induced targeting of profilin and stabilization of dendritic spine morphology. *Nat Neurosci* 6: 1194–1200, 2003.
5. Adasme T, Haeger P, Paula-Lima AC, Espinoza I, Casas-Alarcón MM, Carrasco MA, Hidalgo C. Involvement of ryanodine receptors in neurotrophin-induced hippocampal synaptic plasticity and spatial memory formation. *Proc Natl Acad Sci USA* 108: 3029–3034, 2011.
6. Akbik FV, Bhagat SM, Patel PR, Cafferty WB, Strittmatter SM. Anatomical plasticity of adult brain is titrated by Nogo Receptor 1. *Neuron* 77: 859–866, 2013.

7. Allen D, Bond CT, Luján R, Ballesteros-Merino C, Lin MT, Wang K, Klett N, Watanabe M, Shigemoto R, Stackman RW, Maylie J, Adelman JP. The SK2-long isoform directs synaptic localization and function of SK2-containing channels. *Nat Neurosci* 14: 744–749, 2011.
8. Allen KM, Gleeson JG, Bagrodia S, Partington MW, MacMillan JC, Cerione RA, Mulley JC, Walsh CA. PAK3 mutation in nonsyndromic X-linked mental retardation. *Nat Genet* 20: 25–30, 1998.
9. Alonso M, Medina JH, Pozzo-Miller L. ERK1/2 activation is necessary for BDNF to increase dendritic spine density in hippocampal CA1 pyramidal neurons. *Learn Mem* 11: 172–178, 2004.
10. Alvarez VA, Ridenour DA, Sabatini BL. Distinct structural and ionotropic roles of NMDA receptors in controlling spine and synapse stability. *J Neurosci* 27: 7365–7376, 2007.
11. Amaral MD, Pozzo-Miller L. TRPC3 channels are necessary for brain-derived neurotrophic factor to activate a nonselective cationic current and to induce dendritic spine formation. *J Neurosci* 27: 5179–5189, 2007.
12. An JJ, Gharami K, Liao GY, Woo NH, Lau AG, Vanevski F, Torre ER, Jones KR, Feng Y, Lu B, Xu B. Distinct role of long 3' UTR BDNF mRNA in spine morphology and synaptic plasticity in hippocampal neurons. *Cell* 134: 175–187, 2008.
13. Andrásfalvy BK, Makara JK, Johnston D, Magee JC. Altered synaptic and non-synaptic properties of CA1 pyramidal neurons in Kv4.2 knockout mice. *J Physiol* 586: 3881–3892, 2008.
14. Ango F, Robbe D, Tu JC, Xiao B, Worley PF, Pin JP, Bockaert J, Fagni L. Homer-dependent cell surface expression of metabotropic glutamate receptor type 5 in neurons. *Mol Cell Neurosci* 20: 323–329, 2002.
15. Antar LN, Afroz R, Dictenberg JB, Carroll RC, Bassell GJ. Metabotropic glutamate receptor activation regulates fragile X mental retardation protein and FMR1 mRNA localization differentially in dendrites and at synapses. *J Neurosci* 24: 2648–2655, 2004.
16. Antar LN, Li C, Zhang H, Carroll RC, Bassell GJ. Local functions for FMRP in axon growth cone motility and activity-dependent regulation of filopodia and spine synapses. *Mol Cell Neurosci* 32: 37–48, 2006.
17. Aoki C, Kojima N, Sabaliauskas N, Shah L, Ahmed TH, Oakford J, Ahmed T, Yamazaki H, Hanamura K, Shirao T. Drebrin a knockout eliminates the rapid form of homeostatic synaptic plasticity at excitatory synapses of intact adult cerebral cortex. *J Comp Neurol* 517: 105–121, 2009.
18. Araya R, Jiang J, Eiselthal KB, Yuste R. The spine neck filters membrane potentials. *Proc Natl Acad Sci USA* 103: 17961–17966, 2006.
19. Araya R, Nikolenko V, Eiselthal KB, Yuste R. Sodium channels amplify spine potentials. *Proc Natl Acad Sci USA* 104: 12347–12352, 2007.
20. Arendt T. Synaptic degeneration in Alzheimer's disease. *Acta Neuropathol* 118: 167–179, 2009.
21. Arstikaitis P, Gauthier-Campbell C, Huang K, El-Husseini A, Murphy TH. Proteins that promote filopodia stability, but not number, lead to more axonal-dendritic contacts. *PLoS One* 6: e16998, 2011.
22. Ashby MC, Maier SR, Nishimune A, Henley JM. Lateral diffusion drives constitutive exchange of AMPA receptors at dendritic spines and is regulated by spine morphology. *J Neurosci* 26: 7046–7055, 2006.
23. Ashe KH, Zahs KR. Probing the biology of Alzheimer's disease in mice. *Neuron* 66: 631–645, 2010.
24. Ayhan Y, Abazyan B, Nomura J, Kim R, Ladenheim B, Krasnova IN, Sawa A, Margolis RL, Cadet JL, Mori S, Vogel MW, Ross CA, Pletnikov MV. Differential effects of prenatal and postnatal expressions of mutant human DISC1 on neurobehavioral phenotypes in transgenic mice: evidence for neurodevelopmental origin of major psychiatric disorders. *Mol Psychiatry* 16: 293–306, 2011.
25. Bacchelli E, Blasi F, Biondolillo M, Lamb JA, Bonora E, Barnby G, Parr J, Beyer KS, Klauck SM, Poustka A, Bailey AJ, Monaco AP, Maestrini E, (IMGSAC) IMGSAC. Screening of nine candidate genes for autism on chromosome 2q reveals rare non-synonymous variants in the cAMP-GEFII gene. *Mol Psychiatry* 8: 916–924, 2003.
26. Bae J, Sung BH, Cho IH, Kim SM, Song WK. NESH regulates dendritic spine morphology and synapse formation. *PLoS One* 7: e34677, 2012.
27. Bagni C, Greenough WT. From mRNP trafficking to spine dysmorphogenesis: the roots of fragile X syndrome. *Nat Rev Neurosci* 6: 376–387, 2005.
28. Balamotis MA, Tamberg N, Woo YJ, Li J, Davy B, Kohwi-Shigematsu T, Kohwi Y. Satb1 ablation alters temporal expression of immediate early genes and reduces dendritic spine density during postnatal brain development. *Mol Cell Biol* 32: 333–347, 2012.
29. Ballesteros-Yáñez I, Benavides-Piccione R, Bourgeois JP, Changeux JP, DeFelipe J. Alterations of cortical pyramidal neurons in mice lacking high-affinity nicotinic receptors. *Proc Natl Acad Sci USA* 107: 11567–11572, 2010.
30. Barros CS, Calabrese B, Chamero P, Roberts AJ, Korzus E, Lloyd K, Stowers L, Mayford M, Halpain S, Müller U. Impaired maturation of dendritic spines without disorganization of cortical cell layers in mice lacking NRG1/ErbB signaling in the central nervous system. *Proc Natl Acad Sci USA* 106: 4507–4512, 2009.
31. Bassani S, Cingolani LA, Valnegri P, Folci A, Zapata J, Gianfelice A, Sala C, Goda Y, Passafaro M. The X-linked intellectual disability protein TSPAN7 regulates excitatory synapse development and AMPAR trafficking. *Neuron* 73: 1143–1158, 2012.
32. Bassell GJ, Warren ST. Fragile X syndrome: loss of local mRNA regulation alters synaptic development and function. *Neuron* 60: 201–214, 2008.
33. Beaudoin GM, Schofield CM, Nuwal T, Zang K, Ullian EM, Huang B, Reichardt LF. Afadin, a Ras/Rap effector that controls cadherin function, promotes spine and excitatory synapse density in the hippocampus. *J Neurosci* 32: 99–110, 2012.
34. Becker N, Wierenga CJ, Fonseca R, Bonhoeffer T, Nägerl UV. LTD induction causes morphological changes of presynaptic boutons and reduces their contacts with spines. *Neuron* 60: 590–597, 2008.
35. Benavides-Piccione R, Feraud-Espinosa I, Robles V, Yuste R, Defelipe J. Age-based comparison of human dendritic spine structure using complete three-dimensional reconstructions. *Cereb Cortex* 23: 1798–1810, 2013.
36. Berkel S, Marshall CR, Weiss B, Howe J, Roeth R, Moog U, Endris V, Roberts W, Szatmari P, Pinto D, Bonin M, Riess A, Engels H, Sprengel R, Scherer SW, Rappold GA. Mutations in the SHANK2 synaptic scaffolding gene in autism spectrum disorder and mental retardation. *Nat Genet* 42: 489–491, 2010.
37. Bertram L, Lill CM, Tanzi RE. The genetics of Alzheimer disease: back to the future. *Neuron* 68: 270–281, 2010.
38. Bertram L, Tanzi RE. Thirty years of Alzheimer's disease genetics: the implications of systematic meta-analyses. *Nat Rev Neurosci* 9: 768–778, 2008.
39. Bhakar AL, Dölen G, Bear MF. The pathophysiology of Fragile X (and what it teaches us about synapses). *Annu Rev Neurosci* 35: 417–443, 2012.
40. Bienvenu T, des Portes V, McDonnell N, Carrié A, Zemni R, Couvert P, Ropers HH, Moraine C, van Bokhoven H, Fryns JP, Allen K, Walsh CA, Boué J, Kahn A, Chelly J, Beldjord C. Missense mutation in PAK3, R67C, causes X-linked nonspecific mental retardation. *Am J Med Genet* 93: 294–298, 2000.
41. Billuart P, Bienvenu T, Ronce N, des Portes V, Vinet MC, Zemni R, Roest Croliis H, Carrié A, Fauchereau F, Cherry M, Briault S, Hamel B, Fryns JP, Beldjord C, Kahn A, Moraine C, Chelly J. Oligophrenin-1 encodes a rhoGAP protein involved in X-linked mental retardation. *Nature* 392: 923–926, 1998.
42. Biou V, Brinkhaus H, Malenka RC, Matus A. Interactions between drebrin and Ras regulate dendritic spine plasticity. *Eur J Neurosci* 27: 2847–2859, 2008.
43. Bloodgood BL, Sabatini BL. Neuronal activity regulates diffusion across the neck of dendritic spines. *Science* 310: 866–869, 2005.
44. Bloodgood BL, Sabatini BL. Nonlinear regulation of unitary synaptic signals by CaV(2.3) voltage-sensitive calcium channels located in dendritic spines. *Neuron* 53: 249–260, 2007.
45. Blundell J, Blais CA, Etherton MR, Espinosa F, Tabuchi K, Walz C, Bolliger MF, Südhof TC, Powell CM. Neuroligin-1 deletion results in impaired spatial memory and increased repetitive behavior. *J Neurosci* 30: 2115–2129, 2010.
46. Boda B, Alberi S, Nikonenko I, Node-Langlois R, Jourdain P, Moosmayer M, Paris-Jourdain L, Müller D. The mental retardation protein PAK3 contributes to synapse formation and plasticity in hippocampus. *J Neurosci* 24: 10816–10825, 2004.
47. Boda B, Dubos A, Müller D. Signaling mechanisms regulating synapse formation and function in mental retardation. *Curr Opin Neurobiol* 20: 519–527, 2010.

48. Boda B, Jourdain L, Muller D. Distinct, but compensatory roles of PAK1 and PAK3 in spine morphogenesis. *Hippocampus* 18: 857–861, 2008.
50. Boda B, Mas C, Muller D. Activity-dependent regulation of genes implicated in X-linked non-specific mental retardation. *Neuroscience* 114: 13–17, 2002.
51. Bonaglia MC, Giorda R, Borgatti R, Felisari G, Gagliardi C, Selicorni A, Zuffardi O. Disruption of the ProSAP2 gene in a t(12;22)(q24.1;q133) is associated with the 22q133 deletion syndrome. *Am J Hum Genet* 69: 261–268, 2001.
52. Bongmba OY, Martinez LA, Elhardt ME, Butler K, Tejada-Simon MV. Modulation of dendritic spines and synaptic function by Rac1: a possible link to Fragile X syndrome pathology. *Brain Res* 1399: 79–95, 2011.
53. Bosch M, Hayashi Y. Structural plasticity of dendritic spines. *Curr Opin Neurobiol* 22: 383–388, 2012.
54. Boulware MI, Kordasiewicz H, Mermelstein PG. Caveolin proteins are essential for distinct effects of membrane estrogen receptors in neurons. *J Neurosci* 27: 9941–9950, 2007.
55. Bourgin C, Murai KK, Richter M, Pasquale EB. The EphA4 receptor regulates dendritic spine remodeling by affecting beta1-integrin signaling pathways. *J Cell Biol* 178: 1295–1307, 2007.
56. Bozdagi O, Sakurai T, Papapetrou D, Wang X, Dickstein DL, Takahashi N, Kajiwara Y, Yang M, Katz AM, Scattoni ML, Harris MJ, Saxena R, Silverman JL, Crawley JN, Zhou Q, Hof PR, Buxbaum JD. Haploinsufficiency of the autism-associated Shank3 gene leads to deficits in synaptic function, social interaction, and social communication. *Mol Autism* 1: 15, 2010.
57. Brandon NJ, Sawa A. Linking neurodevelopmental and synaptic theories of mental illness through DISC1. *Nat Rev Neurosci* 12: 707–722, 2011.
58. Bresler T, Ramati Y, Zamorano PL, Zhai R, Garner CC, Ziv NE. The dynamics of SAP90/PSD-95 recruitment to new synaptic junctions. *Mol Cell Neurosci* 18: 149–167, 2001.
59. Brigman JL, Wright T, Talani G, Prasad-Mulcare S, Jinde S, Seabold GK, Mathur P, Davis MI, Bock R, Gustin RM, Colbran RJ, Alvarez VA, Nakazawa K, Delpire E, Lovinger DM, Holmes A. Loss of GluN2B-containing NMDA receptors in CA1 hippocampus and cortex impairs long-term depression, reduces dendritic spine density, and disrupts learning. *J Neurosci* 30: 4590–4600, 2010.
60. Bryan B, Kumar V, Stafford LJ, Cai Y, Wu G, Liu M. GEFT, a Rho family guanine nucleotide exchange factor, regulates neurite outgrowth and dendritic spine formation. *J Biol Chem* 279: 45824–45832, 2004.
61. Brüning I, Kaech S, Brinkhaus H, Oertner TG, Matus A. Influx of extracellular calcium regulates actin-dependent morphological plasticity in dendritic spines. *Neuropharmacology* 47: 669–676, 2004.
62. Burette AC, Strehler EE, Weinberg RJ. A plasma membrane Ca^{2+} ATPase isoform at the postsynaptic density. *Neuroscience* 169: 987–993, 2010.
63. Butler MG, Dasouki MJ, Zhou XP, Talebizadeh Z, Brown M, Takahashi TN, Miles JH, Wang CH, Stratton R, Pilarski R, Eng C. Subset of individuals with autism spectrum disorders and extreme macrocephaly associated with germline PTEN tumour suppressor gene mutations. *J Med Genet* 42: 318–321, 2005.
64. Cahill ME, Xie Z, Day M, Photowala H, Barbolina MV, Miller CA, Weiss C, Radulovic J, Sweatt JD, Disterhoft JF, Surmeier DJ, Penzes P. Kalirin regulates cortical spine morphogenesis and disease-related behavioral phenotypes. *Proc Natl Acad Sci USA* 106: 13058–13063, 2009.
65. Carlisle HJ, Manzer P, Marcora E, Kennedy MB. SynGAP regulates steady-state and activity-dependent phosphorylation of cofilin. *J Neurosci* 28: 13673–13683, 2008.
66. Carmona MA, Murai KK, Wang L, Roberts AJ, Pasquale EB. Glial ephrin-A3 regulates hippocampal dendritic spine morphology and glutamate transport. *Proc Natl Acad Sci USA* 106: 12524–12529, 2009.
67. Carrie A, Jun L, Bienvenu T, Vinet MC, McDonell N, Couvert P, Zemni R, Cardona A, Van Buggenhout G, Fris S, Hamel B, Moraine C, Ropers HH, Strom T, Howell GR, Whittaker A, Ross MT, Kahn A, Fryns JP, Beldjord C, Marynen P, Chelly J. A new member of the IL-1 receptor family highly expressed in hippocampus and involved in X-linked mental retardation. *Nat Genet* 23: 25–31, 1999.
68. Chakravarthy S, Saiepour MH, Bence M, Perry S, Hartman R, Couey JJ, Mansvelder HD, Levett CN. Postsynaptic TrkB signaling has distinct roles in spine maintenance in adult visual cortex and hippocampus. *Proc Natl Acad Sci USA* 103: 1071–1076, 2006.
69. Chao HW, Hong CJ, Huang TN, Lin YL, Hsueh YP. SUMOylation of the MAGUK protein CASK regulates dendritic spinogenesis. *J Cell Biol* 182: 141–155, 2008.
70. Chapple CA, Calfa GD, Lane MC, Albertson AJ, Larimore JL, Kudo S, Armstrong DL, Percy AK, Pozzo-Miller L. Dendritic spine pathologies in hippocampal pyramidal neurons from Rett syndrome brain and after expression of Rett-associated MECP2 mutations. *Neurobiol Dis* 35: 219–233, 2009.
71. Charrier C, Joshi K, Coutinho-Budd J, Kim JE, Lambert N, de Marchena J, Jin WL, Vanderhaeghen P, Ghosh A, Sassa T, Polleux F. Inhibition of SRGAP2 function by its human-specific paralogs induces neoteny during spine maturation. *Cell* 149: 923–935, 2012.
72. Chen BS, Thomas EV, Sanz-Clemente A, Roche KW. NMDA receptor-dependent regulation of dendritic spine morphology by SAPI02 splice variants. *J Neurosci* 31: 89–96, 2011.
73. Chen TW, Wardill TJ, Sun Y, Pulver SR, Renninger SL, Baohan A, Schreier ER, Kerr RA, Orger MB, Jayaraman V, Looger LL, Svoboda K, Kim DS. Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature* 499: 295–300, 2013.
74. Chen Y, Andres AL, Frotscher M, Baram TZ. Tuning synaptic transmission in the hippocampus by stress: the CRH system. *Front Cell Neurosci* 6: 13, 2012.
75. Chen YK, Chen CY, Hu HT, Hsueh YP. CTTNBP2, but not CTTNBP2NL, regulates dendritic spinogenesis and synaptic distribution of the striatin-PP2A complex. *Mol Biol Cell* 23: 4383–4392, 2012.
76. Chen YK, Hsueh YP. Cortactin-binding protein 2 modulates the mobility of cortactin and regulates dendritic spine formation and maintenance. *J Neurosci* 32: 1043–1055, 2012.
77. Cheng A, Wan R, Yang JL, Kamimura N, Son TG, Ouyang X, Luo Y, Okun E, Mattson MP. Involvement of PGC-1 α in the formation and maintenance of neuronal dendritic spines. *Nat Commun* 3: 1250, 2012.
78. Chih B, Afridi SK, Clark L, Scheiffele P. Disorder-associated mutations lead to functional inactivation of neuroligins. *Hum Mol Genet* 13: 1471–1477, 2004.
79. Chih B, Engelman H, Scheiffele P. Control of excitatory and inhibitory synapse formation by neuroligins. *Science* 307: 1324–1328, 2005.
80. Chiu CQ, Lur G, Morse TM, Carnevale NT, Ellis-Davies GC, Higley MJ. Compartmentalization of GABAergic inhibition by dendritic spines. *Science* 340: 759–762, 2013.
81. Cho IH, Kim DH, Lee MJ, Bae J, Lee KH, Song WK. SPIN90 phosphorylation modulates spine structure and synaptic function. *PLoS One* 8: e54276, 2013.
82. Choi J, Ko J, Raczy B, Burette A, Lee JR, Kim S, Na M, Lee HW, Kim K, Weinberg RJ, Kim E. Regulation of dendritic spine morphogenesis by insulin receptor substrate 53, a downstream effector of Rac1 and Cdc42 small GTPases. *J Neurosci* 25: 869–879, 2005.
83. Choi S, Ko J, Lee JR, Lee HW, Kim K, Chung HS, Kim H, Kim E. ARF6 and EFA6A regulate the development and maintenance of dendritic spines. *J Neurosci* 26: 4811–4819, 2006.
84. Ciani L, Boyle KA, Dickins E, Sahores M, Anane D, Lopes DM, Gibb AJ, Salinas PC. Wnt7a signaling promotes dendritic spine growth and synaptic strength through Ca^{2+} /Calmodulin-dependent protein kinase II. *Proc Natl Acad Sci USA* 108: 10732–10737, 2011.
85. Clemens LE, Heldmaier G, Exner C. Keep cool: memory is retained during hibernation in Alpine marmots. *Physiol Behav* 98: 78–84, 2009.
86. Clement JP, Aceti M, Creson TK, Ozkan ED, Shi Y, Reish NJ, Almonte AG, Miller BH, Wiltgen BJ, Miller CA, Xu X, Rumbaugh G. Pathogenic SYNGAP1 mutations impair cognitive development by disrupting maturation of dendritic spine synapses. *Cell* 151: 709–723, 2012.
87. Comoletti D, De Jacobo A, Jennings LL, Flynn RE, Gaietta G, Tsigelny I, Ellisman MH, Taylor P. The Arg451Cys-neuroligin-3 mutation associated with autism reveals a defect in protein processing. *J Neurosci* 24: 4889–4893, 2004.

88. Coss RG, Perkel DH. The function of dendritic spines: a review of theoretical issues. *Behav Neural Biol* 44: 151–185, 1985.
89. Coulter DA, Huguenard JR, Prince DA. Calcium currents in rat thalamocortical relay neurones: kinetic properties of the transient, low-threshold current. *J Physiol* 414: 587–604, 1989.
90. Counts SE, He B, Nadeem M, Wu J, Scheff SW, Mufson EJ. Hippocampal drebrin loss in mild cognitive impairment. *Neurodegener Dis* 10: 216–219, 2012.
91. Cubelos B, Sebastián-Serrano A, Beccari L, Calcagnotto ME, Cisneros E, Kim S, Dopazo A, Alvarez-Dolado M, Redondo JM, Bovolenta P, Walsh CA, Nieto M. Cux I and Cux2 regulate dendritic branching, spine morphology, and synapses of the upper layer neurons of the cortex. *Neuron* 66: 523–535, 2010.
92. Cuesto G, Enriquez-Barreto L, Caramés C, Cantarero M, Gasull X, Sandi C, Ferrús A, Acebes Á, Morales M. Phosphoinositide-3-kinase activation controls synaptogenesis and spinogenesis in hippocampal neurons. *J Neurosci* 31: 2721–2733, 2011.
93. Dalla C, Whetstone AS, Hodes GE, Shors TJ. Stressful experience has opposite effects on dendritic spines in the hippocampus of cycling versus masculinized females. *Neurosci Lett* 449: 52–56, 2009.
94. Dalva MB, McClelland AC, Kayser MS. Cell adhesion molecules: signalling functions at the synapse. *Nat Rev Neurosci* 8: 206–220, 2007.
95. Dalva MB, Takasu MA, Lin MZ, Shamah SM, Hu L, Gale NW, Greenberg ME. EphB receptors interact with NMDA receptors and regulate excitatory synapse formation. *Cell* 103: 945–956, 2000.
96. Danglot L, Freret T, Le Roux N, Narboux Nême N, Burgo A, Hyenne V, Roumier A, Contremoulins V, Dauphin F, Bizot JC, Vodjdani G, Gaspar P, Boulouard M, Ponce JC, Galli T, Simmler MC. Vezatin is essential for dendritic spine morphogenesis and functional synaptic maturation. *J Neurosci* 32: 9007–9022, 2012.
97. De Rubeis S, Bagni C. Fragile X mental retardation protein control of neuronal mRNA metabolism: Insights into mRNA stability. *Mol Cell Neurosci* 43: 43–50, 2010.
- 97a. De Vivo L, Landi S, Panniello M, Baroncelli L, Chierzi S, Mariotti L, Spolidoro M, Pizzorusso T, Maffei L, Ratto GM. Extracellular matrix inhibits structural and functional plasticity of dendritic spines in the adult visual cortex. *Nat Commun* 4: 1484, 2013.
98. Dean C, Dresbach T. Neuroligins and neurexins: linking cell adhesion, synapse formation and cognitive function. *Trends Neurosci* 29: 21–29, 2006.
99. DeKosky ST, Scheff SW. Synapse loss in frontal cortex biopsies in Alzheimer's disease: correlation with cognitive severity. *Ann Neurol* 27: 457–464, 1990.
100. Deller T, Orth CB, Del Turco D, Vlachos A, Burbach GJ, Drakew A, Chabanis S, Korte M, Schwegler H, Haas CA, Frotscher M. A role for synaptopodin and the spine apparatus in hippocampal synaptic plasticity. *Ann Anat* 189: 5–16, 2007.
101. Desmond NL, Levy WB. Changes in the postsynaptic density with long-term potentiation in the dentate gyrus. *J Comp Neurol* 253: 476–482, 1986.
102. Dindot SV, Antalffy BA, Bhattacharjee MB, Beaudet AL. The Angelman syndrome ubiquitin ligase localizes to the synapse and nucleus, and maternal deficiency results in abnormal dendritic spine morphology. *Hum Mol Genet* 17: 111–118, 2008.
103. Dubos A, Combeau G, Bernardinelli Y, Barnier JV, Hartley O, Gaertner H, Boda B, Muller D. Alteration of synaptic network dynamics by the intellectual disability protein PAK3. *J Neurosci* 32: 519–527, 2012.
104. Dudanova I, Tabuchi K, Rohlmann A, Südhof TC, Missler M. Deletion of alpha-neurexins does not cause a major impairment of axonal pathfinding or synapse formation. *J Comp Neurol* 502: 261–274, 2007.
105. Dumanis SB, Tesoriero JA, Babus LW, Nguyen MT, Trotter JH, Ladu MJ, Weeber EJ, Turner RS, Xu B, Rebeck GW, Hoe HS. ApoE4 decreases spine density and dendritic complexity in cortical neurons in vivo. *J Neurosci* 29: 15317–15322, 2009.
106. Dunah AW, Hueske E, Wyszynski M, Hoogenraad CC, Jaworski J, Pak DT, Simonetta A, Liu G, Sheng M. LAR receptor protein tyrosine phosphatases in the development and maintenance of excitatory synapses. *Nat Neurosci* 8: 458–467, 2005.
107. Durand CM, Betancur C, Boeckers TM, Bockmann J, Chaste P, Fauchereau F, Nygren G, Rastam M, Gillberg IC, Anckarsater H, Sponheim E, Goubran-Botros H, Delorme R, Chabane N, Mouren-Simeoni MC, de Mas P, Bieth E, Roge B, Heron D, Burglen L, Gillberg C, Leboyer M, Bourgeron T. Mutations in the gene encoding the synaptic scaffolding protein SHANK3 are associated with autism spectrum disorders. *Nat Genet* 39: 25–27, 2007.
108. Durand CM, Perroy J, Loll F, Perraiss D, Fagni L, Bourgeron T, Montcouquiol M, Sans N. SHANK3 mutations identified in autism lead to modification of dendritic spine morphology via an actin-dependent mechanism. *Mol Psychiatry* 17: 71–84, 2012.
109. Edbauer D, Neilson JR, Foster KA, Wang CF, Seeburg DP, Batterton MN, Tada T, Dolan BM, Sharp PA, Sheng M. Regulation of synaptic structure and function by FMRP-associated microRNAs miR-125b and miR-132. *Neuron* 65: 373–384, 2010.
110. Ehrlich I, Malinow R. Postsynaptic density 95 controls AMPA receptor incorporation during long-term potentiation and experience-driven synaptic plasticity. *J Neurosci* 24: 916–927, 2004.
111. El-Husseini AE, Schnell E, Chetkovich DM, Nicoll RA, Brecht DS. PSD-95 involvement in maturation of excitatory synapses. *Science* 290: 1364–1368, 2000.
112. Emptage N, Bliss TV, Fine A. Single synaptic events evoke NMDA receptor-mediated release of calcium from internal stores in hippocampal dendritic spines. *Neuron* 22: 115–124, 1999.
113. Emptage NJ, Reid CA, Fine A, Bliss TV. Optical quantal analysis reveals a presynaptic component of LTP at hippocampal Schaffer-associational synapses. *Neuron* 38: 797–804, 2003.
114. Endris V, Wogatzky B, Leimer U, Bartsch D, Zatyka M, Latif F, Maher ER, Tariverdian G, Kirsch S, Karch D, Rappold GA. The novel Rho-GTPase activating gene MEGAP/srGAP3 has a putative role in severe mental retardation. *Proc Natl Acad Sci USA* 99: 11754–11759, 2002.
115. Engert F, Bonhoeffer T. Dendritic spine changes associated with hippocampal long-term synaptic plasticity. *Nature* 399: 66–70, 1999.
116. Enoki R, Hu YL, Hamilton D, Fine A. Expression of long-term plasticity at individual synapses in hippocampus is graded, bidirectional, and mainly presynaptic: optical quantal analysis. *Neuron* 62: 242–253, 2009.
117. Ethell I, Yamaguchi Y. Cell surface heparan sulfate proteoglycan syndecan-2 induces the maturation of dendritic spines in rat hippocampal neurons. *J Cell Biol* 144: 575–586, 1999.
118. Ethell IM, Irie F, Kalo MS, Couchman JR, Pasquale EB, Yamaguchi Y. EphB/syndecan-2 signaling in dendritic spine morphogenesis. *Neuron* 31: 1001–1013, 2001.
119. Fabian-Fine R, Skehel P, Errington ML, Davies HA, Sher E, Stewart MG, Fine A. Ultrastructural distribution of the alpha7 nicotinic acetylcholine receptor subunit in rat hippocampus. *J Neurosci* 21: 7993–8003, 2001.
120. Fan Y, Tang X, Vitriol E, Chen G, Zheng JQ. Actin capping protein is required for dendritic spine development and synapse formation. *J Neurosci* 31: 10228–10233, 2011.
121. Feldman DE. The spike-timing dependence of plasticity. *Neuron* 75: 556–571, 2012.
122. Feng J, Yan Z, Ferreira A, Tomizawa K, Liauw JA, Zhuo M, Allen PB, Ouimet CC, Greengard P. Spinophilin regulates the formation and function of dendritic spines. *Proc Natl Acad Sci USA* 97: 9287–9292, 2000.
123. Fernández de Sevilla D, Buño W. The muscarinic long-term enhancement of NMDA and AMPA receptor-mediated transmission at Schaffer collateral synapses develop through different intracellular mechanisms. *J Neurosci* 30: 11032–11042, 2010.
124. Ferrari F, Mercaldo V, Piccoli G, Sala C, Cannata S, Achsel T, Bagni C. The fragile X mental retardation protein-RNP granules show an mGluR-dependent localization in the post-synaptic spines. *Mol Cell Neurosci* 34: 343–354, 2007.
125. Fester L, Prange-Kiel J, Zhou L, Blittersdorf BV, Böhm J, Jarry H, Schumacher M, Rune GM. Estrogen-regulated synaptogenesis in the hippocampus: sexual dimorphism in vivo but not in vitro. *J Steroid Biochem Mol Biol* 131: 24–29, 2012.
126. Fiala JC, Feinberg M, Popov V, Harris KM. Synaptogenesis via dendritic filopodia in developing hippocampal area CA1. *J Neurosci* 18: 8900–8911, 1998.
127. Fiala JC, Spacek J, Harris KM. Dendritic spine pathology: cause or consequence of neurological disorders? *Brain Res* 39: 29–54, 2002.

128. Fifkova E, Van Harreveld A. Long-lasting morphological changes in dendritic spines of dentate granular cells following stimulation of the entorhinal area. *J Neurocytol* 6: 211–230, 1977.
129. Fortin DA, Srivastava T, Soderling TR. Structural modulation of dendritic spines during synaptic plasticity. *Neuroscientist* 18: 326–341, 2012.
130. Fraser MM, Bayazitov IT, Zakharenko SS, Baker SJ. Phosphatase and tensin homolog, deleted on chromosome 10 deficiency in brain causes defects in synaptic structure, transmission and plasticity, and myelination abnormalities. *Neuroscience* 151: 476–488, 2008.
131. Frick A, Magee J, Johnston D. LTP is accompanied by an enhanced local excitability of pyramidal neuron dendrites. *Nat Neurosci* 7: 126–135, 2004.
132. Frotscher M, Drakew A, Heimrich B. Role of afferent innervation and neuronal activity in dendritic development and spine maturation of fascia dentata granule cells. *Cereb Cortex* 10: 946–951, 2000.
133. Fu M, Yu X, Lu J, Zuo Y. Repetitive motor learning induces coordinated formation of clustered dendritic spines in vivo. *Nature* 483: 92–95, 2012.
134. Fujii R, Okabe S, Urushido T, Inoue K, Yoshimura A, Tachibana T, Nishikawa T, Hicks GG, Takumi T. The RNA binding protein TLS is translocated to dendritic spines by mGluR5 activation and regulates spine morphology. *Curr Biol* 15: 587–593, 2005.
135. Fujii R, Takumi T. TLS facilitates transport of mRNA encoding an actin-stabilizing protein to dendritic spines. *J Cell Sci* 118: 5755–5765, 2005.
136. Fukazawa Y, Saitoh Y, Ozawa F, Ohta Y, Mizuno K, Inokuchi K. Hippocampal LTP is accompanied by enhanced F-actin content within the dendritic spine that is essential for late LTP maintenance in vivo. *Neuron* 38: 447–460, 2003.
137. Gao C, Frausto SF, Guede AL, Tronson NC, Jovasevic V, Leaderbrand K, Corcoran KA, Guzmán YF, Swanson GT, Radulovic J. IQGAP1 regulates NR2A signaling, spine density, and cognitive processes. *J Neurosci* 31: 8533–8542, 2011.
138. Gartner U, Alpar A, Behrbohm J, Heumann R, Arendt T. Enhanced Ras activity promotes spine formation in synRas mice neocortex. *Neuroreport* 16: 149–152, 2005.
139. Gauthier J, Champagne N, Lafrenière RG, Xiong L, Spiegelman D, Brustein E, Lapointe M, Peng H, Côté M, Noreau A, Hamdan FF, Addington AM, Rapoport JL, Delisi LE, Krebs MO, Joobar R, Fathalli F, Mouaffak F, Haghighi AP, Néri C, Dubé MP, Samuels ME, Marineau C, Stone EA, Awadalla P, Barker PA, Carbonetto S, Drapeau P, Rouleau GA, Team SD. De novo mutations in the gene encoding the synaptic scaffolding protein SHANK3 in patients ascertained for schizophrenia. *Proc Natl Acad Sci USA* 107: 7863–7868, 2010.
140. Gauthier J, Spiegelman D, Piton A, Lafreniere RG, Laurent S, St-Onge J, Lapointe L, Hamdan FF, Cossette P, Mottron L, Fombonne E, Joobar R, Marineau C, Drapeau P, Rouleau GA. Novel de novo SHANK3 mutation in autistic patients. *Am J Med Genet B Neuropsychiatr Genet* 150B: 421–424, 2009.
141. Geinisman Y, Disterhoft JF, Gundersen HJ, McEchron MD, Persina IS, Power JM, van der Zee EA, West MJ. Remodeling of hippocampal synapses after hippocampus-dependent associative learning. *J Comp Neurol* 417: 49–59, 2000.
142. Gerrow K, Romorini S, Nabi SM, Colicos MA, Sala C, El-Husseini A. A preformed complex of postsynaptic proteins is involved in excitatory synapse development. *Neuron* 49: 547–562, 2006.
143. Ghirelli AE, Paradis S. The GTPase Rem2 regulates synapse development and dendritic morphology. *Dev Neurobiol* 71: 374–389, 2011.
144. Giannone G, Hosy E, Levat F, Constals A, Schulze K, Sobolevsky AI, Rosconi MP, Gouaux E, Tampé R, Choquet D, Cognet L. Dynamic superresolution imaging of endogenous proteins on living cells at ultra-high density. *Biophys J* 99: 1303–1310, 2010.
145. Giessel AJ, Sabatini BL. M1 muscarinic receptors boost synaptic potentials and calcium influx in dendritic spines by inhibiting postsynaptic SK channels. *Neuron* 68: 936–947, 2010.
146. Glantz LA, Lewis DA. Decreased dendritic spine density on prefrontal cortical pyramidal neurons in schizophrenia. *Arch Gen Psychiatry* 57: 65–73, 2000.
147. Globus A, Rosenzweig MR, Bennett EL, Diamond MC. Effects of differential experience on dendritic spine counts in rat cerebral cortex. *J Comp Physiol Psychol* 82: 175–181, 1973.
148. Goetze B, Tuebing F, Xie Y, Dorostkar MM, Thomas S, Pehl U, Boehm S, Macchi P, Kiebler MA. The brain-specific double-stranded RNA-binding protein Staufen2 is required for dendritic spine morphogenesis. *J Cell Biol* 172: 221–231, 2006.
149. Gogtay N. Cortical brain development in schizophrenia: insights from neuroimaging studies in childhood-onset schizophrenia. *Schizophr Bull* 34: 30–36, 2008.
150. Goldin M, Segal M, Avignone E. Functional plasticity triggers formation and pruning of dendritic spines in cultured hippocampal networks. *J Neurosci* 21: 186–193, 2001.
151. Goto J, Mikoshiba K. Inositol 1,4,5-trisphosphate receptor-mediated calcium release in Purkinje cells: from molecular mechanism to behavior. *Cerebellum* 10: 820–833, 2011.
152. Gould E, Woolley CS, Frankfurt M, McEwen BS. Gonadal steroids regulate dendritic spine density in hippocampal pyramidal cells in adulthood. *J Neurosci* 10: 1286–1291, 1990.
153. Govek EE, Newey SE, Akerman CJ, Cross JR, Van der Veken L, Van Aelst L. The X-linked mental retardation protein oligophrenin-1 is required for dendritic spine morphogenesis. *Nat Neurosci* 7: 364–372, 2004.
154. Graf ER, Zhang X, Jin SX, Linhoff MW, Craig AM. Neurexins induce differentiation of GABA and glutamate postsynaptic specializations via neuroligins. *Cell* 119: 1013–1026, 2004.
155. Grove M, Demyanenko G, Echarri A, Zipfel PA, Quiroz ME, Rodriguez RM, Playford M, Martensen SA, Robinson MR, Wetsel WC, Maness PF, Pendergast AM. ABI2-deficient mice exhibit defective cell migration, aberrant dendritic spine morphogenesis, and deficits in learning and memory. *Mol Cell Biol* 24: 10905–10922, 2004.
156. Grover LM, Teyler TJ. Two components of long-term potentiation induced by different patterns of afferent activation. *Nature* 347: 477–479, 1990.
157. Grueber WB, Jan LY, Jan YN. Different levels of the homeodomain protein cut regulate distinct dendrite branching patterns of *Drosophila* multidendritic neurons. *Cell* 112: 805–818, 2003.
158. Grunditz A, Holbro N, Tian L, Zuo Y, Oertner TG. Spine neck plasticity controls postsynaptic calcium signals through electrical compartmentalization. *J Neurosci* 28: 13457–13466, 2008.
159. Gu J, Firestein BL, Zheng JQ. Microtubules in dendritic spine development. *J Neurosci* 28: 12120–12124, 2008.
160. Guthrie PB, Segal M, Kater SB. Independent regulation of calcium revealed by imaging dendritic spines. *Nature* 354: 76–80, 1991.
161. Hale CF, Dietz KC, Varela JA, Wood CB, Zirlin BC, Leverich LS, Greene RW, Cowan CW. Essential role for vav Guanine nucleotide exchange factors in brain-derived neurotrophic factor-induced dendritic spine growth and synapse plasticity. *J Neurosci* 31: 12426–12436, 2011.
162. Hamdan FF, Gauthier J, Spiegelman D, Noreau A, Yang Y, Pellerin S, Dobrzyniecka S, Côté M, Perreault-Linck E, Perreault-Linck E, Carmant L, D'Anjou G, Fombonne E, Addington AM, Rapoport JL, Delisi LE, Krebs MO, Mouaffak F, Joobar R, Mottron L, Drapeau P, Marineau C, Lafrenière RG, Lacaille JC, Rouleau GA, Michaud JL, Group StD. Mutations in SYNGAP1 in autosomal nonsyndromic mental retardation. *N Engl J Med* 360: 599–605, 2009.
163. Hamilton AM, Oh WC, Vega-Ramirez H, Stein IS, Hell JW, Patrick GN, Zito K. Activity-dependent growth of new dendritic spines is regulated by the proteasome. *Neuron* 74: 1023–1030, 2012.
164. Han S, Nam J, Li Y, Kim S, Cho SH, Cho YS, Choi SY, Choi J, Han K, Kim Y, Na M, Kim H, Bae YC, Kim E. Regulation of dendritic spines, spatial memory, and embryonic development by the TANC family of PSD-95-interacting proteins. *J Neurosci* 30: 15102–15112, 2010.
165. Hanley JG. PICK1: a multi-talented modulator of AMPA receptor trafficking. *Pharmacol Ther* 118: 152–160, 2008.
166. Hansen KF, Sakamoto K, Wayman GA, Impey S, Obrietan K. Transgenic miR132 alters neuronal spine density and impairs novel object recognition memory. *PLoS One* 5: e15497, 2010.
167. Harnett MT, Makara JK, Spruston N, Kath WL, Magee JC. Synaptic amplification by dendritic spines enhances input cooperativity. *Nature* 491: 599–602, 2012.

168. Harris EW, Ganong AH, Cotman CW. Long-term potentiation in the hippocampus involves activation of *N*-methyl-D-aspartate receptors. *Brain Res* 323: 132–137, 1984.
169. Harris K. Structure, development, and plasticity of dendritic spines. *Curr Opin Neurobiol* 9: 343–348, 1999.
170. Harris KM. Calcium from internal stores modifies dendritic spine shape. *Proc Natl Acad Sci USA* 96: 12213–12215, 1999.
171. Harris KM, Stevens JK. Dendritic spines of CA I pyramidal cells in the rat hippocampus: serial electron microscopy with reference to their biophysical characteristics. *J Neurosci* 9: 2982–2997, 1989.
172. Harvey CD, Svoboda K. Locally dynamic synaptic learning rules in pyramidal neuron dendrites. *Nature* 450: 1195–1200, 2007.
173. Harvey CD, Yasuda R, Zhong H, Svoboda K. The spread of Ras activity triggered by activation of a single dendritic spine. *Science* 321: 136–140, 2008.
174. Hasbani MJ, Schlieff ML, Fisher DA, Goldberg MP. Dendritic spines lost during glutamate receptor activation reemerge at original sites of synaptic contact. *J Neurosci* 21: 2393–2403, 2001.
175. Hayashi K, Ohshima T, Hashimoto M, Mikoshiba K. Pak1 regulates dendritic branching and spine formation. *Dev Neurobiol* 67: 655–669, 2007.
176. Hayashi K, Shirao T. Change in the shape of dendritic spines caused by overexpression of drebrin in cultured cortical neurons. *J Neurosci* 19: 3918–3925, 1999.
177. Hayashi K, Suzuki A, Hirai S, Kurihara Y, Hoogenraad CC, Ohno S. Maintenance of dendritic spine morphology by partitioning-defective 1b through regulation of microtubule growth. *J Neurosci* 31: 12094–12103, 2011.
178. Hayashi M, Tang C, Verpelli C, Narayanan R, Stearns M, Xu RM, Li H, Sala C, Hayashi Y. The postsynaptic density proteins homer and shank form a polymeric network structure. *Neurosci Res* 68: E339, 2010.
179. Hayashi MK, Tang C, Verpelli C, Narayanan R, Stearns MH, Xu RM, Li H, Sala C, Hayashi Y. The postsynaptic density proteins Homer and Shank form a polymeric network structure. *Cell* 137: 159–171, 2009.
180. Hayashi ML, Choi SY, Rao BS, Jung HY, Lee HK, Zhang D, Chattarji S, Kirkwood A, Tonegawa S. Altered cortical synaptic morphology and impaired memory consolidation in forebrain-specific dominant-negative PAK transgenic mice. *Neuron* 42: 773–787, 2004.
181. Hayashi-Takagi A, Takaki M, Graziane N, Seshadri S, Murdoch H, Dunlop AJ, Makino Y, Seshadri AJ, Ishizuka K, Srivastava DP, Xie Z, Baraban JM, Houslay MD, Tomoda T, Brandon NJ, Kamiya A, Yan Z, Penzes P, Sawa A. Disrupted-in-Schizophrenia 1 (DISC1) regulates spines of the glutamate synapse via Rac1. *Nat Neurosci* 13: 327–332, 2010.
182. Heilemann M. Fluorescence microscopy beyond the diffraction limit. *J Biotechnol* 149: 243–251, 2010.
- 182a. Heine M, Groc L, Frischknecht R, Béique JC, Lounis B, Rumbaugh G, Huganir RL, Cognet L, Choquet D. Surface mobility of postsynaptic AMPARs tunes synaptic transmission. *Science* 320: 201–205, 2008.
183. Heinen K, Baker RE, Spijker S, Rosahl T, van Pelt J, Brussaard AB. Impaired dendritic spine maturation in GABA_A receptor alpha1 subunit knock out mice. *Neuroscience* 122: 699–705, 2003.
184. Henkemeyer M, Itkis OS, Ngo M, Hickmott PW, Ethell IM. Multiple EphB receptor tyrosine kinases shape dendritic spines in the hippocampus. *J Cell Biol* 163: 1313–1326, 2003.
185. Hering H, Sheng M. Activity-dependent redistribution and essential role of cortactin in dendritic spine morphogenesis. *J Neurosci* 23: 11759–11769, 2003.
186. Herron CE, Lester RA, Coan EJ, Collingridge GL. Frequency-dependent involvement of NMDA receptors in the hippocampus: a novel synaptic mechanism. *Nature* 322: 265–268, 1986.
187. Hill JJ, Hashimoto T, Lewis DA. Molecular mechanisms contributing to dendritic spine alterations in the prefrontal cortex of subjects with schizophrenia. *Mol Psychiatry* 11: 557–566, 2006.
188. Hill TC, Zito K. LTP-induced long-term stabilization of individual nascent dendritic spines. *J Neurosci* 33: 678–686, 2013.
189. Hodges JL, Newell-Litwa K, Asmussen H, Vicente-Manzanares M, Horwitz AR. Myosin IIb activity and phosphorylation status determines dendritic spine and post-synaptic density morphology. *PLoS One* 6: e24149, 2011.
190. Hoe HS, Lee JY, Pak DT. Combinatorial morphogenesis of dendritic spines and filopodia by SPAR and alpha-actinin2. *Biochem Biophys Res Commun* 384: 55–60, 2009.
191. Hojo Y, Murakami G, Mukai H, Higo S, Hatanaka Y, Ogiue-Ikeda M, Ishii H, Kimoto T, Kawato S. Estrogen synthesis in the brain—role in synaptic plasticity and memory. *Mol Cell Endocrinol* 290: 31–43, 2008.
192. Holbro N, Grunditz A, Oertner TG. Differential distribution of endoplasmic reticulum controls metabotropic signaling and plasticity at hippocampal synapses. *Proc Natl Acad Sci USA* 106: 15055–15060, 2009.
193. Holtmaat A, Svoboda K. Experience-dependent structural synaptic plasticity in the mammalian brain. *Nat Rev Neurosci* 10: 647–658, 2009.
194. Holtmaat AJ, Trachtenberg JT, Wilbrecht L, Shepherd GM, Zhang X, Knott GW, Svoboda K. Transient and persistent dendritic spines in the neocortex in vivo. *Neuron* 45: 279–291, 2005.
195. Hotulainen P, Llano O, Smirnov S, Tanhuanpää K, Faix J, Rivera C, Lappalainen P. Defining mechanisms of actin polymerization and depolymerization during dendritic spine morphogenesis. *J Cell Biol* 185: 323–339, 2009.
196. Hozumi Y, Fukaya M, Adachi N, Saito N, Otani K, Kondo H, Watanabe M, Goto K. Diacylglycerol kinase beta accumulates on the perisynaptic site of medium spiny neurons in the striatum. *Eur J Neurosci* 28: 2409–2422, 2008.
197. Hsieh H, Boehm J, Sato C, Iwatsubo T, Tomita T, Sisodia S, Malinow R. AMPAR removal underlies Abeta-induced synaptic depression and dendritic spine loss. *Neuron* 52: 831–843, 2006.
198. Hu H, Qin Y, Bochorishvili G, Zhu Y, van Aelst L, Zhu JJ. Ras signaling mechanisms underlying impaired GluR1-dependent plasticity associated with fragile X syndrome. *J Neurosci* 28: 7847–7862, 2008.
199. Huber KM, Gallagher SM, Warren ST, Bear MF. Altered synaptic plasticity in a mouse model of fragile X mental retardation. *Proc Natl Acad Sci USA* 99: 7746–7750, 2002.
200. Hung AY, Futai K, Sala C, Valtchanoff JG, Ryu J, Woodworth MA, Kidd FL, Sung CC, Miyakawa T, Bear MF, Weinberg RJ, Sheng M. Smaller dendritic spines, weaker synaptic transmission, but enhanced spatial learning in mice lacking Shank1. *J Neurosci* 28: 1697–1708, 2008.
201. Hutsler JJ, Zhang H. Increased dendritic spine densities on cortical projection neurons in autism spectrum disorders. *Brain Res* 1309: 83–94, 2010.
202. Impey S, Davare M, Lesiak A, Lasiek A, Fortin D, Ando H, Varlamova O, Obrietan K, Soderling TR, Goodman RH, Wayman GA. An activity-induced microRNA controls dendritic spine formation by regulating Rac1-PAK signaling. *Mol Cell Neurosci* 43: 146–156, 2010.
203. Irie F, Yamaguchi Y. EphB receptors regulate dendritic spine development via intersectin, Cdc42 and N-WASP. *Nat Neurosci* 5: 1117–1118, 2002.
204. Irwin SA, Idupulapati M, Gilbert ME, Harris JB, Chakravarti AB, Rogers EJ, Crisostomo RA, Larsen BP, Mehta A, Alcantara CJ, Patel B, Swain RA, Weiler IJ, Oostra BA, Greenough WT. Dendritic spine and dendritic field characteristics of layer V pyramidal neurons in the visual cortex of fragile-X knockout mice. *Am J Med Genet* 111: 140–146, 2002.
205. Irwin SA, Patel B, Idupulapati M, Harris JB, Crisostomo RA, Larsen BP, Kooy F, Willemis PJ, Cras P, Kozlowski PB, Swain RA, Weiler IJ, Greenough WT. Abnormal dendritic spine characteristics in the temporal and visual cortices of patients with fragile-X syndrome: a quantitative examination. *Am J Med Genet* 98: 161–167, 2001.
206. Ishikawa Y, Katoh H, Negishi M. A role of Rnd1 GTPase in dendritic spine formation in hippocampal neurons. *J Neurosci* 23: 11065–11072, 2003.
207. Jacobsen JS, Wu CC, Redwine JM, Comery TA, Arias R, Bowlby M, Martone R, Morrison JH, Pangalos MN, Reinhart PH, Bloom FE. Early-onset behavioral and synaptic deficits in a mouse model of Alzheimer's disease. *Proc Natl Acad Sci USA* 103: 5161–5166, 2006.
208. Jamain S, Radyushkin K, Hammerschmidt K, Granon S, Boretius S, Varoqueaux F, Ramanantsoa N, Gallego J, Ronnenberg A, Winter D, Frahm J, Fischer J, Bourgeron T, Ehrenreich H, Brose N. Reduced social interaction and ultrasonic communication in a

- mouse model of monogenic heritable autism. *Proc Natl Acad Sci USA* 105: 1710–1715, 2008.
209. Janssens RM, Brus L, Cahill DJ, Huirne JA, Schoemaker J, Lambalk CB. Direct ovarian effects and safety aspects of GnRH agonists and antagonists. *Hum Reprod Update* 6: 505–518, 2000.
 210. Jaworski J, Kapitein LC, Gouveia SM, Dortland BR, Wulf PS, Grigoriev I, Camera P, Spangler SA, Di Stefano P, Demmers J, Krugers H, Defilippi P, Akhmanova A, Hoogenraad CC. Dynamic microtubules regulate dendritic spine morphology and synaptic plasticity. *Neuron* 61: 85–100, 2009.
 211. Jaworski J, Spangler S, Seeburg DP, Hoogenraad CC, Sheng M. Control of dendritic arborization by the phosphoinositide-3'-kinase-Akt-mammalian target of rapamycin pathway. *J Neurosci* 25: 11300–11312, 2005.
 212. Jensen CS, Rasmussen HB, Misonou H. Neuronal trafficking of voltage-gated potassium channels. *Mol Cell Neurosci* 48: 288–297, 2011.
 213. Ji Y, Gong Y, Gan W, Beach T, Holtzman DM, Wisniewski T. Apolipoprotein E isoform-specific regulation of dendritic spine morphology in apolipoprotein E transgenic mice and Alzheimer's disease patients. *Neuroscience* 122: 305–315, 2003.
 214. Jinushi-Nakao S, Arvind R, Amikura R, Kinameri E, Liu AW, Moore AW. Knot/Collier and cut control different aspects of dendrite cytoskeleton and synergize to define final arbor shape. *Neuron* 56: 963–978, 2007.
 215. Johnson OL, Ouimet CC. A regulatory role for actin in dendritic spine proliferation. *Brain Res* 1113: 1–9, 2006.
 216. Johnston MV, Blue ME, Naidu S. Rett syndrome and neuronal development. *J Child Neurol* 20: 759–763, 2005.
 217. Jones KA, Srivastava DP, Allen JA, Strachan RT, Roth BL, Penzes P. Rapid modulation of spine morphology by the 5-HT_{2A} serotonin receptor through kalirin-7 signaling. *Proc Natl Acad Sci USA* 106: 19575–19580, 2009.
 218. Jourdain P, Fukunaga K, Muller D. Calcium/calmodulin-dependent protein kinase II contributes to activity-dependent filopodia growth and spine formation. *J Neurosci* 23: 10645–10649, 2003.
 219. Jung CK, Herms J. Structural dynamics of dendritic spines are influenced by an environmental enrichment: an in vivo imaging study. *Cereb Cortex* 2012.
 220. Kaech S, Fischer M, Doll T, Matus A. Isoform specificity in the relationship of actin to dendritic spines. *J Neurosci* 17: 9565–9572, 1997.
 221. Kamikubo Y, Egashira Y, Tanaka T, Shinoda Y, Tominaga-Yoshino K, Ogura A. Long-lasting synaptic loss after repeated induction of LTD: independence to the means of LTD induction. *Eur J Neurosci* 24: 1606–1616, 2006.
 222. Kang MG, Guo Y, Hugarir RL. AMPA receptor and GEF-H1/Lfc complex regulates dendritic spine development through RhoA signaling cascade. *Proc Natl Acad Sci USA* 106: 3549–3554, 2009.
 223. Kasai H, Fukuda M, Watanabe S, Hayashi-Takagi A, Noguchi J. Structural dynamics of dendritic spines in memory and cognition. *Trends Neurosci* 33: 121–129, 2010.
 224. Kayser MS, Lee AC, Hruska M, Dalva MB. Preferential control of basal dendritic protrusions by EphB2. *PLoS One* 6: e17417, 2011.
 225. Kayser MS, Nolt MJ, Dalva MB. EphB receptors couple dendritic filopodia motility to synapse formation. *Neuron* 59: 56–69, 2008.
 226. Keck T, Mrcic-Flogel TD, Vaz Afonso M, Eysel UT, Bonhoeffer T, Hübener M. Massive restructuring of neuronal circuits during functional reorganization of adult visual cortex. *Nat Neurosci* 11: 1162–1167, 2008.
 227. Kerrisk ME, Greer CA, Koleske AJ. Integrin $\alpha 3$ is required for late postnatal stability of dendrite arbors, dendritic spines and synapses, and mouse behavior. *J Neurosci* 33: 6742–6752, 2013.
 228. Kerti K, Lorincz A, Nusser Z. Unique somato-dendritic distribution pattern of Kv4.2 channels on hippocampal CA1 pyramidal cells. *Eur J Neurosci* 35: 66–75, 2012.
 229. Khelifaoui M, Denis C, van Galen E, de Bock F, Schmitt A, Houbroun C, Morice E, Geros B, Ramakers G, Fagni L, Chelly J, Nosten-Bertrand M, Billuart P. Loss of X-linked mental retardation gene oligophrenin I in mice impairs spatial memory and leads to ventricular enlargement and dendritic spine immaturity. *J Neurosci* 27: 9439–9450, 2007.
 230. Khelifaoui M, Pavlowsky A, Powell AD, Valnegri P, Cheong KW, Blandin Y, Passafaro M, Jefferys JG, Chelly J, Billuart P. Inhibition of RhoA pathway rescues the endocytosis defects in Oligophrenin I mouse model of mental retardation. *Hum Mol Genet* 18: 2575–2583, 2009.
 231. Kim IH, Raciz B, Wang H, Burianek L, Weinberg R, Yasuda R, Wetsel WC, Soderling SH. Disruption of Arp2/3 results in asymmetric structural plasticity of dendritic spines and progressive synaptic and behavioral abnormalities. *J Neurosci* 33: 6081–6092, 2013.
 232. Kim JY, Oh MH, Bernard LP, Macara IG, Zhang H. The RhoG/ELMO1/Dock180 signaling module is required for spine morphogenesis in hippocampal neurons. *J Biol Chem* 286: 37615–37624, 2011.
 233. Kim K, Yang J, Zhong XP, Kim MH, Kim YS, Lee HW, Han S, Choi J, Han K, Seo J, Prescott SM, Topham MK, Bae YC, Koretzky G, Choi SY, Kim E. Synaptic removal of diacylglycerol by DGK ζ and PSD-95 regulates dendritic spine maintenance. *EMBO J* 28: 1170–1179, 2009.
 234. Kim MH, Choi J, Yang J, Chung W, Kim JH, Paik SK, Kim K, Han S, Won H, Bae YS, Cho SH, Seo J, Bae YC, Choi SY, Kim E. Enhanced NMDA receptor-mediated synaptic transmission, enhanced long-term potentiation, and impaired learning and memory in mice lacking IRSp53. *J Neurosci* 29: 1586–1595, 2009.
 235. Kim S, Burette A, Chung HS, Kwon SK, Woo J, Lee HW, Kim K, Kim H, Weinberg RJ, Kim E. NGL family PSD-95-interacting adhesion molecules regulate excitatory synapse formation. *Nat Neurosci* 9: 1294–1301, 2006.
 236. Kim SM, Choi KY, Cho IH, Rhy JH, Kim SH, Park CS, Kim E, Song WK. Regulation of dendritic spine morphology by SPIN90, a novel Shank binding partner. *J Neurochem* 109: 1106–1117, 2009.
 237. Kim Y, Sung JY, Ceglia I, Lee KW, Ahn JH, Halford JM, Kim AM, Kwak SP, Park JB, Ho Ryu S, Schenck A, Bardoni B, Scott JD, Nairn AC, Greengard P. Phosphorylation of WAVE1 regulates actin polymerization and dendritic spine morphology. *Nature* 442: 814–817, 2006.
 238. Kirov SA, Petrak LJ, Fiala JC, Harris KM. Dendritic spines disappear with chilling but proliferate excessively upon rewarming of mature hippocampus. *Neuroscience* 127: 69–80, 2004.
 - 238a. Kleindienst T, Winnubst J, Roth-Alpermann C, Bonhoeffer T, Lohmann C. Activity-dependent clustering of functional synaptic inputs on developing hippocampal dendrites. *Neuron* 72: 1012–1024, 2011.
 239. Knafo S, Alonso-Nanclares L, Gonzalez-Soriano J, Merino-Serrais P, Feraud-Espinosa I, Ferrer I, DeFelipe J. Widespread changes in dendritic spines in a model of Alzheimer's disease. *Cereb Cortex* 19: 586–592, 2009.
 240. Knobloch M, Mansuy IM. Dendritic spine loss and synaptic alterations in Alzheimer's disease. *Mol Neurobiol* 37: 73–82, 2008.
 241. Knott GW, Holtmaat A, Wilbrecht L, Welker E, Svoboda K. Spine growth precedes synapse formation in the adult neocortex in vivo. *Nat Neurosci* 9: 1117–1124, 2006.
 242. Ko J, Kim S, Chung HS, Kim K, Han K, Kim H, Jun H, Kaang BK, Kim E. SALM synaptic cell adhesion-like molecules regulate the differentiation of excitatory synapses. *Neuron* 50: 233–245, 2006.
 243. Koleske AJ. Molecular mechanisms of dendrite stability. *Nat Rev Neurosci* 14: 536–550, 2013.
 244. Kolluri N, Sun Z, Sampson AR, Lewis DA. Lamina-specific reductions in dendritic spine density in the prefrontal cortex of subjects with schizophrenia. *Am J Psychiatry* 162: 1200–1202, 2005.
 245. Kolomeets NS, Orlovskaya DD, Rachmanova VI, Uranova NA. Ultrastructural alterations in hippocampal mossy fiber synapses in schizophrenia: a postmortem morphometric study. *Synapse* 57: 47–55, 2005.
 246. Komatsuzaki Y, Hatanaka Y, Murakami G, Mukai H, Hojo Y, Saito M, Kimoto T, Kawato S. Corticosterone induces rapid spinogenesis via synaptic glucocorticoid receptors and kinase networks in hippocampus. *PLoS One* 7: e34124, 2012.
 247. Komiyama T, Luo L. Intrinsic control of precise dendritic targeting by an ensemble of transcription factors. *Curr Biol* 17: 278–285, 2007.
 248. Kopec CD, Real E, Kessels HW, Malinow R. GluRI links structural and functional plasticity at excitatory synapses. *J Neurosci* 27: 13706–13718, 2007.

249. Korkotian E, Holzman D, Segal M. Dynamic regulation of spine-dendrite coupling in cultured hippocampal neurons. *Eur J Neurosci* 20: 2649–2663, 2004.
250. Korkotian E, Segal M. Morphological constraints on calcium dependent glutamate receptor trafficking into individual dendritic spine. *Cell Calcium* 42: 41–57, 2007.
251. Korkotian E, Segal M. Release of calcium from stores alters the morphology of dendritic spines in cultured hippocampal neurons. *Proc Natl Acad Sci USA* 96: 12068–12072, 1999.
252. Korkotian E, Segal M. Spike-associated fast contraction of dendritic spines in cultured hippocampal neurons. *Neuron* 30: 751–758, 2001.
253. Korkotian E, Segal M. Synaptopodin regulates release of calcium from stores in dendritic spines of cultured hippocampal neurons. *J Physiol* 589: 5987–5995, 2011.
254. Kovalchuk Y, Eilers J, Lisman J, Konnerth A. NMDA receptor-mediated subthreshold Ca^{2+} signals in spines of hippocampal neurons. *J Neurosci* 20: 1791–1799, 2000.
255. Kramár EA, Babayan AH, Gavin CF, Cox CD, Jafari M, Gall CM, Rumbaugh G, Lynch G. Synaptic evidence for the efficacy of spaced learning. *Proc Natl Acad Sci USA* 109: 5121–5126, 2012.
256. Kreis P, Thévenot E, Rousseau V, Boda B, Muller D, Barnier JV. The p21-activated kinase 3 implicated in mental retardation regulates spine morphogenesis through a Cdc42-dependent pathway. *J Biol Chem* 282: 21497–21506, 2007.
257. Kretz O, Fester L, Wehrenberg U, Zhou L, Brauckmann S, Zhao S, Prange-Kiel J, Naumann T, Jarry H, Frotscher M, Rune GM. Hippocampal synapses depend on hippocampal estrogen synthesis. *J Neurosci* 24: 5913–5921, 2004.
258. Kulkarni VA, Firestein BL. The dendritic tree and brain disorders. *Mol Cell Neurosci* 50: 10–20, 2012.
259. Kumar V, Zhang MX, Swank MW, Kunz J, Wu GY. Regulation of dendritic morphogenesis by Ras-PI3K-Akt-mTOR and Ras-MAPK signaling pathways. *J Neurosci* 25: 11288–11299, 2005.
260. Kushima I, Nakamura Y, Aleksic B, Ikeda M, Ito Y, Shiino T, Okochi T, Fukuo Y, Ujike H, Suzuki M, Inada T, Hashimoto R, Takeda M, Kaibuchi K, Iwata N, Ozaki N. Resequencing and association analysis of the KALRN and EPHB1 genes and their contribution to schizophrenia susceptibility. *Schizophr Bull* 38: 552–560, 2012.
261. Kutsche K, Yntema H, Brandt A, Jantke I, Nothwang HG, Orth U, Boavida MG, David D, Chelly J, Fryns JP, Moraine C, Ropers HH, Hamel BC, van Bokhoven H, Gal A. Mutations in ARHGEF6, encoding a guanine nucleotide exchange factor for Rho GTPases, in patients with X-linked mental retardation. *Nat Genet* 26: 247–250, 2000.
262. Kvaajo M, McKellar H, Arguello PA, Drew LJ, Moore H, MacDermott AB, Karayiorgou M, Gogos JA. A mutation in mouse Disc1 that models a schizophrenia risk allele leads to specific alterations in neuronal architecture and cognition. *Proc Natl Acad Sci USA* 105: 7076–7081, 2008.
263. Kwon CH, Luikart BW, Powell CM, Zhou J, Matheny SA, Zhang W, Li Y, Baker SJ, Parada LF. Pten regulates neuronal arborization and social interaction in mice. *Neuron* 50: 377–388, 2006.
264. Kwon HB, Sabatini BL. Glutamate induces de novo growth of functional spines in developing cortex. *Nature* 474: 100–104, 2011.
265. Lacor PN, Buniel MC, Furlow PW, Clemente AS, Velasco PT, Wood M, Viola KL, Klein WL. Abeta oligomer-induced aberrations in synapse composition, shape, and density provide a molecular basis for loss of connectivity in Alzheimer's disease. *J Neurosci* 27: 796–807, 2007.
266. Laggenbauer B, Ostareck D, Keidel EM, Ostareck-Lederer A, Fischer U. Evidence that fragile X mental retardation protein is a negative regulator of translation. *Hum Mol Genet* 10: 329–338, 2001.
267. Lai CS, Franke TF, Gan WB. Opposite effects of fear conditioning and extinction on dendritic spine remodelling. *Nature* 483: 87–91, 2012.
268. Lamprecht R, Farb CR, Rodrigues SM, LeDoux JE. Fear conditioning drives profilin into amygdala dendritic spines. *Nat Neurosci* 9: 481–483, 2006.
269. Landi S, Putignano E, Boggio EM, Giustetto M, Pizzorusso T, Ratto GM. The short-term structural plasticity of dendritic spines is altered in a model of Rett syndrome. *Sci Rep* 1: 45, 2011.
270. Lanz TA, Carter DB, Merchant KM. Dendritic spine loss in the hippocampus of young PDAPP and Tg2576 mice and its prevention by the ApoE2 genotype. *Neurobiol Dis* 13: 246–253, 2003.
271. Lauri SE, Lamsa K, Pavlov I, Riekkari R, Johnson BE, Molnar E, Rauvala H, Taira T. Activity blockade increases the number of functional synapses in the hippocampus of newborn rats. *Mol Cell Neurosci* 22: 107–117, 2003.
272. Lee H, Raiker SJ, Venkatesh K, Geary R, Robak LA, Zhang Y, Yeh HH, Shrager P, Giger RJ. Synaptic function for the Nogo-66 receptor NgR1: regulation of dendritic spine morphology and activity-dependent synaptic strength. *J Neurosci* 28: 2753–2765, 2008.
273. Lee HW, Choi J, Shin H, Kim K, Yang J, Na M, Choi SY, Kang GB, Eom SH, Kim H, Kim E. Preso, a novel PSD-95-interacting FERM and PDZ domain protein that regulates dendritic spine morphogenesis. *J Neurosci* 28: 14546–14556, 2008.
274. Lee KJ, Lee Y, Rozeboom A, Lee JY, Udagawa N, Hoe HS, Pak DT. Requirement for PIK2 in orchestrated ras and rap signaling, homeostatic structural plasticity, and memory. *Neuron* 69: 957–973, 2011.
- 274a. Lee KJ, Park IS, Kim H, Greenough WT, Pak DT, Rhyu IJ. Motor skill training induces coordinated strengthening and weakening between neighboring synapses. *J Neurosci* 33: 9794–9799, 2013.
275. Lee S, Lee K, Hwang S, Kim SH, Song WK, Park ZY, Chang S. SPIN90/WISH interacts with PSD-95 and regulates dendritic spinogenesis via an N-WASP-independent mechanism. *EMBO J* 25: 4983–4995, 2006.
276. Lee SJ, Escobedo-Lozoya Y, Szatmari EM, Yasuda R. Activation of CaMKII in single dendritic spines during long-term potentiation. *Nature* 458: 299–304, 2009.
277. Leitch B, Szostek A, Lin R, Shevtsova O. Subcellular distribution of L-type calcium channel subtypes in rat hippocampal neurons. *Neuroscience* 164: 641–657, 2009.
278. Li B, Woo RS, Mei L, Malinow R. The neuregulin-1 receptor erbB4 controls glutamatergic synapse maturation and plasticity. *Neuron* 54: 583–597, 2007.
279. Li C, Brake WG, Romeo RD, Dunlop JC, Gordon M, Buzescu R, Magarinos AM, Allen PB, Greengard P, Luine V, McEwen BS. Estrogen alters hippocampal dendritic spine shape and enhances synaptic protein immunoreactivity and spatial memory in female mice. *Proc Natl Acad Sci USA* 101: 2185–2190, 2004.
280. Li Z, Okamoto K, Hayashi Y, Sheng M. The importance of dendritic mitochondria in the morphogenesis and plasticity of spines and synapses. *Cell* 119: 873–887, 2004.
281. Li Z, Zhang Y, Ku L, Wilkinson KD, Warren ST, Feng Y. The fragile X mental retardation protein inhibits translation via interacting with mRNA. *Nucleic Acids Res* 29: 2276–2283, 2001.
282. Lin WH, Nebhan CA, Anderson BR, Webb DJ. Vasodilator-stimulated phosphoprotein (VASP) induces actin assembly in dendritic spines to promote their development and potentiate synaptic strength. *J Biol Chem* 285: 36010–36020, 2010.
283. Lin YL, Lei YT, Hong CJ, Hsueh YP. Syndecan-2 induces filopodia and dendritic spine formation via the neurofibromin-PAK-Ena/VASP pathway. *J Cell Biol* 177: 829–841, 2007.
284. Lippi G, Steinert JR, Marczylo EL, D'Oro S, Fiore R, Forsythe ID, Schmitt G, Zoli M, Nicotera P, Young KW. Targeting of the Arp3 actin nucleation factor by miR-29a/b regulates dendritic spine morphology. *J Cell Biol* 194: 889–904, 2011.
285. Lipska BK, Peters T, Hyde TM, Halim N, Horowitz C, Mitkus S, Weickert CS, Matsumoto M, Sawa A, Straub RE, Vakkalanka R, Herman MM, Weinberger DR, Kleinman JE. Expression of DISC1 binding partners is reduced in schizophrenia and associated with DISC1 SNPs. *Hum Mol Genet* 15: 1245–1258, 2006.
286. Liston C, Gan WB. Glucocorticoids are critical regulators of dendritic spine development and plasticity in vivo. *Proc Natl Acad Sci USA* 108: 16074–16079, 2011.
287. Liu F, Day M, Muñiz LC, Bitran D, Arias R, Revilla-Sanchez R, Grauer S, Zhang G, Kelley C, Pulito V, Sung A, Mervis RF, Navarra R, Hirst WD, Reinhart PH, Marquis KL, Moss SJ, Pangalos MN, Brandon NJ. Activation of estrogen receptor-beta regulates hippocampal synaptic plasticity and improves memory. *Nat Neurosci* 11: 334–343, 2008.
288. Lorincz A, Nusser Z. Molecular identity of dendritic voltage-gated sodium channels. *Science* 328: 906–909, 2010.

289. Lozada AF, Wang X, Gounko NV, Massey KA, Duan J, Liu Z, Berg DK. Induction of dendritic spines by $\beta 2$ -containing nicotinic receptors. *J Neurosci* 32: 8391–8400, 2012.
290. Lu R, Wang H, Liang Z, Ku L, O'donnell WT, Li W, Warren ST, Feng Y. The fragile X protein controls microtubule-associated protein 1B translation and microtubule stability in brain neuron development. *Proc Natl Acad Sci USA* 101: 15201–15206, 2004.
291. Lu Y, Christian K, Lu B. BDNF: a key regulator for protein synthesis-dependent LTP and long-term memory? *Neurobiol Learn Mem* 89: 312–323, 2008.
292. Luikart BW, Nef S, Virmani T, Lush ME, Liu Y, Kavalali ET, Parada LF. TrkB has a cell-autonomous role in the establishment of hippocampal Schaffer collateral synapses. *J Neurosci* 25: 3774–3786, 2005.
293. Luikart BW, Parada LF. Receptor tyrosine kinase B-mediated excitatory synaptogenesis. *Prog Brain Res* 157: 15–24, 2006.
294. Luján R, Maylie J, Adelman JP. New sites of action for GIRK and SK channels. *Nat Rev Neurosci* 10: 475–480, 2009.
295. Ma XM, Huang J, Wang Y, Eipper BA, Mains RE. Kalirin, a multifunctional Rho guanine nucleotide exchange factor, is necessary for maintenance of hippocampal pyramidal neuron dendrites and dendritic spines. *J Neurosci* 23: 10593–10603, 2003.
296. Ma XM, Kiraly DD, Gaier ED, Wang Y, Kim EJ, Levine ES, Eipper BA, Mains RE. Kalirin-7 is required for synaptic structure and function. *J Neurosci* 28: 12368–12382, 2008.
297. Mabb AM, Judson MC, Zylka MJ, Philpot BD. Angelman syndrome: insights into genomic imprinting and neurodevelopmental phenotypes. *Trends Neurosci* 34: 293–303, 2011.
298. Maglione M, Sigrist SJ. Seeing the forest tree by tree: super-resolution light microscopy meets the neurosciences. *Nat Neurosci* 16: 790–797, 2013.
299. Mainen ZF, Malinow R, Svoboda K. Synaptic calcium transients in single spines indicate that NMDA receptors are not saturated. *Nature* 399: 151–155, 1999.
300. Makino H, Malinow R. Compartmentalized versus global synaptic plasticity on dendrites controlled by experience. *Neuron* 72: 1001–1011, 2011.
301. Malenka RC, Nicoll RA. Silent synapses speak up. *Neuron* 19: 473–476, 1997.
302. Maloney MT, Bamberg JR. Cofilin-mediated neurodegeneration in Alzheimer's disease and other amyloidopathies. *Mol Neurobiol* 35: 21–44, 2007.
303. Mammoto A, Sasaki T, Asakura T, Hotta I, Imamura H, Takahashi K, Matsuura Y, Shirao T, Takai Y. Interactions of drebrin and gephyrin with profilin. *Biochem Biophys Res Commun* 243: 86–89, 1998.
304. Manley S, Gillette JM, Patterson GH, Shroff H, Hess HF, Betzig E, Lippincott-Schwartz J. High-density mapping of single-molecule trajectories with photoactivated localization microscopy. *Nat Methods* 5: 155–157, 2008.
305. Manning MA, Cassidy SB, Clericuzio C, Cherry AM, Schwartz S, Hudgins L, Enns GM, Hoyne HE. Terminal 22q deletion syndrome: a newly recognized cause of speech and language disability in the autism spectrum. *Pediatrics* 114: 451–457, 2004.
306. Maras PM, Baram TZ. Sculpting the hippocampus from within: stress, spines, and CRH. *Trends Neurosci* 35: 315–324, 2012.
307. Marcello E, Epis R, Saraceno C, Di Luca M. Synaptic dysfunction in Alzheimer's disease. *Adv Exp Med Biol* 700: 573–601, 2012.
308. Marquet P, Depeursinge C, Magistretti PJ. Exploring neural cell dynamics with digital holographic microscopy. *Annu Rev Biomed Eng* 15: 407–431, 2013.
309. Mateos JM, Lüthi A, Savic N, Stierli B, Streif P, Gähwiler BH, McKinney RA. Synaptic modifications at the CA3-CA1 synapse after chronic AMPA receptor blockade in rat hippocampal slices. *J Physiol* 581: 129–138, 2007.
310. Matsuo N, Reijmers L, Mayford M. Spine-type-specific recruitment of newly synthesized AMPA receptors with learning. *Science* 319: 1104–1107, 2008.
311. Matsuzaki M, Honkura N, Ellis-Davies GC, Kasai H. Structural basis of long-term potentiation in single dendritic spines. *Nature* 429: 761–766, 2004.
312. Maynard KR, Stein E. DSCAM contributes to dendrite arborization and spine formation in the developing cerebral cortex. *J Neurosci* 32: 16637–16650, 2012.
313. McAvoy T, Zhou MM, Greengard P, Nairn AC. Phosphorylation of Rap1GAP, a striatally enriched protein, by protein kinase A controls Rap1 activity and dendritic spine morphology. *Proc Natl Acad Sci USA* 106: 3531–3536, 2009.
314. Mei L, Xiong WC. Neuregulin I in neural development, synaptic plasticity and schizophrenia. *Nat Rev Neurosci* 9: 437–452, 2008.
315. Mellios N, Sugihara H, Castro J, Banerjee A, Le C, Kumar A, Crawford B, Strathmann J, Tropea D, Levine SS, Edbauer D, Sur M. miR-132, an experience-dependent microRNA, is essential for visual cortex plasticity. *Nat Neurosci* 14: 1240–1242, 2011.
316. Mendez P, De Roo M, Poglia L, Klausner P, Muller D. N-cadherin mediates plasticity-induced long-term spine stabilization. *J Cell Biol* 189: 589–600, 2010.
317. Meng J, Meng Y, Hanna A, Janus C, Jia Z. Abnormal long-lasting synaptic plasticity and cognition in mice lacking the mental retardation gene Pak3. *J Neurosci* 25: 6641–6650, 2005.
318. Meng Y, Zhang Y, Tregoubov V, Janus C, Cruz L, Jackson M, Lu WY, MacDonald JF, Wang JY, Falls DL, Jia Z. Abnormal spine morphology and enhanced LTP in LIMK-1 knockout mice. *Neuron* 35: 121–133, 2002.
320. Menna E, Zambetti S, Morini R, Donzelli A, Disanza A, Calvigioni D, Braida D, Nicolini C, Orlando M, Fossati G, Cristina Regondi M, Pattini L, Frassoni C, Francolini M, Scita G, Sala M, Fahnstock M, Matteoli M. Eps8 controls dendritic spine density and synaptic plasticity through its actin-capping activity. *EMBO J* 32: 1730–1744, 2013.
321. Menon P, Deane R, Sagare A, Lane SM, Zarccone TJ, O'Dell MR, Yan C, Zlokovic BV, Berk BC. Impaired spine formation and learning in GPCR kinase 2 interacting protein-1 (GIT1) knockout mice. *Brain Res* 1317: 218–226, 2010.
322. Mezey S, Doyère V, De Souza I, Harrison E, Cambon K, Kendal CE, Davies H, Laroche S, Stewart MG. Long-term synaptic morphometry changes after induction of long-term potentiation and long-term depression in the dentate gyrus of awake rats are not simply mirror phenomena. *Eur J Neurosci* 19: 2310–2318, 2004.
323. Michaelsen K, Murk K, Zagrebelsky M, Drezniak A, Jockusch BM, Rothkegel M, Korte M. Fine-tuning of neuronal architecture requires two profilin isoforms. *Proc Natl Acad Sci USA* 107: 15780–15785, 2010.
324. Middei S, Restivo L, Caprioli A, Aceti M, Ammassari-Teule M. Region-specific changes in the microanatomy of single dendritic spines over time might account for selective memory alterations in ageing hAPPswTg2576 mice, a mouse model for Alzheimer disease. *Neurobiol Learn Mem* 90: 467–471, 2008.
325. Miller JP, Rall W, Rinzel J. Synaptic amplification by active membrane in dendritic spines. *Brain Res* 325: 325–330, 1985.
326. Milner B, Squire LR, Kandel ER. Cognitive neuroscience and the study of memory. *Neuron* 20: 445–468, 1998.
327. Mitra SW, Hoskin E, Yudkovitz J, Pear L, Wilkinson HA, Hayashi S, Pfaff DW, Ogawa S, Rohrer SP, Schaeffer JM, McEwen BS, Alves SE. Immunolocalization of estrogen receptor beta in the mouse brain: comparison with estrogen receptor alpha. *Endocrinology* 144: 2055–2067, 2003.
328. Miyamoto Y, Yamauchi J, Tanoue A, Wu C, Mobley WC. TrkB binds and tyrosine-phosphorylates Tiam1, leading to activation of Rac1 and induction of changes in cellular morphology. *Proc Natl Acad Sci USA* 103: 10444–10449, 2006.
329. Miyazaki H, Yamazaki M, Watanabe H, Maehama T, Yokozeki T, Kanaho Y. The small GTPase ADP-ribosylation factor 6 negatively regulates dendritic spine formation. *FEBS Lett* 579: 6834–6838, 2005.
330. Mizui T, Takahashi H, Sekino Y, Shirao T. Overexpression of drebrin A in immature neurons induces the accumulation of F-actin and PSD-95 into dendritic filopodia, and the formation of large abnormal protrusions. *Mol Cell Neurosci* 30: 149–157, 2005.
331. Moeller ML, Shi Y, Reichardt LF, Ethell IM. EphB receptors regulate dendritic spine morphogenesis through the recruitment/phosphorylation of focal adhesion kinase and RhoA activation. *J Biol Chem* 281: 1587–1598, 2006.
332. Moessner R, Marshall CR, Sutcliffe JS, Skaug J, Pinto D, Vincent J, Zwaigenbaum L, Fernandez B, Roberts W, Szatmari P, Scherer SW. Contribution of SHANK3 mutations to autism spectrum disorder. *Am J Hum Genet* 81: 1289–1297, 2007.
333. Moore CD, Thacker EE, Larimore J, Gaston D, Underwood A, Kearns B, Patterson SI, Jackson T, Chapleau C, Pozzo-Miller L, Theibert A. The neuronal Arf GAP centaurin alpha1 modulates dendritic differentiation. *J Cell Sci* 120: 2683–2693, 2007.

334. Muddashetty RS, Kelić S, Gross C, Xu M, Bassell GJ. Dysregulated metabotropic glutamate receptor-dependent translation of AMPA receptor and postsynaptic density-95 mRNAs at synapses in a mouse model of fragile X syndrome. *J Neurosci* 27: 5338–5348, 2007.
335. Muddashetty RS, Nalavadi VC, Gross C, Yao X, Xing L, Laur O, Warren ST, Bassell GJ. Reversible inhibition of PSD-95 mRNA translation by miR-125a, FMRP phosphorylation, and mGluR signaling. *Mol Cell* 42: 673–688, 2011.
336. Mukai H, Tsurugizawa T, Murakami G, Kominami S, Ishii H, Ogiue-Ikeda M, Takata N, Tanabe N, Furukawa A, Hojo Y, Ooishi Y, Morrison JH, Janssen WG, Rose JA, Chambon P, Kato S, Izumi S, Yamazaki T, Kimoto T, Kawato S. Rapid modulation of long-term depression and spinogenesis via synaptic estrogen receptors in hippocampal principal neurons. *J Neurochem* 100: 950–967, 2007.
337. Mukai J, Dhilla A, Drew LJ, Stark KL, Cao L, MacDermott AB, Karayiorgou M, Gogos JA. Palmitoylation-dependent neurodevelopmental deficits in a mouse model of 22q11 microdeletion. *Nat Neurosci* 11: 1302–1310, 2008.
338. Mundel P, Heid HW, Mundel TM, Kruger M, Reiser J, Kriz W. Synaptopodin: an actin-associated protein in telencephalic dendrites and renal podocytes. *J Cell Biol* 139: 193–204, 1997.
339. Murai KK, Nguyen LN, Irie F, Yamaguchi Y, Pasquale EB. Control of hippocampal dendritic spine morphology through ephrin-A3/EphA4 signaling. *Nat Neurosci* 6: 153–160, 2003.
340. Murakami G, Tsurugizawa T, Hatanaka Y, Komatsuzaki Y, Tanabe N, Mukai H, Hojo Y, Kominami S, Yamazaki T, Kimoto T, Kawato S. Comparison between basal and apical dendritic spines in estrogen-induced rapid spinogenesis of CA1 principal neurons in the adult hippocampus. *Biochem Biophys Res Commun* 351: 553–558, 2006.
341. Murakoshi H, Wang H, Yasuda R. Local, persistent activation of Rho GTPases during plasticity of single dendritic spines. *Nature* 472: 100–104, 2011.
342. Murphy DD, Cole NB, Greenberger V, Segal M. Estradiol increases dendritic spine density by reducing GABA neurotransmission in hippocampal neurons. *J Neurosci* 18: 2550–2559, 1998.
343. Müller W, Connor JA. Dendritic spines as individual neuronal compartments for synaptic Ca^{2+} responses. *Nature* 354: 73–76, 1991.
344. Nägerl UV, Kostinger G, Anderson JC, Martin KA, Bonhoeffer T. Protracted synaptogenesis after activity-dependent spinogenesis in hippocampal neurons. *J Neurosci* 27: 8149–8156, 2007.
345. Nair D, Hosy E, Petersen JD, Constals A, Giannone G, Choquet D, Sibarita JB. Super-resolution imaging reveals that AMPA receptors inside synapses are dynamically organized in nanodomains regulated by PSD95. *J Neurosci* 33: 13204–13224, 2013.
346. Naisbitt S, Kim E, Tu JC, Xiao B, Sala C, Valtschanoff J, Weinberg RJ, Worley PF, Sheng M. Shank, a novel family of postsynaptic density proteins that binds to the NMDA receptor/PSD-95/GKAP complex and cortactin. *Neuron* 23: 569–582, 1999.
347. Nakamura Y, Wood CL, Patton AP, Jaafari N, Henley JM, Mellor JR, Hanley JG. PICK1 inhibition of the Arp2/3 complex controls dendritic spine size and synaptic plasticity. *EMBO J* 30: 719–730, 2011.
348. Nakata H, Nakamura S. Brain-derived neurotrophic factor regulates AMPA receptor trafficking to post-synaptic densities via IP_3R and TRPC calcium signaling. *FEBS Lett* 581: 2047–2054, 2007.
349. Nakazawa T, Kuriu T, Tezuka T, Umemori H, Okabe S, Yamamoto T. Regulation of dendritic spine morphology by an NMDA receptor-associated Rho GTPase-activating protein, p250GAP. *J Neurochem* 105: 1384–1393, 2008.
350. Napoli I, Mercaldo V, Boyl PP, Eleuteri B, Zalfa F, De Rubeis S, Di Marino D, Mohr E, Massimi M, Falconi M, Witke W, Costa-Mattioli M, Sonenberg N, Achsel T, Bagni C. The fragile X syndrome protein represses activity-dependent translation through CYFIP1, a new 4E-BP. *Cell* 134: 1042–1054, 2008.
351. Nassirpour R, Bahima L, Lalive AL, Lüscher C, Luján R, Slesinger PA. Morphine- and CaMKII-dependent enhancement of GIRK channel signaling in hippocampal neurons. *J Neurosci* 30: 13419–13430, 2010.
352. Navarrete M, Perea G, Fernandez de Sevilla D, Gómez-Gonzalo M, Núñez A, Martín ED, Araque A. Astrocytes mediate in vivo cholinergic-induced synaptic plasticity. *PLoS Biol* 10: e1001259, 2012.
353. Newey SE, Velamoor V, Govek EE, Van Aelst L. Rho GTPases, dendritic structure, and mental retardation. *J Neurobiol* 64: 58–74, 2005.
354. Nikonenko I, Boda B, Steen S, Knott G, Welker E, Müller D. PSD-95 promotes synaptogenesis and multi-innervated spine formation through nitric oxide signaling. *J Cell Biol* 183: 1115–1127, 2008.
355. Nilius B, Owsianik G. The transient receptor potential family of ion channels. *Genome Biol* 12: 218, 2011.
356. Nishimura T, Yamaguchi T, Tokunaga A, Hara A, Hamaguchi T, Kato K, Iwamatsu A, Okano H, Kaibuchi K. Role of numb in dendritic spine development with a Cdc42 GEF intersectin and EphB2. *Mol Biol Cell* 17: 1273–1285, 2006.
- 356a. Niu S, Yabut O, D'Arcangelo G. The Reelin signaling pathway promotes dendritic spine development in hippocampal neurons. *J Neurosci* 28: 10339–10348, 2008.
357. Noam Y, Phan L, McClelland S, Manders EM, Ehrengreuer MU, Wadman WJ, Baram TZ, Chen Y. Distinct regional and subcellular localization of the actin-binding protein filamin A in the mature rat brain. *J Comp Neurol* 520: 3013–3034, 2012.
359. Nodé-Langlois R, Müller D, Boda B. Sequential implication of the mental retardation proteins ARHGAP6 and PAK3 in spine morphogenesis. *J Cell Sci* 119: 4986–4993, 2006.
360. Nuriya M, Jiang J, Nemet B, Eisenthal KB, Yuste R. Imaging membrane potential in dendritic spines. *Proc Natl Acad Sci USA* 103: 786–790, 2006.
361. Nägerl UV, Eberhorn N, Cambridge SB, Bonhoeffer T. Bidirectional activity-dependent morphological plasticity in hippocampal neurons. *Neuron* 44: 759–767, 2004.
362. O'Donnell C, Nolan MF, van Rossum MC. Dendritic spine dynamics regulate the long-term stability of synaptic plasticity. *J Neurosci* 31: 16142–16156, 2011.
363. Oh Y, Waxman SG. The $\beta 1$ subunit of mRNA of the rat brain Na^+ channel is expressed in glial cells. *Proc Natl Acad Sci USA* 91: 9985–9989, 1994.
364. Okabe S. Molecular dynamics of the excitatory synapse. *Adv Exp Med Biol* 970: 131–152, 2012.
365. Okabe S, Miwa A, Okado H. Spine formation and correlated assembly of presynaptic and postsynaptic molecules. *J Neurosci* 21: 6105–6114, 2001.
366. Okamoto K, Narayanan R, Lee SH, Murata K, Hayashi Y. The role of CaMKII as an F-actin-bundling protein crucial for maintenance of dendritic spine structure. *Proc Natl Acad Sci USA* 104: 6418–6423, 2007.
367. Okamura K, Tanaka H, Yagita Y, Saeki Y, Taguchi A, Hiraoka Y, Zeng LH, Colman DR, Miki N. Cadherin activity is required for activity-induced spine remodeling. *J Cell Biol* 167: 961–972, 2004.
368. Okubo-Suzuki R, Okada D, Sekiguchi M, Inokuchi K. Synaptopodin maintains the neural activity-dependent enlargement of dendritic spines in hippocampal neurons. *Mol Cell Neurosci* 38: 266–276, 2008.
369. Oliver CJ, Terry-Lorenzo RT, Elliott E, Bloomer WA, Li S, Brautigan DL, Colbran RJ, Shenolikar S. Targeting protein phosphatase 1 (PP1) to the actin cytoskeleton: the neurabin I/PP1 complex regulates cell morphology. *Mol Cell Biol* 22: 4690–4701, 2002.
370. Ooishi Y, Kawato S, Hojo Y, Hatanaka Y, Higo S, Murakami G, Komatsuzaki Y, Ogiue-Ikeda M, Kimoto T, Mukai H. Modulation of synaptic plasticity in the hippocampus by hippocampus-derived estrogen and androgen. *J Steroid Biochem Mol Biol* 131: 37–51, 2012.
- 370a. Oray S, Majewska A, Sur M. Dendritic spine dynamics are regulated by monocular deprivation and extracellular matrix degradation. *Neuron* 44: 1021–1030, 2004.
371. Osterweil E, Wells DG, Mooseker MS. A role for myosin VI in postsynaptic structure and glutamate receptor endocytosis. *J Cell Biol* 168: 329–338, 2005.
372. Ostroff LE, Fiala JC, Allwardt B, Harris KM. Polyribosomes redistribute from dendritic shafts into spines with enlarged synapses during LTP in developing rat hippocampal slices. *Neuron* 35: 535–545, 2002.
373. Ovtcharoff W, Segal M, Goldin M, Helmeke C, Kreher U, Greenberger V, Herzog A, Michaelis B, Braun K. Electron microscopic 3D-reconstruction of dendritic spines in cultured hippocampal neurons undergoing synaptic plasticity. *Dev Neurobiol* 68: 870–876, 2008.

374. Packer AM, Peterka DS, Hirtz JJ, Prakash R, Deisseroth K, Yuste R. Two-photon optogenetics of dendritic spines and neural circuits. *Nat Methods* 9: 1202–1205, 2012.
375. Pak DT, Sheng M. Targeted protein degradation and synapse remodeling by an inducible protein kinase. *Science* 302: 1368–1373, 2003.
376. Pak DT, Yang S, Rudolph-Correia S, Kim E, Sheng M. Regulation of dendritic spine morphology by SPAR, a PSD-95-associated RapGAP. *Neuron* 31: 289–303, 2001.
377. Palmer LM, Stuart GJ. Membrane potential changes in dendritic spines during action potentials and synaptic input. *J Neurosci* 29: 6897–6903, 2009.
378. Park E, Chi S, Park D. Activity-dependent modulation of the interaction between CaMKII α and Abi1 and its involvement in spine maturation. *J Neurosci* 32: 13177–13188, 2012.
379. Park E, Na M, Choi J, Kim S, Lee JR, Yoon J, Park D, Sheng M, Kim E. The Shank family of postsynaptic density proteins interacts with and promotes synaptic accumulation of the beta PIX guanine nucleotide exchange factor for Rac1 and Cdc42. *J Biol Chem* 278: 19220–19229, 2003.
380. Passafium M, Nakagawa T, Sala C, Sheng M. Induction of dendritic spines by an extracellular domain of AMPA receptor subunit GluR2. *Nature* 424: 677–681, 2003.
381. Pathania M, Torres-Reveron J, Yan L, Kimura T, Lin TV, Gordon V, Teng ZQ, Zhao X, Fulga TA, Van Vactor D, Border A. miR-132 enhances dendritic morphogenesis, spine density, synaptic integration, and survival of newborn olfactory bulb neurons. *PLoS One* 7: e38174, 2012.
382. Pavlowsky A, Gianfelice A, Pallotto M, Zanchi A, Vara H, Khelifaoui M, Valnegri P, Rezaei X, Bassani S, Brambilla D, Kumpost J, Blahos J, Roux MJ, Humeau Y, Chelly J, Passafium M, Giustetto M, Billuart P, Sala C. A postsynaptic signaling pathway that may account for the cognitive defect due to ILIRAPL1 mutation. *Curr Biol* 20: 103–115, 2010.
383. Penzes P, Beeser A, Chernoff J, Schiller MR, Eipper BA, Mains RE, Huganir RL. Rapid induction of dendritic spine morphogenesis by trans-synaptic ephrinB-EphB receptor activation of the Rho-GEF kalirin. *Neuron* 37: 263–274, 2003.
384. Penzes P, Cahill ME, Jones KA, VanLeeuwen JE, Woolfrey KM. Dendritic spine pathology in neuropsychiatric disorders. *Nat Neurosci* 14: 285–293, 2011.
385. Penzes P, Johnson RC, Sattler R, Zhang X, Huganir RL, Kambampati V, Mains RE, Eipper BA. The neuronal Rho-GEF Kalirin-7 interacts with PDZ domain-containing proteins and regulates dendritic morphogenesis. *Neuron* 29: 229–242, 2001.
386. Penzes P, Rafalovich I. Regulation of the actin cytoskeleton in dendritic spines. *Adv Exp Med Biol* 970: 81–95, 2012.
387. Perkel DH, Perkel DJ. Dendritic spines: role of active membrane in modulating synaptic efficacy. *Brain Res* 325: 331–335, 1985.
388. Petrecca K, Miller DM, Shrier A. Localization and enhanced current density of the Kv4.2 potassium channel by interaction with the actin-binding protein filamin. *J Neurosci* 20: 8736–8744, 2000.
389. Peça J, Feliciano C, Ting JT, Wang W, Wells MF, Venkatraman TN, Lascola CD, Fu Z, Feng G. Shank3 mutant mice display autistic-like behaviours and striatal dysfunction. *Nature* 472: 437–442, 2011.
390. Phelan MC, Rogers RC, Saul RA, Stapleton GA, Sweet K, McDermid H, Shaw SR, Claytor J, Willis J, Kelly DP. 22q13 deletion syndrome. *Am J Med Genet* 101: 91–99, 2001.
391. Pilpel Y, Segal M. Rapid WAVE dynamics in dendritic spines of cultured hippocampal neurons is mediated by actin polymerization. *J Neurochem* 95: 1401–1410, 2005.
392. Pontrello CG, Sun MY, Lin A, Fiocco TA, DeFea KA, Ethell IM. Cofilin under control of β -arrestin-2 in NMDA-dependent dendritic spine plasticity, long-term depression (LTD), and learning. *Proc Natl Acad Sci USA* 109: E442–451, 2012.
393. Pop AS, Levenga J, de Esch CE, Buijsen RA, Nieuwenhuizen IM, Li T, Isaacs A, Gasparini F, Oostra BA, Willemsen R. Rescue of dendritic spine phenotype in Fmr1 KO mice with the mGluR5 antagonist AFQ056/Mavoglurant. *Psychopharmacology* 2012.
394. Popov VI, Bocharova LS. Hibernation-induced structural changes in synaptic contacts between mossy fibres and hippocampal pyramidal neurons. *Neuroscience* 48: 53–62, 1992.
395. Popov VI, Davies HA, Rogachevsky VV, Patrushev IV, Errington ML, Gabbott PL, Bliss TV, Stewart MG. Remodelling of synaptic morphology but unchanged synaptic density during late phase long-term potentiation (LTP): a serial section electron micrograph study in the dentate gyrus in the anaesthetised rat. *Neuroscience* 128: 251–262, 2004.
396. Popov VI, Medvedev NI, Patrushev IV, Ignat'ev DA, Morenkov ED, Stewart MG. Reversible reduction in dendritic spines in CA1 of rat and ground squirrel subjected to hypothermia-normothermia in vivo: a three-dimensional electron microscope study. *Neuroscience* 149: 549–560, 2007.
397. Portera-Cailliau C, Pan DT, Yuste R. Activity-regulated dynamic behavior of early dendritic protrusions: evidence for different types of dendritic filopodia. *J Neurosci* 23: 7129–7142, 2003.
398. Prange O, Murphy TH. Modular transport of postsynaptic density-95 clusters and association with stable spine precursors during early development of cortical neurons. *J Neurosci* 21: 9325–9333, 2001.
399. Prange O, Wong TP, Gerrow K, Wang YT, El-Husseini A. A balance between excitatory and inhibitory synapses is controlled by PSD-95 and neuroligin. *Proc Natl Acad Sci USA* 101: 13915–13920, 2004.
400. Prange-Kiel J, Jarry H, Schoen M, Kohlmann P, Lohse C, Zhou L, Rune GM. Gonadotropin-releasing hormone regulates spine density via its regulatory role in hippocampal estrogen synthesis. *J Cell Biol* 180: 417–426, 2008.
401. Proepper C, Johannsen S, Liebau S, Dahl J, Vaida B, Bockmann J, Kreutz MR, Gundelfinger ED, Boeckers TM. Abelson interacting protein 1 (Abi-1) is essential for dendrite morphogenesis and synapse formation. *EMBO J* 26: 1397–1409, 2007.
402. Proepper C, Steinestel K, Schmeisser MJ, Heinrich J, Steinestel J, Bockmann J, Liebau S, Boeckers TM. Heterogeneous nuclear ribonucleoprotein k interacts with Abi-1 at postsynaptic sites and modulates dendritic spine morphology. *PLoS One* 6: e27045, 2011.
403. Qualmann B, Boeckers TM, Jeromin M, Gundelfinger ED, Kessels MM. Linkage of the actin cytoskeleton to the postsynaptic density via direct interactions of Abp1 with the ProSAP/Shank family. *J Neurosci* 24: 2481–2495, 2004.
404. Raemaekers T, Peric A, Baatsen P, Sannerud R, Declerck I, Baert V, Michiels C, Annaert W. ARF6-mediated endosomal transport of telencephalin affects dendritic filopodia-to-spine maturation. *EMBO J* 31: 3252–3269, 2012.
405. Rall W. Distributions of potential in cylindrical coordinates and time constants for a membrane cylinder. *Biophys J* 9: 1509–1541, 1969.
406. Ramakers GJ, Wolfer D, Rosenberger G, Kuchenbecker K, Kreienkamp HJ, Prange-Kiel J, Rune G, Richter K, Langnaese K, Masneuf S, Bösl MR, Fischer KD, Krugers HJ, Lipp HP, van Galen E, Kutsche K. Dysregulation of Rho GTPases in the α Pix/Arhgef6 mouse model of X-linked intellectual disability is paralleled by impaired structural and synaptic plasticity and cognitive deficits. *Hum Mol Genet* 21: 268–286, 2012.
407. Rami G, Caillard O, Medina I, Pellegrino C, Fattoum A, Ben-Ari Y, Ferhat L. Change in the shape and density of dendritic spines caused by overexpression of acidic calponin in cultured hippocampal neurons. *Hippocampus* 16: 183–197, 2006.
408. Ramon y Cajal S. Sur la structure de l'écorce cérébrale de quelques mammifères. *La Cellule* 7: 125–176, 1891.
409. Rauch A, Hoyer J, Guth S, Zweier C, Kraus C, Becker C, Zenker M, Hüffmeier U, Thiel C, Rüschenhoff F, Nürnberg P, Reis A, Trautmann U. Diagnostic yield of various genetic approaches in patients with unexplained developmental delay or mental retardation. *Am J Med Genet A* 140: 2063–2074, 2006.
410. Raymond CR, Redman SJ. Spatial segregation of neuronal calcium signals encodes different forms of LTP in rat hippocampus. *J Physiol* 570: 97–111, 2006.
411. Reeve SP, Bassetto L, Genova GK, Kleyner Y, Leyssen M, Jackson FR, Hassan BA. The *Drosophila* fragile X mental retardation protein controls actin dynamics by directly regulating profilin in the brain. *Curr Biol* 15: 1156–1163, 2005.
412. Reilly JE, Hanson HH, Phillips GR. Persistence of excitatory shaft synapses adjacent to newly emerged dendritic protrusions. *Mol Cell Neurosci* 48: 129–136, 2011.
413. Restivo L, Ferrari F, Passino E, Sgobio C, Bock J, Oostra BA, Bagni C, Ammassari-Teule M. Enriched environment promotes behavioral and morphological recovery in a mouse model for the fragile X syndrome. *Proc Natl Acad Sci USA* 102: 11557–11562, 2005.

414. Ricciardi S, Ungaro F, Hambrock M, Rademacher N, Stefanelli G, Brambilla D, Sessa A, Magagnotti C, Bachi A, Giarda E, Verpelli C, Kilstrup-Nielsen C, Sala C, Kalscheuer VM, Broccoli V. CDKL5 ensures excitatory synapse stability by reinforcing NGL-1-PSD95 interaction in the postsynaptic compartment and is impaired in patient iPSC-derived neurons. *Nat Cell Biol* 14: 911–923, 2012.
415. Richards DA, Mateos JM, Hugel S, de Paola V, Caroni P, Gähwiler BH, McKinney RA. Glutamate induces the rapid formation of spine head protrusions in hippocampal slice cultures. *Proc Natl Acad Sci USA* 102: 6166–6171, 2005.
416. Rochefort NL, Konnerth A. Dendritic spines: from structure to in vivo function. *EMBO Rep* 13: 699–708, 2012.
417. Ron S, Dudai Y, Segal M. Overexpression of PKM ζ alters morphology and function of dendritic spines in cultured cortical neurons. *Cereb Cortex* 22: 2519–2528, 2012.
418. Rose CR, Kovalchuk Y, Eilers J, Konnerth A. Two-photon Na⁺ imaging in spines and fine dendrites of central neurons. *Pflügers Arch* 439: 201–207, 1999.
419. Roselli F, Hutzler P, Wegerich Y, Livrea P, Almeida OF. Disassembly of shank and homer synaptic clusters is driven by soluble beta-amyloid(1–40) through divergent NMDAR-dependent signalling pathways. *PLoS One* 4: e6011, 2009.
420. Roussignol G, Ango F, Romorini S, Tu JC, Sala C, Worley PF, Bockeaert J, Fagni L. Shank expression is sufficient to induce functional dendritic spine synapses in aspinous neurons. *J Neurosci* 25: 3560–3570, 2005.
421. Roussignol G, Ango F, Romorini S, Tu JC, Sala C, Worley PF, Bockeaert JRL, Fagni L. Shank expression is sufficient to induce functional dendritic spine synapses in aspinous neurons. *J Neurosci* 25: 3560–3570, 2005.
422. Rubio MD, Johnson R, Miller CA, Haganir RL, Rumbaugh G. Regulation of synapse structure and function by distinct myosin II motors. *J Neurosci* 31: 1448–1460, 2011.
423. Rune GM, Wehrenberg U, Prange-Kiel J, Zhou L, Adelman G, Frotscher M. Estrogen up-regulates estrogen receptor alpha and synaptophysin in slice cultures of rat hippocampus. *Neuroscience* 113: 167–175, 2002.
424. Ryan XP, Alldritt J, Svenningsson P, Allen PB, Wu GY, Nairn AC, Greengard P. The Rho-Specific GEF Lfc Interacts with Neurabin and Spinophilin to Regulate Dendritic Spine Morphology. *Neuron* 47: 85–100, 2005.
425. Ryu J, Futai K, Feliu M, Weinberg R, Sheng M. Constitutively active Rap2 transgenic mice display fewer dendritic spines, reduced extracellular signal-regulated kinase signaling, enhanced long-term depression, and impaired spatial learning and fear extinction. *J Neurosci* 28: 8178–8188, 2008.
426. Ryu J, Liu L, Wong TP, Wu DC, Burette A, Weinberg R, Wang YT, Sheng M. A critical role for myosin IIb in dendritic spine morphology and synaptic function. *Neuron* 49: 175–182, 2006.
427. Rác B, Weinberg RJ. Microdomains in forebrain spines: an ultrastructural perspective. *Mol Neurobiol* 47: 77–89, 2013.
428. Sabatini BL, Oertner TG, Svoboda K. The life cycle of Ca²⁺ ions in dendritic spines. *Neuron* 33: 439–452, 2002.
429. Sacktor TC. PKMzeta, LTP maintenance, and the dynamic molecular biology of memory storage. *Prog Brain Res* 169: 27–40, 2008.
430. Saglietti L, Dequidt C, Kamieniarz K, Rousset MC, Valnegri P, Thoumine O, Beretta F, Fagni L, Choquet D, Sala C, Sheng M, Passafaro M. Extracellular interactions between GluR2 and N-cadherin in spine regulation. *Neuron* 54: 461–477, 2007.
431. Sakamoto H, Mezaki Y, Shikimi H, Ukena K, Tsutsui K. Dendritic growth and spine formation in response to estrogen in the developing Purkinje cell. *Endocrinology* 144: 4466–4477, 2003.
432. Sala C, Futai K, Yamamoto K, Worley PF, Hayashi Y, Sheng M. Inhibition of dendritic spine morphogenesis and synaptic transmission by activity-inducible protein Homer 1a. *J Neurosci* 23: 6327–6337, 2003.
433. Sala C, Piech V, Wilson NR, Passafaro M, Liu G, Sheng M. Regulation of dendritic spine morphology and synaptic function by Shank and Homer. *Neuron* 31: 115–130, 2001.
435. Sala C, Roussignol C, Meldolesi J, Fagni L. Key role of the postsynaptic density scaffold proteins shank and homer in the functional architecture of Ca²⁺ homeostasis at dendritic spines in hippocampal neurons. *J Neurosci* 25: 4587–4592, 2005.
436. Salih DA, Rashid AJ, Colas D, de la Torre-Ubieta L, Zhu RP, Morgan AA, Santo EE, Ucar D, Devarajan K, Cole CJ, Madison DV, Shamloo M, Butte AJ, Bonni A, Josselyn SA, Brunet A. FoxO6 regulates memory consolidation and synaptic function. *Genes Dev* 26: 2780–2801, 2012.
437. Salinas PC, Price SR. Cadherins and catenins in synapse development. *Curr Opin Neurobiol* 15: 73–80, 2005.
438. Sanchez AM, Flamini MI, Baldacci C, Goglia L, Genazzani AR, Simoncini T. Estrogen receptor-alpha promotes breast cancer cell motility and invasion via focal adhesion kinase and N-WASP. *Mol Endocrinol* 24: 2114–2125, 2010.
439. Sanchez AM, Flamini MI, Fu XD, Mannella P, Giretti MS, Goglia L, Genazzani AR, Simoncini T. Rapid signaling of estrogen to WAVE1 and moesin controls neuronal spine formation via the actin cytoskeleton. *Mol Endocrinol* 23: 1193–1202, 2009.
440. Sanders J, Cowansage K, Baumgärtel K, Mayford M. Elimination of dendritic spines with long-term memory is specific to active circuits. *J Neurosci* 32: 12570–12578, 2012.
441. Saneyoshi T, Wayman G, Fortin D, Davare M, Hoshi N, Nozaki N, Natsume T, Soderling TR. Activity-dependent synaptogenesis: regulation by a CaM-kinase kinase/CaM-kinase I/betaPIX signaling complex. *Neuron* 57: 94–107, 2008.
442. Sapoznik S, Ivenshitz M, Segal M. Age-dependent glutamate induction of synaptic plasticity in cultured hippocampal neurons. *Learn Mem* 13: 719–727, 2006.
443. Sawallisch C, Berhörster K, Disanza A, Mantoani S, Kintscher M, Stoenica L, Dityatev A, Sieber S, Kindler S, Morellini F, Schweizer M, Boeckers TM, Korte M, Scita G, Kreienkamp HJ. The insulin receptor substrate of 53 kDa (IRS53) limits hippocampal synaptic plasticity. *J Biol Chem* 284: 9225–9236, 2009.
444. Schmeisser MJ, Ey E, Wegener S, Bockmann J, Stempel AV, Kuebler A, Janssen AL, Udvardi PT, Shiban E, Spilker C, Balschun D, Skryabin BV, Dieck S, Smalla KH, Montag D, Leblond CS, Faure P, Torquet N, Le Sourd AM, Toro R, Grabrucker AM, Shiochet SA, Schmitz D, Kreutz MR, Bourgeron T, Gundelfinger ED, Boeckers TM. Autistic-like behaviours and hyperactivity in mice lacking ProSAP1/Shank2. *Nature* 486: 256–260, 2012.
445. Schratz GM, Tuebing F, Nigh EA, Kane CG, Sabatini ME, Kiebler M, Greenberg ME. A brain-specific microRNA regulates dendritic spine development. *Nature* 439: 283–289, 2006.
446. Schumacher J, Laje G, Abou Jamra R, Becker T, Mühleisen TW, Vasilescu C, Mattheisen M, Herms S, Hoffmann P, Hillmer AM, Georgi A, Herold C, Schulze TG, Thompson P, Rietschel M, McMahon FJ, Nöthen MM, Cichon S. The DISC locus and schizophrenia: evidence from an association study in a central European sample and from a meta-analysis across different European populations. *Hum Mol Genet* 18: 2719–2727, 2009.
447. Schwarz JM, Liang SL, Thompson SM, McCarthy MM. Estradiol induces hypothalamic dendritic spines by enhancing glutamate release: a mechanism for organizational sex differences. *Neuron* 58: 584–598, 2008.
448. Segal M, Vlachos A, Korkotian E. The spine apparatus, synaptopodin, and dendritic spine plasticity. *Neuroscientist* 16: 125–131, 2010.
449. Segev I, Friedman A, White EL, Gutnick MJ. Electrical consequences of spine dimensions in a model of a cortical spiny stellate cell completely reconstructed from serial thin sections. *J Comput Neurosci* 2: 117–130, 1995.
450. Sfakianos MK, Eisman A, Gourley SL, Bradley WD, Scheetz AJ, Settleman J, Taylor JR, Greer CA, Williamson A, Koleske AJ. Inhibition of Rho via Arg and p190RhoGAP in the postnatal mouse hippocampus regulates dendritic spine maturation, synapse and dendrite stability, and behavior. *J Neurosci* 27: 10982–10992, 2007.
451. Shamir A, Kwon OB, Karavanova I, Vullhorst D, Leiva-Salcedo E, Janssen MJ, Buonanno A. The importance of the NRG-1/ErbB4 pathway for synaptic plasticity and behaviors associated with psychiatric disorders. *J Neurosci* 32: 2988–2997, 2012.
452. Shankar GM, Bloodgood BL, Townsend M, Walsh DM, Selkoe DJ, Sabatini BL. Natural oligomers of the Alzheimer amyloid-beta protein induce reversible synapse loss by modulating an NMDA-type glutamate receptor-dependent signaling pathway. *J Neurosci* 27: 2866–2875, 2007.
453. Sharp AH, McPherson PS, Dawson TM, Aoki C, Campbell KP, Snyder SH. Differential immunohistochemical localization of inositol 1,4,5-trisphosphate- and ryanodine-sensitive Ca²⁺ release channels in rat brain. *J Neurosci* 13: 3051–3063, 1993.

454. Sheng M, Kim E. The Shank family of scaffold proteins. *J Cell Sci* 113: 1851–1856, 2000.
455. Sheng M, Sala C. PDZ domains and the organization of supramolecular complexes. *Annu Rev Neurosci* 24: 1–29, 2001.
456. Shepherd GM, Brayton RK, Miller JP, Segev I, Rinzel J, Rall W. Signal enhancement in distal cortical dendrites by means of interactions between active dendritic spines. *Proc Natl Acad Sci USA* 82: 2192–2195, 1985.
457. Sherren N, Pappas BA. Selective acetylcholine and dopamine lesions in neonatal rats produce distinct patterns of cortical dendritic atrophy in adulthood. *Neuroscience* 136: 445–456, 2005.
458. Shi SH, Hayashi Y, Petralia RS, Zaman SH, Wenthold RJ, Svoboda K, Malinow R. Rapid spine delivery and redistribution of AMPA receptors after synaptic NMDA receptor activation. *Science* 284: 1811–1816, 1999.
459. Shi Y, Pontrello CG, DeFea KA, Reichardt LF, Ethell IM. Focal adhesion kinase acts downstream of EphB receptors to maintain mature dendritic spines by regulating cofilin activity. *J Neurosci* 29: 8129–8142, 2009.
460. Shin E, Kashiwagi Y, Kuriu T, Iwasaki H, Tanaka T, Koizumi H, Gleeson JG, Okabe S. Doublecortin-like kinase enhances dendritic remodeling and negatively regulates synapse maturation. *Nat Commun* 4: 1440, 2013.
461. Shipman SL, Herring BE, Suh YH, Roche KW, Nicoll RA. Distance-dependent scaling of AMPARs is cell-autonomous and GluA2 dependent. *J Neurosci* 33: 13312–13319, 2013.
462. Shors TJ, Chua C, Falduto J. Sex differences and opposite effects of stress on dendritic spine density in the male versus female hippocampus. *J Neurosci* 21: 6292–6297, 2001.
463. Siegel G, Obernosterer G, Fiore R, Oehmen M, Bicker S, Christensen M, Khudayberdiev S, Leuschner PF, Busch CJ, Kane C, Hübel K, Dekker F, Hedberg C, Rengasajan B, Drepper C, Waldmann H, Kauppinen S, Greenberg ME, Draguhn A, Rehmsmeier M, Martinez J, Schratt GM. A functional screen implicates microRNA-138-dependent regulation of the palmitoylation enzyme APT1 in dendritic spine morphogenesis. *Nat Cell Biol* 11: 705–716, 2009.
464. Sloniewski S, Ethell IM. Looking forward to EphB signaling in synapses. *Semin Cell Dev Biol* 23: 75–82, 2012.
465. Smith MA, Ellis-Davies GC, Magee JC. Mechanism of the distance-dependent scaling of Schaffer collateral synapses in rat CA1 pyramidal neurons. *J Physiol* 548: 245–258, 2003.
466. Snyder EM, Nong Y, Almeida CG, Paul S, Moran T, Choi EY, Nairn AC, Salter MW, Lombroso PJ, Gouras GK, Greengard P. Regulation of NMDA receptor trafficking by amyloid-beta. *Nat Neurosci* 8: 1051–1058, 2005.
467. Soares C, Lee KF, Nassrallah W, Béique JC. Differential subcellular targeting of glutamate receptor subtypes during homeostatic synaptic plasticity. *J Neurosci* 33: 13547–13559, 2013.
468. Soderling SH, Guire ES, Kaech S, White J, Zhang F, Schutz K, Langeberg LK, Banker G, Raber J, Scott JD. A WAVE-1 and WRP signaling complex regulates spine density, synaptic plasticity, and memory. *J Neurosci* 27: 355–365, 2007.
469. Soderling SH, Guire ES, Kaech S, White J, Zhang F, Schutz K, Langeberg LK, Banker G, Raber J, Scott JD. A WAVE-1 and WRP signaling complex regulates spine density, synaptic plasticity, and memory. *J Neurosci* 27: 355–365, 2007.
470. Soderling TR. CaM-kinases: modulators of synaptic plasticity. *Curr Opin Neurobiol* 10: 375–380, 2000.
471. Solís O, Limón DI, Flores-Hernández J, Flores G. Alterations in dendritic morphology of the prefrontal cortical and striatum neurons in the unilateral 6-OHDA-rat model of Parkinson's disease. *Synapse* 61: 450–458, 2007.
472. Soltau M, Richter D, Kreienkamp HJ. The insulin receptor substrate IRSp53 links postsynaptic shank1 to the small G-protein cdc42. *Mol Cell Neurosci* 21: 575–583, 2002.
473. Sorra KE, Harris KM. Overview on the structure, composition, function, development, and plasticity of hippocampal dendritic spines. *Hippocampus* 10: 501–511, 2000.
474. Sorra KE, Harris KM. Stability in synapse number and size at 2 hr after long-term potentiation in hippocampal area CA1. *J Neurosci* 18: 658–671, 1998.
475. Spencer JL, Waters EM, Romeo RD, Wood GE, Milner TA, McEwen BS. Uncovering the mechanisms of estrogen effects on hippocampal function. *Front Neuroendocrinol* 29: 219–237, 2008.
476. Spilker C, Acuna Sanhuesa GA, Bockers TM, Kreutz MR, Gundelfinger ED. SPAR2, a novel SPAR-related protein with GAP activity for Rap1 and Rap2. *J Neurochem* 104: 187–201, 2008.
477. Srivastava DP, Jones KA, Woolfrey KM, Burgdorf J, Russell TA, Kalmbach A, Lee H, Yang C, Bradberry MM, Wokosin D, Moskal JR, Casanova MF, Waters J, Penzes P. Social, communication, and cortical structural impairments in Epac2-deficient mice. *J Neurosci* 32: 11864–11878, 2012.
478. Srivastava DP, Woolfrey KM, Liu F, Brandon NJ, Penzes P. Estrogen receptor β activity modulates synaptic signaling and structure. *J Neurosci* 30: 13454–13460, 2010.
479. St Clair D, Blackwood D, Muir W, Carothers A, Walker M, Spowart G, Gosden C, Evans HJ. Association within a family of a balanced autosomal translocation with major mental illness. *Lancet* 336: 13–16, 1990.
480. Stamatakou E, Marzo A, Gibb A, Salinas PC. Activity-dependent spine morphogenesis: a role for the actin-capping protein Eps8. *J Neurosci* 33: 2661–2670, 2013.
481. Stark KL, Xu B, Bagchi A, Lai WS, Liu H, Hsu R, Wan X, Pavlidis P, Mills AA, Karayiorgou M, Gogos JA. Altered brain microRNA biogenesis contributes to phenotypic deficits in a 22q11-deletion mouse model. *Nat Genet* 40: 751–760, 2008.
482. Steen RG, Mull C, McClure R, Hamer RM, Lieberman JA. Brain volume in first-episode schizophrenia: systematic review and meta-analysis of magnetic resonance imaging studies. *Br J Psychiatry* 188: 510–518, 2006.
483. Steiner P, Higley MJ, Xu W, Czervionke BL, Malenka RC, Sabatini BL. Destabilization of the postsynaptic density by PSD-95 serine 73 phosphorylation inhibits spine growth and synaptic plasticity. *Neuron* 60: 788–802, 2008.
484. Stewart MG, Medvedev NI, Popov VI, Schoepfer R, Davies HA, Murphy K, Dallérac GM, Kraev IV, Rodríguez JJ. Chemically induced long-term potentiation increases the number of perforated and complex postsynaptic densities but does not alter dendritic spine volume in CA1 of adult mouse hippocampal slices. *Eur J Neurosci* 21: 3368–3378, 2005.
485. Stroth O, Freichel M, Kretz O, Birnbaumer L, Hartmann J, Egger V. NMDA receptor-dependent synaptic activation of TRPC channels in olfactory bulb granule cells. *J Neurosci* 32: 5737–5746, 2012.
486. Stutzmann GE, Mattson MP. Endoplasmic reticulum Ca^{2+} handling in excitable cells in health and disease. *Pharmacol Rev* 63: 700–727, 2011.
487. Südhof TC. Neuroligins and neuroligins link synaptic function to cognitive disease. *Nature* 455: 903–911, 2008.
488. Sung JY, Engmann O, Teylan MA, Nairn AC, Greengard P, Kim Y. WAVE1 controls neuronal activity-induced mitochondrial distribution in dendritic spines. *Proc Natl Acad Sci USA* 105: 3112–3116, 2008.
489. Suzuki K, Hayashi Y, Nakahara S, Kumazaki H, Prox J, Horiuchi K, Zeng M, Tanimura S, Nishiyama Y, Osawa S, Sehara-Fujisawa A, Saftig P, Yokoshima S, Fukuyama T, Matsuki N, Koyama R, Tomita T, Iwatsubo T. Activity-dependent proteolytic cleavage of neuroligin-1. *Neuron* 76: 410–422, 2012.
490. Svoboda K, Tank DW, Denk W. Direct measurement of coupling between dendritic spines and shafts. *Science* 272: 716–719, 1996.
491. Sweet RA, Henteloff RA, Zhang W, Sampson AR, Lewis DA. Reduced dendritic spine density in auditory cortex of subjects with schizophrenia. *Neuropsychopharmacology* 34: 374–389, 2009.
492. Szatmari EM, Oliveira AF, Sumner EJ, Yasuda R. Centaurin- α 1-Ras-Elk-1 signaling at mitochondria mediates β -amyloid-induced synaptic dysfunction. *J Neurosci* 33: 5367–5374, 2013.
493. Tabuchi K, Blundell J, Ethernell MR, Hammer RE, Liu X, Powell CM, Südhof TC. A neuroligin-3 mutation implicated in autism increases inhibitory synaptic transmission in mice. *Science* 318: 71–76, 2007.
494. Takahashi H, Sekino Y, Tanaka S, Mizui T, Kishi S, Shirao T. Drebrin-dependent actin clustering in dendritic filopodia governs synaptic targeting of postsynaptic density-95 and dendritic spine morphogenesis. *J Neurosci* 23: 6586–6595, 2003.

495. Takahashi N, Kitamura K, Matsuo N, Mayford M, Kano M, Matsuki N, Ikegaya Y. Locally synchronized synaptic inputs. *Science* 335: 353–356, 2012.
496. Tan AM, Chang YW, Zhao P, Hains BC, Waxman SG. Rac1-regulated dendritic spine remodeling contributes to neuropathic pain after peripheral nerve injury. *Exp Neurol* 232: 222–233, 2011.
497. Tan HY, Callicott JH, Weinberger DR. Dysfunctional and compensatory prefrontal cortical systems, genes and the pathogenesis of schizophrenia. *Cereb Cortex* 17 Suppl 1: i171–181, 2007.
498. Tan XJ, Dai YB, Wu WF, Kim HJ, Barros RP, Richardson TI, Yaden BC, Warner M, McKinzie DL, Krishnan V, Gustafsson J. Reduction of dendritic spines and elevation of GABAergic signaling in the brains of mice treated with an estrogen receptor β ligand. *Proc Natl Acad Sci USA* 109: 1708–1712, 2012.
499. Tanaka J, Horiike Y, Matsuzaki M, Miyazaki T, Ellis-Davies GC, Kasai H. Protein synthesis and neurotrophin-dependent structural plasticity of single dendritic spines. *Science* 319: 1683–1687, 2008.
500. Tanokashira D, Morita T, Hayashi K, Mayanagi T, Fukumoto K, Kubota Y, Yamashita T, Sobue K. Glucocorticoid suppresses dendritic spine development mediated by down-regulation of caldesmon expression. *J Neurosci* 32: 14583–14591, 2012.
501. Tashiro A, Minden A, Yuste R. Regulation of dendritic spine morphology by the rho family of small GTPases: antagonistic roles of Rac and Rho. *Cereb Cortex* 10: 927–938, 2000.
502. Tashiro A, Yuste R. Regulation of dendritic spine motility and stability by Rac1 and Rho kinase: evidence for two forms of spine motility. *Mol Cell Neurosci* 26: 429–440, 2004.
503. Tavaoie SF, Alvarez VA, Ridenour DA, Kwiatkowski DJ, Sabatini BL. Regulation of neuronal morphology and function by the tumor suppressors Tsc1 and Tsc2. *Nat Neurosci* 8: 1727–1734, 2005.
504. Thomas A, Kim DS, Fields RL, Chin H, Gainer H. Quantitative analysis of gene expression in organotypic slice-explant cultures by particle-mediated gene transfer. *J Neurosci Methods* 84: 181–191, 1998.
505. Thomas S, Ritter B, Verbich D, Sanson C, Bourbonniere L, McKinney RA, McPherson PS. Intersectin regulates dendritic spine development and somatodendritic endocytosis but not synaptic vesicle recycling in hippocampal neurons. *J Biol Chem* 284: 12410–12419, 2009.
506. Thévenot E, Moreau AW, Rousseau V, Combeau G, Domenichini F, Jacquet C, Goupille O, Amar M, Kreis P, Fossier P, Barnier JV. p21-Activated kinase 3 (PAK3) protein regulates synaptic transmission through its interaction with the Nck2/Grb4 protein adaptor. *J Biol Chem* 286: 40044–40059, 2011.
507. Tolias KF, Bikoff JB, Burette A, Paradis S, Harrar D, Tavaoie S, Weinberg RJ, Greenberg ME. The Rac1-GEF Tiam1 couples the NMDA receptor to the activity-dependent development of dendritic arbors and spines. *Neuron* 45: 525–538, 2005.
508. Tolias KF, Bikoff JB, Kane CG, Tolias CS, Hu L, Greenberg ME. The Rac1 guanine nucleotide exchange factor Tiam1 mediates EphB receptor-dependent dendritic spine development. *Proc Natl Acad Sci USA* 104: 7265–7270, 2007.
509. Toro R, Konyukh M, Delorme R, Leblond C, Chaste P, Fauchereau F, Coleman M, Leboyer M, Gillberg C, Bourgeron T. Key role for gene dosage and synaptic homeostasis in autism spectrum disorders. *Trends Genet* 26: 363–372, 2010.
510. Tortosa E, Montenegro-Venegas C, Benoist M, Härtel S, González-Billault C, Esteban JA, Avila J. Microtubule-associated protein 1B (MAP1B) is required for dendritic spine development and synaptic maturation. *J Biol Chem* 286: 40638–40648, 2011.
511. Trachtenberg JT, Chen BE, Knott GW, Feng G, Sanes JR, Welker E, Svoboda K. Long-term in vivo imaging of experience-dependent synaptic plasticity in adult cortex. *Nature* 420: 788–794, 2002.
512. Tran TS, Rubio ME, Clem RL, Johnson D, Case L, Tessier-Lavigne M, Huganir RL, Ginty DD, Kolodkin AL. Secreted semaphorins control spine distribution and morphogenesis in the postnatal CNS. *Nature* 462: 1065–1069, 2009.
513. Tropea D, Giacometti E, Wilson NR, Beard C, McCurry C, Fu DD, Flannery R, Jaenisch R, Sur M. Partial reversal of Rett Syndrome-like symptoms in MeCP2 mutant mice. *Proc Natl Acad Sci USA* 106: 2029–2034, 2009.
514. Tsai J, Grutzendler J, Duff K, Gan WB. Fibrillar amyloid deposition leads to local synaptic abnormalities and breakage of neuronal branches. *Nat Neurosci* 7: 1181–1183, 2004.
515. Tsai SY, Hayashi T, Harvey BK, Wang Y, Wu WW, Shen RF, Zhang Y, Becker KG, Hoffer BJ, Su TP. Sigma-1 receptors regulate hippocampal dendritic spine formation via a free radical-sensitive mechanism involving Rac1 xGTP pathway. *Proc Natl Acad Sci USA* 106: 22468–22473, 2009.
516. Tseng CY, Firestein BL. The role of PSD-95 and cypin in morphological changes in dendrites following sublethal NMDA exposure. *J Neurosci* 31: 15468–15480, 2011.
517. Tuckwell HC. Quantitative aspects of L-type Ca^{2+} currents. *Prog Neurobiol* 96: 1–31, 2012.
518. Turrigiano G. Homeostatic signaling: the positive side of negative feedback. *Curr Opin Neurobiol* 17: 318–324, 2007.
519. Tyler WJ, Pozzo-Miller L. Miniature synaptic transmission and BDNF modulate dendritic spine growth and form in rat CA1 neurones. *J Physiol* 553: 497–509, 2003.
520. Ultanir SK, Hertz NT, Li G, Ge WP, Burlingame AL, Pleasure SJ, Shokat KM, Jan LY, Jan YN. Chemical genetic identification of NDR1/2 kinase substrates AAK1 and Rabin8 Uncovers their roles in dendrite arborization and spine development. *Neuron* 73: 1127–1142, 2012.
521. Ultanir SK, Kim JE, Hall BJ, Deerinck T, Ellisman M, Ghosh A. Regulation of spine morphology and spine density by NMDA receptor signaling in vivo. *Proc Natl Acad Sci USA* 104: 19553–19558, 2007.
522. Vadodaria KC, Brakebusch C, Suter U, Jessberger S. Stage-specific functions of the small Rho GTPases Cdc42 and Rac1 for adult hippocampal neurogenesis. *J Neurosci* 33: 1179–1189, 2013.
523. Valnegri P, Montrasio C, Brambilla D, Ko J, Passafium M, Sala C. The X-linked intellectual disability protein ILIRAPL1 regulates excitatory synapse formation by binding PTP delta and RhoGAP2. *Hum Mol Genet* 20: 4797–4809, 2011.
524. Valnegri P, Montrasio C, Brambilla D, Ko J, Passafium M, Sala C. The X-linked intellectual disability protein ILIRAPL1 regulates excitatory synapse formation by binding PTP δ and RhoGAP2. *Hum Mol Genet* 20: 4797–4809, 2011.
525. Valnegri P, Sala C, Passafium M. Synaptic dysfunction and intellectual disability. *Adv Exp Med Biol* 970: 433–449, 2012.
526. Van de Ven TJ, VanDongen HM, VanDongen AM. The nonkinase phorbol ester receptor alpha 1-chimerin binds the NMDA receptor NR2A subunit and regulates dendritic spine density. *J Neurosci* 25: 9488–9496, 2005.
527. Vastagh C, Gardoni F, Bagetta V, Stanic J, Zianni E, Giampà C, Picconi B, Calabresi P, Di Luca M. N-methyl-D-aspartate (NMDA) receptor composition modulates dendritic spine morphology in striatal medium spiny neurons. *J Biol Chem* 287: 18103–18114, 2012.
528. Vazquez LE, Chen HJ, Sokolova I, Knuesel I, Kennedy MB. SynGAP regulates spine formation. *J Neurosci* 24: 8862–8872, 2004.
529. Verkhratsky A. Physiology and pathophysiology of the calcium store in the endoplasmic reticulum of neurons. *Physiol Rev* 85: 201–279, 2005.
530. Verpelli C, Dvoretzskova E, Vicidomini C, Rossi F, Chiappalone M, Schoen M, Di Stefano B, Mantegazza R, Broccoli V, Boeckers TM, Dityatev A, Sala C. Importance of shank3 in regulating metabotropic glutamate receptor 5 (mGluR5) expression and signaling at synapses. *J Biol Chem* 286: 34839–34850, 2011.
531. Verpelli C, Piccoli G, Zanchi A, Gardoni F, Huang K, Brambilla D, Di Luca M, Battaglioli E, Sala C. Synaptic activity controls dendritic spine morphology by modulating eEF2-dependent BDNF synthesis. *J Neurosci* 30: 5830–5842, 2010.
532. Verpelli C, Sala C. Molecular and synaptic defects in intellectual disability syndromes. *Curr Opin Neurobiol* 22: 530–536, 2012.
533. Vickers CA, Stephens B, Bowen J, Arbuthnott GW, Grant SG, Ingham CA. Neurone specific regulation of dendritic spines in vivo by post synaptic density 95 protein (PSD-95). *Brain Res* 1090: 89–98, 2006.
534. Vlachos A, Korkotian E, Schonfeld E, Copanaki E, Deller T, Segal M. Synaptopodin regulates plasticity of dendritic spines in hippocampal neurons. *J Neurosci* 29: 1017–1033, 2009.
535. Von Bohlen und Halbach O, Minichiello L, Unsicker K. TrkB but not trkC receptors are necessary for postnatal maintenance of hippocampal spines. *Neurobiol Aging* 29: 1247–1255, 2008.

536. Von der Ohe CG, Darian-Smith C, Garner CC, Heller HC. Ubiquitous and temperature-dependent neural plasticity in hibernators. *J Neurosci* 26: 10590–10598, 2006.
537. Walsh DM, Selkoe DJ. Deciphering the molecular basis of memory failure in Alzheimer's disease. *Neuron* 44: 181–193, 2004.
538. Walsh T, McClellan JM, McCarthy SE, Addington AM, Pierce SB, Cooper GM, Nord AS, Kusenda M, Malhotra D, Bhandari A, Stray SM, Rippey CF, Rocanova P, Makarov V, Lakshmi B, Findling RL, Sikich N, Stromberg T, Merriman B, Gogtay N, Butler P, Eckstrand K, Noory L, Gochman P, Long R, Chen Z, Davis S, Baker C, Eichler EE, Meltzer PS, Nelson SF, Singleton AB, Lee MK, Rapoport JL, King MC, Sebat J. Rare structural variants disrupt multiple genes in neurodevelopmental pathways in schizophrenia. *Science* 320: 539–543, 2008.
539. Waltereit R, Leimer U, von Bohlen Und Halbach O., Panke J., Höltter SM, Garrett L, Wittig K, Schneider M, Schmitt C, Calzada-Wack J, Neff F, Becker L, Pohn C, Kutschera-Jawy S, Endris V, Bacon C, Fuchs H, Gailus-Durner V, Berger S, Schönicg K, Adamski J, Klopstock T, Esposito I, Wurst W, de Angelis MH, Rappold G, Wieland T, Bartsch D. Srgap3^{-/-} mice present a neurodevelopmental disorder with schizophrenia-related intermediate phenotypes. *FASEB J* 26: 4418–4428, 2012.
540. Wang HD, Deutch AY. Dopamine depletion of the prefrontal cortex induces dendritic spine loss: reversal by atypical antipsychotic drug treatment. *Neuropsychopharmacology* 33: 1276–1286, 2008.
541. Wang X, McCoy PA, Rodriguiz RM, Pan Y, Je HS, Roberts AC, Kim CJ, Berrios J, Colvin JS, Bousquet-Moore D, Lorenzo I, Wu G, Weinberg RJ, Ehlers MD, Philpot BD, Beaudet AL, Wetsel WC, Jiang YH. Synaptic dysfunction and abnormal behaviors in mice lacking major isoforms of Shank3. *Hum Mol Genet* 20: 3093–3108, 2011.
542. Wayman GA, Lee YS, Tokumitsu H, Silva AJ, Silva A, Soderling TR. Calmodulin-kinases: modulators of neuronal development and plasticity. *Neuron* 59: 914–931, 2008.
543. Webb DJ, Zhang H, Majumdar D, Horwitz AF. alpha5 integrin signaling regulates the formation of spines and synapses in hippocampal neurons. *J Biol Chem* 282: 6929–6935, 2007.
544. Wegner AM, Nebhan CA, Hu L, Majumdar D, Meier KM, Weaver AM, Webb DJ. N-wasp and the arp2/3 complex are critical regulators of actin in the development of dendritic spines and synapses. *J Biol Chem* 283: 15912–15920, 2008.
545. Weltzin MM, Zhao HW, Drew KL, Bucci DJ. Arousal from hibernation alters contextual learning and memory. *Behav Brain Res* 167: 128–133, 2006.
546. Wigström H, Gustafsson B. Postsynaptic control of hippocampal long-term potentiation. *J Physiol* 81: 228–236, 1986.
547. Willig KI, Kellner RR, Medda R, Hein B, Jakobs S, Hell SW. Nanoscale resolution in GFP-based microscopy. *Nat Methods* 3: 721–723, 2006.
548. Wilson HL, Wong AC, Shaw SR, Tse WY, Stapleton GA, Phelan MC, Hu S, Marshall J, McDermid HE. Molecular characterisation of the 22q13 deletion syndrome supports the role of haploinsufficiency of SHANK3/PROSAP2 in the major neurological symptoms. *J Med Genet* 40: 575–584, 2003.
549. Won H, Lee HR, Gee HY, Mah W, Kim JI, Lee J, Ha S, Chung C, Jung ES, Cho YS, Park SG, Lee JS, Lee K, Kim D, Bae YC, Kaang BK, Lee MG, Kim E. Autistic-like social behaviour in Shank2-mutant mice improved by restoring NMDA receptor function. *Nature* 486: 261–265, 2012.
550. Woolfrey KM, Srivastava DP, Photowala H, Yamashita M, Barbolina MV, Cahill ME, Xie Z, Jones KA, Quilliam LA, Prakriya M, Penzes P. Epac2 induces synapse remodeling and depression and its disease-associated forms alter spines. *Nat Neurosci* 12: 1275–1284, 2009.
551. Woolley CS, Gould E, Frankfurt M, McEwen BS. Naturally occurring fluctuation in dendritic spine density on adult hippocampal pyramidal neurons. *J Neurosci* 10: 4035–4039, 1990.
552. Woolley CS, McEwen BS. Roles of estradiol and progesterone in regulation of hippocampal dendritic spine density during the estrous cycle in the rat. *J Comp Neurol* 336: 293–306, 1993.
553. Wu GY, Deisseroth K, Tsien RW. Spaced stimuli stabilize MAPK pathway activation and its effects on dendritic morphology. *Nat Neurosci* 4: 151–158, 2001.
554. Wu HY, Hudry E, Hashimoto T, Kuchibhotla K, Rozkalne A, Fan Z, Spires-Jones T, Xie H, Arbel-Ornath M, Grosskreutz CL, Bacskai BJ, Hyman BT. Amyloid beta induces the morphological neurodegenerative triad of spine loss, dendritic simplification, and neuritic dystrophies through calcineurin activation. *J Neurosci* 30: 2636–2649, 2010.
555. Xiao B, Tu JC, Worley PF. Homer: a link between neural activity and glutamate receptor function. *Curr Opin Neurobiol* 10: 370–374, 2000.
556. Xie Z, Cahill ME, Penzes P. Kalirin loss results in cortical morphological alterations. *Mol Cell Neurosci* 43: 81–89, 2010.
557. Xie Z, Hugarir RL, Penzes P. Activity-dependent dendritic spine structural plasticity is regulated by small GTPase Rap1 and its target AF-6. *Neuron* 48: 605–618, 2005.
558. Xie Z, Srivastava DP, Photowala H, Kai L, Cahill ME, Woolfrey KM, Shum CY, Surmeier DJ, Penzes P. Kalirin-7 controls activity-dependent structural and functional plasticity of dendritic spines. *Neuron* 56: 640–656, 2007.
559. Xu B, Hsu PK, Stark KL, Karayiorgou M, Gogos JA. Derepression of a neuronal inhibitor due to miRNA dysregulation in a schizophrenia-related microdeletion. *Cell* 152: 262–275, 2013.
560. Xu HT, Pan F, Yang G, Gan WB. Choice of cranial window type for in vivo imaging affects dendritic spine turnover in the cortex. *Nat Neurosci* 10: 549–551, 2007.
561. Xu T, Yu X, Perlik AJ, Tobin WF, Zweig JA, Tennant K, Jones T, Zuo Y. Rapid formation and selective stabilization of synapses for enduring motor memories. *Nature* 462: 915–919, 2009.
562. Yamagata Y, Kobayashi S, Umeda T, Inoue A, Sakagami H, Fukaya M, Watanabe M, Hatanaka N, Totsuka M, Yagi T, Obata K, Imoto K, Yanagawa Y, Manabe T, Okabe S. Kinase-dead knock-in mouse reveals an essential role of kinase activity of Ca²⁺/calmodulin-dependent protein kinase IIalpha in dendritic spine enlargement, long-term potentiation, and learning. *J Neurosci* 29: 7607–7618, 2009.
563. Yang G, Pan F, Gan WB. Stably maintained dendritic spines are associated with lifelong memories. *Nature* 462: 920–924, 2009.
564. Yang Y, Paspalas CD, Jin LE, Picciotto MR, Arnsten AF, Wang M. Nicotinic $\alpha 7$ receptors enhance NMDA cognitive circuits in dorsolateral prefrontal cortex. *Proc Natl Acad Sci USA* 110: 12078–12083, 2013.
565. Yang Y, Shu X, Liu D, Shang Y, Wu Y, Pei L, Xu X, Tian Q, Zhang J, Qian K, Wang YX, Petralia RS, Tu W, Zhu LQ, Wang JZ, Lu Y. EPAC null mutation impairs learning and social interactions via aberrant regulation of miR-124 and Zif268 translation. *Neuron* 73: 774–788, 2012.
566. Yang Y, Wang XB, Frerking M, Zhou Q. Spine expansion and stabilization associated with long-term potentiation. *J Neurosci* 28: 5740–5751, 2008.
567. Yashiro K, Riday TT, Condon KH, Roberts AC, Bernardo DR, Prakash R, Weinberg RJ, Ehlers MD, Philpot BD. Ube3a is required for experience-dependent maturation of the neocortex. *Nat Neurosci* 12: 777–783, 2009.
568. Yasuda S, Tanaka H, Sugiura H, Okamura K, Sakaguchi T, Tran U, Takemiya T, Mizoguchi A, Yagita Y, Sakurai T, De Robertis EM, Yamagata K. Activity-induced protocadherin arcadin regulates dendritic spine number by triggering N-cadherin endocytosis via TAO2beta and p38 MAP kinases. *Neuron* 56: 456–471, 2007.
569. Yoshida T, McCarley RW, Nakamura M, Lee K, Koo MS, Bouix S, Salisbury DF, Morra L, Shenton ME, Niznikiewicz MA. A prospective longitudinal volumetric MRI study of superior temporal gyrus gray matter and amygdala-hippocampal complex in chronic schizophrenia. *Schizophr Res* 113: 84–94, 2009.
570. Youn H, Jeoung M, Koo Y, Ji H, Markesbery WR, Ji I, Ji TH. Kalirin is under-expressed in Alzheimer's disease hippocampus. *J Alzheimers Dis* 11: 385–397, 2007.
571. Youn H, Ji I, Ji HP, Markesbery WR, Ji TH. Under-expression of Kalirin-7 increases iNOS activity in cultured cells and correlates to elevated iNOS activity in Alzheimer's disease hippocampus. *J Alzheimers Dis* 12: 271–281, 2007.
572. Yu W, Lu B. Synapses and dendritic spines as pathogenic targets in Alzheimer's disease. *Neural Plast* 2012: 247150, 2012.
573. Yu W, Polepalli J, Wagh D, Rajadas J, Malenka R, Lu B. A critical role for the PAR-1/MARK-tau axis in mediating the toxic effects of A β on synapses and dendritic spines. *Hum Mol Genet* 21: 1384–1390, 2012.
574. Yu W, Sun Y, Guo S, Lu B. The PINK1/Parkin pathway regulates mitochondrial dynamics and function in mammalian hippocampal and dopaminergic neurons. *Hum Mol Genet* 20: 3227–3240, 2011.

575. Yuan JP, Kiselyov K, Shin DM, Chen J, Shcheynikov N, Kang SH, Dehoff MH, Schwarz MK, Seeburg PH, Muallem S, Worley PF. Homer binds TRPC family channels and is required for gating of TRPC1 by IP₃ receptors. *Cell* 114: 777–789, 2003.
576. Zagrebelsky M, Holz A, Dechant G, Barde YA, Bonhoeffer T, Korte M. The p75 neurotrophin receptor negatively modulates dendrite complexity and spine density in hippocampal neurons. *J Neurosci* 25: 9989–9999, 2005.
577. Zalfa F, Eleuteri B, Dickson KS, Mercaldo V, De Rubeis S, di Penta A, Tabolacci E, Chiurazzi P, Neri G, Grant SG, Bagni C. A new function for the fragile X mental retardation protein in regulation of PSD-95 mRNA stability. *Nat Neurosci* 10: 578–587, 2007.
578. Zalfa F, Giorgi M, Primerano B, Moro A, Di Penta A, Reis S, Oostra B, Bagni C. The fragile X syndrome protein FMRP associates with BCL RNA and regulates the translation of specific mRNAs at synapses. *Cell* 112: 317–327, 2003.
579. Zemni R, Bienvu T, Vinet MC, Sefiani A, Carrié A, Billuart P, McDonnell N, Couvert P, Francis F, Chafey P, Fauchereau F, Friocourt G, des Portes V, Cardona A, Frints S, Meindl A, Brandau O, Ronce N, Moraine C, van Bokhoven H, Ropers HH, Sudbrak R, Kahn A, Fryns JP, Beldjord C, Chelly J. A new gene involved in X-linked mental retardation identified by analysis of an X;2 balanced translocation. *Nat Genet* 24: 167–170, 2000.
580. Zhang H, Macara IG. The PAR-6 polarity protein regulates dendritic spine morphogenesis through p190 RhoGAP and the Rho GTPase. *Dev Cell* 14: 216–226, 2008.
581. Zhang H, Macara IG. The polarity protein PAR-3 and TIAM1 cooperate in dendritic spine morphogenesis. *Nat Cell Biol* 8: 227–237, 2006.
582. Zhang H, Webb DJ, Asmussen H, Niu S, Horwitz AF. A GIT1/PIX/Rac/PAK signaling module regulates spine morphogenesis and synapse formation through MLC. *J Neurosci* 25: 3379–3388, 2005.
583. Zhang XL, Pöschel B, Faul C, Upreti C, Stanton PK, Mundel P. Essential role for synaptopodin in dendritic spine plasticity of the developing hippocampus. *J Neurosci* 33: 12510–12518, 2013.
584. Zhao L, Ma QL, Calon F, Harris-White ME, Yang F, Lim GP, Morihara T, Ubeda OJ, Ambegaokar S, Hansen JE, Weisbart RH, Teter B, Frautschy SA, Cole GM. Role of p21-activated kinase pathway defects in the cognitive deficits of Alzheimer disease. *Nat Neurosci* 9: 234–242, 2006.
585. Zhou J, Du W, Zhou K, Tai Y, Yao H, Jia Y, Ding Y, Wang Y. Critical role of TRPC6 channels in the formation of excitatory synapses. *Nat Neurosci* 11: 741–743, 2008.
586. Zhou L, Lehan N, Wehrenberg U, Disteldorf E, von Lossow R, Mares U, Jarry H, Rune GM. Neuroprotection by estradiol: a role of aromatase against spine synapse loss after blockade of GABA(A) receptors. *Exp Neurol* 203: 72–81, 2007.
587. Zhou L, Martinez SJ, Haber M, Jones EV, Bouvier D, Doucet G, Corera AT, Fon EA, Zisch AH, Murai KK. EphA4 signaling regulates phospholipase Cγ1 activation, cofilin membrane association, and dendritic spine morphology. *J Neurosci* 27: 5127–5138, 2007.
588. Zhou Q, Homma KJ, Poo MM. Shrinkage of dendritic spines associated with long-term depression of hippocampal synapses. *Neuron* 44: 749–757, 2004.
589. Zhu Y, Pak D, Qin Y, McCormack SG, Kim MJ, Baumgart JP, Velamoor V, Auberson YP, Osten P, van Aelst L, Sheng M, Zhu JJ. Rap2-JNK removes synaptic AMPA receptors during depotentiation. *Neuron* 46: 905–916, 2005.
590. Zito K, Knott G, Shepherd GM, Shenolikar S, Svoboda K. Induction of spine growth and synapse formation by regulation of the spine actin cytoskeleton. *Neuron* 44: 321–334, 2004.
591. Zito K, Scheuss V, Knott G, Hill T, Svoboda K. Rapid functional maturation of nascent dendritic spines. *Neuron* 61: 247–258, 2009.
592. Zuo Y, Yang G, Kwon E, Gan WB. Long-term sensory deprivation prevents dendritic spine loss in primary somatosensory cortex. *Nature* 436: 261–265, 2005.

Dendritic Spines: The Locus of Structural and Functional Plasticity

Carlo Sala and Menahem Segal

Physiol Rev 94:141-188, 2014. doi:10.1152/physrev.00012.2013

You might find this additional info useful...

This article cites 592 articles, 260 of which can be accessed free at:

</content/94/1/141.full.html#ref-list-1>

This article has been cited by 2 other HighWire hosted articles

Learning and behavioral deficits associated with the absence of the fragile X mental retardation protein: what a fly and mouse model can teach us

Ana Rita Santos, Alexandros K. Kanellopoulos and Claudia Bagni

Learn. Mem., October, 2014; 21 (10): 543-555.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Synaptopodin Regulates Spine Plasticity: Mediation by Calcium Stores

Eduard Korkotian, Michael Frotscher and Menahem Segal

J. Neurosci., August 27, 2014; 34 (35): 11641-11651.

[\[Abstract\]](#) [\[Full Text\]](#) [\[PDF\]](#)

Updated information and services including high resolution figures, can be found at:

</content/94/1/141.full.html>

Additional material and information about *Physiological Reviews* can be found at:

<http://www.the-aps.org/publications/prv>

This information is current as of January 24, 2015.

Physiological Reviews provides state of the art coverage of timely issues in the physiological and biomedical sciences. It is published quarterly in January, April, July, and October by the American Physiological Society, 9650 Rockville Pike, Bethesda MD 20814-3991. Copyright © 2014 by the American Physiological Society. ISSN: 0031-9333, ESSN: 1522-1210. Visit our website at <http://www.the-aps.org/>.