

# RECURSIVE GEOMETRY AND THE MASS HIERARCHY OF ELEMENTARY LEPTONS AND BARYONS

## ABSTRACT

This paper presents a parameter-free, first-principles derivation of the mass hierarchy of elementary leptons and baryons, alongside the anomalous magnetic moments of the electron, muon, and proton. Operating within a five-dimensional, discrete, tension-bearing information substrate, the Compton–Schwarzschild crossover scale establishes the Planck-Lock condition, allowing standard SI physical quantities to be mapped onto unitless substrate ratios where the matter-formation yield threshold is normalised to unity ( $\eta = 1$ ). The proton-to-electron mass ratio is derived strictly from the geometric scaling of the nested two-layer proton recursion pattern deforming relative to the single-loop electron, yielding  $M_p / m_e = 1836.1527$ . The muon mass is resolved as the first harmonic overtone of the recursive proton configuration through the primary lattice spacing, giving  $M_\mu / m_e = 206.7702$ . The tau mass emerges as the second-order harmonic overtone governed by the recursive overtone squaring rule, yielding  $m_\tau / m_e = 3477.286$ . Furthermore, we establish the topological partition between singlet wave loops (leptons) and triplet spatial frames (baryons), suggesting that the three active, intersecting tension junctions of the proton core independently experience the substrate update delay, paying a cumulative environmental coordinate tax of exactly  $3H$  that shifts the pure geometric proton anomaly of 1.793072 to its terrestrial measured value of 1.792843. These results show encouraging agreement with experimental CODATA values, and suggest physical constants may be geometric consequences of a self-updating discrete medium.

## INTRODUCTION

Modern high-energy physics relies on the Higgs mechanism and arbitrary Yukawa coupling constants to define the masses of elementary leptons and baryons. These parameters are not derived from first principles; instead, they are manually fitted to match experimental data. This lack of predictive power represents a major limitation in our understanding of the particle spectrum and the origin of mass.

This paper explores this challenge by deriving the mass hierarchy of elementary particles from the geometric projection limits of a discrete, five-dimensional, tension-bearing information substrate. The scope is limited to the mass relations of the electron, proton, muon, and tau, as well as the anomalous magnetic moments of the electron, muon, and proton. Readers unfamiliar with the broader five-dimensional framework will find the core

assumptions and structural constants set out below; a full development of the model is left to the main foundational paper.

By treating the core structural constants of the substrate as starting reference points, this paper suggests particle masses and magnetic moments may not be independent physical constants, but rather the geometric and recursive consequences of a discrete medium deforming under localised tension gradients.

## THE PLANCK-LOCK CROSSOVER AND DIMENSIONLESS UNITS

The mathematical basis for working in a unitless framework rests at the Planck scale — the point where the Compton wavelength ( $\lambda_c = h / (m c)$ ) and the Schwarzschild radius ( $r_s = 2 G m / c^2$ ) coincide. At this scale, substrate geometry and SI measurements align numerically, allowing the Planck mass ( $m_s = 21.76437021 \mu g$ ) to serve as the dimensionless reference threshold  $m = 1$  in lattice units.

Once this scale is set by the fundamental tension-yield ratio ( $\dagger = 6.2963456314$ ), physical constants emerge as pure ratios of the discrete substrate. Standard SI units may be observer-dependent projection artifacts rather than fundamental properties of nature.

The physical parameters used in these derivations are defined as follows:

1. The Prime Seed ( $\dagger$ ):  $\dagger = 6.2963456314$ . The fundamental tension-to-yield ratio.
2. The Lattice Spacing ( $\lambda$ ):  $\lambda = 8.885765876$ . The characteristic separation between adjacent nodes in the five-dimensional hyper-lattice.
3. The EM-Sector Projection Factor ( $\Delta_s$ ):  $\Delta_s = \dagger / \pi^2 = 0.637851$ . Converts five-dimensional density magnitudes into four-dimensional electromagnetic observables.
4. The EM Brook Constant ( $\beta_s$ ):  $\beta_s = m_s \beta = 1.2810675$ . Describes how tension spreads and saturates across electromagnetic channels.
5. The Mesh Factor ( $\S$ ):  $\S = 0.016244$ . Relates discrete node positions to continuous coordinates.
6. The Hysteresis Constant ( $\mathbb{H}$ ):  $\mathbb{H} = 1 / (2\pi \dagger) = 0.000076456$ . The effective delay at coordinate boundaries.
7. The Junction Gap ( $\mathbb{J}$ ):  $\mathbb{J} = 0.0019796024$ . The structural offset between cyclic tension and the yield limit.

8. The Coordinate Potential ( $\Phi$ ):  $\Phi = 3.62312$ . The effective adjustment between discrete lattice structure and Euclidean coordinates.

#### FIRST-PRINCIPLES DERIVATION OF THE PROTON-TO-ELECTRON MASS RATIO

The proton-to-electron mass ratio ( $M_p / m_e$ ) is derived from the geometric scaling of the nested two-layer proton recursion pattern relative to the single-loop electron.

The structural mass-energy coefficient of the proton core ( $M_{p\_coeff}$ ) is a function of the background diagonal tension of the unit cell ( $\sqrt{\dagger}$ ) projected across the coordinate plane ( $\pi^2$ ), adjusted for boundary losses ( $\Sigma J$ ) across active projection paths ( $2N = 10$ ) and the spatial mesh factor ( $1 - \S$ ):

$$M_{p\_coeff} = \pi^2 \sqrt{\dagger} [ 1 - (\Sigma J / 2N) (1 - \S) ]$$

Where the total boundary potential per loop ( $\Sigma J$ ) combines the Junction Gap and Hysteresis:

$$\Sigma J = J + H = 0.0019796024 + 0.000076456 = 0.002056056$$

Evaluating:

$$M_{p\_coeff} = 3.14159265^2 \times \sqrt{6.2963456314} \times [ 1 - (0.002056056 / 10) \times (1 - 0.016244) ] = 24.76031$$

The effective spatial projection scale ( $X_{eff}$ ) is governed by the characteristic matter-decay scale ( $\lambda_{char} = 0.21178$ ), representing the effective boundary layer of spacetime. To relate this to electromagnetic mass, the square root of the characteristic scale is scaled by the Planck reference ( $m_s$ ) and the hysteresis of active channels:

$$X_{eff} = (\sqrt{\lambda_{char}} / m_s) (1 - 1.5H)$$

$$\text{Where } \lambda_{char} = \sqrt{[1 / (\lambda \sqrt{\dagger})]} = 0.21178:$$

$$X_{eff} = (\sqrt{0.21178} / 21.76437021) \times (1 - 1.5 \times 0.000076456) = 0.021142$$

The proton-to-electron mass ratio then follows directly from these quantities:

$$M_p / m_e = [ \pi^2 \sqrt{\dagger} (1 - (\Sigma J / 2N) (1 - \S)) ] / [ \Delta_s X_{eff} ]$$

$$M_p / m_e = 24.76031 / (0.637851 \times 0.021142) = 24.76031 / 0.0134848 = 1836.1527$$

This value is close to the experimental CODATA value of  $1836.15267343 \pm 0.0000011$ . In the normalised  $\eta = 1$  system, the ratio emerges as a structural relation without additional SI conversion.

#### LEPTONIC HARMONIC OVERTONES (MUON AND TAU MASS RATIOS)

Lepton generations may correspond to stable harmonic overtones of the lattice recursion. The electron occupies the ground state ( $\Xi^0 = 1$ ). The muon and tau are treated as the first and second overtones.

##### 4.1 The Muon Mass Ratio

The muon is treated as the projection of triplet tension into the four-dimensional frame at the primary lattice spacing  $\lambda$ :

$$M_{\mu} / m_e = (M_p / m_e) / \lambda \times (1 + j / \pi)$$

$$M_{\mu} / m_e = (1836.1527 / 8.885765876) \times (1 + 0.0019796024 / 3.14159265)$$

$$M_{\mu} / m_e = 206.6400 \times 1.000630 = 206.7702$$

This compares well with the CODATA value of  $206.7682830 \pm 0.0000046$ .

##### 4.2 The Tau Mass Ratio via Recursive Squaring

The tau is treated as the second-order overtone. The structural mass scales with the square of the first overtone:

$$M_{\tau}(\text{struct}) = M_{\mu}^2 = 8.0486^2 = 64.780$$

Projection into four dimensions introduces an effective coordinate resistance:

$$\begin{aligned} \text{Projection Viscosity} &= \Delta \Phi = 0.029305 \times 3.62312 \\ &= 0.106176 \end{aligned}$$

Attenuating the full capacity ( $\Xi = 2.0$ ) yields:

$$m_{\tau} / m_e(\text{pure}) = (M_p / m_e) (2 - \Delta \Phi) = 1836.1527 \times (2 - 0.106176)$$

$$= 3477.3509$$

A small environmental adjustment is applied for local laboratory conditions:

$$\Delta_{\text{drift}} = -0.5 \text{ H} (1 - \Delta \Phi + \Delta) (M_p / m_e)$$

$$\Delta_{\text{drift}} = -0.5 \times 0.000076456 \times (1 - 0.106176 + 0.029305) \times 1836.1527$$

$$= -0.0648$$

$$m_{\tau} / m_e(\text{observed}) = 3477.3509 - 0.0648$$

$$= 3477.286$$

This falls within the range of the experimental CODATA value of  $3477.23 \pm 0.23$ .

## TRIPLET HYSTERESIS LOADING AND THE PROTON MAGNETIC MOMENT

The number of intersecting tension junctions may determine how many times the hysteresis delay applies. This distinguishes singlet leptons from triplet baryons.

### 5.1 Leptons (Singlet Hysteresis)

Leptons are single-loop configurations with one effective update delay:

$$\Sigma J = J + \text{H} = 0.002056056$$

The electron anomaly is derived from the projected potential:

$$a_e = [ \Sigma J \times (\ell_g / \Xi^3) \times (1 - \S_s) ] / 2\Delta_s$$

$$a_e = [ 0.002056056 \times (8.903507 / 8) \times 0.646447 ] / 1.275702 = 0.00115965218$$

$$g_e = 2(1 + a_e) = 2.00231930436$$

Both values match the CODATA averages closely.

### 5.2 Baryons (Triplet Hysteresis)

Baryons have three intersecting junctions, suggesting three separate contributions to the effective delay. The pure geometric anomaly is:

$$a_{p\_pure} = (M_p / m_e) / (\Xi^{(2N)}) (1 - \mathbb{H})^{(1/3)}$$

$$a_{p\_pure} = (1836.1527 / 1024) \times (1 - 0.000076456)^{(1/3)} = 1.793072$$

Each junction then contributes the environmental delay, giving a total shift of  $3\mathbb{H}$ :

$$a_{p\_terrestrial} = a_{p\_pure} - 3\mathbb{H}$$

$$a_{p\_terrestrial} = 1.793072 - 3 \times 0.000076456 = 1.792843$$

This is very close to the measured value of  $1.79284734462 \pm 0.00000000096$ . The difference may reflect the effect of the medium on triplet structures in real laboratory conditions.

## CONCLUSION

This work suggests the mass hierarchy and magnetic moments of leptons and baryons may follow from the recursive geometry and projection properties of a discrete substrate. The proton-to-electron ratio, lepton mass scales, and magnetic anomalies all align closely with experimental values.

This paper focuses specifically on the mass and magnetic anomaly sectors. How these same structural ideas extend to general relativity, quantum field theory, and cosmology is developed fully in the main foundational paper.

## REFERENCES

- [0] J. Fenn, "The Mechaniverse: Foundations of Emergent Physical Law," Zenodo Working Paper, 2026.
- [1] CODATA, "Fundamental physical constants," National Institute of Standards and Technology, Gaithersburg, MD, USA, 2018.
- [2] International Bureau of Weights and Measures, "SI Brochure: The International System of Units (SI)," 9th ed., BIPM, Sèvres, France, 2019.
- [3] P. J. Mohr, D. B. Newell, and B. N. Taylor, "The Planck constant and the revision of the International System of Units," Rev. Mod. Phys., vol. 88, no. 3, p. 035009, 2016.

[4] S. Adams, "The equivalence of Compton wavelength and Schwarzschild radius for the Planck mass," *Am. J. Phys.*, vol. 87, no. 11, pp. 833–836, 2019.

[5] Particle Data Group, "Review of particle physics," *Prog. Theor. Exp. Phys.*, vol. 2024, no. 8, p. 083C01, 2024.