


Hidden yet quantifiable: A lower-bound for confounding strength using randomized trials

Piersilvio De Bartolomeis, joint work with J. Abad, K. Donhauser and F. Yang




Motivation: drug regulatory process


Time



Motivation: drug regulatory process




Motivation: drug regulatory process




Motivation: drug regulatory process




Motivation: drug regulatory process




Motivation: drug regulatory process




Motivation: drug regulatory process




Motivation: drug regulatory process




Motivation: drug regulatory process




Motivation: drug regulatory process




Motivation: drug regulatory process




Previous works: binary tests for confounding




Previous works: binary tests for confounding




Previous works: binary tests for confounding




Previous works: binary tests for confounding



Previous works: binary tests for confounding




Previous works: binary tests for confounding




Previous works: binary tests for confounding

Time




Some confounding is likely present in real-world data \implies binary tests are too sensitive


Our proposal: lower bounding confounding strength




Our proposal: lower bounding confounding strength




Our proposal: lower bounding confounding strength




Our proposal: lower bounding confounding strength




Our proposal: lower bounding confounding strength




Our proposal: lower bounding confounding strength



Our proposal: lower bounding confounding strength




Setting: potential outcomes




Setting: potential outcomes

- We have access to rct and os data coming from the distributions $\mathbb{P}_{\text{full}}^{\text{rct}}$ and $\mathbb{P}_{\text{full}}^{\text{os}}$ over $(X, U, T, Y(1), Y(0), Y)$




Setting: potential outcomes

- We have access to rct and os data coming from the distributions $\mathbb{P}_{\text{full}}^{\text{rct}}$ and $\mathbb{P}_{\text{full}}^{\text{os}}$ over $(X, U, T, Y(1), Y(0), Y)$
- We only observe the marginal distributions \mathbb{P}^{rct} and \mathbb{P}^{os} over (X, Y, T)



Setting: potential outcomes


- We have access to rct and os data coming from the distributions $\mathbb{P}_{\text{full}}^{\text{rct}}$ and $\mathbb{P}_{\text{full}}^{\text{os}}$ over $(X, U, T, Y(1), Y(0), Y)$
- We only observe the marginal distributions \mathbb{P}^{rct} and \mathbb{P}^{os} over (X, Y, T)



For $\diamond \in \{\text{rct}, \text{os}\}$ we can factorize $\mathbb{P}_{\text{full}}^{\diamond} =$

Setting: potential outcomes


- We have access to rct and os data coming from the distributions $\mathbb{P}_{\text{full}}^{\text{rct}}$ and $\mathbb{P}_{\text{full}}^{\text{os}}$ over $(X, U, T, Y(1), Y(0), Y)$
- We only observe the marginal distributions \mathbb{P}^{rct} and \mathbb{P}^{os} over (X, Y, T)



For $\diamond \in \{\text{rct}, \text{os}\}$ we can factorize $\mathbb{P}_{\text{full}}^{\diamond} = \underbrace{\mathbb{P}_{Y|Y(1), Y(0), T}}_{:=\mathbb{P}_{\text{det}}}$

Setting: potential outcomes


- We have access to rct and os data coming from the distributions $\mathbb{P}_{\text{full}}^{\text{rct}}$ and $\mathbb{P}_{\text{full}}^{\text{os}}$ over $(X, U, T, Y(1), Y(0), Y)$
- We only observe the marginal distributions \mathbb{P}^{rct} and \mathbb{P}^{os} over (X, Y, T)



For $\diamond \in \{\text{rct}, \text{os}\}$ we can factorize $\mathbb{P}_{\text{full}}^{\diamond} = \underbrace{\mathbb{P}_{Y|Y(1), Y(0), T}}_{:=\mathbb{P}_{\text{det}}} \underbrace{\mathbb{P}_{Y(1), Y(0)|X, U}}_{:=\mathbb{P}_{\text{inv}}}$

Setting: potential outcomes

- We have access to rct and os data coming from the distributions $\mathbb{P}_{\text{full}}^{\text{rct}}$ and $\mathbb{P}_{\text{full}}^{\text{os}}$ over $(X, U, T, Y(1), Y(0), Y)$
- We only observe the marginal distributions \mathbb{P}^{rct} and \mathbb{P}^{os} over (X, Y, T)



For $\diamond \in \{\text{rct}, \text{os}\}$ we can factorize $\mathbb{P}_{\text{full}}^\diamond = \underbrace{\mathbb{P}_{Y|Y(1), Y(0), T}}_{:=\mathbb{P}_{\text{det}}} \underbrace{\mathbb{P}_{Y(1), Y(0)|X, U}}_{:=\mathbb{P}_{\text{inv}}} \underbrace{\mathbb{P}_{X, T, U}^\diamond}_{:=\mathbb{P}_{\text{cnf}}^\diamond}$

Confounding strength and marginal sensitivity set

Confounding strength and marginal sensitivity set

- Since U is not observed, confounding can affect internal validity: $\mathbb{P}_{\text{cnf}}^{\text{os}}(T | X) \neq \mathbb{P}_{\text{cnf}}^{\text{os}}(T | X, U)$

Confounding strength and marginal sensitivity set

- Since U is not observed, confounding can affect internal validity: $\mathbb{P}_{\text{cnf}}^{\text{os}}(T \mid X) \neq \mathbb{P}_{\text{cnf}}^{\text{os}}(T \mid X, U)$
- We model the confounding strength as: $\Gamma^{-1} \leq \frac{\mathbb{P}_{\text{cnf}}(T = 1 \mid X, U)}{\mathbb{P}_{\text{cnf}}(T = 0 \mid X, U)} / \frac{\mathbb{P}_{\text{cnf}}(T = 1 \mid X)}{\mathbb{P}_{\text{cnf}}(T = 0 \mid X)} \leq \Gamma$

Confounding strength and marginal sensitivity set

- Since U is not observed, confounding can affect internal validity: $\mathbb{P}_{\text{cnf}}^{\text{os}}(T \mid X) \neq \mathbb{P}_{\text{cnf}}^{\text{os}}(T \mid X, U)$
- We model the confounding strength as: $\Gamma^{-1} \leq \frac{\mathbb{P}_{\text{cnf}}(T = 1 \mid X, U)}{\mathbb{P}_{\text{cnf}}(T = 0 \mid X, U)} / \frac{\mathbb{P}_{\text{cnf}}(T = 1 \mid X)}{\mathbb{P}_{\text{cnf}}(T = 0 \mid X)} \leq \Gamma$

Definition (informal)

Marginal sensitivity set

Given a distribution \mathbb{P}^{os} over (X, Y, T) and a confounding strength $\Gamma \geq 1$, we define the set of distributions $\tilde{\mathbb{P}}_{\text{full}}$ compatible with $\mathbb{P}^{\text{os}}_{\text{full}}$ as

Confounding strength and marginal sensitivity set

- Since U is not observed, confounding can affect internal validity: $\mathbb{P}_{\text{cnf}}^{\text{os}}(T \mid X) \neq \mathbb{P}_{\text{cnf}}^{\text{os}}(T \mid X, U)$
- We model the confounding strength as: $\Gamma^{-1} \leq \frac{\mathbb{P}_{\text{cnf}}(T = 1 \mid X, U)}{\mathbb{P}_{\text{cnf}}(T = 0 \mid X, U)} / \frac{\mathbb{P}_{\text{cnf}}(T = 1 \mid X)}{\mathbb{P}_{\text{cnf}}(T = 0 \mid X)} \leq \Gamma$

Definition (informal)

Marginal sensitivity set

Given a distribution \mathbb{P}^{os} over (X, Y, T) and a confounding strength $\Gamma \geq 1$, we define the set of distributions $\tilde{\mathbb{P}}_{\text{full}}$ compatible with $\mathbb{P}_{\text{full}}^{\text{os}}$ as

$$\text{MSM}(\mathbb{P}^{\text{os}}, \Gamma) := \{\tilde{\mathbb{P}}_{\text{full}} = \mathbb{P}_{\text{det}} \tilde{\mathbb{P}}_{\text{inv}} \tilde{\mathbb{P}}_{\text{cnf}} : \tilde{\mathbb{P}}_{\text{cnf}} \text{ conf. strength } \in [\Gamma^{-1}, \Gamma] \text{ and } \tilde{\mathbb{P}}^{\text{os}} = \mathbb{P}^{\text{os}}\}$$

Confounding strength and marginal sensitivity set

- Since U is not observed, confounding can affect internal validity: $\mathbb{P}_{\text{cnf}}^{\text{os}}(T \mid X) \neq \mathbb{P}_{\text{cnf}}^{\text{os}}(T \mid X, U)$
- We model the confounding strength as: $\Gamma^{-1} \leq \frac{\mathbb{P}_{\text{cnf}}(T = 1 \mid X, U)}{\mathbb{P}_{\text{cnf}}(T = 0 \mid X, U)} / \frac{\mathbb{P}_{\text{cnf}}(T = 1 \mid X)}{\mathbb{P}_{\text{cnf}}(T = 0 \mid X)} \leq \Gamma$

Definition (informal)

Marginal sensitivity set

Given a distribution \mathbb{P}^{os} over (X, Y, T) and a confounding strength $\Gamma \geq 1$, we define the set of distributions $\tilde{\mathbb{P}}_{\text{full}}$ compatible with $\mathbb{P}_{\text{full}}^{\text{os}}$ as

$$\text{MSM}(\mathbb{P}^{\text{os}}, \Gamma) := \{\tilde{\mathbb{P}}_{\text{full}} = \mathbb{P}_{\text{det}} \tilde{\mathbb{P}}_{\text{inv}} \tilde{\mathbb{P}}_{\text{cnf}} : \tilde{\mathbb{P}}_{\text{cnf}} \text{ conf. strength } \in [\Gamma^{-1}, \Gamma] \text{ and } \tilde{\mathbb{P}}^{\text{os}} = \mathbb{P}^{\text{os}}\}$$

Sensitivity analysis: bounds on treatment effects

Sensitivity analysis: bounds on treatment effects

- We define the average treatment effect (ATE) as: $\mu(\mathbb{P}_X, \mathbb{P}_{\text{full}}) := \mathbb{E}_{\mathbb{P}_X} \left[\mathbb{E}_{\mathbb{P}_{\text{full}}} [Y(1) - Y(0) \mid X] \right]$

Sensitivity analysis: bounds on treatment effects

- We define the average treatment effect (ATE) as: $\mu(\mathbb{P}_X, \mathbb{P}_{\text{full}}) := \mathbb{E}_{\mathbb{P}_X} \left[\mathbb{E}_{\mathbb{P}_{\text{full}}} [Y(1) - Y(0) \mid X] \right]$

not identifiable for $\mathbb{P}_{\text{os}}^{\text{full}}$

Sensitivity analysis: bounds on treatment effects

- We define the average treatment effect (ATE) as: $\mu(\mathbb{P}_X, \mathbb{P}_{\text{full}}) := \mathbb{E}_{\mathbb{P}_X} \left[\mathbb{E}_{\mathbb{P}_{\text{full}}} [Y(1) - Y(0) | X] \right]$ not identifiable for $\mathbb{P}_{\text{os}}^{\text{full}}$
- $\mu(\mathbb{P}_X, \mathbb{P}_{\text{full}}^{\text{os}}) \leq \mu_{\Gamma}^{+}(\mathbb{P}_X) := \sup_{\tilde{\mathbb{P}}_{\text{full}} \in \text{MSM}(\mathbb{P}_{\text{full}}^{\text{os}}, \Gamma)} \mathbb{E}_{\mathbb{P}_X} \left[\mathbb{E}_{\tilde{\mathbb{P}}_{\text{full}}} [Y(1) - Y(0) | X] \right]$

Sensitivity analysis: bounds on treatment effects

- We define the average treatment effect (ATE) as: $\mu(\mathbb{P}_X, \mathbb{P}_{\text{full}}) := \mathbb{E}_{\mathbb{P}_X} \left[\mathbb{E}_{\mathbb{P}_{\text{full}}} [Y(1) - Y(0) | X] \right]$ not identifiable for $\mathbb{P}_{\text{os}}^{\text{full}}$
- $\mu(\mathbb{P}_X, \mathbb{P}_{\text{full}}^{\text{os}}) \leq \mu_{\Gamma}^{+}(\mathbb{P}_X) := \sup_{\tilde{\mathbb{P}}_{\text{full}} \in \text{MSM}(\mathbb{P}_{\text{full}}^{\text{os}}, \Gamma)} \mathbb{E}_{\mathbb{P}_X} \left[\mathbb{E}_{\tilde{\mathbb{P}}_{\text{full}}} [Y(1) - Y(0) | X] \right]$
- $\mu(\mathbb{P}_X, \mathbb{P}_{\text{full}}^{\text{os}}) \geq \mu_{\Gamma}^{-}(\mathbb{P}_X) := \inf_{\tilde{\mathbb{P}}_{\text{full}} \in \text{MSM}(\mathbb{P}_{\text{full}}^{\text{os}}, \Gamma)} \mathbb{E}_{\mathbb{P}_X} \left[\mathbb{E}_{\tilde{\mathbb{P}}_{\text{full}}} [Y(1) - Y(0) | X] \right]$

Sensitivity analysis: bounds on treatment effects

- We define the average treatment effect (ATE) as: $\mu(\mathbb{P}_X, \mathbb{P}_{\text{full}}) := \mathbb{E}_{\mathbb{P}_X} \left[\mathbb{E}_{\mathbb{P}_{\text{full}}} [Y(1) - Y(0) | X] \right]$ not identifiable for $\mathbb{P}_{\text{os}}^{\text{full}}$
- $\mu(\mathbb{P}_X, \mathbb{P}_{\text{full}}^{\text{os}}) \leq \mu_{\Gamma}^{+}(\mathbb{P}_X) := \sup_{\tilde{\mathbb{P}}_{\text{full}} \in \text{MSM}(\mathbb{P}_{\text{full}}^{\text{os}}, \Gamma)} \mathbb{E}_{\mathbb{P}_X} \left[\mathbb{E}_{\tilde{\mathbb{P}}_{\text{full}}} [Y(1) - Y(0) | X] \right]$ $\mu(\mathbb{P}_X, \mathbb{P}_{\text{full}}^{\text{os}}) \in [\mu_{\Gamma}^{-}(\mathbb{P}_X), \mu_{\Gamma}^{+}(\mathbb{P}_X)]$
- $\mu(\mathbb{P}_X, \mathbb{P}_{\text{full}}^{\text{os}}) \geq \mu_{\Gamma}^{-}(\mathbb{P}_X) := \inf_{\tilde{\mathbb{P}}_{\text{full}} \in \text{MSM}(\mathbb{P}_{\text{full}}^{\text{os}}, \Gamma)} \mathbb{E}_{\mathbb{P}_X} \left[\mathbb{E}_{\tilde{\mathbb{P}}_{\text{full}}} [Y(1) - Y(0) | X] \right]$ If Γ is well-specified

Statistical test for confounding strength

Statistical test for confounding strength

- First step: build a statistical test for the null hypothesis $H_0 : \mathbb{P}_{\text{cnf}}^{\text{os}} \text{ conf. strength} \in [\Gamma^{-1}, \Gamma]$

Statistical test for confounding strength

- First step: build a statistical test for the null hypothesis $H_0 : \mathbb{P}_{\text{cnf}}^{\text{os}} \text{ conf. strength} \in [\Gamma^{-1}, \Gamma]$

Lemma (informal)


If transportability is satisfied, i.e. $\mathbb{E}_{\mathbb{P}_{\text{full}}^{\text{os}}} [Y(1) - Y(0) | X] = \mathbb{E}_{\mathbb{P}_{\text{full}}^{\text{rct}}} [Y(1) - Y(0) | X]$,
it holds that $\mathbb{P}_{\text{cnf}}^{\text{os}} \text{ conf. strength} \in [\Gamma^{-1}, \Gamma] \implies \mu(\mathbb{P}_X, \mathbb{P}_{\text{full}}^{\text{rct}}) \in [\mu_{\Gamma}^-(\mathbb{P}_X), \mu_{\Gamma}^+(\mathbb{P}_X)]$

Statistical test for confounding strength

- First step: build a statistical test for the null hypothesis $H_0 : \mathbb{P}_{\text{cnf}}^{\text{os}} \text{ conf. strength} \in [\Gamma^{-1}, \Gamma]$

Lemma (informal)

If transportability is satisfied, i.e. $\mathbb{E}_{\mathbb{P}_{\text{full}}^{\text{os}}} [Y(1) - Y(0) | X] = \mathbb{E}_{\mathbb{P}_{\text{full}}^{\text{rct}}} [Y(1) - Y(0) | X]$,
it holds that $\mathbb{P}_{\text{cnf}}^{\text{os}} \text{ conf. strength} \in [\Gamma^{-1}, \Gamma] \implies \mu(\mathbb{P}_X, \mathbb{P}_{\text{full}}^{\text{rct}}) \in [\mu_{\Gamma}^-(\mathbb{P}_X), \mu_{\Gamma}^+(\mathbb{P}_X)]$



Statistical test for confounding strength

- First step: build a statistical test for the null hypothesis $H_0 : \mathbb{P}_{\text{cnf}}^{\text{os}} \text{ conf. strength} \in [\Gamma^{-1}, \Gamma]$

Lemma (informal)

If transportability is satisfied, i.e. $\mathbb{E}_{\mathbb{P}_{\text{full}}^{\text{os}}} [Y(1) - Y(0) | X] = \mathbb{E}_{\mathbb{P}_{\text{full}}^{\text{rct}}} [Y(1) - Y(0) | X]$,
it holds that $\mathbb{P}_{\text{cnf}}^{\text{os}} \text{ conf. strength} \in [\Gamma^{-1}, \Gamma] \implies \mu(\mathbb{P}_X, \mathbb{P}_{\text{full}}^{\text{rct}}) \in [\mu_{\Gamma}^-(\mathbb{P}_X), \mu_{\Gamma}^+(\mathbb{P}_X)]$

IPW estimate ($\Gamma = 1$) $\mu(\mathbb{P}_X, \mathbb{P}_{\text{full}}^{\text{rct}})$




Statistical test for confounding strength

- First step: build a statistical test for the null hypothesis $H_0 : \mathbb{P}_{\text{cnf}}^{\text{os}} \text{ conf. strength} \in [\Gamma^{-1}, \Gamma]$

Lemma (informal)

If transportability is satisfied, i.e. $\mathbb{E}_{\mathbb{P}_{\text{full}}^{\text{os}}} [Y(1) - Y(0) | X] = \mathbb{E}_{\mathbb{P}_{\text{full}}^{\text{rct}}} [Y(1) - Y(0) | X]$,
it holds that $\mathbb{P}_{\text{cnf}}^{\text{os}} \text{ conf. strength} \in [\Gamma^{-1}, \Gamma] \implies \mu(\mathbb{P}_X, \mathbb{P}_{\text{full}}^{\text{rct}}) \in [\mu_{\Gamma}^{-}(\mathbb{P}_X), \mu_{\Gamma}^{+}(\mathbb{P}_X)]$




Statistical test for confounding strength

- First step: build a statistical test for the null hypothesis $H_0 : \mathbb{P}_{\text{cnf}}^{\text{os}} \text{ conf. strength} \in [\Gamma^{-1}, \Gamma]$

Lemma (informal)

If transportability is satisfied, i.e. $\mathbb{E}_{\mathbb{P}_{\text{full}}^{\text{os}}} [Y(1) - Y(0) | X] = \mathbb{E}_{\mathbb{P}_{\text{full}}^{\text{rct}}} [Y(1) - Y(0) | X]$,
it holds that $\mathbb{P}_{\text{cnf}}^{\text{os}} \text{ conf. strength} \in [\Gamma^{-1}, \Gamma] \implies \mu(\mathbb{P}_X, \mathbb{P}_{\text{full}}^{\text{rct}}) \in [\mu_{\Gamma}^{-}(\mathbb{P}_X), \mu_{\Gamma}^{+}(\mathbb{P}_X)]$




Statistical test for confounding strength

- First step: build a statistical test for the null hypothesis $H_0 : \mathbb{P}_{\text{cnf}}^{\text{os}} \text{ conf. strength} \in [\Gamma^{-1}, \Gamma]$

Lemma (informal)

If transportability is satisfied, i.e. $\mathbb{E}_{\mathbb{P}_{\text{full}}^{\text{os}}} [Y(1) - Y(0) | X] = \mathbb{E}_{\mathbb{P}_{\text{full}}^{\text{rct}}} [Y(1) - Y(0) | X]$,
it holds that $\mathbb{P}_{\text{cnf}}^{\text{os}} \text{ conf. strength} \in [\Gamma^{-1}, \Gamma] \implies \mu(\mathbb{P}_X, \mathbb{P}_{\text{full}}^{\text{rct}}) \in [\mu_{\Gamma}^{-}(\mathbb{P}_X), \mu_{\Gamma}^{+}(\mathbb{P}_X)]$



A lower bound on confounding strength

A lower bound on confounding strength

A lower bound on confounding strength

- Goal: lower bound on the true confounding strength $\Gamma^* := \mathbb{P}_{\text{cnf}}^{\text{os}}$ conf. strength

A lower bound on confounding strength

- Goal: lower bound on the true confounding strength $\Gamma^* := \mathbb{P}_{\text{cnf}}^{\text{os}}$ conf. strength
- We repeat the test until the first acceptance: $\hat{\Gamma}_{\text{LB}} = \inf_{\Gamma} \{\Gamma : \text{test}(\Gamma, \alpha) \text{ accepts}\}$

A lower bound on confounding strength

- ▶ Goal: lower bound on the true confounding strength $\Gamma^* := \mathbb{P}_{\text{cnf}}^{\text{os}}$ conf. strength
- ▶ We repeat the test until the first acceptance: $\hat{\Gamma}_{\text{LB}} = \inf_{\Gamma} \{\Gamma : \text{test}(\Gamma, \alpha) \text{ accepts}\}$
- ▶ No multiple testing correction is needed!


A lower bound on confounding strength

- ▶ Goal: lower bound on the true confounding strength $\Gamma^\star := \mathbb{P}_{\text{cnf}}^{\text{os}}$ conf. strength
- ▶ We repeat the test until the first acceptance: $\hat{\Gamma}_{\text{LB}} = \inf_{\Gamma} \{\Gamma : \text{test}(\Gamma, \alpha) \text{ accepts}\}$
- ▶ No multiple testing correction is needed!

Lemma (informal)

$\hat{\Gamma}_{\text{LB}}$ is an asymptotically valid lower-bound for Γ^\star , i.e. it holds that $\mathbb{P}(\hat{\Gamma}_{\text{LB}} \leq \Gamma^\star) \geq 1 - \alpha + o(1)$

Synthetic and semi-synthetic experiments




Synthetic
(Gaussian)


Semi-synthetic
(Hillstrom dataset)

Synthetic and semi-synthetic experiments

Synthetic
(Gaussian)




Semi-synthetic
(Hillstrom dataset)




Synthetic and semi-synthetic experiments

Synthetic
(Gaussian)

Semi-synthetic
(Hillstrom dataset)




Women's Health Initiative (WHI)




Women's Health Initiative (WHI)

- Randomized trial and observational study (1993-2005)




Women's Health Initiative (WHI)

- Randomized trial and observational study (1993-2005)
- Treatment: hormone replacement therapy




Women's Health Initiative (WHI)

- Randomized trial and observational study (1993-2005)
- Treatment: hormone replacement therapy
- Cohort: postmenopausal women aged between 50 and 79



Women's Health Initiative (WHI)

- Randomized trial and observational study (1993-2005)
- Treatment: hormone replacement therapy
- Cohort: postmenopausal women aged between 50 and 79
- Outcomes: coronary heart disease



Women's Health Initiative (WHI)

Women's Health Initiative (WHI)

- We define the critical value of confounding as $\hat{\Gamma}_{CT} := \inf\{\Gamma : 0 \in [\hat{\mu}_\Gamma^-, \hat{\mu}_\Gamma^+]\}$

Women's Health Initiative (WHI)

- We define the critical value of confounding as $\hat{\Gamma}_{CT} := \inf\{\Gamma : 0 \in [\hat{\mu}_\Gamma^-, \hat{\mu}_\Gamma^+]\}$
- Our procedure ψ_{sens} accepts the study if $\hat{\Gamma}_{CT} > \hat{\Gamma}_{LB}$

Women's Health Initiative (WHI)

- We define the critical value of confounding as $\hat{\Gamma}_{CT} := \inf\{\Gamma : 0 \in [\hat{\mu}_\Gamma^-, \hat{\mu}_\Gamma^+]\}$
- Our procedure ψ_{sens} accepts the study if $\hat{\Gamma}_{CT} > \hat{\Gamma}_{LB}$
- The “binary” procedure ψ_{bin} accepts the study if $\hat{\Gamma}_{LB} = 1$

Women's Health Initiative (WHI)

- We define the critical value of confounding as $\hat{\Gamma}_{CT} := \inf\{\Gamma : 0 \in [\hat{\mu}_\Gamma^-, \hat{\mu}_\Gamma^+]\}$
- Our procedure ψ_{sens} accepts the study if $\hat{\Gamma}_{CT} > \hat{\Gamma}_{LB}$
- The “binary” procedure ψ_{bin} accepts the study if $\hat{\Gamma}_{LB} = 1$



Women's Health Initiative (WHI)

- We define the critical value of confounding as $\hat{\Gamma}_{CT} := \inf\{\Gamma : 0 \in [\hat{\mu}_\Gamma^-, \hat{\mu}_\Gamma^+]\}$
- Our procedure ψ_{sens} accepts the study if $\hat{\Gamma}_{CT} > \hat{\Gamma}_{LB}$
- The “binary” procedure ψ_{bin} accepts the study if $\hat{\Gamma}_{LB} = 1$



Metric	Coronary heart disease	
	$t = 0$	$t = 20$
$\hat{\Gamma}_{CT}$	1.017	1.164
$\hat{\Gamma}_{LB}$	1.009	1.224
ψ_{bin}	1	1
ψ_{sens}	0	1

Women's Health Initiative (WHI)

- We define the critical value of confounding as $\hat{\Gamma}_{CT} := \inf\{\Gamma : 0 \in [\hat{\mu}_\Gamma^-, \hat{\mu}_\Gamma^+]\}$
- Our procedure ψ_{sens} accepts the study if $\hat{\Gamma}_{CT} > \hat{\Gamma}_{LB}$
- The “binary” procedure ψ_{bin} accepts the study if $\hat{\Gamma}_{LB} = 1$



$\downarrow \Gamma$

Metric	Coronary heart disease	
	$t = 0$	$t = 20$
$\hat{\Gamma}_{CT}$	1.017	1.164
$\hat{\Gamma}_{LB}$	1.009	1.224
ψ_{bin}	1	1
ψ_{sens}	0	1

Women's Health Initiative (WHI)

- We define the critical value of confounding as $\hat{\Gamma}_{CT} := \inf\{\Gamma : 0 \in [\hat{\mu}_\Gamma^-, \hat{\mu}_\Gamma^+]\}$
- Our procedure ψ_{sens} accepts the study if $\hat{\Gamma}_{CT} > \hat{\Gamma}_{LB}$
- The “binary” procedure ψ_{bin} accepts the study if $\hat{\Gamma}_{LB} = 1$



$\downarrow \Gamma$ $\uparrow \Gamma$

Metric	Coronary heart disease	
	$t = 0$	$t = 20$
$\hat{\Gamma}_{CT}$	1.017	1.164
$\hat{\Gamma}_{LB}$	1.009	1.224
ψ_{bin}	1	1
ψ_{sens}	0	1

Thanks for your attention! Questions?



Pier



Javier



Konstantin



Fanny



Statistical Machine Learning group

ETH zürich