Refinement

H atoms were positioned geometrically, with C—H = 0.95–0.98 Å, and constrained to ride on their parent atoms, with U_{iso}(H) = 1.2–1.5U_{eq}(C).

Figures




Fig. 1. The molecular structure of (II) showing the atomic labelling scheme and displacement ellipsoids drawn at the 50% probability level.

Pentacarbonyl-2κ^5*C*-chlorido-1κ*Cl*-bis[1(η^5)-\ cyclopentadienyl][μ$_2$-oxido(methyl)methylene-1:2κ^2*O*:*C*]\ tungsten(IV)zirconium(0)

Crystal data

$[ZrW(C_5H_5)_2(C_2H_3O)Cl(CO)_5]$	D_x = 2.065 Mg m^{-3}
M_r = 623.79	Mo *K*α radiation λ = 0.71073 Å
Orthorhombic, *Pnma*	Cell parameters from 25 reflections
a = 22.3794 (8) Å	θ = 2–17º
b = 12.3852 (7) Å	μ = 6.41 mm^{-1}
c = 7.2404 (3) Å	*T* = 273 (2) K
V = 2006.85 (16) Å^3	Prism, red
Z = 4	0.34 × 0.31 × 0.29 mm
F_{000} = 1176	

Data collection

Philips PW1100 diffractometer	R_{int} = 0.042
Radiation source: fine-focus sealed tube	θ_{max} = 25.0º
Monochromator: graphite	θ_{min} = 2.5º
T = 273(2) K	*h* = 0→26
ω/2θ scans	*k* = 0→14
Absorption correction: ψ scan (North *et al.*, 1968)	*l* = −8→0
T_{min} = 0.219, T_{max} = 0.258	3 standard reflections
2070 measured reflections	every 50 reflections
1851 independent reflections	intensity decay: none
1521 reflections with *I* > 2σ(*I*)	