

Organophosphate Poisoning

Section 1: Case Summary

Scenario Title:	Organophosphate poisoning
Keywords:	Toxicology, organophosphate
Brief Description of Case:	60-year old farmer presents to the ED with vomiting, rhinorrhea and dyspnea. He is placed into the resuscitation bay due to his appearance and hypoxia. Initial vital signs show tachypnea, hypoxia, and bradycardia. The team will need to identify the potential for poisoning as early as possible while proceeding with work-up and treatment. They must then initiate decontamination procedures of the patient and all staff who provided direct patient care. After appropriate treatment, the patient improves and is admitted to the ICU.

Goals and Objectives	
Educational Goal:	To expose learners to the identification and management of the uncommon, but clinically and academically important toxidrome of organophosphate poisoning.
Objectives: (Medical and CRM)	CRM: 1. Clear communication with emphasis on task prioritization and closed loop communication Medical 1. Effective resuscitation and treatment an organophosphate poisoning while maintaining patient and staff safety. 2. Recognize the need to protect staff from contamination. 3. Appropriately initiate patient and staff decontamination. 4. Administer appropriate antidotes and treatments for the patient.
EPAs Assessed:	

Learners, Setting and Personnel					
Target Learners:	☒ Junior Learners		☒ Senior Learners		☐ Staff
	☒ Physicians	☐ Nurses	☐ RTs	☐ Inter-professional	
	☐ Other Learners:				
Location:	☒ Sim Lab		☐ In Situ		☐ Other:
Recommended Number of Facilitators:	Instructors: 1				
	Sim Actors: 1				
	Sim Techs: 1				

Scenario Development	
Date of Development:	09/07/2021
Scenario Developer(s):	Phillip Lunsford, MD
Affiliations/Institutions(s):	Guthrie Emergency Medicine Residency Program
Contact E-mail:	Phillip.lunsford@guthrie.org
Last Revision Date:	
Revised By:	
Version Number:	1.0



Organophosphate Poisoning

Section 2A: Initial Patient Information

A. Patient Chart					
Patient Name: John Watts		Age: 60		Gender: M	Weight: 100 kg
Presenting complaint: Vomiting and shortness of breath					
Temp: 37.6°C	HR: 48	BP: 88/50	RR: 28	O ₂ Sat: 87%	FiO ₂ : RA
Cap glucose: 5.8			GCS: 15		
Triage note: Patient has been unwell since this morning. Runny nose, shortness of breath and vomiting.					
Allergies: None					
Past Medical History: CAD DM II HLD			Current Medications: Metoprolol Lisinopril Metformin Atorvastatin		

Section 2B: Extra Patient Information

A. Further History	
<i>Include any relevant history not included in triage note above. What information will only be given to learners if they ask? Who will provide this information (mannequin's voice, confederate, SP, etc.)?</i>	
When asked, also complains of excessive drooling, diffuse abdominal pain, diarrhea.	
When asked, patient runs the family berry farm. He has been treating crops with insecticide himself because the supplier has become too expensive. Not sure of the name. Has used it the last 3 days, but became ill last night and worse today.	
B. Physical Exam	
<i>List any pertinent positive and negative findings</i>	
Cardio: Bradycardic, no murmur	Neuro: Alert, GCS 15
Resp: Diffuse crackles bilaterally	Head & Neck: Heavy salivation, rhinorrhea
Abdo: Diffuse, mild tenderness	MSK/skin: Diaphoretic
Other: Pupils 2 and minimally reactive	

Organophosphate Poisoning

Section 3: Technical Requirements/Room Vision

A. Patient
☒ Mannequin (<i>specify type and whether infant/child/adult</i>)
☐ Standardized Patient
☐ Task Trainer
☐ Hybrid
B. Special Equipment Required
Crash cart/atropine Intubation supplies, including direct and video laryngoscopy Suction set-up PPE for providers
C. Required Medications
Atropine Pralidoxime
D. Moulage
Diaphoretic, wet clothes/gown
E. Monitors at Case Onset
☐ Patient on monitor with vitals displayed
☒ Patient not yet on monitor
F. Patient Reactions and Exam
<i>Include any relevant physical exam findings that require mannequin programming or cues from patient (e.g. – abnormal breath sounds, moaning when RUQ palpated, etc.) May be helpful to frame in ABCDE format.</i> A: Drooling, Vomiting B: Increased rate and work of breathing C: Lightheaded D: Miotic pupils, minimally reactive E: Sweaty, Moaning with diffuse palpation of abdomen

Organophosphate Poisoning

Section 4: Sim Actors and Standardized Patients

Sim Actors and Standardized Patient Roles and Scripts	
Role	Description of role, expected behavior, and key moments to intervene/prompt learners. Include any script required (including conveying patient information if patient is unable)
Bedside RN	RN in the room for scenario flow, nursing tasks, and cues when needed. RN can also complain of early symptoms of exposure if no PPE is used.
Wife	Wife can check in for more mild symptoms, also bring pesticide bottle/answer questions for decontamination cues.
Pharmacy/ Poison Control	Available by phone to answer questions of dosing and administration of antidotes



Organophosphate Poisoning

Section 5: Scenario Progression

Scenario States, Modifiers and Triggers				
Patient State/Vitals	Patient Status	Learner Actions, Modifiers & Triggers to Move to Next State		Facilitator Notes
1. Baseline State Rhythm: Sinus brady HR: 48 BP: 88/50 RR: 28 O ₂ SAT: 87% T: 37.6°C GCS: 15	Alert, answering questions. Dyspneic, vomiting, diaphoretic.	<u>Expected Learner Actions</u> ☐ IV, access, monitors ☐ ABCs, supp O2 ☐ Focused history ☐ Focused physical exam ☐ Verbalize possible poisoning ☐ Verbalize need for PPE ☐ Decontamination (pt/staff) ☐ ECG ☐ CXR	<u>Modifiers</u> - No PPE → RN states that he/she is starting to get nauseated - Supp O2 → 92% <u>Triggers</u> - Identification and treatment with atropine → 2. Treatment - No treatment for organophosphate by 5 min → 3. Decompensation	If learner does not request decontamination and PPE, nurse can comment an additional time about symptoms, but case will otherwise continue. Recognition for decontamination to be discussed at debrief.
2. Treatment for organophosphate HR: 48 BP: 88/50 RR: 24 O ₂ SAT: 92% GCS: 14	Unchanged	<u>Expected Learner Actions</u> ☐ Atropine IV (start with 1 mg, double dose q3-5m) ☐ Pralidoxime IV (30 mg/kg) ☐ Consider HF nasal cannula ☐ Consult pharmacist ☐ Consult poison control	<u>Modifiers</u> - First dose of atropine will improve symptoms, but patient will require multiple doses for clinical stability. <u>Triggers</u> - If only one dose of atropine and stop → 3. Decompensation - Multiple doses of atropine + pralidoxime given → 4. Resolution	Poison control confirms likely toxidrome and recommends continued atropine for symptom management and pralidoxime.
3. Decompensation HR: 36 BP: 74/47 RR: 36 O2 Sat: 84% GCS: 7 (E1V2M4)	Obtunded, urinary incontinence	<u>Expected Learner Actions</u> ☐ Obtain collateral info (wife) ☐ Consider intubation ☐ Consider vasopressors ☐ Atropine IV (start with 1 mg, double dose q3-5m) ☐ Pralidoxime IV (30 mg/kg)	<u>Modifiers</u> - RN may prompt that mental status changed <u>Triggers</u> - Multiple doses of atropine + pralidoxime given → 4. Resolution	Intubation would be high risk due to both anatomical and physiological challenges. Leader should be cued to the arrival of wife with pesticide bottle and similar but mild symptoms.



Organophosphate Poisoning

4. Resolution HR: 56 BP: 104/75 RR: 20 O2: 98% GCS 14		<u>Expected Learner Actions</u> ☐ Titrate atropine to dry secretions ☐ Labs/CXR if not obtained ☐ Pralidoxime infusion (8 mg/kg/h) ☐ Prepare diazepam ☐ Consult poison control (if not already completed)	<u>Modifiers</u> <u>Triggers</u> - Discuss with ICU → end case	
---	--	---	---	--




Organophosphate Poisoning

Appendix A: Laboratory Results


<u>VBG</u> pH 7.28 pCO ₂ 49 pO ₂ 39 HCO ₃ 21 Lactate 3.4	
--	--

Organophosphate Poisoning

Appendix B: ECGs, X-rays, Ultrasounds and Pictures



[Case courtesy of Frank Gaillard, Radiopaedia.org, rID: 3011](#)



https://en.wikipedia.org/wiki/Sinus_bradycardia#/media/File:ECG_Sinus_Bradycardia_43_bpm.jpg

Organophosphate Poisoning

Appendix C: Facilitator Cheat Sheet & Debriefing Tips

Sample questions:

1. What was on your initial differential for this patient?
2. How did you manage the ABCs?
3. How did you identify the toxidrome?
4. How and when do you decontaminate staff?
5. What did you consider for airway management? NIPPV?

Questions specific to organophosphate poisoning:

1. What are the clinical manifestations of organophosphate poisoning?
 - DUMBELS, SLUDGE/BBB
 - Cardiac manifestations?
 - Breathing manifestations?
 - Neurologic manifestations?
2. What are the management options for airway management?
 - NIPPV – Risks with copious nasal and oral secretions?
 - Intubation – Anatomic and physiologic challenges?
 - Atropine/pralidoxime effects
 - Neuromuscular fatigue from the poison may require intubations as well
 - Succinylcholine effects will be exaggerated and prolonged because its metabolized by Ach , Rocuronium will likely require higher dosing to overcome competitive inhibition at the NMJ
3. What are the treatments/antidotes for organophosphate poisoning?
 - Atropine – usually large amounts – for secretions (muscarinic receptors)
 - Pralidoxime – bolus, may require drip as well – for neuromuscular symptoms (nicotinic receptors)
4. Seizure management
 - Responds to benzos
 - Consider prophylactic diazepam
 - Will not respond to phenytoin or levetiracetam
5. Decontamination
 - Remove clothes
 - Copious skin irrigation +/- wash with soap
 - Gastric lavage generally not recommended (considered in ingestions <1 hour)
 - Charcoal recommended for ingestions (within 1 hour)
6. Prognosis
 - GCS < 13 at presentation generally poor prognosis
 - However exact agent involved plays a key role

Organophosphate Poisoning

Organophosphate and carbamate poisoning: Rapid overview

To obtain emergency consultation with a medical toxicologist, in the United States, call 1-800-222-1222, or the nearest international regional poison center. Contact information for regional poison centers around the world is available at the website referenced below.^[1]

Clinical syndromes

Acute toxicity

Generally manifests in minutes to hours

Evidence of cholinergic excess

SLUDGE = Salivation, Lacrimation, Urination, Defecation, Gastric Emptying

BBB = Bradycardia, Bronchorrhea, Bronchospasm

Respiratory insufficiency can result from muscle weakness, decreased central drive, increased secretions, and bronchospasm

Intermediate syndrome

Occurs 24-96 hours after exposure

Bulbar, respiratory, and proximal muscle weakness are prominent features

Generally resolves in 1-3 weeks

Organophosphorus Agent-Induced Delayed Peripheral Neuropathy (OPIDN)

Usually occurs several weeks after exposure

Primarily motor involvement

May resolve spontaneously, but can result in permanent neurologic dysfunction

Diagnostic evaluation of acute toxicity

Atropine challenge if diagnosis is in doubt (1 mg IV in adults, 0.01 to 0.02 mg/kg in children)

Absence of anticholinergic signs (tachycardia, mydriasis, decreased bowel sounds, dry skin) strongly suggests poisoning with organophosphate or carbamate

Draw blood sample for measurement of RBC acetylcholinesterase activity to confirm diagnosis

Treatment of acute toxicity

Deliver 100 percent oxygen via facemask; early intubation often required; avoid succinylcholine

Decontamination if ingestion within 1 hour give single dose activated charcoal, adult 50 g (1 g/kg in children) unless airway not protected or other contraindication. Aggressive dermal and ocular irrigation as needed. Bag/discard clothing.

Atropine 2 to 5 mg IV/IM/IO bolus (0.05 mg/kg IV in children)

Escalate (double) dose every 3-5 minutes until bronchial secretions and wheezing stop

TACHYCARDIA AND MYDRIASIS ARE NOT CONTRAINDICATIONS TO ATROPINE USE

Hundreds of milligrams may be needed over several days in severe poisonings

Inhaled ipratropium 0.5 mg with parenteral atropine may be helpful for bronchospasm; may repeat

Pralidoxime (2-PAM) 2 g (25 mg/kg in children) IV over 30 minutes; may repeat after 30 minutes or give continuous infusion if severe

Continuous infusion at 8 mg/kg/hour in adults (10 mg/kg/hour in children)

If no IV access, give pralidoxime 600 mg IM (15 mg/kg in children <40 kg). Rapidly repeat as needed to total of 1800 mg or 45 mg/kg in children.

Pralidoxime is given with atropine

Benzodiazepine therapy

Diazepam 10 mg IV (0.1 to 0.2 mg/kg in children), repeat as necessary if seizures occur. Do not give phenytoin.

Reference:

1. Poison emergency center contact numbers. Liquid Glass Nanotech. Available at: <https://www.liquidglassnanotech.com/poison-emergency-center-contact-numbers/> (Accessed on May 25, 2021)

UpToDate®



Organophosphate Poisoning

References

1. https://www.uptodate.com/contents/organophosphate-and-carbamate-poisoning?search=organophosphate%20poisoning&source=search_result&selectedTitle=1~28&usage_type=default&display_rank=1#H9 (Accessed 09/07/2021)
2. Nelson, Lewis S and Goldfrank, Lewis R. *Goldfrank's Toxicologic Emergencies*. New York: McGraw-Hill, 2019. Pages 1486-1513.

