

Exploring the Central Limit Theorem: How Many Groups for Normality?

Pearl Bipin Pulickal
pearlbipin@gmail.com

January 14, 2025

Abstract

This short technical note explores the Central Limit Theorem (CLT) and addresses a common query regarding its application. While it is well known that 30 samples are sufficient for the CLT to ensure the sample mean distribution approaches normality, the question remains: how many groups of 30 samples do we need to create a reliable plot of these sample means? This note explains that applying the Central Limit Theorem twice—once for a single group of 30 samples and again for multiple groups of 30 samples—results in a distribution that approximates normality and accurately represents the population mean.

1 Introduction

The Central Limit Theorem (CLT) is a fundamental concept in statistics that states that, given a sufficiently large sample size, the distribution of the sample mean will tend toward a normal distribution regardless of the original population distribution. It is widely accepted that 30 samples in a group are sufficient for CLT to apply and for the sample mean distribution to approach normality. However, a common question arises: how many groups of 30 samples are required to obtain a clear normal distribution of the sample means?

This technical note seeks to answer this question by explaining that the CLT must be applied twice: first to a group of 30 samples, and then again to 30 groups of 30 samples.

2 Application of the Central Limit Theorem

2.1 First Application of CLT

When the Central Limit Theorem is applied to a sample of size 30, the mean of these 30 samples will, by the CLT, follow a distribution that approaches normality as the sample size increases. However, a single group of 30 samples is insufficient to plot the distribution of sample means. To form a meaningful and reliable distribution of sample means, it is essential to take multiple groups of samples.

2.2 Second Application of CLT

To obtain a well-formed normal distribution of sample means, the Central Limit Theorem must be applied a second time. Instead of relying on just one group of 30 samples, we take 30 separate groups, each consisting of 30 samples. After calculating the mean of each of these 30 groups, the distribution of the means of these groups will begin to approximate a normal distribution, as predicted by the Central Limit Theorem.

Thus, to create a plot of sample means that truly resembles a normal distribution, we need to apply CLT to 30 groups of 30 samples, for a total of 900 samples. The distribution of these 30 group means will be close to normal, and the mean of these sample means will be very close to the population mean.

3 Conclusion

The Central Limit Theorem is a powerful tool in statistics, but to obtain a normal distribution of sample means, it must be applied twice. First, 30 samples are taken from the population, and their mean distribution approaches normality. Second, by taking 30 groups of 30 samples, the distribution of the means of those groups approximates a normal distribution, with the sample means clustering around the population mean. Therefore, the answer to the question of how many groups of 30 samples are needed to obtain a reliable normal distribution is 30 groups of 30 samples.

4 References

- B. S. Everitt, *The Cambridge Dictionary of Statistics*, Cambridge University Press, 2010.
- W. Feller, “An Introduction to Probability Theory and Its Applications,” *Vol. 1*, Wiley, 1968.
- J. L. Doob, “The Central Limit Theorem,” *Transactions of the American Mathematical Society*, 1949.
- M. L. Stein, *Interpolation of Random Processes*, Springer-Verlag, 1972.