

When the Imaginary Becomes Real: A Rigorous Exploration of Complex Exponentiation

Pearl Bipin Pulickal

October 7, 2025

Abstract

The expressions i^i and $i^{1/i}$ are remarkable examples in complex analysis where operations on purely imaginary numbers yield a set of exclusively real values. This paper provides a rigorous, pedagogically-focused derivation of these results, grounded in the formal definition of complex exponentiation, $z^w = e^{w \log z}$. We explore the multi-valued nature of the complex logarithm, the concept of branch cuts, and the distinction between the principal value and the infinite family of possible solutions. We demonstrate that all possible values of i^i are real and positive, and we examine the elegant symmetry between i^i and its reciprocal, i^{-i} . Finally, we extend Euler's formula to complex arguments, revealing its role as a profound bridge between circular and hyperbolic functions.

1. Introduction

The imaginary unit $i = \sqrt{-1}$ is a cornerstone of complex numbers. Its geometric interpretation as the point $(0, 1)$ in the complex plane, particularly through Euler's identity $e^{i\pi} + 1 = 0$, connects algebra to the geometry of rotations. This foundation, however, hints at deeper and less intuitive behaviors when exponentiation involves complex numbers in both the base and the exponent.

A canonical example that surprises many is the expression i^i . Is it real, imaginary, or complex? As we will demonstrate, not only is i^i real, but it represents an infinite set of real numbers. This paper aims to provide a clear and rigorous explanation of this phenomenon. To do so, we must first establish the formal machinery of the complex logarithm and exponentiation, which are essential for navigating these seemingly paradoxical results.

2. The Machinery of Complex Exponentiation

To evaluate an expression like z^w where $z, w \in \mathbb{C}$, we must define it in terms of functions we understand, namely the exponential and logarithm.

2.1. The Multi-Valued Complex Logarithm

For any non-zero complex number $z = x + iy$, we can write it in polar form as $z = re^{i\theta}$, where $r = |z| = \sqrt{x^2 + y^2}$ and $\theta = \arg(z)$ is the argument or angle. Because adding any integer multiple of 2π to the angle θ results in the same complex number, the argument is multi-valued.

$$z = re^{i(\theta+2\pi k)}, \quad k \in \mathbb{Z}$$

The complex logarithm, $\log z$, is defined as the inverse of the exponential function. Therefore,

$$\log z = \log(re^{i(\theta+2\pi k)}) = \ln(r) + i(\theta + 2\pi k).$$

Here, $\ln(r)$ is the standard real natural logarithm of the modulus of z . The multi-valued nature of the argument directly imparts a multi-valued nature to the logarithm.

2.2. Branch Cuts and the Principal Value

To work with a single-valued function, we typically restrict the argument θ to a specific interval of length 2π . The standard choice is $(-\pi, \pi]$, and the value of the argument in this interval is called the **principal argument**, denoted $\text{Arg}(z)$. This restriction defines the **principal value** of the logarithm, denoted $\text{Log}(z)$:

$$\text{Log}(z) = \ln |z| + i \text{Arg}(z).$$

To make $\text{Log}(z)$ continuous and analytic, we must introduce a **branch cut**—a curve in the complex plane where the function is discontinuous. For the principal branch, the cut is typically made along the negative real axis, including the origin.

The general logarithm can then be expressed in terms of the principal value:

$$\log z = \text{Log}(z) + i(2\pi k), \quad k \in \mathbb{Z}.$$

2.3. Defining Complex Exponentiation

Using the complex logarithm, we can now unambiguously define complex exponentiation for $z, w \in \mathbb{C}$ (with $z \neq 0$):

$$z^w := e^{w \log z} = e^{w(\text{Log}(z) + i(2\pi k))}.$$

This definition shows that, in general, z^w is multi-valued. The **principal value** of z^w is obtained by using the principal value of the logarithm ($k = 0$):

$$\text{P.V. } z^w = e^{w \text{Log}(z)}.$$

3. Evaluating i^i : An Infinite Family of Real Numbers

We now apply this framework to evaluate i^i .

3.1. The Logarithm of i

First, we find the logarithm of i . In polar form, $i = 1 \cdot e^{i\pi/2}$.

- The modulus is $|i| = 1$.
- The principal argument is $\text{Arg}(i) = \pi/2$.

Therefore, the principal value of the logarithm of i is:

$$\text{Log}(i) = \ln(1) + i\frac{\pi}{2} = i\frac{\pi}{2}.$$

The general, multi-valued logarithm is:

$$\log i = i\left(\frac{\pi}{2} + 2\pi k\right), \quad k \in \mathbb{Z}.$$

3.2. The Values of i^i

Using the definition $i^i = e^{i \log i}$, we substitute our expression for $\log i$:

$$i^i = e^{i \cdot [i(\frac{\pi}{2} + 2\pi k)]} = e^{i^2(\frac{\pi}{2} + 2\pi k)} = e^{-(\frac{\pi}{2} + 2\pi k)}.$$

Since k can be any integer, we have found that i^i is not a single number but an infinite set of real, positive values.

The **principal value** occurs when $k = 0$:

$$\text{P.V. } i^i = e^{-\pi/2} \approx 0.207879.$$

Other values include:

- For $k = 1$: $e^{-(\pi/2 + 2\pi)} = e^{-5\pi/2} \approx 0.000388$.
- For $k = -1$: $e^{-(\pi/2 - 2\pi)} = e^{3\pi/2} \approx 111.31778$.

All values are real, demonstrating that an imaginary number raised to an imaginary power can indeed be purely real.

4. Evaluating the Symmetric Counterpart: $i^{1/i}$

A similar analysis can be applied to $i^{1/i}$. Noting that $1/i = -i$, the expression is equivalent to i^{-i} .

Using the definition $i^{-i} = e^{-i \log i}$:

$$i^{-i} = e^{-i \cdot [i(\frac{\pi}{2} + 2\pi k)]} = e^{-i^2(\frac{\pi}{2} + 2\pi k)} = e^{\frac{\pi}{2} + 2\pi k}.$$

Once again, we find an infinite set of real, positive numbers. The principal value (for $k = 0$) is:

$$\text{P.V. } i^{1/i} = e^{\pi/2} \approx 4.810477.$$

It is immediately apparent that the set of values for $i^{1/i}$ is the reciprocal of the set of values for i^i . Specifically, the principal value of $i^{1/i}$ is the reciprocal of the principal value of i^i .

5. The Bridge to Hyperbolic Functions

The rigorous framework of complex analysis allows us to extend familiar functions, like sine and cosine, to complex arguments, revealing deep connections to other areas of mathematics. Euler's formula, $e^{i\theta} = \cos \theta + i \sin \theta$, is the key.

5.1. Trigonometric Functions with Complex Arguments

For any complex number z , the trigonometric functions are defined via the exponential function:

$$\cos z = \frac{e^{iz} + e^{-iz}}{2}, \quad \sin z = \frac{e^{iz} - e^{-iz}}{2i}.$$

These definitions are consistent with Euler's formula and extend the functions from the real line to the entire complex plane. Let's explore what happens when the argument is purely imaginary, say $z = ix$ for some $x \in \mathbb{R}$.

$$\begin{aligned} \cos(ix) &= \frac{e^{i(ix)} + e^{-i(ix)}}{2} = \frac{e^{-x} + e^x}{2} = \cosh(x). \\ \sin(ix) &= \frac{e^{i(ix)} - e^{-i(ix)}}{2i} = \frac{e^{-x} - e^x}{2i} = \frac{-(e^x - e^{-x})}{2i} = i \frac{e^x - e^{-x}}{2} = i \sinh(x). \end{aligned}$$

These fundamental identities, $\cos(ix) = \cosh(x)$ and $\sin(ix) = i \sinh(x)$, form a bridge between circular and hyperbolic trigonometry.

5.2. Euler's Formula Revisited

Let's verify Euler's formula for the imaginary argument ix :

$$\cos(ix) + i \sin(ix) = \cosh(x) + i(i \sinh(x)) = \cosh(x) - \sinh(x).$$

From the definition of hyperbolic functions, we know $\cosh(x) - \sinh(x) = e^{-x}$. On the other hand, the left side of Euler's formula gives:

$$e^{i(ix)} = e^{-x}.$$

The identity holds perfectly. This demonstrates that Euler's formula is not merely a statement about points on the unit circle but a universal identity in the complex plane. It shows that an "imaginary rotation" (multiplication by i) of a real angle corresponds to exponential scaling (hyperbolic motion).

6. Conclusion

The study of expressions like i^i offers more than just a mathematical curiosity; it serves as a gateway to a deeper understanding of the structures underlying complex analysis. Our rigorous exploration has revealed several key insights:

1. **Complex exponentiation is multi-valued:** Due to its definition via the complex logarithm, z^w generally represents an infinite set of values. Understanding this is crucial for correct interpretation.

2. **The imaginary can become real:** All values of i^i and $i^{1/i}$ are real numbers, a non-intuitive result that is demystified by the application of Euler's formula.
3. **Structure and symmetry:** The principal values of i^i and $i^{1/i}$ are reciprocals, a direct consequence of the sign change in the exponent.
4. **Unification of functions:** Extending trigonometric functions to the complex plane reveals that they are intrinsically linked to hyperbolic functions through Euler's formula, which acts as a fundamental bridge between the two.

These examples beautifully illustrate the consistency, elegance, and interconnectedness of mathematical concepts in the complex plane, turning what seems paradoxical into an exhibition of profound structural harmony.

References

1. Ahlfors, L. V. *Complex Analysis: An Introduction to the Theory of Analytic Functions of One Complex Variable*. McGraw-Hill, 1979.
2. Churchill, R. V., and Brown, J. W. *Complex Variables and Applications*. McGraw-Hill, 2009.
3. Euler, L. *Introductio in analysin infinitorum* (1748).
4. Nahin, P. J. *An Imaginary Tale: The Story of $\sqrt{-1}$* . Princeton University Press, 1998.