

## A New Hypothesis for Electrical Transmission in the Mammalian Central Nervous System

H. HILLMAN

*Unity Laboratory of Applied Neurobiology, University of Surrey, Guildford GU2 5XH, UK*

**Abstract** — It has been previously shown that the anatomical synapse does not exist in the living central nervous system, and that receptors are unlikely to be unique molecules sited on particular locations in or on the cell membrane. These conclusions and the rapidity of action of intravenously administered substances necessitated the adumbration of a new electrical hypothesis for the mechanism whereby signals pass from one part of the brain and spinal cord to another. The hypothesis proposed implies some testable predictions.

### **Brief historical summary on transmission**

In 1897, Sherrington proposed the physiological concept of the synapse (1, 2, 3), and soon afterwards Held saw 'end-fusse' on neurons in the trapezoid nucleus of cat, dog and rabbit. 'I shall designate (them) physiologically simply as zones for the transmission of stimuli' (4). Bernard demonstrated the dissociation of nerve and muscle activity by curarising frogs (5), and this is regarded as one of the key experiments in the development of the concept of chemical transmission. The observations that the actions of adrenalin and sympathetic stimulation were similar (6), as were acetylcholine and parasympathetic stimulation (7, 8) further supported this idea.

During the first half of the 20th Century, the alternative view, that transmission was electrical, enjoyed wide support, mainly because the electrical properties of what are now called conduction

and transmission were seen to be similar (9, 10, 11). At about the same time, the belief gradually gained currency that findings from the neuromuscular function were relevant not only to autonomic, but also to central synaptic, transmission. 'The original hypothesis was made as general as possible by applying it to the neuromuscular functions of skeletal muscle, and to the synapses of the sympathetic ganglia as well as of the central nervous system', wrote Eccles (11). This assumption has been almost universally accepted.

Katz and his collaborators studied the frog neuromuscular function by intracellular recording and electron microscopy. They concluded that transmission across it was chemical (12), except for some invertebrate synapses (13, 14). Both the current chemical hypothesis and the previous electrical hypothesis (11, 15) start off with the belief in the existence of the anatomical synapse in the living mammalian central nervous system. It should

be stressed that the hypothesis to be presented addresses the question of transmission in the central nervous system only, and does not share the assumption that its properties are the same as in other junctional tissues.

### *Introduction to new hypothesis*

Detailed evidence has been published that: (a) the central anatomical synapse probably does not exist in life (16), although there is no reasonable doubt about the physiological properties attributed to synapses; (b) receptors are probably not unique molecules located at discrete regions of the neural, synaptic or target cell membranes (17); (c) the central nervous system is largely composed of neuroglia, or 'nerve-glue' in the original sense of Virchow (18), that is, it is a syncytium composed of a ground substance enclosing relatively few neurons (19, 20, 21); and (d) many of the experiments designed to localise receptors by histochemical, electron microscopical, immunocytochemical and subcellular fractionation techniques, have never been controlled in respect of the possible effects of the procedures on the properties of the organelles being studied. Therefore, such experiments are incomplete, and it would be premature to draw conclusions from them (22, 23, 24).

Nevertheless, there is overwhelming evidence that signals do pass from one part of the central nervous system to another in the living animal. Therefore, serious reservations about important elements of current views induce an obligation to formulate alternative, testable hypotheses.

### *Previous evidence*

The following additional findings have also been used as the bases for the new hypotheses: (e) the concept of the blood brain barrier arose from the delay in arrival of dyes, ions, sugars and aminoacids in the brain (25–30); (f) anaesthetics, analgesics, analeptics, tranquillisers and other neuroactive substances, when injected intravenously, spread through the circulation in 0–12 s, and act on the central nervous system within seconds to minutes later (31–35); (g) cerebral and spinal capillaries are spaced 60  $\mu\text{m}$  or more apart in adults, as measured in stained tissues, which are shrunk (36, 37), so that the real intercapillary distance may be much higher. Thus substances entering the brain from the blood would have to

pass through the cerebral or spinal capillary membranes, and then the choroid plexus or the full length of the neuroglial cells before reaching the excitable cells (38–41); (h) cells presumed to be neurons are excited or inhibited by the extracellular application of transmitters and aminoacids (42–45); (i) lack of oxygen or substrate depolarises cells (46–50); (j) changes in the direct current potential differences across parts of the nervous system or individual cell membranes affect their excitability (51, 52, 53); (k) there is a potential difference between the cerebral blood vessels and the brain substance (54–58); (l) the conductance of the intact brain is greater than that of the membranes of the cells within it (59, 60, 61); (m) the concentrations of ions in the interstitial fluid of the brain are remarkably constant (62, 63); (n) non-synaptic transmission has been reported in a wide variety of mainly invertebrate tissues (13, 14). This has usually been assumed to occur at 'tight' junctions (64, 65), but this localisation cannot be proved directly; (o) electrical signals can be conducted through biological tissues without crossing synapses, for example, when an electrocorticograph or an electroencephalograph is recorded, or when extracellular stimulation or recording is made in relation to a neuron or a peripheral nerve; and (p) many drugs in clinical concentrations have high affinities for plasma proteins (66).

### *Assumptions of the proposed hypothesis*


1. Electrical signals can be conducted more rapidly through tissues than molecules can move by diffusion, passage across membranes, or as a consequence of enzyme reactions.
2. The capillaries penetrating throughout the central nervous system are the sites at which all rapidly acting substances have their effects.
3. There are no membranes between the capillaries and the neuroglia, or 'interstitial' compartment.
4. Rapidly acting neuroactive substances affect the electrical properties of the capillary membrane, and thus initiate an electrical impulse, which 'broadcasts' rapidly in all directions across the neuroglia to the nearest dendrites and somas, acting as aerials (Fig. 1).
5. The electrical impulse gradually dissipates in

the conductive neuroglia, with a speed and to an extent depending upon its chemical composition at that particular moment. The response of excitable cells to a stimulus depends upon the sensitivity of their membranes at that particular moment.

6. The particular routes of physiological signals between neurons is a consequence of their anatomical relationship to each other in a topographical sense, similar to the localisation of the sensory and motor cortices in the tracts and brain.

*Train of events following intravenous injection of a rapidly acting neuroactive substance, T*

T circulates within 9–12 s. Meanwhile, it partitions between the plasma, the blood cells and the endothelia of the blood vessels on the way, generally in favour of the plasma (Fig. 1).




**Fig. 1** Diagram showing the route of passage of the electrical excitability caused by the neuroactive substance acting on the membrane of the cerebral or spinal capillary, on the left. The electrical signal passes across the neuroglia to the dendrites or somas of the nearest neurons, on the right. For a description, please see the text.

T spreads rapidly through the total network of capillary membranes of the central nervous system. The network possesses a considerable resistance, impedance and capacitance. The capillary membranes are extremely thin in life, even when pericyte nuclei adhere to them, so that it is likely that the potential difference of the blood-brain

barrier is across the total capillary wall thickness, rather than between the blood and outer capillary membrane into the capillary cytoplasm, with a further potential difference between the cytoplasm across the inner capillary membrane into the neuroglia.

If T is of the right chemical configuration, it affects the resistance, impedance or capacitance of the capillary membrane. A transient electrical signal is induced at the capillary membrane-neuroglial interface (Fig. 2). The electrical signal is conducted in every direction through the low resistance neuroglia. The transient is not self-regenerative, and so it decrements within the volume of the neuroglia.



**Fig. 2** The equivalent circuit showing the passage of the electrical excitability from the capillary, on the left, across the capillary membrane and the neuroglia to the excitable cell membrane, on the right. The capacitances of the plasma, the neuroglia and the cytoplasm are assumed to be small, and are not shown.

The direct current potential difference across the tissue determines the excitability of the particular region of the brain or spinal cord. It could be the cause of the 'central excitatory' and 'central inhibitory' state of Sherrington (67, 68).

The 'broadcast' impulse is picked up by all the dendrites and somas in the region and decrements in the low resistance neuroglia (Fig. 1). Large enough, or several, impulses excite the dendrites or cell bodies in the region, and an orthodromic stimulus is initiated down the axon. Preformed discrete receptors do not exist on the soma or dendrites, but the whole, presumably soma-dendritic, membrane, is sensitive to electrical activity and substances which act rapidly on the central nervous system. According to a hypothesis previously proposed (17), the inherent characteristic of the excitable membrane is the range of substances to which it reacts by altering physico-chemical properties.

*Predictions of the proposed hypothesis*

1. T is found in relatively high concentration in the capillaries, but in extremely low or undetectable concentration adjacent to cells — upon which it acts if added iontophoretically, — as soon as it produces an effect.
2. The cerebral and spinal capillaries have a high affinity for T.
3. T affects the electrical, but not the chemical properties of the capillary membranes, before it can have any effect on excitable cells in the brain or spinal cord. Therefore, an electrical signal should be detected between the capillary and the excitable cell, before that cell fires.
4. Brief periods of natural physiological stimulation *in vivo* do not produce detectable biochemical effects.
5. Substances acting within a few seconds of administration cause their effects by reacting within the capillary membrane. Substances taking longer could affect the chemical or electrical properties of the neuroglia.
6. The properties of the blood brain barrier are relevant to slower events in the central nervous system, such as partitioning of T within the different compartments, affinity for the chemical constituents of the latter, the reaction of T with the latter (including its metabolism), and excretion of T and its metabolic products.

**References**

1. Sherrington C.S. The central nervous system, pps 915 – 1252, in *Textbook of Physiology*, by Foster M. and Sherrington C.S. Macmillan, London, 1897.
2. Fulton J.S. *Textbook of Neurophysiology*. Oxford University Press, New York, pps 70 – 104, 1949.
3. Eccles J.C. *The Physiology of Synapses*. Springer, Berlin, pps 3 – 26, 1963.
4. Held H. Beiträge zur Struktur der Nervenzellen und ihrer Fortsätze Zweite Abhandlung. *Arch Anat Physiol (Lpz.)* 21: 204 – 294, 1897.
5. Bernard C. Leçons sur les Effets des Substances Toxiques et Médicamenteuses. Baillière, Paris, pps 351 – 353, 1857.
6. Elliott T.R. The action of adrenalin. *J Physiol Lond* 32: 401 – 467, 1905.
7. Loewi O. Über humorale Übertragbarkeit der Herzherzenwirkung II. Mitteilung. *Pflügers Arch* 193: 201 – 213, 1922.
8. Dale H.H. published in, *Nobel Lectures in Physiology or Medicine, 1922 – 1941*. Elsevier, Amsterdam, pps 402 – 413, 1956.
9. Erlanger J. The Initiation of impulses in axons. *J Neurophysiol* 2: 370 – 379, 1939.
10. Gasser H.S. Axons as samples of nervous tissue. *J Neurophysiol* 2: 361 – 369, 1939.
11. Eccles J.C. An electrical hypothesis of synaptic and neuromuscular transmission. *Ann NY Acad Sci* 47: 429 – 455, 1946.
12. Katz B. *The Release of Neural Transmitter Substances*. Sherrington Lecture. Liverpool University Press, Liverpool, 1969.
13. Bennett M.V.L. Electrical transmission: a functional analysis and comparison to chemical transmission, in *Handbook of Physiology. The Nervous System. Vol. 1. Part 1*, ed. by Kandel E. The American Physiological Society, Bethesda, pps 357 – 416, 1977.
14. Vizi E.S. *Non-Synaptic Actions Between Neurons: Modulation of Neurochemical Transmission*. Wiley, New York, pps 43 – 62; 143 – 169, 1984.
15. Eccles J.C. A review and restatement of the electrical hypotheses of synaptic excitatory and inhibitory action. *Arch Sci Physiol* 3: 567 – 584, 1949.
16. Hillman H. The anatomical synapse in mammals by light and electron microscopy. *Medical Hypotheses* 17: 1 – 32, 1985a.
17. Hillman H. Critique of current views on receptors, and a new general theory of receptor location and action *in vivo*. *Medical Hypotheses* 26: 193 – 308, 1988.
18. Virchow R. Ueber das granulirte Aussehen der Wandungen des Gehirnvventrikels. *Allg Z Psychiatrie, Berlin* 3: 242 – 250, 1846.
19. Hess W.R. The ground substance of the central nervous system. *J Comp Neurol* 98: 69 – 91, 1953.
20. Hillman H. *Cellular Structure of the Mammalian System*. MTP Press, Lancaster, 1986.
21. Hillman H. Jarman D. *Atlas of the Cellular Structure of the Human Nervous System*. Academic Press, London 1990.
22. Hillman H. *Certainty and Uncertainty in Biochemical Techniques*. Surrey University Press, Blackie, Glasgow, 1972.
23. Hillman H. Some fundamental theoretical and practical problems associated with neurochemical techniques in mammalian studies. *Neurochem Internat* 5: 1 – 13, 1983.
24. Hillman H. Uncertainties, artifacts, uncontrolled experiments and incomplete evidence in modern biology. *Physiol Chem & Phys & Med NMR* 21: 145 – 164, 1989.
25. Lewandowsky M. Zur Lehre der Cerebrospinalflüssigkeit. *Z Klin Med* 40: 480 – 494, 1900.
26. Goldman E.E. Der aussere und innere Sekretion des gesunden und Kranken Organismus in Lichte der 'Vitalen Färbung'. *Beitr Klin Chirurg* 64: 192 – 265, 1900.
27. Goldman E.E. Vitalfärbung am Zentralnervensystem. *Abh Press Akad Wiss, Phys Math Kl* 1, 1 – 60, 1913.
28. Rapoport S.I. *Blood-Brain Barrier in Physiology and Medicine*. Raven Press, New York, 1976.
29. Suckling A.J. Rumsby M.G. Bradbury A.W.B. (eds.) *The Blood-Brain Barrier in Health and Disease*. Horwood, Chichester, 1986.
30. Davson H. Keasley W. Segal M. *The Physiology and Pathology of the Blood-Brain Barrier*. Churchill Livingstone, Edinburgh 1987.
31. Oldendorf W.H. Measurement of the mean transit time of cerebral circulation by external detection of an intravenously injected isotope. *J Nuc Med* 3: 382 – 398, 1962.

32. Dundee J W, Wyant G M. Intravenous Anaesthesia. Churchill Livingstone, Edinburgh pps 8–39, 1974.
33. Goldstein A. Aronow L. Kalman S.M. Principles of Drug Action. Second Edition, Harper International, New York, pps 131–134, 1974.
34. Ghonesin M, Korttila K. Pharmacokinetics of intravenous anaesthetics: implications for clinical use. *Clin Pharmacol* 2: 344–372, 1977.
35. Lake C.L. Cardiovascular Anaesthesia. Springer Verlag, New York, pps 96–126, 1985.
36. Lodin Z. Mares V, Karasek J.S. Skrivanova. Studies on the effect of fixation on nervous tissue. II Changes of size of nuclei of nervous cells after fixation and further histological treatment of the nervous tissue. *Acta Histochem* 28: 297–310, 1967.
37. Chughtai I, Hillman H, Jarman D. The effects of haematoxylin and eosin, Palmgren's and osmic acid procedures on the dimensions and appearance of isolated rabbit medullary neurones. *Microscopy* 35: 652–659, 1987.
38. Craigie E.H. Post natal changes in vascularity in the cerebral cortex of the male albino rat. *J Comp Neurol* 39: 301–324, 1925.
39. Diemer K. Capillarisation and oxygen supply to the brain, in *Oxygen Transport in Blood and Tissue*, ed. by Lubbers D.W. Luft U.C. Theus G. Witzleb E. Thieme, Stuttgart, pps 118–123, 1968.
40. Ohno K. Pettigrew K.D. Rapoport S.I. Lower limits of cerebrovascular permeability to non-electrolytes in the conscious rat. *Amer J Physiol* 235: H229–307, 1978.
41. Weiss H.R. Buchweitz E. Martha T.J. Auletta M. Quantitative regional determination of morphometric indices of total and perfused capillary network in the rat brain. *Circ Res* 51: 495–503, 1982.
42. Curtis D.R. Watkins J.C. Acidic aminoacids with strong excitatory action on mammalian neurones. *J Physiol* 166: 1–14, 1963.
43. Krnjevic K. Chemical nature of synaptic transmission in vertebrates. *Physiol Rev* 51: 418–540, 1974.
44. Obata K. Biochemistry and physiology of aminoacid transmitters, in the *Nervous System Vol. 1 Cell Biology of Neurons Pt 1*, ed by Kandel E.R. American Physiological Society, Bethesda, pps 625–650, 1977.
45. Roberts P.J. Storm-Mathieson J. Bradford H.F. (eds) *Excitatory Amino-Acids*. Macmillan Press, London (many papers), 1986.
46. Kolmodin G.M. Skoglund C.R. Influence of asphyxia on membrane potentials of spinal moto- and interneurones. *Acta Physiol Scand* 45: 1–18, 1959.
47. Hillman H. Membrane potentials in guinea pig cerebral cortex slices in vitro. Their dependence on substrate and oxygen. *J Neurochem* 8: 257–261, 1961.
48. Glaser G.H. Sodium and seizures, *Epilepsia* 5: 97–111, 1964.
49. Hillman H. Hyden H. Membrane potentials in isolated neurones in vitro from Deiters' nucleus of rabbit. *J Physiol* 177: 398–410, 1965.
50. Meldrum B. Brierley J.B. Prolonged epileptic seizures in primates: ischaemic cell change and its relation to local physiological events. *Arch Neurol, Chicago* 28: 10–17, 1973.
51. Li C.L. Jasper H.H. Microelectrode studies of the electrical activity of the cerebral cortex in the cat. *J Physiol* 121: 117–140, 1953.
52. Creutzfeldt D.D. Fromm G.H. Kapp H. Influence of transcortical d-c currents on cortical neuronal activity. *Exp Neurol* 5: 436–452, 1962.
53. Bindman L. Lippold O. The Neurophysiology of the Cerebral Cortex. Arnold, London, pps 113–130, 1981.
54. Loeschke H.H. Über den Einfluss von CO<sub>2</sub> auf die Bestandpotentiale der Hirnhäute. *Pflug Arch ges Physiol* 262: 532–536, 1956.
55. Tschirgi R.D. Taylor J.L. Slowly changing bioelectric potentials associated with the blood brain barrier. *Amer J Physiol* 195: 7–22, 1958.
56. O'Leary J.L. Goldring S. D-c potentials of the brain. *Physiol Rev* 44: 99–125, 1964.
57. Crone C. Olesen S.P. Electrical resistance of brain microvascular endothelium. *Brain Res* 241: 49–55, 1981.
58. Butt A.M. Jones H.C. Electrical resistance across the developing blood brain barrier in anaesthetised rats. *J Physiol* 423: 32P, 1989.
59. Freygang W.H. Landau W.M. Some relations between resistivity and electrical activity in the cerebral cortex of the cat. *J Cell Comp Physiol* 45: 377–392, 1955.
60. Ranck J.B. Specific impedance of rabbit cerebral cortex. *Exptl Neurol* 7: 144–152, 1963.
61. Van Harreveld A. *Brain Tissue Electrolytes*, Butterworth, London pps 55–94.
62. Jones H.C. Keep R. The control of potassium concentration in the cerebrospinal fluid and brain interstitial fluid of developing rats. *J Physiol* 383: 441–453, 1987.
63. Pardridge W.M. Brain metabolism: a perspective from the blood brain barrier. *Physiol Rev* 63: 1481–1535, 1981.
64. Brightman M.W.L. Reese T.S. Junctions between intimately opposed cell membranes in the vertebrate brain. *J Cell Biol* 40: 648–677, 1969.
65. Crone C. The blood brain barrier: a modified tight epithelium in the Blood Brain Barrier in Health and Disease, ed by Suckling A.J., Rumsby M.G., Bradbury M.W.B. Horwood, Chichester pps 17–40, 1986.
66. Lancaster R. *Pharmacology in Clinical Practice*. Heinemann, London, pps 13–14, 1980.
67. Sherrington C.S. Remarks on some aspects of reflex inhibition. *Proc Roy Soc B97*: 519–545, 1925.
68. Lorente de No, R. Facilitation of motor neurones. *Amer J Physiol* 113: 505–523, 1935.