

The Cellular Structure of the Mammalian Nervous System

By the same author

Certainty and Uncertainty in Biochemical Techniques
(1972). Surrey University Press, Blackie,
London.

(With P. Sartory)

The Living Cell - A Re-examination of its Fine
Structure (1980). Packard Publishing Limited,
Chichester.

The Cellular Structure of the Mammalian Nervous System

A re-examination, and some consequences for neurobiology

by Harold Hillman

*The Reader in Physiology and
the Director of the Unity Laboratory of Applied Neurobiology,
University of Surrey*



MTP PRESS LIMITED
a member of the KLUWER ACADEMIC PUBLISHERS GROUP
LANCASTER / BOSTON / THE HAGUE / DORDRECHT



Published in the UK and Europe by
MTP Press Limited
Falcon House
Lancaster, England

British Library Cataloguing in Publication Data

Hillman, H.

The cellular structure of the mammalian nervous system : a re-examination and some consequences for neurobiology.

1. Nervous system—Mammals

I. Title

599.01'88 QL923

Published in the USA by
MTP Press
A division of Kluwer Boston Inc
190 Old Derby Street
Hingham, MA 02043, USA

Library of Congress Cataloging-in-Publication Data

Hillman, Harold.

The cellular structure of the mammalian nervous system.

Includes bibliographies and index.

1. Mammals—Cytology. 2. Mammals—Cytology—Technique.

3. Nervous system—Mammals. 4. Nerve tissue—Cytology.

5. Neurons—Identification. 6. Neuroglia—Identification.

I. Title.

[DNLM: 1. Histological Techniques. 2. Mammals—anatomy & histology. 3. Nervous System—cytology. 4. Nervous System—ultrastructure. WL 101 H654c]

QL739.15.H55 1985 599'.0188 85-16570

ISBN-13: 978-94-010-8681-3

e-ISBN-13: 978-94-009-4922-5

DOI: 10.1007/978-94-009-4922-5

Copyright © 1986 MTP Press Limited

Softcover reprint of the hardcover 1st edition 1986

All rights reserved. No part of this publication may be produced, stored in a retrieval system, or transmitted in any form, or by any means, electronic, mechanical, photocopying, recording or otherwise, without prior permission from the publishers.

Contents

Foreword	1
Acknowledgements	3
Philosophy	4
1 Landmarks in the cell biology of the nervous system	6
<i>Description of the cell. Fixation and cutting sections. Achromatic objectives. Early findings in neurohistology. Neuroglia described. Staining procedures. Control experiments. Brain as a syncytium. The reticular versus neuronal doctrine. Axons and dendrites. Mitochondria. Golgi body. Neuromuscular junction. Synapses. Transmission. Neurons and neuroglia. Nissl body. Granules. Myelinated fibres. The Schwann cell. Nucleoli and nucleolar membrane. Nucleolulus. Cytoarchitectonics and myeloarchitectonics. White matter. Physiological experiments on tracts. Microscopical techniques. Tissue culture. Cellular and intracellular movements. The cell membrane. Electron microscopy.</i>	
2 Methods of studying neural structure	27
<i>Microdissection. Histological techniques. Tissue culture. Electron microscopy. Subcellular fractionation.</i>	
3 Cellular identification – general comments	49
<i>Appearances of cells. Geometry of cells. Specificity of stains. Physiological properties. Specific cell markers. Appearances in tissue culture. The current view of different kinds of cells as described in the literature.</i>	
4 Specific identification	82
<i>Light microscopy of histological sections. Appearances by electron microscopy. Appearances in tissue culture. Specific staining. Markers. Conclusions from the literature.</i>	
5 Experimental studies	92
<i>Histology. Examination of unstained tissues. Cells with apparently thin films of cytoplasm. Serial sections. Quantimetry. Electron microscopical studies. Dorsal root ganglia. White matter. Axons.</i>	
6 Critique of these experiments	119

7	The fine structure of the nervous system	126
	<i>The neuron. The synapses. Dendrites. Neuromuscular junctions.</i>	
	<i>Cytoplasm of neurons. The nuclear membrane of neurons.</i>	
	<i>Nuclear pores. Nucleoplasm. Nucleoli of neurons. Astrocytes, oligodendrocytes and microglia. The syncytium. Naked nuclei.</i>	
	<i>White matter. Ganglion cells. Ependymal cells and endothelial cells. Myelin sheath. The Geren model. The Schwann cell.</i>	
8	Other evidence supporting these conclusions	183
	<i>Tissue between neurons. Resistance and impedance of brain and spinal cord. Proportion of brain in which abrupt direct current potentials can be recorded. Viscosity of brain and spinal cord. Astrocytes and oligodendrocytes.</i>	
9	Summary of conclusions about nervous tissue in living mammals	189
	<i>Types of cells. Syncytium. Neurons. Absence of cell contacts.</i>	
	<i>Cytoplasm. Nuclear membranes. Mitochondria. Nucleoli.</i>	
	<i>Nucleolar membrane. Naked nuclei. Ganglia. Unmyelinated fibres. Nature of myelin sheath. Fibre tracts in white matter.</i>	
	<i>Transmitters. Blood-brain barrier. The Geren model. The nature of axoplasmic flow.</i>	
10	Problems arising from the latter conclusions	193
	<i>Previous holders of these views. Identification of astrocytes.</i>	
	<i>Synapses and transmission. The nature of the structures described as synapses seen by light and by electron microscopy. 'Unit' membranes. Cytoskeleton or endoskeleton. The integrity of the brain. The central nervous system as a syncytium. Evidence other than standard transmission electron microscopy. Gliosis.</i>	
11	Properties of the nervous system in the light of present conclusions	205
	<i>Classification of cell types, normal and pathological. Membrane transport. The extracellular space. Cerebral oedema. Blood-brain barrier. The syncytium and the extracellular fluid. The vesicle hypothesis of transmission. A general theory of receptors. Does neuroglia have a high sodium ion content? Epilepsy. Myelin and demyelinating diseases. Biochemical studies of the central nervous system.</i>	
12	Practical recommendations	242
13	Postscript	244
Appendices		
1	Specificity of staining procedures	245
2	Specificity of markers	248
3	Typical quantimetric counts	250
4	Biological membranes and measurement of their thickness	251
5	The tissue in the electron microscope	255
6	Unsatisfactory terms	256
7	A few naive questions	258
References		
Author index		295
Subject index		306

List of Figures

Cover, Horizontal section of skull (Vesalius)	
1. Dr R. Virchow	5
2. Peripheral nerve (Leeuwenhoek)	7
3. Cerebellar neuron (Valentin)	8
4. Dr J.E. Purkinje	9
5. Isolated neurons (Deiters)	9
6. Brain adjacent to ventricle (Virchow)	10
7. Nervous system as syncytium (Bauer)	12
8. The Golgi apparatus (Golgi)	14
9. 'End-fusse' on neurons (Held)	16
10. Albumin stained with haematoxylin and eosin	19
11. The Schwann cell (Schwann)	20
12. 'Lipoid double-layer' model of cell membrane (Davson and Danielli)	25
13. Fluid mosaic model of membrane (Singer and Nicholson)	26
14. Professor H. Hyden	28
15. Hyden's technique for isolating neuron cell bodies	29
16. Isolated neuron fresh, and after osmium staining and Araldite embedding	30
17. Dorsal root ganglion, stained with Palmgren's stain	31
18. Isolated unfixed rabbit dorsal root ganglion cell	31
19. Myelinated axons of sciatic nerves unfixed	32
20. Neuroglial clumps (Hamberger)	33
21. Sections of three eggs of the same size	51
22. Cells in the central nervous system (Peters, Palay and Webster)	63
23. Neuroglial cells (Penfield)	79
24. Mitochondria in frontal lobe tissue	94
25. Rat kidney, liver, brain stained simultaneously with haematoxylin and eosin	96
26. Item, stained with cresyl violet	97
27. Item, stained with Weil and Davenport	98
28. Paraffin-embedded unstained sections of frontal cortex, viewed by phase-contrast microscopy	99

LIST OF FIGURES

29.	Osmium-fixed, Araldite-embedded sections of frontal, parietal and occipital cortex, viewed by phase-contrast microscopy	100
30.	Serial sections of rat septal region stained by five different procedures	103-104
31.	Electron micrograph of rat parietal cortex	109
32.	Electron micrograph of neuron in rat parietal cortex	110
33.	Electron micrograph of neuron plus 'satellite' in rat parietal cortex	111
34.	Ventral horn cells of rat stained with cresyl violet, Palmgren's, haematoxylin and eosin, and phosphotungstic acid haematoxylin	112
35.	Cerebellar Purkinje cells stained with cresyl violet, haematoxylin and eosin, Palmgren's stain, and phosphotungstic acid haematoxylin	113
36.	Frontal cortex of rat stained with Marsland, Glees and Erikson, haematoxylin and eosin, Patay's stain and phosphotungstic acid haematoxylin	114
37.	Dorsal root ganglion of human stained with haematoxylin and eosin	116
38.	Commissure in human ventral white matter stained with Palmgren's stain	116
39.	White matter of human spinal cord stained with Palmgren's stain	117
40.	White matter of human spinal cord stained with Weigert - Pal's stain	117
41.	Rat sciatic nerve viewed by dark-ground illumination	118
42.	Mitochondria in unfixed neurons and neuroglia	122
43.	Isolated rabbit neurons stained with phosphotungstic acid haematoxylin; Marsland, Glees and Erikson; Gallyas's; and Weil and Davenport stains	124
44.	Intracellular movements in isolated rabbit neuron	128-130
45.	Diagram showing origin of 'trilaminar' membrane	133
46.	Synapses on ventral motoneurons of cat (Wyckoff and Young)	137
47.	Diagram of synapses on neuron soma (de Robertis)	139
48.	Synapses on rat reticular neuron and ventral motoneuron (Curtis, Jacobson and Marcus)	140
49.	Diagram of expected appearances of attachments of synaptic knobs	141
50.	Visual cortex or cat stained by Golgi-Cox method (Sholl)	142
51.	Silver preparations of human cortex showing dendritic arborization (Conel)	143
52.	Neurons in retina (Wassle)	144
53.	Expected views of synapses by light and electron microscopy	145
54.	Diagram to differentiate if granules are on the cell surface or in the cytoplasm	148
55.	Purkinje cells in 25-day-old culture (Hild and Tasaki) and 42-day-old culture of cat cerebellum (Pomerat and Costero), seen by phase-contrast microscopy	149
56.	Human neuromuscular junction (Ralis, Beasley and Ralis)	157
57.	Drawing of intranucleolar structures in rabbit neurons	160
58.	Cells in cat cerebral cortex (Ramon-Moliner)	161
59.	Human astrocyte	162
60.	Oligodendroglia in white matter (Lewis)	162
61.	Electron micrograph of oligodendrocyte from rat lateral geniculate body (Glees)	163

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

62. Microglial cell (Bouman and Bok)	164
63. Electron micrograph of astrocytic foot (Peters, Palay and Webster)	166
64. Diagram of expected appearance of astrocytic foot	167
65. Schwann cell nuclei (Bloom and Fawcett)	168
66. Cortico-cortical connections (Bailey and Bonin)	169
67. Nuclei in subcellular fractions of rat brain (Nurnberg and Gordon)	170
68. Transverse section of gorilla spinal cord (Waldeyer)	171
69. Anterior horn of lateral ventricle of rat showing ependyma	172
70. The Geren model of myelination (Peters)	174
71. Electron micrograph of myelinated nerve (Fawcett)	175
72. 'Myelin figure' in rabbit kidney	178
73. Myelinated fibres in culture of kitten cerebellum (Hild)	180
74. Diagram of migration of cells in spinal cord, cerebellum and isocortex from ependyma, during development (Jacobson)	185
75. Penetration of guinea pig cerebral slice in vitro with intracellular electrode (Hillman, Campbell and McIlwain)	188
76. Diagram of neuron	190
77. Diagram of brain adjacent to cerebral ventricle	191
78. Diagram of high-power micrograph of central nervous system	192
79. Electron micrograph of cat cerebral cortex (Jones and Powell)	195
80. Cerebral astrocytoma of human being	207
81. A molecule of Na^+/K^+ ATPase protruding from the cell membrane (Repke)	211
82. Computer-assisted tomogram of brain of young man with hydrocephalus and high intelligence (Lorber)	227

Unattributed illustrations are original

There is no remembrance of former things; neither shall there be any remembrance of things that are to come with those that shall come after.

Ecclesiastes (200 B.C.)

But it still moves.

Galileo Galilei (1633)

I admit that the judgement can be biased in many ways, and to an almost incredible degree, so that while exempt from external control, it may be so dependent on another man's words, that it may fitly be said to be ruled by him; but although this influence is carried to great lengths, it has never gone so far as to invalidate the statement, that every man's understanding is his own.

B. Spinoza (1760)

I propose to describe briefly what I have seen during fifty years of work, and what an investigator can verify for himself.

S. Ramon y Cajal (1954)

DEDICATION

This monograph is dedicated to:

- the pursuit of scientific truth
- my wife Elizabeth
- Alexander, Annie Rachel, Benedict and Sophia, whose eyes look into the future.

List of Tables

(Names of tissues listed originate from references)

1. Studies dependent upon characterization of cells by light microscopy	1
2. Light microscopic studies whose authors say that they cannot distinguish neurons from neuroglia	37
3. Electron microscopic studies indicating that authors cannot distinguish different types of neuroglia or they use the term neuroglia	38
4. Light microscopic studies whose authors say that they cannot differentiate neuroglia or they use the term neuroglia	39
5. Light microscopic studies of tissue cultures pointing to inability to distinguish neurons and different kinds of neuroglia	42
6. Light microscopic studies of tissue cultures giving no anatomical criteria for identification	43
7. Ratios of activities of 51 different enzymes in neurons compared with neuroglia	44-45
8. Appearances by light microscopy of allegedly different cellular elements	52-61
9. Appearances by electron microscopy of allegedly different elements	64-71
10. Staining procedures, their specificity indicated by authors	74-77
11. Appearances in tissue culture of allegedly different elements	85
12. Enzymes, proteins, antibodies, transmitters, a toxin, a lipid, used as markers for cell types in tissue culture	88-91
13. Staining procedures used in our experiments	93
14. Cells stained in one section identified in the next serial sections	102
15. Percentage of area of brain stained with six stains in four areas of brain	106
16. Mean percentages of areas stained added up	107
17. Artifacts and their likely origin	127
18. Illustrations showing neurofilaments, etc. not occurring in bundles	131
19. Illustrations of knobs apparently of same size as synapses but seen outside neuronal somas	138
20. Neurons in tissue culture by light microscopy in which the nuclei appear clearer than cytoplasm	148

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

21. Width of synaptic clefts by electron microscopy	150
22. Number of synapses per soma seen by light microscopy	152
23. Number of synapses per soma seen by electron microscopy	153
24. Diameters of synapses by light microscopy stated by authors	153
25. Diameters of synapses by electron microscopy measured from illustrations	154
26. Diameters of synaptic vesicles by electron microscopy	155
27. Substances proposed to be involved in communication between neurons (transmitters)	155
28. Myelin sheaths at different orientations but similar spacing	176
29. Large molecules believed to be present in the membranes of nerve cells	213-216
30. Early measurements of the extracellular space in intact animals by markers	223

Foreword

It would seem an appropriate time to re-examine the cellular structure of the mammalian nervous system for the following reasons. Firstly, there is considerable confusion in the literature about the appearance of the different kinds of neuroglia by light and by electron microscopy, and this is complemented by widespread disagreements among distinguished neuropathologists about the international classification of tumours of the central nervous system. Secondly, there is an increasing volume of experiments on the physiology and biochemistry of tissue cultures of neurons and different kinds of neuroglia, whose validity depends upon the accurate identification of both the parent tissue and also of the cells subsequently growing in culture. The biochemical classification in recent years has often tended to become independent of the cellular identification, which makes the use of the neuroglial cell names doubtful and the significance of the biochemical properties of the cells difficult to relate to the physiological properties *in vitro* or *in vivo* (Table 1).

Table 1 Studies dependent upon the characterization of cell types in the nervous system by light microscopy

- Classification and prognosis of tumours and infections in biopsies and histological sections
 - Identification of cells in electron micrographs
 - Identification of cells in tissue culture
 - Histochemical localizations in sections
 - Immunocytological localizations
 - Identification of cells in developing nervous system
 - Localization of physiological and biochemical properties in whole animals
 - Localization of transmitters and receptors
 - Characterization of cell separation by bulk separation methods
 - Identification of sites of action of drugs
 - Identification of biochemical properties of subcellular fractions from different cell types
 - Identification of microdissected cells
 - Identification of sites of action of bacterial and viral infections
-

FOREWORD

Thirdly, histologists have not been sufficiently concerned with the distortions of the shapes, volumes or appearances of the cellular elements, resulting from the procedures they employ (Hillman, 1982b), nor how much the properties of the stained sections reflect the properties of the living tissue. Fourthly, most stains intended to be specific require a considerable art and skill to obtain the appearance which is agreed to be characteristic of the particular kind of cell. However, more seriously, most of the classical stains used for the central nervous system were developed before the concept of control experiments became as important as it is now agreed to be. Most stains which were said to be 'specific' were judged to be so by well-known neuropathologists, who did not generally test rigorously whether or not their particular stain showed up, for example, neurons as well as a particular kind of neuroglial cell. Fifthly, electron microscopists have attached much importance to whether they were cutting relatively thick (100 nm) or relatively thin (10 nm) sections, but virtually no attention has been given to the multitude of different geometrical shapes which must result from cutting a neuron and its processes in a number of different orientations (please see page 51). Sixthly, I believe that research in the causation and mechanism of neurological diseases such as multiple sclerosis, cerebral tumours and cerebral thrombosis, and psychological illnesses, such as manic-depression and schizophrenia, has been remarkably unsuccessful. I have previously suggested that this is partly due to the widespread use of biochemical procedures which themselves may change the tissues excessively and therefore obscure differences between normal and pathological tissue, or may destroy the unstable factors causing disease which they are intending to identify (Hillman, 1972). We have further suggested that the lack of success is also partly due to the considerable effort and resources devoted to the examination of structures apparent after electron and light microscopical preparation, which could not exist in living cells (Hillman and Sartory, 1980a).

Acknowledgements

I have great pleasure in acknowledging the careful and meticulous contribution to this work of Mrs Dalveen Taylor, in respect of assistance with the experiments and the literature. Mr James Kirby of St Peter and St Paul's Hospital, London, carried out considerable histological work using many staining procedures, and Miss Julie Howarth of the Biochemistry Department, Surrey University, also helped with histology. Two kind electron microscopists, who prefer to remain anonymous, nevertheless helped with some preparations. Mrs Nita Foale, Mr Kevin Shaughnessy and the whole Audio-Visual Aids Unit, directed by Mr Russell Towns, aided me very much with the photography and illustrations. The Librarians of Surrey University, particularly Mr Denis Hewlett and Mr Glyn Davis, and the many Librarians of the British Medical Association, the University of London, University College London and the Science Museum Library, could not have been more helpful with the bibliography, especially of older source material. Miss Elizabeth Bruce and Mrs. Linda James typed out the various drafts and corrections of the manuscript, which was an awesome task. Mr. David Jarman helped with the manuscript. Dr. Mary Hearns and Mr. Robert Calderwood helped with the literature. To all these colleagues, I express my profound gratitude.

I will be grateful to any reader who wishes to discuss, or take issue with, the views expressed, - about which I would be willing to take part in any dialogue, private or public.

Philosophy

The philosophy behind this monograph may be summarized: firstly, we are seeking to elucidate properties of living cells by direct examination or extrapolation. A corollary of this is that information from living or unfixed tissue is to be regarded as more valuable than that obtained from tissues subjected to treatments such as killing, fixation, freezing, dehydration, staining, homogenization, centrifugation, sonication, addition of inhibitors, or the application of powerful non-physiological chemicals, when the data derived from these two kinds of experiments are mutually incompatible. For example, information about the structure of unfixed neurons derived from phase-contrast microscopy is considered more valuable than that found by freeze etching (Hillman, 1976). Secondly, Occam's Razor is a useful instrument. Thirdly, Popper's insistence on the necessity to seek to falsify one's own hypothesis is regarded as a valuable scientific modus operandi. Fourthly, geometry has to be respected, irrespective of the size of the objects being examined. Fifthly, one should not adopt an agnostic position about an important theory in one's own discipline, which one is propounding; either one concludes that sufficient evidence has been proffered to accept the validity of a particular theory, or one regards the evidence as being insufficient to be persuasive - in which case one should reject the particular viewpoint. One should not espouse a theory in a subject central to one's discipline, and then hide behind one's own ignorance of the evidence upon which it is critically based. Sixthly, findings, or a theory based upon them, cannot be accepted as true if major assumptions inherent in the use of the technique have been shown experimentally to be wrong, or contradict the laws of optics, physics or thermodynamics - unless the effects of the assumptions being wrong have been calculated (or shown) to make no significant difference to the results or interpretation of the experiments. Seventhly, an experiment is only as good as the control observations made on the effect of the technique itself on the system under study. Obviously, no amount of repetition of experiments in which major assumptions have been ignored or the control observations have not been made can make them valid or justify deriving conclusions from them.

I do not consider that these philosophical attitudes are in any way controversial, but would be interested to hear from anyone who does. Certain reviewers have said or implied that they do not agree with some of the elements of the philosophy summarized here, but they have not stated precisely or at all which are the particular ones to

PHILOSOPHY

which they take objection, or why they do (Lucy, 1973; Robards, 1981; Gregson, 1981).

The writing of the monograph arose from the re-examination of the structure of the living cell (Hillman and Sartory, 1980a), and the application of the same philosophy and geometrical considerations to the examination of the mammalian nervous system. However, it coincided with an attempt to answer a question which has been facing me daily for nearly 20 years. During this period I have been using the method of Hyden (1959) to dissect out by hand neurons from the cranial nerve nuclei of rabbits, rats, guinea pigs and mice. In this technique one cuts transverse sections about 1 mm thick of the fresh medulla, and stains it slightly with weak methylene blue in 0.25 mol/l sucrose solution. The sections are laid flat on a wax block, and under a dissecting microscope the neurons appear marine blue on a background of intensely white tissue. One is struck by the relatively small area that the somas of these neurons - defined as the cells which stain blue - occupy as a proportion of the total area of the tissue. At the same time, one frequently asks oneself 'What is the cellular nature of the tissue between the neurons?' One may surmise that this question was in Virchow's mind (1846), when he gave the name neuroglia to this tissue (Figures 1 and 6).

The present monograph examines the normal mammalian nervous system, although its implications for neuropathology will be considered later (please see page 205). Much of the literature on neuroglia has been derived from studies on sections from brains which have infections or tumours. In my opinion the demonstration that cell types described as neuroglia in healthy tissue are identical with those given the same names in abnormal tissue or in tissue cultures has not been done rigorously, especially in view of the knowledge that the appearance and biochemistry of cells change as a result of infections, tumours or growth in tissue cultures.




FIGURE 1 Dr Rudolf Virchow, who described the neuroglia as a nerve glue in the 1840s

1

Landmarks in the Cell Biology of the Nervous System

In this chapter the landmarks in the development of current views on the cellular structure of the mammalian nervous system will be reviewed. It is not intended to be a comprehensive disquisition, but a critical review of the thinking about the important advances and their limitations and validity. However, I have indulged in the luxury of employing the more questioning attitudes that have evolved since the epoch of the pioneers, and have quoted findings since then which are relevant. In all cases I have attempted to identify the originators of each technique, finding or theory. Some of the citations refer to findings in other tissues, but they were subsequently regarded as applicable to the nervous system. Agreement on the initiators or the first publications is not always universal.

There is a large literature on the history of the histology of the nervous system. The reader is referred to the authors listed below. However, I would like to emphasize that the critique of these findings is largely my own, and should not be taken as that of the authors cited, even when they have given permission to use illustrations from their works. The following references have been found to be invaluable: Barker, 1899; Robertson, 1899; Stieda, 1899; Singer, 1921; Garrison, 1924; Tiegs, 1927, 1929; Bourne, 1942; Baker, 1948–1953; Fulton, 1949; Brazier, 1951; Bauer, 1953; Ramon y Cajal, 1954; Glees, 1955; Baker, 1958; Hyden, 1961; Zaunick, 1961; Hamberger, 1963; Bradbury, 1967; van der Loos, 1967; Bunge, 1968; Clarke and O'Malley, 1968; Romeis, 1968; Peters, Palay and Webster, 1976; von Kirschbaum, 1977; Rottenberg and Hochberg, 1977; Niessing, Scharrer, Scharrer and Oksche, 1980; Ruska, 1980. Most of these publications were intended to be historical reviews; some of them have served this function incidentally, although they were not written with that intention in mind.

DESCRIPTION OF THE CELL

The description of cells in cork and the juices in them (Hooke, 1667) is often taken as the first description of cell walls and cytoplasm. The understanding that the intra-cellular phase was a liquid was reinforced by the more firmly based observations of Leeuwenhoek (1719) and Fontana (1784) that the axoplasm appeared fluid after it had been cut transversely (Figure 2).

The generalization of the cell theory to plants and animals formulated by Schleiden and Schwann (1838, 1847) was recognized as applicable to nervous tissue but Schwann (his

LANDMARKS IN CELL BIOLOGY OF NERVOUS SYSTEM

page 229) was mistaken about the structure of neurons. He drew them located within the axons, and he was evidently more interested in, and knowledgeable about, peripheral nerves and ganglia than about cerebral neurons.

FIXATION AND CUTTING SECTIONS

The microtome was invented for sectioning wood by Cummings before 1770 (see Hill, 1770), but, of course, wood is a hard tissue, and distorts less than soft tissue when cut. Sectioning of nervous tissue was not very satisfactory until Reil (1809) fixed samples of brain by immersing them in alcohol for several days. He observed that this procedure diminished the volume of the tissue by a quarter, and that small pieces had to be used to permit penetration. Hannover (1840) fixed brain and spinal cord with chromic acid, and noted that their gross appearances were not changed by the reagent. (The cellular structure of the nervous system had not yet been examined.) The microtome was much improved and made suitable for soft animal tissues by Von Gudden (1875).




FIGURE 2 A cross-section of a peripheral nerve of cow or sheep (from Leeuwenhoek, 1719). This is probably the first illustration of a myelinated nerve fibre

ACHROMATIC OBJECTIVES

The great era of light microscopy which lasted more than 100 years - and for whose return we must fervently strive - began with the development of the achromatic objective by Lister (1830). By combining lenses of two kinds of glass he managed to diminish spherical and chromatic aberrations to a degree which made the microscope a useful and accurate device.

EARLY FINDINGS IN NEUROHISTOLOGY

Ehrenberg (1833) and Valentin (1836) described neurons, nuclei and nucleoli (Figure

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

3), and Remak (1838) saw fibres originating from nerve cell bodies. At about the same time, Purkinje (1838) (Figure 4) and later Deiters (1865) were cutting pieces of cerebrum and cerebellum and dissecting out neurons and nerve fibres by hand (Figure 5). This was an important advance in that whole cells could be isolated and drawn. Incidentally, it should be noted that Purkinje did not report knobs on the cell surface, although Deiters did. Parallel with this development, other histologists were cutting sections and attempting to elucidate the three-dimensional appearance of cells by examining them.

The difficulty of such reconstructions was appreciated by Stilling and Wallach (1842), who conceived the idea of cutting serial sections of the spinal cord. With the benefit of hindsight, one must say that those nowadays who extrapolate simply from the two-dimensional histological sections to the three dimensions of life, often fail to appreciate fully the consequences of the shrinkage and distortion of the cells which has already occurred during the preparation of histological slides (Baker, 1958).

NEUROGLIA DESCRIBED

Virchow (1854, 1856) made what must be regarded as the most significant histological observation about the tissue adjacent to the ventricles. It 'becomes continuous with the interstitial matter, the real cement, which binds the nervous elements together, and that in all its properties, it constitutes a tissue different from the other forms of connective tissue; [this] has induced me to give it a new name, that of the neuroglia [nerve cement]' (Figure 6).




FIGURE 3 Drawing from Valentin (1836) of a cerebellar neuron. His nomenclature was a, parenchyma; b, tail-like appendage; c, nucleus; and d, nucleolus. The latter arrow may indicate that he observed a nucleolar membrane. It should be noted that he did not observe a difference in the appearance between the nucleoplasm and the cytoplasm.



FIGURE 4 J.E. Purkinje, who was probably the first person to dissect the nervous system by hand. This is reproduced from Jan Purkyne. *Purkyne-Symposion. Nova Acta Leopoldina. Johann Ambrosius Barth Verlag: Leipzig. Page 5.* 1961




FIGURE 5 Deiters' (1865) drawings of isolated nerve cells. He has not drawn granules (synapses) on them. Please compare with Figure 44

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM




FIGURE 6 Drawing by Virchow (1871) of brain adjacent to the ependyma, E. The fibres are indicated as N, and the tissue between them and the amylaceous corpuscles (ca) is the neuroglia. This illustration is reproduced from Virchow, R. (1971). Cellular Pathology (Dover Publications, New York), page 313, by kind permission of the publishers

His illustration (Figure 94 of Virchow, 1863, republished in 1971, page 313) is reproduced here (Figure 6). Virchow saw a number of concentric bodies, which he thought were starch granules, as well as a number of nuclei. The 'corpora amylacea' may have been neurons observed under suboptimal optical conditions. The nuclei were probably identical with what were subsequently called 'Deiters' cells'; they were also seen by Nissl (1899), and they may also be the cells which del Rio Hortega (1919a) called microglia, and I will call 'naked nuclei' in this monograph (please see page 189).

STAINING PROCEDURES

Gerlach (1858) introduced carmine for staining tissues, and Golgi (1879a,b, 1886) devised his classical bichromate and silver staining procedure which was subsequently modified by Cox (1891). Improvements of the microscope (Clay and Court, 1932; Bradbury, 1967) and the development of the aniline dye industry in the late nineteenth century and early twentieth century, especially in German-speaking central Europe, led to an enormous multiplication of staining procedures (Bolles-Lee, 1885; Sharpey-Schafer, 1885; Bohm and Davidoff, 1895; Mann, 1902; Carleton, 1926; McClung, 1928). Most procedures for neurons, neuroglia and fibres involved depositing on them salts of heavy metals such as silver, gold and osmium, although some of them coloured other elements of the tissues at the same time (Golgi, 1886; Bielschowsky, 1904; Ramon y Cajal, 1913a; Glees, 1946).

CONTROL EXPERIMENTS

The concept of control observations was first used by Darwin (1875) in a study of the bladders of the plant species *Utricularia*, but it did not gain popular currency among experimenters at the time. Hankin (1890) used unimmunized control mice in a study of the effects of tetanus. I would venture to suggest that the empirical nature of the

LANDMARKS IN CELL BIOLOGY OF NERVOUS SYSTEM

relationships between clinical medicine, gross pathology and histopathology, conditioned a philosophy in which control experiments were not thought necessary. Histochemistry has been much less complacent in this respect (Barka and Anderson, 1963; Adams, 1965; Pearse, 1972, 1980), but, unfortunately, to date control experiments have been carried out far too infrequently, in experimental histochemistry (Hillman, 1972, page 52). The considerable improvement of experimental design resulting from Fisher's pioneering work in applied statistics (1935), has until recently failed to make sufficient impact on histochemistry, histology or electron microscopy. This probably can be attributed to the considerable number of observations and analyses which would have to be made in small pieces of tissue to achieve statistically significant results.

BRAIN AS A SYNCYTIUM

In 1854 Virchow wrote a paper whose title in English was 'On a substance presenting the chemical reaction of cellulose found in the brain and spinal cord of man'. Golgi (1875), using his newly described method in 'thick' sections of brain, saw a nerve net. Weigert (1895) showed an apparently random arrangement of interstitial fibres in the neuroglia. There were three senses in which the brain was described as a syncytium. Gerlach (1872) regarded the neurons to be a continuous reticulum (please see below), and this would be a syncytium. Many other authors regarded the syncytium to be present in neuroglial 'nerve-glue' in the sense of Virchow (Virchow, 1846, 1871; Hardesty, 1904; Held, 1909; Holzer, 1923). A third view, whose last major exponent was Bauer (1953), regarded the syncytium as being continuous through the neuron and the neuroglial material (Figure 7 is reproduced from Bauer, op. cit.) The whole idea that there might be a syncytium was vehemently and widely opposed by the majority of neurohistologists (Taft and Ludlum, 1929; Penfield, 1932; Glees, 1955; most authors in Windle, 1958). Both Virchow (1846) and Hess (1953) examined the chemistry of the 'ground-substance'.




FIGURE 7 Bauer's (1953) view of the syncytium of the nervous system. He believed that the fibrils in the cytoplasm were continuous with those outside the cell bodies. This concept of the syncytium is very different from the present one (please see page 11 and Figure 78). This figure is reproduced from Bauer, K.F. (1953) *Organisation des Nervengewebes und Neureencythiumtheorie*, Urban and Schwarzenberg, Munchen, by kind permission of the publishers

THE RETICULAR VERSUS NEURONAL DOCTRINE

On the basis of observations using his gold chloride staining method Gerlach (1872) put forward the reticular theory, which united the whole central nervous system by fibres joining all its elements together. Golgi (1883, 1959) also accepted this view, and did not conceive the central role generally believed nowadays to be played by the neuron and all its processes. The reticularist view was questioned by observations of His (1887, 1888) and Forel (1887), and the neuronal doctrine was formulated by Waldeyer-Hartz (1891), although his contribution to the concept was relatively little. They believed that each neuron with its axon and dendrites was autonomous and had no direct connection with any other. Ramon y Cajal in 1933 (translated in 1954) summed up his views in a strongly worded book, entitled 'Neuron Theory or Reticular Theory', which is generally regarded as the definitive summary of the current view. On pages 8-9, he wrote:

We have made careful investigations of the course and connections of nerve fibres in the cerebral and cerebellar convolutions of man, monkey, dog, etc., and we have never seen an anastomosis between the ramifications of two different protoplasmic expansions, nor have we observed them between filaments emanating

LANDMARKS IN CELL BIOLOGY OF NERVOUS SYSTEM

from the same expansions of Deiters. The fibres intermingle in a most complex manner, producing a thick and intricate plexus, but never a net. It could be said that each element is an absolutely autonomous physiological canton.

Ramon y Cajal indicated that this statement was a summary of his views gathered from observations over a period of 50 years (Ramon y Cajal, 1954). This view was frequently used as an argument for the existence of synapses - that is - neurons are in contact, but not in continuity. I believe that Ramon y Cajal meant what he said, literally, that there are very few connections in the mammalian central nervous system which are visible by the light microscope.

AXONS AND DENDRITES

Deiters (1865) differentiated between the single axon and the several much finer dendrites (Figure 5), a view which is generally accepted nowadays. Golgi (1883) classified neurons into a Type I with a long axon, and a Type II with a short axon. He does not seem to have examined serial sections, nor does he seem to have appreciated that it is difficult to know with certainty the length of an axon from a single section.

Peters, Palay and Webster (1976, page 10) gave thirteen different criteria for distinguishing between axons and dendrites. For example, the axon extends either from the cell body or dendrite, while the dendrite extends from the cell body; the axon is unique in most cells but there are some examples of multiple origin, while the dendrite is usually multiple; the axon may be myelinated or unmyelinated, but the dendrite is rarely myelinated, and - if so - thinly; the axon usually gives rise to branches of the same diameter as the parent stem, but the dendrites usually divide into branches smaller than the parent stem. The only criterion which Peters, Palay and Webster indicated as absolute was that the axons extend outside the central nervous system, which the dendrites do not.

MITOCHONDRIA

Mitochondria were originally seen by La Vallette St George (1867) but Altmann's staining method using picric acid, and his clear illustrations (1890) led to them being called 'Altmann's granules' for a long period subsequently. They were probably seen in neurons for the first time by Held (1897) and Alzheimer (1904 and 1910). The cristae were first seen by electron microscopy by Palade (1953) and Sjostrand (1953a). However, no comment was made at the time on the fact that the cristae appear in the mitochondria of neurons and most other tissues far more frequently in the same plane of section than geometry would permit; see, for example, the latter authors and others (Rhodin, 1975; Peters, Palay and Webster, 1976, page 89; Fawcett, 1981, page 456). Perhaps the failure to notice the inadequacy of the geometry of the cristae arose from the beauty of the mental image of the oxidative enzymes situated in rows along the cristae.

GOLGI BODY

In the early 1890s Golgi described a 'reticular' apparatus in neurons of the barn owl

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

(Figure 8) and then in other cells (Golgi, 1898, 1899). Other authors later described the Golgi 'body', 'apparatus', or 'complex' in neurons as a shell of granules around the nucleus (Holmgren, 1900), or as a single paranuclear mass (Bourne, 1942). There was much discussion about whether the Golgi apparatus was an artifact or not. Walker and Allen (1927) and Walker (1828) made deposits of inorganic materials, which looked like Golgi apparatuses under the microscope, when they had been similarly prepared. Baker (1942) showed a silver grain under the electron microscope which looked disarmingly like a Golgi body. A full symposium of the Royal Microscopic Society in 1955, whose papers were published in 1956, debated its reality. However, its existence came to be accepted generally when the electron microscopists saw a lamellar structure throughout all tissues, which they identified as the Golgi body (Palay, 1958b; Novikoff and Goldfischer, 1961; Beams and Kessel, 1968). Unfortunately, no-one had compared the dimensions or appearances of, on the one hand, the rete diffusa, the shell and the paranuclear mass, with, on the other hand, the shape of a piece of an onion (please see Hillman and Sartory, 1980b).

Before lysosomes were described, research workers generally identified any particles in the cytoplasm, which were evidently not mitochondria, as Golgi bodies, or one of the more than a hundred other names used to describe the structure (Kirkman and Severinghaus, 1938). The relationship of the Golgi body to the endoplasmic reticulum or any other element of the 'cytoskeleton' remains to be clarified.




FIGURE 8 Golgi's view of the Golgi apparatus (1883) in a spinal neuron

NEUROMUSCULAR JUNCTION

The neuromuscular junction was first seen by Kuhne (1862). Since the interest in transmission, it has been classed with the synapse as a junctional tissue. This has enabled much which has been written about the structure and physiology of the neuromuscular junction to be taken as descriptions of the properties of synapses.

SYNAPSES

'End-feet' were seen in human neurons by Held (1897), and in rabbit spinal cord by Auerbach (1898). Illustrations from their papers show granules easily visible by light microscopy but some distance from the neuron somas (Figure 9). At about the same time, Sherrington's study of reflexes led him to postulate connections, which he called 'synapses' (Sherrington, 1897). It was not long before the events and properties which the physiologists described were ascribed to the 'end-feet' or 'end-knobs'.

When the electron microscopists started looking at the mammalian central nervous system, they soon saw 'pre-' and 'post-' synaptic thickenings (Palay, 1956a, 1958b). Synaptic 'vesicles' were seen in the early 1950s, first in retina (Sjostrand, 1953), and subsequently in brain and spinal cord (Palade, 1954; Palay, 1954, 1956b). They appear as circular or slightly oval bodies 20-60 nm in diameter, and are remarkably uniform in their dimensions in a particular tissue (references in Table 26). They were named in frog and earthworm by de Robertis and Bennett (1955), who noted that they predominated on one side of the synapse. Indeed, it is usually said that they occur on the pre-synaptic side, although, one could say that the pre-synaptic side is identified as the one with most vesicles.

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM




FIGURE 9. Drawings of neurons from rabbit trapezoid nucleus showing silver stained end-feet, which are considered one of the first demonstrations by light microscopy of synapses. Note particularly figures 1, 2, 4, 7, 8. These illustrations are from Held (1897, his plate XI)

LANDMARKS IN CELL BIOLOGY OF NERVOUS SYSTEM

Nowadays, electron microscopists use three different criteria for identification of a synapse; firstly, it must be in the central nervous system or ganglia; secondly, it must have a pre- and post-synaptic thickening, and, thirdly, synaptic 'vesicles' must be present, mainly on the 'pre-synaptic' side. Ramon y Cajal (1954, page 11) listed 11 different kinds of synapses, his classification being based entirely on the propinquity of fine fibres from different parts of the neurons. The more recent list from the electron microscopists comprises axo-dendritic, axo-somatic, axo-axonal, dendro-dendritic, somato-dendritic, dendro-somatic, somato-axonic, dendro-axonic, and glomerular, synapses (Peters, Palay and Webster, 1976, pages 159-174); (this does not include neuromuscular junctions). It means that each element of the total neuron can synapse with every other one. The latter authors made the point (page 159) that the 'existence of dendro-axonic and somato-axonic synapses must bring about radical changes in our thoughts about the function of connections between neurons'. Using the above criteria Henrikson and Vaughn (1974) found axo-glial synapses between the 13th and 14th days after gestation in the mouse spinal cord. Similar synapses have been found in the cerebral cortex of turtle (Ebner and Colonnier, 1975).

TRANSMISSION

The idea of a chemical or electrical excitation was first put forward by Du Bois Reymond (1877), but Elliott (1904) was the first person to suggest in a Communication to the Physiological Society that adrenalin might be able to transmit excitation. Incidentally Davson and Eggleton (1968) in a footnote remark that this concept is usually wrongly quoted as having been published by Elliott in a longer paper in the same year as the Proceedings; they noted that 'Apparently he was dissuaded from publishing this suggestion in his large paper, because it was too revolutionary'. From this early idea, and some seminal experiments of Fatt and Katz (1952) on frog nerve-muscle preparations, del Castillo and Katz (1956) proposed the vesicle hypothesis, for neuro-muscular transmission (for summary please see Katz, 1969). It is interesting to note that in the latter publication 13 years after the original concept, he wrote 'I should like to introduce a working hypothesis'.

This hypothesis has gradually been accepted as being applicable to central nervous system synapses, and also as being substantial supportive evidence for interpreting electron microscopic findings about synapses and synaptic vesicles. The criteria for transmitters have been enumerated frequently (see, for example, Hillarp, Fuxe and Dahlstrom, 1966; Werman, 1966; Barchas, Akil, Elliott, Holman and Watson, 1978). The latter authors give a long list, to which Osborne (1981) has added a few to make a total of 50 (reproduced here as Table 27). The vesicle and transmitter hypothesis is dealt with on pages 226-230.

NEURONS AND NEUROGLIA

When Deiters (1865) looked at the brain, he saw neurons and 'cell equivalents'; the latter had little cytoplasm and were subsequently called Deiters' cells. Henle (1871), Boll (1874) and Weigert (1895) all observed the same skein of fibres attached to the neuroglia. However, Golgi (1873), using his own stains, was not certain that the neurons and neuroglia were different cells. In 1894 he showed that the astrocytes

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

were connected to blood vessels and that their incidence increased with age. He also thought of the intercellular substance as a syncytium, in which he noted the fibres. Stricker and Unger (1879) made the point that there were various transition forms between neurons and neuroglia, perhaps implying, somewhat obliquely, that one cannot distinguish between the neurons and the neuroglia unequivocally (please see Tables 2-5).

Deiters (1865) described the microscopic appearance of neurons in detail (reproduced here as Figure 5). Cohnheim (1872) also saw the Deiters' cells without membrane or cytoplasm, and these free neuroglia were considered for many years to be important vehicles for gliosis. However, Golgi (1885), Weigert (1895) and Ramon y Cajal (1893a,b) said that these free nuclei did, indeed, have processes, which earlier staining procedures had not shown up. This then became generally accepted, as it is still today. However, it does leave open a number of important questions. How does one know that the membranes and cytoplasm are present around the naked nuclei if one cannot see them? How does one know that Deiters was staining the same kind of cells as Golgi, Weigert and Ramon y Cajal? What relationship do the nuclei seen by Virchow (Figure 6) or by Deiters, or in any cursory examination of any healthy part of the central nervous system (Figures 36 and 60) bear to the neuroglia which occur in acute or chronic gliosis in infection and injury? Are they the same cells from which glioblastomas and gliomas arise? What relationship, if any, is there between Deiters' nuclei and the amoeboid glia of Alzheimer (1904 and 1910)?

Although neuroglia had been seen and their classification had been debated earlier, it was to Ramon y Cajal (1892) that we largely owe the current classification of neuroglia - star-shaped cells with processes (astrocytes), adendritic cells (oligodendroglia) and those with elongate nuclei (microglia). Further detailed descriptions, especially of the oligodendroglia and microglia, were given by del Rio-Hortega (1919a,b, 1920). The latter author, like Alzheimer before him (1904 and 1910) and Dodgson (1948) after him, suggested that tissue around neurons could move and was elastic.

NISSSL BODY

In the late nineteenth century the Nissl body was described by Key and Retzius (1876) and by Flemming (1882), but it was shown up much more clearly by Nissl, using his own staining system (Nissl, 1894). It then came to be named after him. Hyden (1961), in reviewing the Nissl body, concluded that it was identical with the 'endoplasmic reticulum' (his inverted commas). He then goes on to make the point that

'one must bear in mind that these discrete particles in membranes are observed in fixed, alcohol-treated, and embedded material. The pertinent question, therefore, is whether they really exist and have their counterpart in the living cell. In view of existing studies on the nerve cells there is reason to regard with caution such small structures at high resolution' (Hyden, 1961).

GRANULES

By light microscopy one may see granules. If they appear in the nucleoli they are called nucleololi or nucleolonema (Estable and Sotelo, 1951). In the nucleoplasm granules would

be called nucleoli, specks of dirt or artifacts; in the cytoplasm they would be called Golgi bodies or centromeres by older workers and lysosomes or ribosomes by modern electron microscopists; in the synapses they would be called synaptic vesicles, dense core vesicles, endocytotic or exocytotic vesicles. They all look like granules of different sizes (see Figure 10).




FIGURE 10 Drops of egg albumin were smeared and stained with haematoxylin and eosin. Please note the non-uniformity of the deposit, which could appear as showing fibres, granules, reticula and cytoskeletons. The bar is 20 µm.

MYELINATED FIBRES

In 1717 Leeuwenhoek (see 1719) saw myelinated fibres (Figure 2), but they were more clearly described by Fontana (1781). Remak (1836), and subsequently Koelliker, Dalton and Todd, obviously saw the node (Stirling, 1881), but it fell to Ranvier (1871) to give the definitive description, and consequently his name to the node of Ranvier. Incisures in the peripheral nerves were seen by Schmidt (1874) and Lantermann (1877), whose names were also given to them. Although they can be seen in unfixed peripheral nerves by light microscopy, their existence was not regarded as 'confirmed', until the electron microscopists also saw them in peripheral nerve and spinal cord (Blakemore, 1969; Hildebrand, 1971 and sources quoted there).

The history and appearance of the myelin sheath as revealed before electron microscopy was reviewed by Young (1945). The myelin sheath was examined by low-angle diffraction (Schmitt, Bear and Clark, 1935; Schmidt, 1936, Schmitt and Bear, 1939),

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

and its lamellar structure was described by electron microscopy in the whole nerve (Fernandez-Moran, 1950b; Rozsa, Morgan, Szent-Gyorgyi and Wyckoff, 1950; Sjostrand, 1953a); the same lamellar structure was also seen in homogenates (Fernandez-Moran, 1950a; Sjostrand, 1950). The beautiful concentric appearance is the pride of the literature (for example, Rhodin, 1975; Landon, 1976; Peters, Palay and Webster, 1976, chapter VI, pages 181-230).

THE SCHWANN CELL

This was described by Schwann (1838), and his illustration is reproduced here in Figure 11. From studies of chick embryo, Geren (1954) described how the Schwann cell wraps around the axon to form the myelinated sheath. The central end of the sheath has been called the internal mesaxon, and the peripheral 'entry' point of the axon, the external mesaxon (see, for example, Robertson, 1959, 1969, 1980; Fawcett, 1981, and Figure 71 from the latter book).

NUCLEOLI AND NUCLEOLAR MEMBRANE

Valentin (1836) drew a nucleolus which looked as if it had a membrane around it and was empty (reproduced as Figure 3). He described the nucleus as containing in its centre a 'circular corpuscle', which was both clear and solid (this seems somewhat contradictory). Purkinje (1838) and Deiters (1865) also showed clear nucleoli. 'Nucleolus-associated chromatin' was seen in neurons by Hyden and Hamberger (1945) and Hertl (1957). The nucleolonema was first seen by Estable and Sotelo (1951). In unfixed rabbit neurons it has been shown to be in continual movement, which was stopped by the addition of phenol (Sartory, Fasham and Hillman, 1971). However, the nucleolonema is not seen as well by electron microscopy as it is by phase-contrast (compare Figures 31-33 and 16, 18 and 57), so it is rarely described in modern textbooks. A nucleolar membrane was described in all unfixed neurons of the cranial nerve nuclei and anterior horns of the spinal cord of rabbit, rat and guinea pig (Hussain, Hillman and Sartory, 1974).



FIGURE 11 The myelinated axon as drawn by Schwann (1838). Please note the position of the nucleus

THE NUCLEOLOLUS

Mauthner (1860) first saw a small body in the nucleolus, which he called the nucleolulus. Some authors since then have used the term for one or more minute granules, while others have used it for the appearance of a vacuole (Shantha, Manocha, Bourne and

LANDMARKS IN CELL BIOLOGY OF NERVOUS SYSTEM

Ariens-Kappers, 1969, and authors quoted there). It is difficult to conceive that the appearances of a granule and a vacuole could represent the same object, unless the former were viewed out of focus. It is doubtful whether serious microscopists could make the latter error. The nucleolulus has been seen by both light and electron microscopy (Dutta, Siegesmund and Fox, 1963), but it has not been sighted in the textbooks recently.

CYTOARCHITECTONICS AND MYELOARCHITECTONICS

Cytoarchitectonics is a term used for the types and arrangements of cells in the central nervous system, while myeloarchitectonics refers to the fibres within it. It is generally assumed, implicitly rather than explicitly, that the cells are neurons, and the fibres arise from the neurons. It has also been generally assumed that cells which stain with 'neuroglial' procedures are not excitable, and, therefore, their processes do not conduct action potentials.

Meynert (1867, 1868), Bevan-Lewis and Clarke (1878) and Hammarberg (1895) made systematic histological studies of normal and diseased brains. However, the demonstrations by Fritsch and Hitzig (1870), and Ferrier (1876) that stimulation of particular regions of the parietal cortex could produce movement of limbs, generated an interest in the relationship between the histology of the brain and its physiology. In the early twentieth century the cellular and fibrous arrangements of the brain were recognized as having a pattern. Brodmann (1903, 1909), the Vogts (1903, 1919) and Campbell (1905) published extensive studies of the cellular pattern of the cerebral cortex, including descriptions of the five to eight laminae. The identification, relative thickness, and physiological significance of the layers, were the subject of much discussion. Further anatomical studies were pursued by Ramon y Cajal (1909-1911), von Economo and Koskinas (1925) Von Economo, (1929) and many others; they led to the current view that the cortices of the parietal, temporal and occipital regions have an external lamina, consisting of: I, plexiform; II, small pyramid; III, medium-sized pyramid; IVa, star pyramid; and IVb, star cell, layers; and an internal lamina, consisting of: V, large deep pyramids, and VI, spindle, layers. The four main types of cell seen are: those with long descending axons; those with short axons ramifying near the cell body; cells with ascending axons ramifying in cortical layers; and cells with horizontal axons. Brodmann (1903) carried out the monumental task of mapping 52 areas of the brain, while Campbell (1905) looked at 20 areas.

Both in respect of the cortical layers and of the cell types, authors have no difficulty in indicating 'typical' examples in most histological preparations, but there are many cells - perhaps the majority - which are difficult to identify unequivocally and objectively. Some of this difficulty probably arises from the fact that the thickness of a single section cuts only a fraction of the diameter of a cell soma, and an even smaller fraction of the length of a process.

WHITE MATTER

According to Clarke and O'Malley (1968, page 586), De Vieussens (1684) was the first to examine the white matter; he described it as 'composed of innumerable connected fibres divided up into many bundles so close to one another that there is no perceptible

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

space between them, [and] compose a continuous body, that is, the staff.' Von Gudden (1874) devised the technique of cutting the optic nerve or other cranial nerve to observe the degeneration of the fibres and thus map the tracts, using serial sections. Flechsig (1876, 1881) studied myelination in brain and spinal cord as another method of examining the courses of the tracts. The development of the Weigert (1882) and Marchi and Algeri stains (1885-1886) for degenerating fibres constituted another important histological technique for mapping tracts (Schroder, 1914; Minkowski, 1923; Von Economo and Koskinas, 1925). In discussing 'anatomical' techniques, one should mention the Labanic work of Lassek (1948) in classifying a million fibres and counting those of the different diameters in the human pyramidal tract (please see also Lorente de No, 1934).

A number of workers have counted the number of muscle fibres which are innervated by a single nerve fibre and presumably an anterior horn cell. The ventral roots were cut to allow the sensory fibres to degenerate, so that the remaining fibres were all motor. The following ratios of nerve to muscle fibres were found: soleus, 1:120; extensor longus digitorum 1:165 (Clarke, 1931); sartorius 1:70-117 (Van Harreveld, 1947). Of course, each muscle fibre contains a considerable number of sarcomeres in series, some of them several millimetres away from the end-plates. This multiplicity of sarcomeres served by a single nerve fibre has considerable implications for the anatomy and physiology of the neuromuscular junction (please see pages 156-157, 209, 226-230).

PHYSIOLOGICAL EXPERIMENTS ON TRACTS

Following Fritsch and Hitzig (1870) many authors stimulated cortical areas or extirpated them in a variety of species, including monkey and man (Hitzig, 1874; Ferrier, 1876; Beevor and Horsley, 1890a,b; Bidwell and Sherrington, 1893; Sherrington, 1906; Leyton and Sherrington, 1917). Dusser de Barenne (1910, 1916) applied strychnine to the cortex and recorded the spread of excitability, having reasoned that any region away from the area stimulated, which reacted, must have fibrous connections with it.

The idea that localized parts of the brain had particular 'functions' probably originated from the physiognomists who were active well into the beginning of the twentieth century. It was confirmed, of course, by the experiments on mapping of tracts, electrical and strychnine stimulation, animal ablation experiments, and the surgical excision of epileptic foci by Penfield and his collaborators (Penfield and Erickson, 1941). Yet the very important experiments of Goltz (1881), the arch opponent of localization, should not be overlooked. He showed that decorticated dogs could survive months with minimal motor disturbance (Goltz, 1892). Pathologists know that babies who have had unilateral hemispherectomies subsequently grow up with a remarkable degree of cerebral function. Sometimes, also, patients who have had large volumes of their brains replaced by metastases from tumours in their lungs or stomachs have unexpectedly good neurological and psychological function. This is attributed to 'plasticity', but this really is only another expression for the same phenomenon. Furthermore, the concept of 'centres', demonstrated by experiments showing that particular parts of the brain may be stimulated or extirpated to increase or decrease their activity, is somewhat naive. Such experiments only show that the apparent centre is part of the circuit involved in the motor or sensory response, not that it is

LANDMARKS IN CELL BIOLOGY OF NERVOUS SYSTEM

necessarily the 'centre', nor even that a 'centre' exists for that activity.

MICROSCOPICAL TECHNIQUES

The nineteenth century was an epoch of unprecedented developments in the following aspects of microscopy relevant to the study of cerebral histology: tissue fixation; mounting of slides; illumination systems; microtomy; neural stains; photomicrography; micromanipulation; measurement of dimensions of microscopical objects; illustration; manufacture of model systems illustrating the properties of living cells (colpoids); microscope design; surgical pathology; vital staining; polarizing and dark-field microscopy; control experiments; microspectrography; study of artifacts. In the early twentieth century the following further techniques became practical and common: intracellular injection; tissue culture; warm stages; staining of degenerating fibres; autoradiography; biopsy; cinemicrography; tissue transplantation; microsurgery; histochemistry; time lapse photography; immunocytoLOGY; development of microchemical techniques. Several new kinds of microscopy were developed; inverted; X-ray; reflection; phase contrast; anopteral; interference; ultraviolet; cryomicroscopy; automatic photography; electron microscopy; television; acoustic (Hillman and Richards, 1983).

TISSUE CULTURE

Harrison (1907, 1910) was the first person to culture nervous tissue *in vitro*. He showed that fibrils from frog embryo would grow on clotted lymph. Lewis and Lewis (1915) and Lewis (1923, 1931) observed mitochondria, granules and pinocytosis, and Lewis and McCoy (1922) examined the survival in culture of tissues from rats, guinea pigs, rabbit and man. Weiss and Wang (1936) described fibres in the cytoplasm of chick embryo ganglion cells. Hogue (1950) grew human foetal brain cells from dead embryos, and was able to identify the different kinds of cells in culture.

Phase-contrast microscopy was described by Zernike in 1934. Pomerat and Costero (1956) performed epoch-making observations on intracellular movements using time-lapse photography and phase-contrast microscopy on cat cerebellar cultures. It has also been shown that neurons in culture can fire (Crain, 1956; Hild and Tasaki, 1962). Until 1964, Hild believed that neurons lost their synapses in tissue culture, presumably because he could not see them by light microscopy. However, when he and his colleague (Callas and Hild, 1964) used the electron microscope, they saw membrane thickenings which they regarded as synapses.

Hild, Chang and Tasaki (1958) found electrical responses in cultured cat cells, which they thought were astrocytes, and which were quite different from the rapid action potentials of the cells they identified as neurons (Hild and Tasaki, 1962). Myelination of axons in culture was shown (Peterson and Murray, 1956; Hild, 1957). An explant from the brain stem was demonstrated to join up with one from the spinal cord (Crain, Peterson and Bornstein, 1968), and the same school showed that nerves can activate muscles in culture (Peterson and Crain, 1970).

Intracellular movements seen in tissue culture seem to be incompatible with an 'endoskeleton'. A suggestion has been made that there are 'roads' through the 'endoskeleton', which permit such traffic. Such traffic lanes have been sought by optimistic voyagers (Andres, 1961; Bunge, Bunge, Peterson and Murray, 1967).

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

What should one conclude if intrepid explorers cannot find a way out? (Please see page 198-200).

CELLULAR AND INTRACELLULAR MOVEMENTS

Metchnikoff (1893) described phagocytosis and Nissl (1899) saw microglia which seemed so similar to cells seen around infections of the brain that he supposed that they were motile. Del Rio-Hortega (1919b) was also of the opinion that microglia could move.

The migration of neuroglia in developing embryos was concluded to be occurring after examination of brains of embryos at various stages (Kershman, 1938, 1939). Axonal flow in both directions of proteins, amino acids, nucleotides and transmitter-related enzymes, and its arrest by obstruction of the axons, was described by Weiss and Hiscoe (1948), and has since been studied extensively (Weiss, 1961; Lubinska, 1964; Grafstein and Forman, 1980). Leonhardt (1951) injected particles into the brain, and found them subpially subsequently. Considerable work showing intracerebral movements of granules and cells has been done by Klatzo and his colleagues (Klatzo, Piroux and Laskowski, 1958; Klatzo and Miquel, 1960). Oehmichen has more recently published a review of his own experiments on cellular movements in the nervous system (Oehmichen, 1978).

THE CELL MEMBRANE

Cells have been recognized since the time of Hooke by the walls which surround them, and the chemistry of this wall has fascinated physiologists and biochemists for more than 90 years. Overton (1895, 1902) first suggested that the cell surface was lipid. Gorter and Grendel (1925) extracted the lipids from red blood cells of several species and calculated that there would be enough to make a bimolecular layer around each cell. Fricke (1925), Fricke and Curtis (1835), and Cole (1937), measured the capacity and resistance of several eggs, and Schmitt (1936, 1939) and Schmitt and Bear (1939) examined red cells by birefringence, and concluded that membranes were 1-10 nm thick. Davson and Danielli (1943) reviewed a great deal of evidence of: selective permeability of membranes to ions; oil-water partition coefficients of anaesthetics; high capacitances and resistances of membranes; birefringence studies; chemistry of the membranes, their surface tension and elastic properties, etc.; they then proposed their hypothesis that the cell membrane consisted of two protein molecules sandwiching a lipid layer (Figure 12).

A few comments about this seminal hypothesis should be made. Among the measurements of membrane thickness, the electrical measurements gave maximum values, while the birefringence measurements gave minimal values; all of these had a large range and most of the measurements gave larger values than the simple Davson-Danielli hypothesis would suggest. Most values were derived from membranes of plant cells, erythrocytes and marine eggs, but their results have generally been regarded as applicable to all membranes. Many of the measurements adduced were made on hydrated tissues (Davson and Danielli, 1936; Danielli, 1942). However, measurements from electron micrographs and low angle diffraction - particularly of myelin sheaths - usually made on completely dehydrated tissues, were used to support the Davson-Danielli model, without taking any shrinkage into account.

Robertson (1959) described the 'unit membrane', also known as the 'trilaminar'

LANDMARKS IN CELL BIOLOGY OF NERVOUS SYSTEM

appearance. It was originally thought of as the characteristic appearance of the cell membrane, but, subsequently, it was generalized to include the nuclear membrane, the endoplasmic reticulum, the myelin sheath, and occasionally the lamellae of the Golgi apparatus.




FIGURE 12 Diagram of the 'lipoid double-layer' as proposed by Davson and Danielli (1936). This figure is reproduced from Davson, H. and Danielli, J.F. (1936). *The Permeability of Natural Membranes* (Cambridge University Press, Cambridge), by kind permission of the authors and publisher

The fluid mosaic hypothesis of membrane structure was first put forward by Singer and Nicholson (1972) and is now widely accepted as if it were a finding (Semenza and Carafoli, 1977; Repke, 1980; Kates and Kuksis, 1980). It originated from some thermodynamic considerations, and studies on the distribution of antigens applied to red cell membranes and detected by freeze etching (Figure 13).

ELECTRON MICROSCOPY

The first electron microscope was made by Knoll and Ruska (1932), and the instrument was used to examine biological specimens not long after (Driest and Muller, 1935). During the three decades after the Second World War, neural tissues were the subject of intensive examination. The following structures were detected by the authors indicated in parentheses: endoplasmic reticulum (Porter, Claude and Fullam, 1945); cristae (Sjöstrand, 1953a); ribosomes (Palay and Palade, 1955); 'unit membrane' (Robertson, 1959); lysosomes (de Duve, 1963); neurofilaments (Palay, 1964); microtubules (Porter, 1966); microtrabeculae (Wolosewick and Porter, 1976). The existence of the Golgi body, hitherto seen by light microscopy, was regarded as being 'confirmed', when a body which could be seen in the cytoplasm by electron microscopy was given the same name (Beams, van Breeman, Newfang and Evans, 1952). A 'cytoskeleton' which is claimed to consist of the following elements, whose mean diameters (nm) are given, has been reported: microtrabeculae, 3-6 (Wolosewick and Porter, 1979); actin filaments, 4-6 (Fine and Bray, 1971; Puszkin and Berl, 1972); glial filaments, 7-8 (Wuerker, 1970); neurofilaments, 10 (Palay, 1964; Wuerker and Kirkpatrick, 1972);

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

endoplasmic reticulum, 10 (Porter, Claude and Fullam, 1945); microtubules, 20-26 (Elfvin, 1961); Golgi body, 500-1000 (Hillman and Sartory, 1980a, references indicated on page 83). For reviews, see Wuerker and Kirkpatrick (1972), Shelanski (1973), Lazarides (1980), International Congress (1981).

The original techniques of detection of fluorescence (Williams and Chase, 1967-1977; Nairn, 1969; Shelanski, 1973; Weir, 1973; Sternberger, 1974) have been adapted for use with antibodies, for immunocytochemistry (Franke, Schmid, Osborn and Weber, 1978; Cold Spring Harbor Symposium, 1982). The antibodies are said to be specific, but are sometimes cross-reactive. The various elements of the 'endoskeleton' are viewed by light microscopy, usually in tissue cultures. The cells may have to be 'permeabilized', in addition to being fixed, frozen or stained. The dimensions of the elements of the 'endoskeleton' are measured by electron microscopy although during the preparation much of the lipid is lost.




FIGURE 13 Diagram of the 'fluid mosaic' model of the cell membrane, according to Singer and Nicholson. The solid bodies with stippled surfaces 'represent the globular integral proteins, which at long range are randomly distributed in the plane of the membrane'. This diagram is reproduced from Singer, S.J. and Nicholson, G. (1972). The fluid mosaic model of the structure of cell membranes. *Science*, 175, 720-731, by kind permission of the authors and the American Association for the Advancement of Science

The electron microscope has also been used to detect the nuclear pores or pore apparatuses (Callan and Tomlin, 1950; Hartmann, 1953; Palay and Palade, 1955; Fernandez-Moran, 1957; Franke, 1974).

There have been many important developments in the techniques of electron microscopy. Among these are freeze etching (Meryman and Kafig, 1955; Steere, 1957; Moor, 1969), high-voltage electron microscopy (Dupouy, 1972) and others (for reviews, see Williams and Chase, 1967-1977; Meek, 1976; Stolinski and Breathnach, 1975; Hayat, 1970-1978; Weakley, 1981). All the latter techniques involve dehydration, but Parsons (1974, 1978) has attempted to examine tissues in wet chambers, to obviate the shrinkage. Unfortunately the pressure in his electron microscope is well below the vapour pressure of the water at that temperature so it is difficult to see how the specimens remain fully hydrated.

2

Methods of Studying Neural Structure

The following techniques are used to study the cellular structure of the nervous system:

- (a) microdissection of unfixed tissues;
- (b) light microscopic histology;
- (c) observation of tissue cultures;
- (d) electron microscopy;
- (e) subcellular fractionation.

MICRODISSECTION

Microdissection of the mammalian central nervous system was originally pioneered by Purkinje in the 1840s (Van der Loos, 1967). It was widely used subsequently for many non-neural systems (see, for example, Gray, 1931; Chambers and Chambers, 1961; McClung-Jones, 1967). However, its modern renaissance is due to Hyden (1959, 1961, 1967) (Figure 14). The neurons are dissected out by hand under direct vision from sections of brain (Figures 15 and 16). The value of this method is that neurons can be separated with comparatively little disruption, leaving the soma intact, although, of course, the dendrites and axons are severed. The tension on the processes and placing the cells in unphysiological environments undoubtedly has some effect on them. Even this comparatively simple procedure implies fourteen separate assumptions (Hillman, 1983), but most of these are relevant to biochemical measurements made in them, and probably not to their structure. It is likely that placing them in unphysiological media might alter the osmotic conditions within the cells and therefore the total and relative dimensions of their subcellular organelles. It seems unlikely to alter the shapes of any part significantly. Nevertheless, after isolation these neurons respire linearly for several hours and have resting membrane potentials whose values can be altered reversibly (Epstein and O'Connor, 1965; Hertz, 1966; Hillman and Hyden, 1965). It is not known whether they can generate action potentials, since - to the author's knowledge - no systematic effort has yet been made so far to elicit them. It would be a very worthwhile project upon which to embark, since demonstration of the excitability of these cells would help to justify the very extensive biological studies already carried out on these cells, as well as the technique of isolation.

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM



FIGURE 14 Professor Holger Hyden of Goteborg, modern pioneer of separation by hand of mammalian neurons. This photograph was taken in 1981. He is dissecting out the neurons with stainless steel wires




FIGURE 15 Dissection by hand of neurons of the cranial nerve nuclei, by the method of Hyden (1959). Rabbits have large neurons but any mammalian species can be used. The same technique can be used to separate spinal motor neurons: a, the brain is exposed and the first cut is made between the cerebrum and the cerebellum; b, the second cut is made below the medulla; c, several transverse cuts are made through the medulla with a scalpel; d, the slivers are placed on a wax block, stained slightly with methylene blue in 0.25 mol/l sucrose solution which is then washed off with more sucrose, and the stainless steel wire is used to lift out neurons of the cranial nuclei under direct vision using a dissecting microscope. The cells are then placed in a drop of sucrose solution in a cavity slide. These photographs were taken in collaboration with Dr T.S. Hussain




FIGURE 16 Isolated rabbit neuron, viewed by phase-contrast microscopy. Upper unfixed before staining with osmium tetroxide, lower after embedding in Araldite (it was not placed in the electron microscope or sectioned). The bar is 20 μm

The value of the experiments on isolated neuron cell bodies depends upon the truth of the assumption that the 0.25 mol/l sucrose or 0.15 mol/l saline in which they are dissected (Hyden, 1959; Hussain, Hillman and Sartory, 1974) does not significantly distort their shapes from those *in vivo*. There is no way of testing this. One can only plead that this technique of examining cells probably subjects them to less reagents than any other technique *in vitro*.

The relatively large dimensions of the cell bodies of mammalian cranial nerve nuclei and anterior horn cells has so far permitted a considerable accumulation of data about their structure, biology and pharmacology, which is not matched by findings about their processes or surrounding neuroglia.

In addition to neurons from the cranial nerve nuclei in the medulla, neurons can

METHODS OF STUDYING NEURAL STRUCTURE

also be isolated in large numbers from dorsal and sympathetic ganglia (Figures 17 and 18). Trypsin is often used for softening their capsules (Varon, Raiborn, Seto and Pomerat, 1963) but this powerful proteolytic enzyme may be avoided by the simple expedient of tearing open the capsules using dissecting needles (Hillman, 1966). Mammalian axons can also be teased apart with needles (Figure 19).




FIGURE 17 Human dorsal root ganglion, sectioned near the equator, stained with Palmgren's stain. The bar is 500 μm long




FIGURE 18 An unfixed rabbit dorsal root ganglion cell in normal saline viewed by phase-contrast microscopy. Please note the nucleolar membrane. The maximum diameter of the cell body is about 60 μm




FIGURE 19 Myelinated axons from sciatic nerves, teased out, unfixed, placed in 0.25 mol/l sucrose (left) from a rat by bright-field illumination, showing Schmidt-Lanterman clefts, s; (right) from a rabbit by dark ground, showing nodes of Ranvier, r. The axons are 10-15 µm in diameter

Unfixed neurons, axons and neuroglia can be examined under the best optical conditions using phase-contrast, anopteral, dark-ground, polarizing and interference, as well as bright-field microscopy (Hussain, Hillman and Sartory, 1974). It should be stressed that the simplicity of the preparation, and the paucity of the reagents to which the cells have been subjected before they are observed, make these techniques the most valuable for examining the closest reality which can be achieved to the structure of the living cell *in situ*.

It is of considerable interest that hand dissection of the brain isolates neurons and 'neuroglial clumps' (Hyden, 1959; Hyden and Pigon, 1960). Hyden regarded these clumps as containing oligodendroglia. Hamberger (1963) found in 'vascular' glia a few astrocytes (Figure 20) because some of their cells were connected to blood vessels, which has been regarded as a criterion for astrocytes historically (Ramon y Cajal, 1913).

Mammalian peripheral nerve fibres have the advantage that they can conduct action potentials after isolation which are not significantly different from those *in vivo* (Erlanger and Gasser, 1937).




FIGURE 20 Flake of tissue (left) and capillary glia (right) from the lateral vestibular nucleus of rabbit, stained with methylene blue. The dark nuclei visible in the upper figure are regarded by Professor Hamberger as being oligodendroglial, as are most of the cells visible in the capillary glia. This figure is reproduced by kind permission of the author from Hamberger, A. (1963) Differences between isolated neuronal and vascular glia with respect to respiratory activity. *Acta Physiol. Scand.* 58, Suppl. 203, pages 1-58

HISTOLOGICAL TECHNIQUES

Histology developed as a branch of pathology for the correlation of the appearances of diseased tissues with clinical findings from patients. The correlations are empirical and do not imply that the tissue has the same appearance after staining as it did before. Tissue for histology has come from a dead animal; it has been fixed, dehydrated, embedded in a non-aqueous medium, cut a few μm thick, rehydrated, stained, probably counter stained, dehydrated again, cleared and embedded in a non-aqueous mountant. The effects of these procedures on the size, shapes, textures and colour of the cells and their organelles, are not only everyday observations for histologists, but have been carefully studied over a number of years (Tellyesniczky, 1898; Stowell, 1941; Ross, 1953; Baker, 1958; Gersh, 1959; Hillman and Deutsch, 1978).

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

In a single section one cannot usually know the original dimensions of a cell, the presence, position, dimensions of any of its organelles or any of their shapes (page 51). When one has made a number of serial sections, one can measure the dimensions as seen after preparation for histology. Unfortunately, even reconstructions from serial sections do not show the dimensions of the original cells *in vivo*. The cells and all their organelles have shrunk during dehydration, not all in the same proportion (Hertwig, 1931; Lordin, Mares, Karasek and Skrivanova, 1967; Hillman, Hussain and Sartory, 1976). The simplest explanation for the unequal shrinkage of parts of cells is that they each have different contents of water and are chemically very different. The experiments quoted above showed different shrinkage of nuclei and cytoplasm only. There is little information about the shrinkage of membranes during dehydration, except for the low angle diffraction studies of Finean (1957, 1960). He interpreted the diminution of the repeating period by 2-3 nm as being due to the loss of water, although it has also been proposed that it was due to loss of neutral mucopolysaccharide (Robertson, 1959; Wolman and Hestrin-Lerner, 1960). The myelin sheath has generally been regarded as a model for mammalian membranes of all kinds (Robertson, 1959; Adams and Davison, 1965; Fernandez-Moran, 1967).

The main changes occurring after histological preparation are as follows:

1. Shrinkage of the cytoplasm causes the neuron which was rather spherical in the unfixed state to become more concave between its main dendrites.
2. The cytoplasm of the neuron which is normally a fairly translucent liquid containing a few mitochondria precipitates, and crystalline deposits and granules appear within it.
3. Brownian movement, mitochondrial movement, rotation of the neuronal nuclei, and axonal flow, stop, probably due to denaturation of cytoplasmic, axonal and nucleoplasmic proteins.
4. The nucleoplasm, which was a translucent uniform fluid in both neurons and neuroglial nuclei, precipitates and becomes non-uniform.
5. The nucleolar contents in the neurons (the neuroglial nuclei do not appear to contain nucleoli) precipitate, making the nucleolonema appear amorphous and blurring the distinction between the nucleolonema and the pars amorpha. The precipitation also obscures the nucleolar membrane seen in neurons (Hussain, Hillman and Sartory, 1974). The inability to see the nucleolonema or nucleolar membrane in stained histological sections or in electron micrographs probably explains why their existence has been almost completely ignored. This implies that information derived from the latter preparations is more true than that found by examining unfixed untreated neuronal cell bodies, when a finding is made in the latter preparation, but not found in histological sections (Hillman, 1976).
6. The mitochondria, which appear clearly in the cytoplasm of unfixed cells in tissue culture, and both inside and outside neurons in isolated neurons (Hillman, Deutsch, Allen and Sartory, 1977; Hillman and Deutsch, 1978b), are difficult to see in either of these locations in sections of tissue. The simplest explanation is that they float out of the cytoplasm from the face of the very thin slivers of tissue into some of the reagents between the stages of sectioning and of final mounting. Another possibility is that they disintegrate due to 'osmotic shock' of hypotonic solutions,

METHODS OF STUDYING NEURAL STRUCTURE

or due to chemical attack by other agents. Such misfortunes may not befall the mitochondria in a whole uncut neuron or neuroglial clump.

7. Apparently empty holes or vacuoles may appear, which did not exist in life.
8. Membranes may become closer or more separated.

Thus, on the one hand one may see loss during preparation of some free mitochondria or granules which were present in the living tissue. On the other hand one may see appearing new 'structures', which are dehydrated cytoplasm, nucleoplasm, mitochondrioplasm or extracellular fluid, or are reaction products of any of the reagents used with them, or are composed simply of the reagents themselves. One must continuously try to peer into these glazed windows of our knowledge to perceive the blurred shadows of the shy reality which is the living tissue.

With all these reservations in mind, examinations of histological sections can give valid information about:

1. The presence of pathological states in tissue.
2. The number of a particular kind of cell or subcellular component in a particular section.
3. The relative positions of the components.
4. Their shape after shrinkage - an object which is still spherical after shrinkage has probably preserved its original shape. This is not necessarily true for electron microscopy, in which the metal deposit is exposed to electron bombardment and irradiation under high vacuum, and enough energy is liberated to make particles of nanometre dimensions or less become spherical.
5. The presence or location of a particular feature of the preparation, either if it always occurs in a particular situation, or if it has been shown statistically to occur at a more significant incidence in a particular part of the cell than it is found elsewhere.

Staining with haematoxylin and eosin is probably the most popular procedure used throughout histology and pathology. It is as well to remind ourselves that this technique has 14 steps and employs 11 different reagents. Neurons were isolated by hand dissection, but were not embedded and sectioned; they were fixed, stained with haematoxylin, counter stained with haematoxylin, counter stained with eosin, and the whole cell bodies were embedded in DPX medium. These steps resulted in a shrinkage of the cytoplasm to a mean of 25%, and the nuclei to 15%, of their original unfixed areas, respectively (Hillman, Hussain and Sartory, 1976; please also see Figure 16).

Obviously, different staining systems and different reagents change tissues to different extents, and it may be assumed that the fewer the reagents which are used the more accurately the histological section reflects the structure of the living tissue. In order to make comparison of the appearances of different kinds of cells easier and to diminish the number of reagents, a different approach has been tried. Six different tissues were embedded in paraffin wax and sectioned but not stained. They were examined under phase-contrast microscopy, and their appearances compared with those of sections stained with haematoxylin and eosin and viewed by bright-field illumination (Hillman and Sjuve, 1980).

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

Of course, light microscopy in general has its repertoire of artifacts, such as chromatic aberration, spherical aberration, Arey's discs, glare, coverslip thickness corrections, tube length corrections, errors due to the refractive index of the media, etc. These are well recognized, and can usually be avoided by well-trained microscopists (Hillman, 1982b).

A much more serious source of error are the precipitates, shrinkage artifacts, tears and cracks which result from the dehydration. Among these are the Golgi apparatus (Hillman and Sartory, 1980a, pages 63-66) and the cytoskeleton (please see *ibid*, pages 43-60), as well as many apparent structures seen by electron microscopy (please see Table 17). It is appropriate to stress here two points; firstly, any solute must precipitate during dehydration if it does not diffuse away in, or is not dissolved in, the dehydrating agents; secondly, the constancy of any appearance is no evidence whatsoever, that it is not artifactual, pace many referees of journals.

It is extremely difficult to avoid some degree of subjective judgement in experimental histology. The following steps of the procedure contain subjective elements even in the hands of the most expert research workers: choice of species; choice of anaesthetic; method of killing the animal; selection of the sample; selection of the staining procedure to be used; decision on the precise duration of each step in the procedure; choice of type of microscopy, including the illumination and filters used; decision of magnification to be used to examine the tissue; decision on duration of the observations; decision on the number of observations to satisfy the histologist; choice of area to be photographed; choice of figures for submission for publication; reproduction of figures submitted for publication in the final printed form; decision on the nature of the features to be labelled; decision on which parts are to be labelled. In addition, there is some degree of subjectivity on the identification of the seven different major artifacts of light microscopy (White, 1966).

In considering the staining of neurons, astrocytes, oligodendrocytes and microglia, one has to bear in mind that the specificity of a particular stain for one of these elements rested mainly on the authority of the great histologists, such as Golgi, Cajal, Penfield, Del Rio-Hortega, Mallory and Glees. Most of them carried out their major research before the concept of control experiments had achieved the currency that it has today. These giants established the identifications of particular cells in an empirical fashion, and these have generally been accepted by neurologists and neuro-pathologists as working practices. Yet most authors nowadays, while diagnosing tumours both by the histological identification and by the neurological history of the patient, are well aware of the uncertainty of their identification of the cell types. In textbooks, each author shows an illustration of what he believes undoubtedly to be a clear and unequivocal example of a particular cell type. At the same time, there would be many cells in the brain which could not be identified certainly as being of a particular type, or about which two good histopathologists would disagree in good faith, without any objective way of deciding its identity. Thus one can draw up a table of authors who say quite specifically that one cannot distinguish clearly between any two kinds of cells by light microscopy (Table 2).

This understanding poses a considerable threat to the neurohistologists, who thought that they had been saved by the electron microscopists. Unfortunately, the

METHODS OF STUDYING NEURAL STRUCTURE

electron microscopists cannot be too much help, because they agree on the difficulty of distinguishing beyond any reasonable peradventure on the differences, particularly between the different kinds of neuroglia (Table 3). Also, they are usually using too high a magnification to examine many cells in one field.

Many authors cannot distinguish between astrocytes, oligodendrocytes and, sometimes, microglia, and admit this implicitly by using the term neuroglia, without stating to which variety they believe they are referring (Table 4).

The electron microscopists, the tissue culturists, the biochemists and the immunologists cannot help the light microscopists with the identification for a more general reason - that is - they themselves fundamentally depend upon the light microscopists to whom they refer their own preparations, or who have initially named the tissue of origin from which they took their pieces of tissue to examine or grow in culture. The different kinds of neuroglia were characterized by light microscopy more than two decades before the electron microscope was introduced to biology (Glees, 1955).

TABLE 2 Light microscopic studies in which the authors point to their inability to distinguish neurons and the different kinds of neuroglia

Cannot distinguish between neurons and astrocytes	Cannot distinguish between neurons and oligodendrocytes	Cannot distinguish between neurons and microglia
Deiters(1865)	Deiters(1865)	Deiters(1865)
Gerlach(1871-72)	Gerlach(1871-72)	Gerlach(1871-72)
Boll(1874)	Boll(1874)	Boll(1874)
Golgi(1883)	Golgi(1883)	Golgi(1883)
Spatz(1918)	Spatz(1918)	Brownson(1956)
Glees(1955)	Glees(1955)	Ramon-Moliner(1958)
Lewis(1976)	Ramon-Moliner (1958)	Brownson, Suter and Diller(1963)
Kennedy, Lisak and Raff(1980)	Kennedy, Lisak and Raff(1980)	

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

TABLE 3 Electron microscopic studies in which the authors point to their inability to distinguish between different kinds of neuroglia, or use the generic term 'neuroglia' without indicating a particular type

Cannot distinguish between astrocytes and oligodendrocytes	Cannot distinguish between astrocytes and microglia	Cannot distinguish between oligodendrocytes and microglia	'Neuroglia' used without differentiation into types
Luse(1956)	Farquhar and Hartmann(1957)	Farquhar and Hartmann(1957)	Wendell-Smith and Blunt(1965)
Farquhar and Hartmann(1957)	Palay(1958)	Kruger and Maxwell(1966)	Mori and Leblond(1969)
Maynard, Schultz and Pease(1957)	Schultz(1964)	Glees and Meller(1968)	Glees(1973)
Dempsey and Luse(1958)	Glees and Meller(1968)	Schultz(1964)	Schnapp and Mugnaini(1975)
Palay(1958a)	Orkand(1977)	Cammermeyer(1970)	Kuffler and Nicholls(1976)
Gerschenfeld, Wald, Zadunaisky and De Robertis(1959)	Privat(1978)	Matthews and Kruger(1973)	
	Roots(1978)	Lewis(1976)	
Gray(1959)		Peters, Palay and Webster (1976)	
Hartmann(1953, 1962)		Orkand(1977)	
Schultz(1964)		Privat(1978)	
Davson(1967)		Roots(1978)	
Cammermeyer(1970)			
Orkand(1977)			
Privat(1978)			
Roots(1978)			

TABLE 4 Light microscopic studies in which the authors point to their inability to distinguish between different kinds of neuroglia, or use the generic term 'neuroglia' without indicating a particular type

Cannot distinguish between astrocytes and oligodendrocytes	Cannot distinguish between astrocytes and microglia	Cannot distinguish between oligodendrocytes and microglia	'Neuroglia' used without differentiation into types
Deiters(1865)	Deiters(1865)	Deiters(1865)	Stricker and Unger(1879)
Jastrowitz(1870-71)	Jastrowitz(1870-71)	Jastrowitz(1870-71)	Held(1909)
Gerlach(1871-72)	Gerlach(1871-72)	Gerlach(1871-72)	Hess(1953)
Boll(1874)	Boll(1874)	Boll(1874)	Joseph(1954)
Golgi(1883)	Golgi(1883)	Golgi(1883)	Sholl(1956)
Penfield(1924)	Metz and Spatz(1924)	Del Rio-Hortega(1920)	Bairati(1958)
Rydberg(1932)	Urechia and Elekes(1925)	Metz and Spatz(1924)	Nurnberger(1958)
Bairati(1948;1958)	Bergman(1926)	Penfield(1924)	Lumsden(1953)
Ramon-Moliner(1958)	Creutzfeldt and Metz(1926)	Urechia and Elekes(1925)	Hyden(1961)
Koenig, Bunge and Bunge(1962)	Jakob(1927)	Bergman(1926)	Hamberger(1963)
Sulzmann(1962)	Prujjs(1927)	Creutzfeldt and Metz(1926)	Mori and Leblond(1969)
Schultz(1964)	Besta(1929)	Metz(1926)	Shantha, Manocha, Bourne, and Ariens Kappers(1969)
Truex and Carpenter(1969)	Cammermeyer(1963)	Jakob(1927)	Glees(1973)
Cammermeyer(1970)	Schultz(1964)	Prujjs(1927)	Watson(1974)
Orkand(1977)	Orkand(1977)	King(1937)	Kuffler and Nicholls(1976)
Raff, Fields, Hakamori, Mirsky, Pruss and Winter(1979)	Raff, Fields, Hakamori, Mirsky, Pruss and Winter(1979)	Bauer(1953)	Fischbach and Nelson(1977)
		Clemente and Holst(1954)	Currie and Kelly(1981)
		Pokak(1956)	
		Field(1957)	
		Ramon-Moliner(1958)	
		Cammermeyer(1963)	
		Estable-Puig, De Estable, Tobias and Haymaker(1964)	
		Schultz(1964)	
		Hamberger, Hansson, and Sellstrom(1975)	
		Lewis(1976)	
		Orkand(1977)	

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

The different elements of the 'cytoskeleton' are identified by immunofluorescent antibodies which are said to be more or less specific, and often also cross-reactive (Moll, Franke, Schiller, Geiger and Krepler, 1981). The elements are viewed under ultraviolet microscopy, but measured by electron microscopy. They are often identified in tissue cultures, but, to my knowledge they have never been seen in living cells. The cells may have to be 'permeabilized', that is fixed, frozen or stained for electron microscopy and dehydrated. The biochemistry of each of the elements is worked out after extraction of the tissue with powerful reagents, and then comparing the diameters from electron microscopy, the fluorescence from light microscopy and the biochemical properties of the extracts.

The accuracy of the identification of each element depends upon:

1. the nature of the fixative (including freezing) and any detergent used;
2. the degree to which fixation can prevent diffusion, whether or not 'permeabilization' has been used;
3. the specificity of the antigen, or any first or second antibody;
4. the penetration into the cell of the antibody from incubating medium;
5. the degree of the change of dimensions of each cellular constituent during preparation for histology, histochemistry or electron microscopy;
6. the effect of any of the reagents on the intensity of the antigen-antibody reaction;
7. non-specific uptake of antibody by elements in the tissue;
8. the effect of any of the reagents on the intensity of the fluorescence;
9. when immunofluorescence is used on tissue cultures, the intensity of the fluorescence may not be the same as it would be in the parent tissue from which the culture was taken.

Thus the light microscopist's view remains crucial and central, although apparently by no means certain. It thus behoves us to re-examine the whole fabric of current belief about these cells.

TISSUE CULTURE

Until now, techniques have not yet been developed for seeing living cell bodies in intact adult mammals. The nearest one can approach to knowledge about the living cell *in vivo* is to study preparations which have been minimally interfered with. These include cells in tissue culture, peripheral axons which fire, microdissected unfixed somas, and neural explants in the anterior chamber of the eye. One can also look at large non-mammalian cells either without removing them or in circumstances in which they can be visualized, or in conditions in which they fire after isolation: among such preparations are leech neurons and neuroglia, squid and cuttlefish axons, lobster stretch receptors, goldfish Mauthner cells and puffer supramedullary neurons. It is probably a reasonable assumption that the basic anatomy and the physiology of neurons from non-mammalian species are similar to those from mammals, although their biochemistry may be different.

Cells in tissue culture are normally judged to be living by observing intracellular movements with the non-invasive camera. The time-lapse films of Pomerat and his co-workers and Hansson and Sourander (1964) using retina, have given us new insights

METHODS OF STUDYING NEURAL STRUCTURE

into living cellular anatomy. There has also been an enormous development in the past decade of biochemical studies on the cells in culture (Fedoroff and Hertz, 1977; Schoffeniels, Franck, Tower, and Hertz, 1978; Pevzner, 1979; Raff, Fields, Hakamori, Mirsky, Pruss and Winter, 1979). While interesting in themselves, the biochemical properties of the different kinds of cells, particularly the neuroglia, depend crucially on their identification in the whole tissue and their later certain recognition in culture.

Tissue cultures give one a unique method of observing living cells, and the extrapolation is generally accepted that the findings from its use are true for the whole animal *in vivo*. This is an inescapable assumption. Some years ago we tried to observe living cells on the surface of the rat brain using vertical illumination microscopy, but unfortunately the tissue turned out to be too transparent. Nevertheless, such endeavours are still worthwhile pursuing in the future.

When cells or tissues are first put in culture there is a latent period during which they tend to be spherical and exhibit minimal intracellular movements. Subsequently, they begin to put out processes, which later develop into dendrites. The whole cells then gradually become flatter and may form a monolayer. Cells, which Pomerat and Costero (1856) identified as astrocytes, showed undulating movements. The elegant web-like dendrites are very difficult to differentiate from glial processes, unless one has decided on whether they take origin from a neuron or a neuroglial cell. It is of particular interest also, that the latter authors head one of their descriptions of the culture as 'Cells having some of the characteristic features of neurons'.

From the anatomical viewpoint a few generalizations may be made about the different sort of cells in culture. Firstly, in early cultures most cells, including those from non-neural tissues, look very similar. Tissues for culture must normally be taken from embryos, from very young animals or from tumours. The fact that the living cells can be examined in culture permits us to use them to describe general properties of living cells, which can be seen to be lost when the cells are fixed or killed. Among these properties, some cells show intracytoplasmic movements, clear nuclei, cell pulsation, electrical excitability, conduction, physiological synaptic activity, neuromuscular activity, myelination and intranucleolar movements. It therefore seems reasonable to entertain the assumption that the structures engaging in any of these phenomena are present in culture, and probably represent the minimal machinery which can exhibit them. Thirdly, many authors of papers on tissue cultures cannot distinguish the particular cells (Table 5) or do not describe the anatomy of the cells which they claim to identify (Table 6). A sample of the relatively few descriptions of the cells is given, and they can be seen to be vague and overlapping, both as seen by light and by electron microscopy. The commonest modes of identification are those derived from knowledge of the tissues or tumours from which they originated, or from identification by the use of biochemical 'markers' (Table 7). The greatest danger lies in the latter approach, since the correlations between the biochemical markers and the cells which they are believed to show up are not as good as they should be. When pressed, the research workers often adduce other evidence from classical light microscopy.

Fourthly, as mentioned above, the appearance of cells in culture changes. They 'mature', they de-differentiate - especially if cultured as a cell line - they degenerate, they may autolyse, and sometimes they disappear, presumed to have exploded. Of

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

TABLE 5 Light microscopic studies of tissue culture in which the authors point to their inability to distinguish neurons and the different kinds of neuroglia

Cannot distinguish between astrocytes and oligodendrocytes	Cannot distinguish between astrocytes and neurons	Cannot distinguish between oligodendrocytes and neurons	'Neuroglia' used without differentiation into type
Lumsden(1958)	Lumsden(1958)	Lumsden(1958)	Geiger(1957)
Murray(1965)	Fedoroff(1978)	Fedoroff(1978)	Lumsden(1958)
Lindsay, Barber, Sherwood Zimmer and Raisman(1982)	Lindsay, Barber, Sherwood Zimmer and Raisman(1982)	Lindsay, Barber, Sherwood Zimmer and Raisman(1982)	Bornstein and Appel(1961)
			Hild and Tasaki(1962)
			Varon, Raiborn, Seto and Pomerat(1963)
			Murray(1965)
			Vernadakis and Culver(1979)

METHODS OF STUDYING NEURAL STRUCTURE

TABLE 6 Authors using illustrations of light microscopic studies of tissue cultures who have not indicated anatomical criteria that they use for identification of particular cell types

Tissue	Authors
Human, monkey and rabbit cortex	Geiger(1957)
Kitten corpus callosum	Murray(1958)
Rat cerebellum	Bornstein and Appel(1961)
Rat and cat cerebellum	Hild and Tasaki(1962)
Rabbit cerebral hemispheres and cerebellum	Varon, Raiborn, Seto and Pomerat(1963)
Kitten thalamus and cerebellum	Murray(1965)
Rat cerebral hemispheres	Dittman, Sensenbrenner, Hertz and Mandel(1975)
Rat cerebral hemispheres	Moonen, Cam, Sensenbrenner and Mandel(1975)
Mouse dorsal root ganglion	Shahar, Grunfeld, Spiegelstein and Monzain(1975)
Mouse and rat spinal cord	Crain(1976)
Mouse cerebellum	Seil and Leiman(1977)
Mice brain hemispheres	Fedoroff(1978)
Rat optic nerve	Raff, Mirsky, Fields, Lisak, Dorfman, Silberberg, Gregson, Leibowitz and Kennedy(1978)
Rat brain	Lindsay(1979)
Rat corpus callosum, optic nerve, cerebral cortex, cerebellum	Raff, Fields, Hakamori, Mirsky, Pruss and Winter(1979)
Rat and mouse brain hemispheres	Hansson, Sellstrom, Persson and Ronnback(1980)
Rat brain	Abney, Bartlett and Raff(1981)
Rat cerebellum	Bartlett, Noble, Pruss, Raff, Rattray and Williams(1981)
Rat cerebellum	Cohen and Selvendran(1981)
Rat cerebellum	Currie and Kelly(1981)
Rat brain	Lindsay, Barber, Sherwood, Zimmer and Raisman(1982)

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

TABLE 7 Ratios of the activities in neurons and neuroglia of 51 different enzymes from the literature

Enzyme	Glia	Ratio of activity of neurons to Astrocytes Oligodendroglia	
<u>Group I. Oxidoreductases</u>			
Lactate dehydrogenase	1.8		
Malate dehydrogenase	2.6		
Malate dehydrogenase	0.2		
Isocitrate dehydrogenase	0.7		
6-Phosphogluconate dehydrogenase	0.6		
Glucose-6-phosphate dehydrogenase	0.4		
α -Glycerophosphate dehydrogenase	0.5		
Succinate dehydrogenase	0.8		
Succinate dehydrogenase	0.2		
Glutamate dehydrogenase	1.6		
Glutamate dehydrogenase	2.9		
Monoamine oxidase	1.8		
Monoamine oxidase	1.3		
Monoamine oxidase	1.4		
Monoamine oxidase	2.2		
Monoamine oxidase	2.8		
Monoamine oxidase	2.3		
Monoamine oxidase	1.1		
Monoamine oxidase	2.7		
Cytochrome oxidase	0.4		
Cytochrome oxidase	0.8		
Cytochrome oxidase	0.9		
Cytochrome oxidase	0.4		
Succinate: cytochrome c reductase	0.2		
Succinate oxidase	0.5		
Succinate oxidase	1.9		
Succinate oxidase	0.4		
Succinate oxidase	1.4		
Succinate oxidase	2.8		
<u>Group 2. Transferases</u>			
Choline acetyl transferase	1.1		
Galactosyl transferase		1.2	0.6
Sialyl transferase	2.4		
Aspartate amino transferase	2.0		
Aspartate amino transferase	1.0		
Hexokinase	3.2		
DNA polymerase	1.7		
DNA polymerase	0.2		
Acetylneuraminate cytidyltransferase	1.3		
Cerebroside sulphotransferase	0.1		
<u>Group 3. Hydrolases</u>			
Phospholipase A ₂	5.5		
Butyryl cholinesterase	1.1		
Butyryl cholinesterase	2.5		
Butyryl cholinesterase	3.6		
Acetylcholinesterase	1.7		
Acetylcholinesterase	3.1		
Acetylcholinesterase	2.2		
Acetylcholinesterase	1.0		
Phospholipase A	8.1		
Alkaline phosphatase	6.0		
Acid phosphatase	1.4		
DNase:			
acid	6.1		
alkaline	4.3		

METHODS OF STUDYING NEURAL STRUCTURE

Enzyme	Glia	Ratio of activity of neurons to Astrocytes Oligodendroglia	
Group 3. Hydrolases (cont)			
2',3'-cyclic nucleotide 3'-phosphohydrolase		1.0	0.1
2',3'-cyclic nucleotide 3'-phosphohydrolase		0.2	0.2
2',3'-cyclic nucleotide 3'-phosphohydrolase	0.1		
Aryl sulphatase	0.9		
β -glucosidase	1.1		
β -glucosidase	6.8		
β -galactosidase	0.9		
β -galactosidase	0.9		
β -galactosidase	9.9		
β -mannosidase	0.8		
β -mannosidase	1.3		
β -glucosaminidase	0.9		
β -glucosaminidase	1.2		
β -glucosaminidase	9.2		
β -glucosaminidase		0.3	0.1
β -glucuronidase	1.5		
β -glucuronidase	3.4		
Glucocerebrosidase	1.4		
Glucocerebrosidase	1.1		
Galactocerebrosidase		1.1	0.9
N-acetyl- β -galactosaminidase	1.0		
N-acetyl- β -galactosaminidase	1.0		
N-acetyl- β -galactosaminidase	0.6		
Arabinosidase	3.4		
Cathepsin	6.8		
ATP ases			
Mg ⁺ activated	5.7		
K ⁺ , Na ⁺ activated	0.1		
K ⁺ , Na ⁺ activated	0.4		
K ⁺ , Na ⁺ activated	0.2		
Ouabain-insensitive	0.2		
Group 4. Liases			
Glutamate decarboxylase	1.6		
DOPA decarboxylase	0.9		
Carbonic anhydrase	0.1		
Carbonic anhydrase	0.1		
Carbonic anhydrase	0.2		
Guanylate cyclase	2.1		
Group 5. Isomerases			
Glucosephosphate isomerase	2.0		
Group 6. Ligases			
Glutamine synthetase	1.7		

It should be noted that no ratio exceeds 10, and that few authors distinguish the different types of neuroglia. Pevzner concludes that 'in our opinion, the existence of a strictly neuronal or glial enzyme marker is inherently unlikely'. This table is reproduced from Pevzner, L.Z. (1979) Functional Biochemistry of the Neuroglia (Consultants Bureau, New York and London), pages 51-63, by kind permission of the author and publishers

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

course, de-differentiation represents the greatest difficulty for the microscopist trying to identify cells. Identification of neurons is facilitated by their producing classical action potentials when they are stimulated. Cells which produce slower transients of lower voltage are presumed to be neuroglia (Hild and Tasaki, 1962; Crain, 1976) but another possibility is that they are neurons whose spike-generating machinery has been changed by their isolation or the culture media.

Fifthly, probably the staining properties, the biochemical properties and the antigenicity of the original cells change during the isolation and growth in culture. There seems to be a totally inexplicable and dangerous lack of experiments, in which (a) the classical stains of the different kinds of neuroglia have been used to demonstrate the same kinds of cells in culture, (b) the biochemical properties in culture have been shown in tissue sections in regions where the characteristic cells occur in large numbers (it is argued here that in the tissues, as opposed to the cultures of 'pure' kinds of cells, particular kinds of cell occur all mixed up with others, which would make the biochemistry difficult to disentangle); (c) with a few exceptions, most of the 'specific' markers of cells in culture have not been shown to behave similarly in sections, and never *in vivo*; (d) nearly all immunological techniques used fixed and frequently dehydrated tissue cultures (Pickel, 1981 and sources quoted there). It is argued that the antibody could not enter the living cell. Yet a fixed, dehydrated cell is a dead cell, and antigen-antibody reactions should be expected to work better in living cells. There is a great paucity of experiments in tissue culture, in isolated cells or in whole tissue, showing such properties in live tissues. This is a particularly crucial question for those studying catecholamines, endorphins, enzyme markers, etc., in the nervous system in relation to the possible roles of these substances in mental diseases, or in animal models of those diseases.

ELECTRON MICROSCOPY

The electron microscope has a much higher resolution than the light microscope - in practice better than 0.3 nm (Cosslett, 1982). No biological tissue can survive the electron beam, the high vacuum and the radiation, so the tissue must be stained, that is, have deposited on it salts of heavy metals; this deposit is examined in the electron microscope. If one looks at an unfixed cell in a physiological medium by light microscopy one is looking only at a real tissue; when one looks at a stained histological preparation one is examining tissue minus substances removed during preparation plus added reagents - it thus represents tissue plus artifact. In the electron microscope one is only looking at metal or metal salt, that is, the image is of an artifact. Of course, that does not mean to say that the electron microscopic artifact has no value, but clearly it has less value than the fresh tissue as a source of information.

In light microscopic histology one may be looking at a colour image due to, say, eosin, dichromate or picrate, staining the tissue, or a deposit of salt of silver or osmium. The former solutions impregnate certain parts of cells, while the latter react with and precipitate on the structures; the former are colour stains, the latter are solid deposits and have grain. All the 'stains' used in electron microscopy - commonly, salts of osmium, lead, tungsten - have relatively poor solubility;

METHODS OF STUDYING NEURAL STRUCTURE

they precipitate on reaction with the tissue. Although I have often been told that electron micrographs have been published, or are in the possession of research workers, showing clear detail in tissues without any heavy metal stain, used at any time during the procedure so far no-one has referred me to any such pictures, nor favoured me with samples of their own work. Often, electron microscopists have used osmic acid as a fixative, or 'post-osmicated' a tissue, and have implied that the osmic acid was thus not used as a stain. Does this presume that this expensive salt knows when it is meant to act as a fixative, and when it is meant to act as a stain?

Incidentally, osmic acid, which is generally believed nowadays to stain proteins, has been used in light microscopy as a stain for fats since the nineteenth century (Mann, 1902).

Obviously, a deposit will react with both sides of a membrane of single thickness. Thus, any heavy metal reacting with a real membrane will be seen as two lines if the magnification is high enough. In our view that is one reason for the 'unit' membrane of Robertson (1959). Every single 'membrane' seen by electron microscopy, except that seen around 'lysosomes', appears double. Of course, a 'lysosome' is a solid particle, so the heavy metal deposit has only one ring around it. (For further discussion, please see pages 132-133).

Attempts have been made to examine living or hydrated cells by electron microscopy, either by using very high voltages like 3 MeV (Dupouy, 1972; Glauert, 1979), or by placing the tissues in wet chambers (Parsons, 1974, 1978). One awaits with hope the discovery with these instruments of new findings or greater detail, which so far have eluded classical light microscopy or electron microscopy.

Electron microscopy shares with histology all the difficulties arising out of the thickness of section (please see pages 49-62) and of the preparation techniques (Boyde, Bailey, Jones and Tamarin, 1977; Hillman and Deutsch, 1978a). In addition the technique as applied to biology has further problems:

1. The results of observation on what must be minute samples of the specimen are rarely controlled statistically; Cragg (1972) is exceptional in this respect. Whereas in light microscopy most histologists would agree on the identity of a particular structure, there would not necessarily be such wide agreement by electron microscopists. They have many fewer samples to examine, and they miss the dimension of colour.
2. Many structures seen by electron microscopy, whose existence in life we deny, are not seen in serial sections to fit together, or we have failed to find published evidence of these in the literature (please see pages 126-127). We would be interested to be referred to any such publications or receive copies of electron micrographs of serial sections. The structures in question are listed in Table 17. I would like to reiterate our previous assertion that any alleged structure of which one cannot make a three-dimensional transparent model must be an artifact or an optical illusion. No exercise of intellectual acrobatics can avoid the hurdles of geometry.
3. Attempts have been made to use biochemical evidence to support the electron

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

microscopic findings, but of course, since current biochemical findings are interpreted by reference to the electron microscopic images, the biochemistry cannot be used as evidence in this respect. It would be a circular argument to do so.

4. The difficulty of calibrating the electron microscope has been regarded as both an instrumental problem and as a difficulty of choice of object for calibration (Reimer, 1965; Bahr and Zeitler, 1965; Baumeister, Fringeli, Hahn, Kopp and Seredyński, 1976). Little attention has been paid to dimensional changes in the test objects under the conditions of the electron bombardment, which are almost certainly very different from the dimensional changes of the metal deposits and epoxy resins under the same regime.
5. After these reservations have been fully accounted for in our measurements, calculations and conclusions, electron microscopy can be used to give similar information to that derived by using histological techniques (please see pages 34-35).

SUBCELLULAR FRACTIONATION

If one wishes to draw conclusions about the structure of living cells or their organelles from subcellular fractionation, one is assuming that homogenization, centrifugation, separation and preparation for electron microscopy, have no significant effects on the size, shape and staining properties of the organelles under study (Hillman, 1972).

Biochemists who work with the mitochondrial fraction use the same term for the body in a section of tissue as they do for the same apparent structure in the subcellular fraction of the same name. It is worth noting that in both cases the whole mitochondria appear in the full range of shapes which geometry would dictate. Unfortunately, the cristae do not appear so. It is remarkable how frequently in a 'good' preparation they appear to be normal to the plane of section. A biochemical preparation said to be mitochondrial is seen to be packed with what are undoubtedly mitochondria. However, the preparation supposed to be composed largely of endoplasmic reticulum is called the 'microsomal' fraction; this is because it is also believed to contain cell membranes, as well as other intracellular membranes. There is obviously an assumption implied here that intracellular and extracellular membranes are structurally and biochemically similar. Such an assumption has not been proved. It would obviously not survive acceptance of the view that the endoplasmic reticulum and the cytoskeleton are artifacts. Nonetheless, the use of the term 'microsomal' obscures this assumption, and also permits one to conclude that an enzyme, marker, transmitter or chemical activity found in the microsomal fraction may be intracellular or intramembranous. It also makes the results of such experiments compatible with the findings of other authors, who claim that any such substance is located in either of these regions.

3

Cellular Identification – General Comments

The identification of each particular cell type and of the intercellular tissue, and the description of its structure, depends upon the following characteristics:

1. Their appearances by light and electron microscopy, including their relationship to other cells (Tables 8 and 9).
2. The specific staining systems by which they are demonstrated (Table 10).
3. Their physiological properties.
4. The markers, such as enzymes, proteins and transmitters they contain, and the antibodies and toxins for which they show affinity.
5. Their appearance in tissue culture (Table 11).

APPEARANCE OF CELLS

It cannot be stressed too much that the identification by light microscopy of the neurons, the neuroglia and the ependymal cells, is crucial to the understanding of neurohistology, since this was the technique by which the cells were described originally, and all other identifications of cells are still ultimately derived from these original descriptions (please see Table 1). It must be emphasized that none of the other categories under 1 to 5, nor those listed in Table 1, represent independent categories for classification. It has sometimes been said that when, for example, the identification of a particular cell by electron microscopy has failed to be unequivocal, it can be confirmed independently by the presence of a particular marker shown by immunocytochemical techniques.

GEOOMETRY OF CELLS

In examining the structure and appearance of cells, a number of questions arise. Can one describe the three-dimensional appearances of cells merely by looking at a few sections? The simple answer is that one can only do so either by making and studying serial sections systematically, or by using very thick sections, or by isolating the particular cells. The necessity for serial sections is demonstrated by the following simple model (Figure 21). Three eggs of the same size were taken as models of a neuron whose soma had a maximum diameter of 70 µm. Eleven sections equivalent to 6 µm thickness were cut in three different co-ordinates. If one considers the egg

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

yolk at the nucleus, and the whole egg as the cell, it becomes obvious that, without serial sections or very complex stereological calculations, one cannot describe: the dimensions of the cell, the nucleus or any intracellular body in any plane; the shape of the cell, the nucleus or any intracellular body; the absence of the nucleus or any intracellular body; the number of any small intracellular organelles; the position of the nucleus or any intracellular body; the absolute or relative intensity of staining of any membrane or part of the cell; the number of attachments of synapses to the cell bodies. If one were then to attach prolongations representing axons or dendrites to this model, it would then become clear that, without serial sections, one could also not tell: the number of such processes; their thicknesses; their lengths; their directions; their absolute or relative intensities of staining. Of course, one could not know if any parts of the cells were moving.

In addition to these geometrical difficulties, the total and relative changes in dimensions during histology make it problematic whether one could make real measurements - reflecting the state of the living tissue - of the total or relative dimensions of any membrane or intracellular body without knowing the changes that have occurred.

One can go some way to calculating dimensions:

1. if one assumes that the cells have similar dimensions in a plane at right angles to the section as they do in the plane of the section - (this will not be true for any asymmetrical body which is randomly orientated, or for any body which is not randomly orientated);
2. if one has in a single section a very large number of randomly orientated cells or intracellular bodies;
3. if one has many sections;
4. if one cuts sections whose thickness is much greater than the average maximum dimension of the cells or intracellular bodies;
5. if one examines serial sections, or isolated cells.

The problem of shrinkage can only be solved by measuring the degree of shrinkage of the body in which one is interested as the staining proceeds, and then correcting for it.

Thick sections have considerable advantages from the geometrical viewpoint, but optically they are not so satisfactory. The injection intracellularly of such markers as horseradish peroxidase to define the axon and dendritic trees of cells represents an important advance for cell localization, but unfortunately the tissues are normally subjected to histological procedures subsequently (Muller and McMahan, 1976; Kitai and Bishop, 1981). Isolating cells by hand dissection involves pulling off some of their processes, which makes it difficult to count them unless one can see all the broken ends.




FIGURE 21 The different appearances, dimensions and positions of the nuclei which could be obtained from sections of a cell cut vertically, horizontally and diagonally. Three eggs as models of neurons of 70 µm maximum diameter were sectioned at a thickness equivalent to 6 µm corresponding to that of a histological section

Classical histology can yield information about the numbers of cells in a particular plane, their relative positions in a tissue, and their general appearances. A few examples may be given; they can be used to find out: whether cells have divided recently; whether they are adjacent; if an epithelium is columnar cell or squamous; if cells are secretory; if they are symmetrical; if they originate from particular tissue, such as brain, muscle, liver or kidney; if they are granular; if they are pleiomorphic; if they are phagocytic; if they are lysing; whether there is much connective tissue around them; their approximate shapes if they are relatively symmetrical (see also pages 34-35).

These geometrical and histological reservations make one very careful in interpreting much data in the literature reporting structures or dimensions of cells or their

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

Table 8 The appearance of allegedly different cellular elements in the central nervous system as described by the authors. Lack of information under any heading indicates that it was not supplied in that publication

Form	Size	Processes	Nucleolus
APPEARANCES OF NEURONS BY LIGHT MICROSCOPY			
Irregular		Varying, number of dendrites which repeatedly bifurcate: single large axon from cell body, second system of small axons from dendrites	Present
Polygonal		Four to six, slender, irregular highly branched, originating from angles of cell body Thick rectilinear protoplasmic expansion takes origin from the two poles of the cell body Three divergent, infrequently branching expansions	
Fusiform: markedly elongated, slender ovoid cells with smooth contours			
Triangular cells of greater bulk than the fusiform			
Unipolar fusiform cells. Elongated spindle shaped, more or less uniform			
Spherical, flask-shaped, multi-polar, pyramidal and spindle	4-435 μm diameter	Many branching dendrites, single axon	Present
		One axon and several dendrites unipolar, bipolar or multi-polar	Large and spherical
More or less conical perikaryon or spherical		Fine, smooth axon, fine dendritic processes	
Varies from almost spherical to slightly flattened prolate spheroids	Largest cell, highly variable in size	Vary from a few to thousands	One or more present
	100 μm width and 100-200 μm long	Extend for up to 300 μm and branch several times	
Vast range of shapes	Golgi type I: long axons and large cell bodies Golgi type II short axons and usually small cell bodies	Vary in number from one to a score and each dendrite may branch scantily or into a fantastic bush: also has one axon which may branch Unipolar, bipolar or multi-polar; one axon and one or more dendrites	Generally conspicuous Usually more than one nucleolus present, with a nucleolus rich in RNA
Fusiform, stellate, pyramidal, polyhedral spherical, conical etc.			

CELLULAR IDENTIFICATION – GENERAL COMMENTS

Nucleus	Cytoplasm	Author/tissue
Present	Granular	Deiters (1865) ox - spinal motoneurones
Present		Ramon Y Cajal (1892) brain cells
Present	Fibrillary, Nissl granules present (tigroid masses)	Barker (1899) mammalian
Large and spherical	Contains mitochondria, Golgi apparatus, Nissl granules and neurofibrils, pigment granules	Ranson and Clark (1948) vertebrates
		Sholl (1956) cat - visual cortex
Nuclear membrane appears as a thin linear structure	Nissl substance present; mitochondria	Hyden (1961) rabbit Deiters' neurons
	Presence of lipochrome pigment (yellow granules) grouped at one or both ends of cell body. Cytoplasm was a firm gel	Roots and Johnston (1964) ox - red nuclei and lateral vestibular nuclei
Commonly large, and stains lightly, the chromatin network being composed of fine diffused fibrils	Mitochondria present as tiny, rod-like bodies Nissl bodies appear in cell and dendrites but not the axon, and vary in shape, size arrangement and number	Elliott, (1969)
Single nucleus usually centrally located, nucleoplasm is gelatinous. Cajal's accessory body sometimes present; intranuclear rodlet sometimes seen	Cytoplasm is of a soft gelatinous nature, in which mitochondria, Golgi apparatus, ergastoplasm and vacuoles are suspended; Nissl substance present and neurofibrils, and lysosomes	Shantha, Manocha Bourne and Ariens - Kappers (1969) brain cells

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

Table 8 continued

Form	Size	Processes	Nucleolus
Polygonal, pyramidal spherical or flask-shaped			Prominent
Complex, angular or polygonal with a slight concavity of cytoplasmic surfaces between the sites of the emergence of the processes	Large	Several short radiating dendrites and single long axon; axon may branch; multipolar normally	Single, conspicuous
Varied	Varied, 5-80 μm diameter	One axon, one or more dendritic processes	Large central nucleolus present-spherical
Unipolar, bipolar or multipolar		One axon and a variable number of dendrites	
Multiangular		A number of tapering, twisting and ramifying dendrites	Large, conspicuous and spherical

APPEARANCES OF ASTROCYTES BY LIGHT MICROSCOPY

Star-shaped		Processes radiate in all directions, possess one or more perivascular expansions which are applied to the surface of small vessels	
		In fibrous astrocytes fewer straighter processes than in protoplasmic astrocytes. Both types attach to small blood vessels by one or more perivascular feet	None
		Fewer in number, longer and thinner-fibrous astrocytes one or more end-feet per cell	Present
		Attached to blood vessels by processes terminating in perivascular feet	Absent
Stellate or star-shaped	Protoplasmic astrocytes are generally smaller than fibrous ones (2-35 μm long)	Processes radiate in all directions, numerous, some scarcely branch, some branch a lot	Often absent
		Varying number radiating in all directions; have end-feet to blood vessels; sometimes envelops nerve cells with its processes	

CELLULAR IDENTIFICATION – GENERAL COMMENTS

Nucleus	Cytoplasm	Author/tissue
Single nucleus is well visible, large and round	Cytoplasm can be abundant or small in amount, Nissl substance obvious; neurofibrils, mitochondria and Golgi apparatus are present	Ralis, Beasley and Ralis (1973) human – central nervous system
Large, pale, spherical or lightly ovoid, usually centrally placed in perikaryon; very fine uniformly dispersed chromatin articles	Cytoplasm – perikaryon, contains neurofibrils chromophilic substance or Nissl bodies, Golgi apparatus, mitochondria, a centrosome and various inclusions	Bloom and Fawcett (1975) man, rat – central nervous system
Vesicular nucleus, contains nucleoplasm and chromatin nucleolar satellite present	Nissl granules present, mitochondria and neurofibrils	Greenfield (1976) mammalian – brain and spinal cord
Nonnucleate		Landon and Hall (1976) vertebrate – central and peripheral nervous system
Large, round vesicular, located in centre of cell body	Nissl bodies present in different sizes and shapes (absent in axons but present in perikarya and dendrites); Golgi body present in mitochondria	Peters, Palay and Webster (1976) rat – cortex
	Granular cytoplasm which contains gliosomes	Ramon Y Cajal (1913) human – cerebrum
argest glial nuclei; irregularly oval; small amount of scattered chromatin	Granular or finely reticular; centrosome present; Golgi apparatus visible only in young mammals pigment granules present and gliosomes	Penfield (1932) mammalian – cerebral cells
argest glial nuclei, usually egg shaped	Less perinuclear cytoplasm – fibrous astrocyte; gliosomes, mitochondria, centrosomes fuchsinophile granules, pigment granules all present	Glees (1955)
ame size as small nerve cells irregularly oval, stain lightly, contains granules of chromatin		Ranson and Clark (1959)
oval or egg-shaped. Considerably larger than those of oligodendroglia. Chromatin is dispersed in the karyoplasm	Somewhat granular cytoplasm contains vacuoles sometimes scanty; centrosome, Golgi body, gliosomes, glial fibres, pigment granules all present	Hosokawa (1963)
ound or oval, occasionally elongated; larger than that of oligodendroglia or microglia; it usually only 1 nucleus, but occasionally two or more of different sizes		Mannen (1963) mammalian

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

Table 8 continued

Form	Size	Processes	Nucleolus
		Long wavy processes contain fibres (fibrous astrocytes); numerous, branched feathery; (protoplasmic astrocytes)	
			Conspicuous
		Multiple - called neuroglial fibres; perivascular feet attached to walls of small blood vessels and capillaries	
		Numerous, rather thick processes many being attached to the blood vessels and the pia mater by expanded pedicles; or long relatively thin smooth processes	
		Numerous, branching processes, sometimes attached by vascular end-feet to capillaries	
		Long and branched, occasionally capillary end-feet seen	

APPEARANCES OF OLIGODENDROCYTES BY LIGHT MICROSCOPY

	Smaller than astrocytes	Smaller, more delicate expansions than astrocytes	
	Smaller than astrocytes	Few, slender, relatively free from branches	
Round, oval, triangular spindle-shaped, pear-shaped etc	Much smaller than astrocytes 3-3.5 μm	Few thin fibrous processes furnished with delicate, knot-like swellings	
		Few extensions	
		Few, short processes, or none at all	
Round to pear-shaped		Few, extremely delicate	

CELLULAR IDENTIFICATION – GENERAL COMMENTS

Nucleus	Cytoplasm	Author/tissue
argest glial nuclei; oval with ine, evenly distributed rmatin		Gray (1964)
argest (5-10 μm diameter) and alest of glial nuclei; round/oval/egg-shaped. Fine chromatin etwork		Elliott (1969)
smaller than neurones; larger han oligodendrocytes and less ense	Pale, little or no detail	Curtis, Jacobson and Marcus (1972)
oval; bundles of fibres inside ell body		Ráliš, Beasley and Ráliš (1973) human - central nervous system
airly large, light & indented icear envelope has even rmatin lining		Ham (1974) rabbit brain
arger than oligodendroglia id microglia	Neuroglial fibres present	Bloom and Fawcett (1975) man, rat, central nervous system
arge	Granular, may contain fibres	Culling (1975)
entral, round or ovoid	Cytoplasm, surrounding the nucleus extended as processes of varying length	Hamberger, Hansson and Sellström (1975) rabbit - cortex
aller than astrocytes, and under, and contain more rmatin; dark	Small in amount and granular; appears roughly oblong; no end-feet or fibres; has centrosome, Golgi apparatus, fuchsinophile granules, and gliosomes	Penfield (1932) mammalian - brain and spinal cord
aller than astrocytes, stain ich darker		Ranson and Clark (1959) vertebrates
early oval, more chromatin han in astrocytes	Scanty granular cytoplasm surrounds the nucleus; centrosome, Golgi body, and gliosomes present	Hosokawa (1963)
under and smaller than that of astrocytes	Perinuclear mass, is small; form interfascicular glia	Mannen (1963) mammalian
aller than astrocytes, unded; dense 'blobs' of rmatin, especially ncentrated at nuclear envelope		Gray (1964)
centric, round and smaller han those of astrocytes	Are situated next to blood vessels or at a short distance from them, which they contact with a thickening	Cammermeyer (1966) rabbit - cerebral cortex

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

Table 8 continued

Form	Size	Processes	Nucleolus
Compact, well-outlined bodies with a distinct cell membrane		Few, scantily branching	
Often pear-shaped	Small	Few, delicate processes	
	Smaller than astrocytes	Small number of slender, poorly branching processes	
	Smaller than astrocytes	Few, slender processes with few branches	
	Small	No vascular feet, processes fine and rarely branch	
Oval	Small	Small number	

APPEARANCES OF MICROGLIA BY LIGHT MICROSCOPY

Cytoplasm passes out into expansions or give a bipolar or tripolar appearance; the expansions have numerous branches and terminal spines			
		Vary in number, length and thickness, and number of secondary and tertiary branches	
Majority are multipolar but some are bipolar	Very small	Two, three or more spiny branching processes	

CELLULAR IDENTIFICATION – GENERAL COMMENTS

Nucleus	Cytoplasm	Author/tissue
Generally smaller than astrocytes, (4-5 µm diameter); generally elongated, and oval, rounded, pearshaped or indented; dense clumps of chromatin	Gliosomes and centrosomes may be demonstrated by special preparation	Elliott (1969)
smaller than astrocytes, round and eccentric	Scanty with no fibrils	Truex and Carpenter (1969)
small and darkly staining	Thin rim of cytoplasm around nucleus	Curtis, Jacobson and Marcus (1972)
		Ráliš, Beesley and Ráliš (1973) human - central nervous system
mostly medium sized; spherical or oval with density intermediate to astrocytes and microglia; small number have very light, large nuclei, usually with a large central nucleolus; a fair number have small dark spherical nucleus		Ham (1974) rabbit - brain
smaller than astrocytes	No true neuroglial fibres are related to them	Bloom and Fawcett (1975) man, rat - central nervous system
	Stain intensely - no fibres present	Culling (1975)
	Rounded, darkly stained	Lewis (1976)
small and often irregular or elongated; contains rather heavy chromatin elements		Penfield (1924) cat, rabbit - brain
smallest and darkest in the nervous framework; spheromorphic; numerous granules and a large granule representing the nucleolus	Scanty in cell body, more abundant in prolongations; spongy protoplasm; no Golgi apparatus; specific granules and mitochondria also absent; some inclusions. Argentophile granules and, in some cases, lipoid granulations	del Rio Hortega (1932) rabbit, kitten - brain
nucleus surrounded by scanty cytoplasm; smallest, most darkly staining nuclei		Ranson and Clark (1959)

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

Table 8 continued

Form	Size	Processes	Nucleolus
Rod-like to irregularly shaped; monopolar, bipolar, multipolar	Average cell length including processes 52-62 μm , sometimes applied to capillary walls as perivascular satellites; when near to a nerve cell, it is a perineuronal satellite		Present
		Several with characteristic angular branching	
Stellate, elongated or bulb-like		One, two or more, long tortuous thick processes with delicate secondary branches, which contact numerous neighbouring neurons	
Inconspicuous in normal issue			
	Small cells	Long; of variable thickness angular bends and few branches	
Pleomorphic, elongated to triangular perikaryon		Extremely tortuous, prominent cytoplasmic processes varying in number and size	
	Small	Many	
	Small		
Processes and body covered with considerable number of tiny pointed twigs or spines		Few, rather short and twisted	
Oval	Small	A thick process arises from each end of cell which branches freely	

CELLULAR IDENTIFICATION – GENERAL COMMENTS

Nucleus	Cytoplasm	Author/tissue
irregular, somewhat twisted or round; round, oval, rod-like triangular etc. Rich in chromatin	Argyrophilic granules and sometimes lipoid granulation present; centrosome present, also Golgi body; gliosomes are absent and glial fibres	Hosokawa (1963)
smallest; tend to be elongated		Gray (1964)
void, or, rarely, a rounded nucleus with coarse chromatin granules	Situated juxtaposition to neurons and are rarely found along blood vessels	Cammermeyer (1966) rabbit - cerebral cortex
same size as, or slightly smaller than, in oligodendroglia but more darkly staining	Have centrosomes, and various granular inclusions, but no gliosomes or mitochondria	Elliott (1969)
arrow, elongated, stains less intensely than perikaryon, or processes		Mori and Leblond (1969) rats - corpus callosum
elongated or round	Perikaryon frequently situated next to a neuronal perikaryon or vascular wall; cytoplasm finely vacuolated; presence of PAS red stained lysosomal or lipofuscin-like granules in older animals	Cammermeyer (1970) rabbit - brain
elongated	In diseased conditions end up as 'glitter' cells when they are full of phagocytosed debris	Johnston and Roots (1972)
	Migrate rapidly by amoeboid movement; phagocytic	Ráliš, Beasley and Ráliš (1973) human - central nervous system
small, elongated, with dense chromatin separated by light spaces	Rarely distinguishable	Ham (1974) rabbit - brain
small		Bloom and Fawcett (1975) man, rat - central nervous system
deeply staining		Culling (1975)

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

organelles. All these reservations apply to transmission electron microscopy as well as to light microscopy.

When one searches the classical literature for descriptions of the different kinds of cells, one finds that there are many descriptions by light microscopy of neurons but virtually none of astrocytes (Tables 8 and 9). Is this because astrocytes are rare in healthy tissue? Are they difficult to stain? Are their shapes so variable that they defy description? Are they so obvious in sections that they need no description? Is it possible that they do not exist?

When one finds that respected histologists say that one cannot distinguish between cell types (Tables 2 to 5), one must pose the question, Why then do they consider that they can recognize different kinds of cells? One cannot obviate this problem by quoting other criteria for the identification of the particular cells when one knows that these other criteria themselves are based on the histologists' initial classification of the cell type. The same difficulty exists with respect to the use of cell markers.

A very vivid indication of the difficulty of distinguishing cell types by light microscopy is seen in the illustration from the Frontispiece of Peters, Palay and Webster (1976), which they have kindly permitted us to reproduce (Figure 22). Brief perusal of their drawings leads to the realization that rotation of any of their 'typical' cells would give appearances which would be quite indistinguishable from each other. If one were to make three-dimensional models of their Golgi-stained neurons, astrocytes and oligodendrocytes, and if one were to cut sections of them of thickness equivalent to those used in histology, the sections would have a wide range of overlapping appearances, and so would be even less distinguishable from each other than whole cells. Presumably their drawings are intended to represent typical appearances, so one would be tempted to conclude that the cells' external appearances stained by the Golgi method cannot be differentiated by light microscopy.

Pathologists normally examine sections at relatively low-power magnification, and their diagnosis is based as much on the clinical appearance and natural history of the condition as on the appearance of individual cells at high-power magnification. The pattern of cells in the tumour helps diagnosis of its type, and the polymorphism of their shapes gives some indication of the prognosis, yet this polymorphism or pleiomorphism itself renders the identification of the predominant cell type more difficult. This difficulty is carried on to any identification of cells in culture from the tumour.

In many functional studies the location of the cerebral nuclei and the general myeloarchitectonics indicate the physiological role of the whole nucleus so that their morphology is not so important. For example, recording responses in the sensory cortex to stroking the footpads, or stimulating the motor cortex to produce epileptiform movements, has few implications for the morphology of the neurons involved in these physiological events.




FIGURE 22 The original legend reads: 'Various aspects of nerve cells in light microscopic preparations: A: A small pyramidal cell from the visual cortex, Golgi method. The axon (a) descends from the cell body. B: A small neuron in the dentate nucleus of the cerebellum, Golgi method. The axon (a) is represented only by its initial segment. C: Protoplasmic (velate) astrocyte in the grey matter, Golgi method. D: Oligodendrocyte in the white matter, Golgi method.' Peters, A., Palay, A.L. and Webster, H. de F. (1976) *The Fine Structure of the Nervous System*. Frontispiece. (Philadelphia: W.B. Saunders Company). Please note that rotation of any of these cells, and cutting sections of them in various orientations, could produce pictures which would not distinguish any of them from each other. Please also see Tables 8, 9 and 10.

SPECIFICITY OF STAINS

Staining procedures can either be general ones, such as haematoxylin and eosin or cresyl violet, or they can be 'specific' stains for particular cell types or parts of cells. The identification of cells with both types of stains depends upon their shapes and colours. The colours may vary in intensity or distribution within the cell. Some salts, such as those of silver or gold, can stain a tissue grey or yellow, respectively, or deposit as black or brown granules, depending on the precise chemistry of the extremely complex mixtures of all the reagents and the tissue. The particular chemical milieu in which the reagent tissue mixture exceeds its solubility product is unknown and probably unknowable.

There have been many studies of the effects of reagents on particular compounds extracted from tissues (Mann, 1902; Baker, 1958; McClung-Jones, 1967; Symposium, 1965; Pearse, 1972, 1980). However, in view of the complexity of the chemistry of tissue and the alchemical concoctions of all the chemicals which are used in the procedure, it seems extremely unlikely that the simplified chemical studies are adequate. For example, the examination of the reaction of 10% formalin only with an enzyme preparation (Bruce-Casselman, 1959) is unlikely to represent all the chemical events

Table 9 The appearance of allegedly different cellular elements in the central nervous system as described by the authors. Lack of information under any heading indicates that it was not supplied in that publication

APPEARANCES OF NEURONS BY ELECTRON MICROSCOPY						
Form	Size	Processes	Nucleolus	Nucleus	Cytoplasm	Author/tissue
Axon, thin longitudinally orientated filaments, dendrites present	Nucleoli present eccentrically; vacuolated or solid and always lack a smooth contour; no limiting membrane	Great variety in apparent thickness and structural arrangement of nuclear membrane; indentations appear regularly; nucleoplasm has electron dense irregular clumps of fine granular material, less dense	Composed of minute scattered granules; mitochondria occur as spheres, short rods or elongated and possibly wavy filaments, vary in size from 300 to 3000 Å diameter	Harmann (1953) rat - central nervous system		
		Appears as a tangle or convoluted mass of dense and thick filaments made up of dense aggregations of fine granules; no membrane present	Has double nuclear membrane with perforations	Paley and Palade (1955) rat - dorsal root ganglion, medulla oblongata and cerebellar cortex		
			Cytoplasm crowded with formed elements considerably in size and profile, has endoplasmic reticulum orderly arranged within the Nissl bodies, and small granules in close contact with outer membranes of endoplasmic reticulum or scattered throughout matrix; have an orderly arranged agranular reticulum present, vesicles and a concentrically laminated, membranous structure has been found; also numerous filamentous, rod-like, or spherical mitochondria; neurofilaments - fine long threads 60-100 Å in diameter traverse the matrix, either all in one direction, or criss-cross in all directions, dense rounded bodies 350-660 µm in diameter are present			
			Poorly defined chromatin clumps; nuclear membrane appears double	Schultz, Maynard and Pease (1957) rat - cerebellar cortex and corpus callosum		
			Chromatin granules occur in clusters; fine filaments present; nuclear envelope has pores	Gray (1964)		
		Many dendrites containing many canaliculi, mitochondria and membrane-bound cavities. Axon present	Nuclear envelope appeared intact and with paired membranes, large number of pores in envelope	Roots and Johnston (1964) ox - red nuclei and lateral vestibular nuclei of brain		
				Dense: Nissl bodies - granular endoplasmic reticulum; Golgi apparatus a concentrically arranged group of cisternae; dense cytoplasmic granules; mitochondria generally smaller than in other cells; multivesicular bodies; dense granular inclusions; large heterogeneous bodies; glycogen; lipofuscin pigment and basal bodies and cilia; neurofilaments	Ross (1964) rat - dorsal root ganglion	
		Prominent	Lightly granular			

CELLULAR IDENTIFICATION – GENERAL COMMENTS

		Pale and often irregular	Dense, abundant ribonucleoprotein present; elaborate smooth surfaced membrane systems present; sparse disorderly arranged rough, endoplasmic reticulum ; oval or irregular electron dense bodies present	Nathaniel and Nathaniel (1966) rat – posterior horn of spinal cord
	Axon and dendrites; axons diameter 0.2– 2.2 μm , contain vesicles 400–500 \AA and 600–1400 \AA diameter	(a) Irregular with folded and dilated nuclear envelope (b) More spherical	(a) Dark neuron-dense cytoplasm; numerous free ribosomes, dilation of cisternae of the vacuolar system; large Golgi complexes with dense coret vertices of 100 \AA in diameter close by; large dense bodies 3–4000 \AA diameter also present. (b) Pale neuron-less dense cytoplasm with fewer ribosomes; fewer cisternae in endoplasmic reticulum and Golgi complex less well developed	Echiverri and de Iraldi (1968)
	Characteristic; large, and varying round or irregular	Large, round, dense, situated near centre of nucleus; nucleoloma present	Centrally located; round to oval; has nuclear envelope (double membrane 175–300 \AA apart); nuclear pores present, also fibrils and particles	Rapoport and Stempak (1968) rat – ventral horn cell
Multipolar usually	Varying – small to giant	Isodendritic Off centre, nucleolus present	Usually large, spherical and eccentrically placed; contains fine filaments and associated granules Large, eccentrically placed	Nissl substance, small mitochondria, Golgi apparatus, neurofilaments, microtubules, lysosomes, pigment granules and vesicles of various sizes Ribosomes, endoplasmic reticulum, Golgi complex, lysosomes present, numerous elongated random mitochondria
				Sotelo and Palay (1968)
				Bondareff and Hyden (1969) giant nerve cells of Deiters nucleus
APPEARANCES OF ASTROCYTES BY ELECTRON MICROSCOPY				
	Number variable	Ovoid, larger and less dense than other glial cells; 'vesicular' appearance	Abundant cytoplasm appears relatively 'empty'; few mitochondria, occasionally some Golgi material and a few small vesicles and particles; irregularly- shaped masses of finely granular material are sometimes present	Farguhar and Hartmann (1957) man and rat – cerebrum
Perivascular end-feet present	Large Processes which form sheaths	Not present	Conspicuously, clumped chromatin masses, nucleoplasm rather watery	Schultz, Maynard and Pease (1957) rat – cerebral cortex and corpus callosum
	Processes present, practically filled with filaments (90 \AA)	Oval, densely granular	Relatively thin layer; mitochondria or agranular reticulum present, and ergastoplasm with a high proportion of fine nucleoprotein granules; small dense, irregularly shaped clusters enclosed by single thin membrane	Palay (1958) rat – central nervous system

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

Form	Size	Processes	Nucleus	Cytoplasm	Author/tissue
Vascular feet applied to blood capillaries		Prominent	Ovoid; dense accumulations of chromatin near the nuclear membrane	Low density 'watery'; mitochondria, Golgi material, endoplasmic reticulum and small number ribonucleoprotein particles present	de Robertis, Gerschenfeld and Wald (1960)
Multiple, delicately branched refolding processes		Ovoid or reniform Large and light		Scant perinuclear cytoplasm Light and watery; ollofilaments sometimes present, form part of so-called neuropil	Luse (1962) Hosokawa (1963)
				Pale, 'watery'; extensive bundles of fibrils in fibrous astrocytes; practically no granular endoplasmic reticulum; very few mitochondria	Gray (1964)
				Relatively clear filamentous cytoplasm with little endoplasmic reticulum	Mugnaini and Walberg (1964) cat and dog
				Granular endoplasmic reticulum is scarce but numerous free ribosomes present; number of mitochondria is relatively small; Golgi complex is well developed; osmophilic bodies are frequently found; filaments present scattered or in bundles	Glees and Müller (1968) monkey
		Contain osmophilic bodies, glycogen granules and filaments			Vaughn and Peters (1969) rat - optic nerves
		Many, which form glial sheets between fascicles of axons	Some condensed chromatin adjacent to nuclear material	Less electron-dense than oligodendrocytes; contain characteristic filaments and/or glycogen; microtubules are rare	Curtis, Jacobson and Marcus (1972)
		Pale processes, with glycogen present		Many filaments, few microtubules	Elliott (1969)
			Nuclear membrane is double and seems to be continuous with endoplasmic reticulum; presence of pores	Cytoplasm packed with various inclusions mitochondria present as double walled capsules with ridged inner walls; normally have no centrosome; Nissl bodies formed of numerous flattened vesicles, tightly packed neurofibrils present, endoplasmic reticulum and Golgi apparatus	Johnston and Roots (1972) ox - brain
		Unipolar, bipolar, or multipolar	Usually one or two present and consists of small dense granules and threadlike nucleolomemas	Prominent, enclosed in a nuclear envelope which is a double-membrane; contains chromatins granules	
Smooth perikarya 1. More irregular 2.	Large Large 17-35 Outline, pear-shaped or multipolar	Few spines (3-4 per cell) More spines mm in diameter	Well developed; generally centrally placed in nucleus, associated with it can see a heterochromatic mass and a nuclear rodlet	Round and usually lobulated or dimpled at one pole; nucleoplasm is thin and flocculent and has small clumps of chromatin	Chan-Paley (1973) rat - lateral vestibular nucleus

Table 9 continued

CELLULAR IDENTIFICATION – GENERAL COMMENTS

Smooth or irregular	Small, 10-12 μ in diameter	Prominent and central	Cytoplasm is limited, little Nissl substance - scattered cisternae of endoplasmic reticulum and small collection of polyosomes. A prominent Golgi apparatus rings the nucleus	Chan-Palay (1973) rat - lateral vestibular nucleus
Angular or polygonal	Multipolar, dendrites and the axon	Present	Nuclear envelope and pores	Bloom and Fawcett (1975)
	Multipolar, contains microtubules		Large, spherical, conspicuous eccentric, extremely dense, made up of granules and fine filaments	Aggregations of neurofilaments, granular endoplasmic reticulum and associated ribosomes. Golgi network, agranular endoplasmic reticulum, mitochondria, centrosome containing a pair of centrioles and other inclusions
	The axon contains neurofilaments, sparser in dendrites; branching occurs in both		Large, round, contains fine chromatin filaments; surrounded by the plasma-lemma	Strands of chromatin, multivesicular bodies, lysosomes, Golgi apparatus present; and granular endoplasmic reticulum; occasional microtubules; mitochondrial profiles present; ribosomes and neurofilaments
			Normally single nucleolus present	Nissl substance, neurotubules and neuro-filaments, membrane-bound ribosomes, rough endoplasmic reticulum, Golgi apparatus, mitochondria, microfilaments, microtubules
			Several; long, irregular and branched; vascular end feet	Ribosomal endoplasmic reticulum, and microtubules, Golgi complex, numerous free ribosomes, neuro-tubules, ovoid to round mitochondria
	Fibrous astrocytes are longer and branch less than protoplasmic astrocytes		Fibrous astrocytes are longer and branch less than protoplasmic astrocytes	Unusually lucent matrix; numerous fibrils - thinner than neurofibrillaments (diameter = 60-80 \AA); lysosomes and glycogen granules present
Formation of 'end-feet' processes		Ovoid	Light rather large, often indented	Constantinides (1974)
			Round/ovoid with smooth outline (protoplasmic astrocytes) or irregular and indented (fibrous astrocytes)	Hamberger, Hansson, and Sallström (1975) rabbit - cortex
	May be attached to a capillary	Irregular	Light, poor development of endoplasmic reticulum	Blackwood and Corseilis (1976)
			Light, watery; some mitochondria, scanty endoplasmic reticulum	Lewis (1976)
		Occasionally present	Bean-shaped or irregular	Peters, Palay and Webster (1976) rat - optic nerve and cerebellar cortex
				Jacobson (1978)
				Lucent cytoplasm, sparse cytoplasmic organelles, bundles of microfilaments 79-100 \AA in diameter, may contain glycogen granules

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

Table 9 continued

Form	Size	Processes	Nucleolus	Nucleus	Cytoplasm	Author/tissue
Processes expand at their distal ends and abut onto capillaries -astrocyte feet	Varying number, some with few branches, some extensively branched				Numerous filaments, scattered or in discrete bundles; relatively few other organelles present so has 'watery' appearance, glycogen granules are characteristic inclusions; few microtubules	Roots (1978) vertebrate
Pericapillary end-feet	Numerous processes to less numerous	Medium-size to large	Various shapes, often oval; chromatin evenly dispersed, very few peripheral clumps		Relatively clear; mitochondria present sometimes with matrix filaments and angular cristae; granular endoplasmic reticulum present in cell body, some processes and end-feet; intermediate filaments numerous to scanty; microtubules present but scanty; heterogeneous dense bodies present (probably some specific); glycogen like granular material present in some processes and end-feet; cilium, Golgi apparatus, multivesicular bodies and centrioles present	Mugnaini, Olsen, Dahl, Friedrich and Korte (1980)

APPARANCES OF OLIGODENDROCYTES BY ELECTRON MICROSCOPY

	Seldom seen		Round to oval; smaller and denser than astrocytes; chromatin often aggregated peripherally to form dark rim	Scanty; quite dense and contains an abundance of formed elements i.e., mitochondria, Golgi material, endoplasmic reticulum and small particles	Parquhar and Hartmann (1957) man and rat - cerebral tissues
Small.	Few	Dense	Dense fibrils moderate quantities of endoplasmic reticulum and mitochondria; dense RNA particles; sparse groups of Golgi apparatus	Scant rim, moderately dense; no fibrils moderate quantities of endoplasmic reticulum and mitochondria; dense RNA particles; sparse groups of Golgi apparatus	Schultz, Maynard and Pease (1957) rats - cerebral cortex and corpus callosum
Round	Small.	Few	Nearly spherical	Crowded with dense nucleoprotein granules and small vacuoles	Palay (1958) rat - central nervous system
			Round, with double nuclear envelope; condensed chromatin constituted by fine filaments	Denser than that of astrocytes; Mitochondria present plus large quantity of ribonucleic-protein particles	de Robertis, Gerschenfeld and Wald (1960) rat - spinal cord
		Few broad processes which often branch dichotomously	Round/ovoid	Abundant; a little ergastoplasm; few mitochondria	Luse (1962)
				Nucleus smaller than that of astrocytes and more round and more electron dense	Luse (1962)
				Less cytoplasm; and is denser than that of astrocytes	Hosokawa (1968)
				Chromatin in conspicuous 'blobs'	Gray (1964)

CELLULAR IDENTIFICATION – GENERAL COMMENTS

			granular endoplasmic reticulum	Walberg (1964) cats and dogs
		Present as a constant feature	Round and very seldom lobulated; unevenly filled	Glees and Meiller (1968), monkey
	Few		More electron-dense than astrocytes; usually located near one pole of the cell; irregular clumps of chromatin beneath the nuclear membrane	More electron dense than astrocytes, Golgi apparatus present; short regular profiles of granular endoplasmic reticulum are scattered throughout and diffuse, electron dense material, numerous clusters of free ribosomes, and microtubules are present; no filaments
			Dense, nuclear chromatin, light patches adjacent to the numerous nuclear pores	Large quantities of free ribosomes/ ribosome rosettes; prominent microtubules; dark and light multivesicular bodies; granular inclusions; Golgi apparatus and mitochondria present; no fibrils or glycogen granules
			Located towards pole of cell; chromatin heavily clumped	Denser, but scantier than neurons; many microtubules – few neurofilaments; dense clumps of endoplasmic reticulum; clusters of polyribosomes
		Fewer than astrocytes	Eccentric; clumped chromatin	Abundant, denser than astrocyte cytoplasm; almost no fibrils – large number of microtubules; prominent Golgi vesicles; marked development of endoplasmic reticulum; many granules but no glycogen; numerous ribosomes
			Rounded/ovoid/irregular shape; dense	Dense; microtubules in processes; very few cytoplasmic fibrils of glycogen granules; Golgi apparatus, endoplasmic reticulum, mitochondria, and rosettes of free ribosomes present
		Well marked	Round, oval or irregular	Closely packed ribosomes, polyribosomes and endoplasmic reticulum
	Few		Round, oval, or irregular; chromatin tends to clump, dense	Dense, well-developed Golgi apparatus; inconspicuous mitochondria; few fibrils and glycogen granules; abundant microtubules; short cisternae of endoplasmic reticulum
				Fewer cytoplasmic processes than astrocytes, have abundant endoplasmic reticulum and ribosomes, high density of cytoplasmic organelles, contain prominent Golgi apparatus and vesicles and many cytoplasmic microtubules
		Fewer processes than astrocytes		Large number of organelles especially microtubules, ribosomes and various dense bodies; glycogen is not found
				Roots (1978) vertebrate

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

Table 9 continued

Form	Size	Processes	Nucleolus	Nucleus	Cytoplasm	Author/tissue
	Few to numerous, often thin	Small	Round or oval, chromatin in clumps; swollen nuclear envelope	Usually dense; mitochondria few and small; granular endoplasmic reticulum abundant in cell body but not in distal processes; short cisterns with wide transparent lumen; intermediate filaments usually not present; numerous microtubules both in cell bodies and processes; heterogenous dense bodies present (some specific); glycogen-like granular material not observed or cilium; Golgi apparatus, multi-vesicular bodies and centrioles present	Magnaini, Osen, Dahl, Friedrich and Korte (1980); see also Magnaini and Walberg (1964)	

APPEARANCES OF MICROGLIA BY ELECTRON MICROSCOPY

	Several processes with many branches	Great density and variable shape, chromatin evenly distributed and is uniformly dense	Quite dense and contains abundant formed elements (i.e. mitochondria etc); amount is intermediate between that of astrocytes and that of oligodendroglia, frequently there is a large amount of cytoplasm near one pole of the cell	Farguhar and Hartmann (1957) man and rat - cerebrum
		Dense, angular and irregular	Extremely dense; conspicuous Golgi apparatus; moderate amount of endoplasmic reticulum; few, small mitochondria	Schultz, Maynard and Pease (1957) rat - cerebral cortex and corpus callosum
	Very small with no mitochondria inside them	Elongated nuclei	Dense, heterogeneous, cytoplasmic masses present, containing fine granules, rounded larger masses and packets of thin, parallel, sheet-like material	Palay (1958) rat - central nervous system
Irregularly indented	Small	Longish and irregularly shaped, finely granular and of high electron density	Dense, and dark inclusion bodies are often present	Palay (1958) rat - central nervous system
		Dense, highly crenated	Densely packed granules, pale endoplasmic reticulum present	Hosokawa (1963)
	Processes extend regularly	Unique!	Thin rim around nucleus - of same density as interchromatic nucleoplasm; no fibres; infrequent Golgi, endoplasmic reticulum and ribosomes; mitochondria and lysosomes present;	Gray (1964)
				Hori and Leblond (1969) rat - corpus callosum

CELLULAR IDENTIFICATION – GENERAL COMMENTS

		cultes, more than astrocytes; few or no tubules or filaments; long, stringy cisterns; fat droplets and laminar dense bodies present	Marcus (1972)
	Dense chromatin	Sparse; no fibrils or microtubules; no glycogen; few microvilli, some dense bodies	Constantinides (1974)
	Small, shape relates to number of processes; extremely dense	Little rough endoplasmic, but fair amount of dense bodies	Han (1974)
	Oval, or elongate with prominent chromatin clumps	Extremely dense; moderate amounts of 'angular reticulum', endoplasmic reticulum and mitochondria	Greenfield (1976)
	Small, containing dense chromatin clumps and light nucleoplasm.	Thin rim around nucleus, and forms broad processes; long narrow cisternae of endoplasmic reticulum; Golgi apparatus present and large numbers of inclusion bodies	Peters, Palay and Webster (1976) rat – cerebellar cortex and spinal cord
		Scanty cytoplasm containing lipid inclusions, lysosomes and abundant vesicles but sparse endoplasmic reticulum, scattered ribosomes, and few or no microfilaments or microtubules	Jacobson (1978)
		No abundance of filaments or microtubules, but other inclusions are numerous and much more common than in either oligodendrocytes or astrocytes	Roots (1978) vertebrate
	Few, both thin and large, with thin side arms	Oval or indented, chromatin in large clumps, small to large nucleolus	Mugnaini, Osen-Dahl, Friedrich and Korte (1980); see also Mugnaini and Walberg (1964)

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

which occur to enzymes in tissue when it is fixed. Most modern electron microscopists believe that osmium salts stain proteins specifically, while light microscopists for a long time have considered them as a suitable stain for lipids (Romeis, 1968, pages 260-261).

Sometimes the authors of new staining procedures or modifiers of well-accepted procedures claim that they are 'specific' for one cell type (Table 10). It is extremely rare that they are quite so definite about the specificity. The following degrees of certainty are expressed about the specificity of a staining procedure for a particular cell type:

1. it is specific; it does not stain other cell types;
2. it is specific, but may stain other cell types, for example, if permitted to 'overstain', or 'by accident', or occasionally;
3. it is said to be specific but no information is given about whether or not it has been tested on other cell types;
4. it is not explicitly claimed to be specific but is listed as staining one cell type, or is 'recommended' for staining that cell type, and is not listed under staining procedures for other cell types;
5. it stains one type of cell with a range of colours which overlaps grossly the range of colours which it also stains another type of cell;
6. one has to assess subjectively by exercise of judgement, skill or experience at what moment, or with what concentration of reagents at a particular stage in the procedure, the particular cells in which one is interested are stained adequately.

The literature is replete with examples of this practice. In discussing the demonstration of protoplasmic astrocytes in the cerebral cortex by the method of Cajal, Gray (1954, page 537) says: 'The successful application of this technique depends on the ability to judge the exact shade of purple which indicates a satisfactory termination of the staining process.' Carleton and Drury (1962, page 221) comment: 'The complexity of the intervention of factors as yet unknown make it often difficult to stain particular elements of the nervous system in the same way on different occasions.' These reservations, regrettably, are much more frequently expressed in the older literature. The more recent literature has tended to regard the specificity of cells as being defined more accurately by the electron microscopist than by the light microscopist, forgetting that the electron microscopist takes his cue from the light microscopist. (Please see also Appendixes 1 and 2).

In assessing the literature on specificity of staining, one frequently comes across the use of the term 'neuroglia', without a more precise indication of whether the author is referring to astrocytes, oligodendrocytes or microglia. This can only be interpreted as meaning either that the author believes that the particular staining procedure does not differentiate between the types of neuroglia, or that it stains all of them, or that he cannot distinguish between them. So far histologists have not indulged in the luxury of wondering whether, indeed, they are different types of cells.

The study of different kinds of neuroglial cells - if indeed they do exist - is made much more difficult when one compares the specificity of particular stains as listed in authoritative standard textbooks. Frequently, different authors, using the more or less

CELLULAR IDENTIFICATION – GENERAL COMMENTS

explicit expressions listed above either assign rather different specificities to a particular staining procedure than do their equally distinguished colleagues, or describe them as being much less specific. Examples of this occur commonly in the literature, but are most easily appreciated by comparing the texts (Table 10). (Please see also Appendix 1.)

In addition to using the term 'neuroglia', other generic terms used include 'nervous system stain', 'neural stain', 'central nervous system stain', 'neural element stain', 'spinal cord stain', 'cellular element stain in the nervous system'. All these terms carry the implication that one cannot distinguish between the different cellular elements of the nervous system. Dare one also comment that most 'specific' stains also show up capillaries in the nervous system?

If one accepts that the staining procedures are not 'specific' for particular kinds of cells, one could still say that it shows them up, even if in a number of defined and undefined circumstances, it may also stain other cells, to which at the time one was not directing one's attention. Unfortunately, this does not help the histologist who is attempting to classify the cells in the nervous system, since it begs the important question of whether there are any criteria for distinguishing the different kinds of cells; that is, what is the evidence for believing that neurons, astrocytes, oligodendrocytes and microglia are different cells?

In order to claim that a particular stain shows up a specific cell type, one has to show also that, firstly, it does not show up any of the other cell types in that tissue, and secondly, that staining procedures which are believed to be specific for other cell types do not also stain the one for whom the specific staining is claimed. These requirements for specificity apply equally to any claim for the specificity of any marker for a particular cell type.

The specificity of the stains depends heavily on these 'control' experiments, but the great era for the development of stains preceded that in which the concept of control experiments, of testing of statistical significance, or even of questioning upon what evidence any authority had claimed specificity for his staining procedure, had arrived. Since that era, the belief in specificity has apparently continued to be accepted without question.

PHYSIOLOGICAL PROPERTIES

In general, neurophysiologists penetrate cells with intracellular electrodes, or record extracellularly. It is easier to encounter a large cell body than a small one, and it is less damaged by the electrode. Therefore, if a stable membrane potential can be recorded and the cell body can be excited, that cell is usually described by the neurophysiologists as a neuron. If a large resting potential is recorded, but surface, orthodromic, antidromic or intracellular stimulation fails to excite that cell, it is generally felt that it is likely to be a neuroglial cell. From such reasoning, and the fact that neuroglia in culture give very slow potential changes, which look quite different from those in neurons (Hild and Tasaki, 1962), it is generally believed either that neuroglial cells do not fire or that they are too small or too difficult to penetrate with intracellular electrodes. It might be advantageous for those who believe that astrocytes and oligodendrocytes are different cell types to try to give a definitive

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

Table 10 The cell types or parts of cells believed by different authors to be stained by particular procedures

Bielschowsky's silver	Cajal's gold sublimate	Cresyl violet	Colgi-Cox silver	Haematoxylin and eosin	Holzer	Marsland, Glees and Erikson	Palmgren	Patay
				Nerve cells and processes - dark brown Background - pale brown Other elements not reported	General Nuclei - blue Cytoplasm - pink Neuroglia fibres - pink			
Peripheral nerve fibres Axis cylinders - black Myelin - red-violet Axis cylinders Connective tissue - violet or blue-violet Other elements not reported	Protoplasmic neuroglia Neuroglia cells } dark Astrocytes } purple Third element - unstained Background - light purple Other elements not reported		Axis cylinders and dendrites Nerve cells and processes Neuroglia Connective tissue Fibres	silver impregnation	General Neuroglia Other elements not reported			
Cell bodies, axon and dendrites - black Neurofibrils Other elements not reported	Protoplasmic and fibrous astrocytes Microglia Oligodendroglia Neuroglia cells - purple red Nerve cells - pinkish Nerve fibres - nearly or quite unstained		Nerve cells and processes Neuroglia	dense black	General			
Neurofibrils - brown Neuroglia - may be slight colouring Other elements not reported	Astrocytes - black Other elements not reported			General Nuclei - blue Cytoplasm - pink	Glial fibres - blue Other elements not reported			
	Astrocytes Other elements not reported				Glial fibres Other elements not reported			
			General					
Nerve cells and processes Other elements not reported	Neuroglia - dark purple Other elements not reported	Nervous tissue Nissl granules	Nerve cells and processes Other elements not reported	Nerve cells and processes Other elements not reported	Neuroglia Other elements not reported		General Nervous tis grey	
Axons and dendrites - black Neurofibrils - black Neuroglia - may be lightly stained Other elements not reported	Astrocytes - opaque purplish-black Other elements not reported	Nissl substance - deep purple-blue Nuclei - blue Some cytoplasmic processes of neurones - blue Other elements not reported	Neurons - black Neuroglia - black	Nuclei - blue-black Neuroglial fibres Axons } Not stained Nerve endings Cytoplasm - pink	Fibrous glia - deep blue Other elements not reported		Nerve fibres - black Other elements not reported	
Neurofibrils in cell body, axons and dendrites of neurons - black Reticulum fibres - black Other elements not reported	Astrocytes and processes - black Nerve cells - pale red Nerve fibres - not stained Other elements not reported	Nissl substance - purple Other elements not reported	Nerve cells and processes Other elements not reported					
Neurofibrils, axon and dendrites Neuroglia Nerve endings Other elements not reported	Astrocytes - black Neuron bodies } also Microglial nuclei } stained Nerve endings Oligodendroglia } not Axons and myelin } stained sheath	Nervous tissue Nuclei - violet Plasma - blue Amyloid, mucin and mast cells granules - red	Neurons - black Neuroglia - black	General Fibrogliar fibrils - not stained Nuclei - blue				
Nerve cells and processes - black Neurofibrils Neuroglial processes Other elements not reported	Astrocytes - black Oligodendrocytes } black Background - light pinkish purple Other elements not reported			Nerve cells - pink Nuclei of nerve cells } blue Oligodendroglia Astrocytes Microglia Other elements not reported				

CELLULAR IDENTIFICATION – GENERAL COMMENTS

old's silver ate	Phosphotungstic acid haematoxylin	del Rio-Hortega's silver	Weigert-Pal	Weil and Davenport	Author	Comments of authors cited
			Myelinated nerves – blue Nerve cells and large axis cylinders – brown Neuroglia Fibrin Nuclei } blue		Mann (1902)	
			Medullated nerve fibres – blue-black Other elements not reported		Hartridge and Haynes (1930)	
dendroglia – black glia – light brown elements not sorted	Nuclei – blue Intercellular substances – pink	Protoplasmic and fibrous neuroglia Microglia Other elements not reported	* Myelin sheaths – blue; background – colourless Other elements not reported * Glia fibres and nuclei – blue Cytoplasm – colourless Other elements not reported	Oligodendroglia – brown Microglia – brown Other elements not reported	Gatenby and Painter (1937)	Bielschowsky – "According to our experience good results are rarely obtained". Golgi – "and of the fact that the impregnation may be limited sometimes to certain elements, sometimes to others, care should be exercised in the interpretation of the results obtained. A further source of possible error is found in the formation of precipitates which may, up to a point, simulate dendrites and other structures".
		Microglia Other elements not reported	Myelinated nerve fibres – black Other elements not reported		Carleton and Short (1960)	
	Pathologic astrocytes Mitochondria Nuclei Fibrin Fibroglia Microglia } blue Other elements not reported	Neurofibrils – black Other elements not reported	Myelin sheaths Other elements not reported		McManus and Mowry (1960)	
dendroglia and microglia elements not sorted			Glial fibres		Hosokawa (1963)	
					Mannen (1963)	Golgi – "As this technique is not specific for nerve cells and glial cells, both can be seen".
dendroglia] purplish-glia } blue-grey cytes	General	Microglia – dark grey Oligodendroglia Other elements not reported	Myelin sheaths Nerve cells and processes All nervous tissue	Microglia – brown Other elements not reported	Gray (1964)	Penfield – to show astrocytes as well – "simply leave the sections too long in the silver stain" del Rio Hortega – "nor does it stain oligodendroglia except by an occasional accident." Author uses term "recommended for"
dendroglia] black glia cytes – lightly lined elements not reported	Astrocytic glia } deep blue Nuclei Fibrin } blue Myelin – light blue Other elements not reported	* Astrocytes – brown-black * Oligodendroglia – black * Microglia – black Neurons not reported	Myelin sheaths – deep blue-black to blue Other elements not reported	Pathological glial – grey-black Other elements not reported	Drury and Wallington (1976)	
dendroglia] dark grey glia } to black elements not reported					Humason (1967)	Penfield's method – "long fixation tends to increase staining of microglia and astrocytes"
dendroglia] brown glia cytes is not reported	Pibrillar fibrils Myelial fibrils Neuroglial fibrils } blue Fibrin Other elements not reported	Astrocytes Oligodendroglia } grey Microglia Neurons not reported	Myelinated nerve fibres – dark blue Other elements not reported		McClung Jones (1967)	Golgi – "best staining procedure for classical neuroglia"
	Glial fibres – blue Other elements not reported	Microglia – black Other elements not reported	Myelinated nerve fibres – blue Other elements not reported		Cruickshank, Dodds and Gardner (1968)	

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

Table 10 continued

Bielschowsky's silver	Cajal's gold sublimate	Cresyl violet	Golgi-Cox silver	Haematoxylin and eosin	Holzer	Marsland, Glees and Erikson	Palmgren	Patay
					Astrocytes Other elements not reported			
Nerve cells and processes; neurofibrils Untoned: nerve fibres and neurofibrils - brown to black After gold toning - fibres - grey-black Background - pinkish grey to purplish Other elements not reported	Protoplasmic and fibrous astrocytes Astrocytes - black Background - unstained or light brownish purple Nerve cells - red Nerve fibres - not stained	Nissl granules and nuclei of neuroglia cells Nissl granules - purple to violet Nuclei of neuroglia and endothelial cells - slightly bluer than Nissl granules Other elements not reported	Nerve cells, axons and dendrons Nerve cells and processes - black Neuroglia - astrocytes sometimes stained black Background - dull yellow	Nuclei - blue Other elements not reported				
Neurofibrils in cell body, axons and dendrites of neurons - black	Astrocytes and processes - deep ruby to dark purple Neurons and axons - deep ruby to dark purple	Nissl substance - purple Glial cells - lilac Background - colourless	Nerve cells and fibres - black Neuroglia - sometimes blue	General Nuclei - blue-black		Nerve fibres - dark brown-black Other elements not reported	Nerve fibres - brown-black Other elements not reported	
Neurofibrils Other elements not reported		Nissl bodies in neurons Other elements not reported	Nerve cells and processes Other elements not reported	Nuclei - dark purple or blue Cytoplasm - pink				
Neurons, neurofibrils axons and dendrites - black	Astrocytes - red-black			Nervous system		Neurons, axons and dendrites - black		
Other elements not reported	Other elements not reported					Other elements not reported		
Nerve fibres - black Nuclei - brown Other elements not reported	Astrocytes Other elements not reported			Nerve cells Reactive microglia Astrocytes Microglia Other elements not reported				
				Nuclei - blue-black Cytoplasm - pink-red		axons and dendrites - black Other tissues light grey to yellowish brown		
		Nerve cell bodies and dendrites - blue Nuclei of oligodendrocytes - blue Nissl bodies in nerve cells - blue Other elements not reported	Nerve cells and processes - brown Nuclei of oligodendrocytes - brown Thin nerve fibres - brown-black Other elements not reported	Nerve cells and processes - purple Nuclei of oligodendrocytes - blue Cytoplasm - pink Small nerve cells - purple				
Cell bodies, axon and dendrites - black Neurofibrils Other elements not reported	Protoplasmic and fibrous astrocytes Microglia not Oligodendroglia stained Neuroglia cells - purple-red Nerve cells - pinkish Nerve fibres - nearly or quite unstained		Nerve cells and processes - black Neuroglia cells and processes - black	Nuclei - blue Karyosomes - dark blue Plasmosomes - red Cytoplasm - shades of pink				
Bielschowsky (1904) neurofibrils - black						Marsland, Glees and Erikson (1954) Rabbit neurons, anterior horn cells and fibres, boutons terminaux and free terminals - black precipitate Other elements not reported	Palmgren (1948) Nerve fibres and endings Cell bodies - red Nerve fibres - black "Hence within certain limits the selectivity of the nerve fibre staining is directly proportional to the delay of the precipitation"	

Cell types in bold indicate that the authors stated explicitly that the stain is specific for that cell type. Other cell types not in bold have been indicated to be stained by the procedure, but not specifically. Absence of an entry under a particular heading means that the author did not mention that procedure, sometimes because it had not yet been described. 'Other elements not reported' means that the authors have not mentioned whether or not other cell types or organelles are stained by that procedure. 'General' means that all neural elements are stained by that procedure, although not necessarily the same colour or intensity. An asterisk(*) means that the author mentions slight modifications of the procedure. Please see also Appendix 1.

CELLULAR IDENTIFICATION – GENERAL COMMENTS

Author's silver mate	Phosphotungstic acid haematoxylin	del Rio-Hortega's silver	Weigert-Pal	Weil and Davenport	Author	Comments of authors cited
	Astrocytes Other elements not reported				Uyeda, Eng and Bignami (1972)	
dendrocytes dark grey glia to black round - pale grey elements not reported	Tissue elements in general Neuroglia fibres Fibroglia fibres Myoglia Nuclei Fibrin	blue	*Protoplasmic and fibrous astrocytes Astrocytes - black Background - unstained or light brownish purple Nerve cells - red Nerve fibres - unstained Not reported *Oligodendrocytes Processes and cytoplasm - black Nuclei - practically unstained Background - grey Not reported *Microglia - black Other glial cells dark grey to black Background - pale	Myelin sheath - dark blue Other structures - unstained	Clark (1973)	Hortega's method for microglia is very similar to his method for oligodendroglia. "Although this method is probably the most selective for oligoden- drocytes, other types of cells may be stained particularly microglia. To secure staining of the former only, the time of fixation should be kept within 2 days, since long fixation tends to cause staining of microglia, astrocytes and even nerve cells".
glia black dendroglia elements not ported	Astrocytes - blue Neurons - pink Neuroglia Nuclei, fibrin blue Glial fibres Microglia		*Microglia - black Oligodendroglia - black *Astrocytes - black to dark grey Neurofibrils - black Neurons not reported	Myelin sheaths - dark blue-black Other elements not reported	Ralis, Beasley, and Ralis (1973)	"Cresyl violet can be used to show general cytoarchitecture of brain and spinal cord. Weil and Davenport's method may give some non-specific staining. Hortega's method for astrocytes may give some non-specific staining. Holzer-normal glia are unstained".
dendroglia - black glia - black glia elements not ported	Nervous tissue Nuclei, centrioles, fibroglia, myoglia, blue and neuroglia, fibres, fibrin			Myelin Other elements not reported	Bloom and Fawcett (1975)	
	Astrocytes and fibres Microglia Nerve cells Other elements not not reported		Nerve cells Neurofibrils Oligodendroglia Other elements not reported	Myelin sheaths - deep blue-black Not reported Other elements not reported	Oligodendroglia - black Microglia - black Neuroglia Other elements not reported	Culling (1975)
				Myelinated nerve fibres Other elements not reported	Greenfield, Blackwood and Corstellis (1976)	
	Nuclei Fibroglia Myoglia Neuroglia fibres light blue Centrioles Mitochondria Myelin Fibrin Other elements not not reported	light blue dark blue			Smith and Bruton (1977)	"Silver impregnations for nerve fibres are capricious and frequently fail". Phosphotungstic acid haematoxylin - "The fact that a number of structures are demonstrated by one colour in no way detracts from its usefulness, as the colour is present in many shades and the structure identification is helped by its morphology". "The length of staining time will determine the elements stained and degree of intensity".
	Nerve cells - blue Nerve fibres - grey Myelin - pale Other elements not reported			White matter, i.e. nerve fibres in tracts, stain dark grey. Nerve cells in grey matter stain lighter. Other elements not reported	Leeson and Leeson (1979)	
dendroglia bluish- grey elements not ported	Neuroglia - blue Fibrin - blue Other elements not reported		Myelin - blue-black Other elements not reported	*Protoplasmic astrocytes - violet *Fibrous astrocytes *Oligodendrocytes - grey *Neuroglia - deeply stained Astrocytes faintly Oligodendrocytes stained Glia fibres	Carleton, Drury and Wallington (1980)	Haematoxylin and eosin - "Specialized structures such as neuroglial fibres, dendrites etc. are not brought into evidence" Weigert-Pal - "A simple and rapid not specific method". Penfield "Differentiation between the two is easy on morphological grounds"
				Neurons not reported	Millhouse (1981)	
old (1930) dendroglia purple- glia grey elements not ported	Mallory (1900) Neuroglia fibres Nuclei, neuroglia fibres and fibrin - blue Axis cylinders and ganglion cells - pale pink Connective tissue - deep pink "The blue colour is a little sensitive to strong light, and on prolonged exposure will fade to pink" Other elements not reported		del Rio Hortega (1971) Neuroglia and connective tissue; Microglia and oligodendroglia - black Neurons not reported	Weigert (1885) Medullated fibres - dark blue to blue- black. Has a variety uses Other elements not reported	Weil and Davenport (1933) Oligodendroglia and microglia - brown Particularly good for "pathologic glial proliferations" Other elements not reported	

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

answer to the question, 'What are the electrical properties of neuroglial cells in the living intact adult mammalian central nervous system?' Obviously, this question is not answered by dubbing those cells as neuroglia, which one cannot see being penetrated by the electrode or subsequently marked histologically. Quite explicitly, the statement that an unexcitable cell is a neuroglial cell, followed by the deduction that neuroglial cells are not excitable, is a circular argument. It will be concluded later in this monograph that besides neurons, the only kinds of neuroglia are the naked nuclei free in the ubiquitous syncytium, and that what are generally called astrocytes and oligodendrocytes are neurons (page 189), so this problem does not arise for us.

Neuroglia have been widely characterized as cells 'supporting' or 'satellite' to neurons (Muller, 1930; Krypsin-Exner, 1952; Brownson, 1955, 1956; Hyden, 1962; Koenig, 1964; Friede, 1965; Peters, Palay and Webster, 1976, pages 231-266; Pevzner, 1979). Prima facie, it is difficult to know what is meant by this implicitly teleological term. It is highly likely that every sort of cell in any tissue interacts with its neighbouring extracellular fluids and cells. In this sense all tissue fluids and cells are interdependent, and are therefore mutually supporting. One cannot do the experiment of withdrawing particular cellular elements from any living tissue and examining the behaviour of the remaining tissue afterwards, as one does classically in ablation experiments of cerebral nuclei or endocrine organs.

In the middle 1960s at a meeting of the Brain Research Association in London, the distinguished neurobiologist, J.Z. Young, was talking about memory in the neurons of octopuses. He was asked by a questioner in what way his experiments could be interpreted as locating the memory to neurons rather than to the nuclei upon which he was experimenting. He agreed immediately that one could not attribute the properties to the neurons rather than to the central nervous nuclei as a whole. Nevertheless, it is widely believed that memory and many other probably cerebral functions are located in neurons, although there is no unequivocal evidence for this belief; in many senses this view is complementary to the view that neuroglia are 'supporting' cells.

The idea that the neuroglial cells support the neurons, or are 'satellites' of them, probably originated from the fact that some small neuroglia adjacent to neuron somata look like 'satellites' (Figure 23). Many biochemical experiments have been carried out on the cytochrome oxidase, succinic dehydrogenase, and ATPase in isolated neurons and their enzyme activities have been compared with those of 'neuroglial clumps' (Cummins and Hyden, 1962; Hamberger, 1963; Hyden, 1967). Their oxygen uptake was different, and the clumps responded to high potassium ions by increasing their oxygen uptake, while the neurons did not (Hertz, 1966, 1979). Of course, it is not at all surprising that, since all the cellular elements in the nervous system stain differently from the tissue between them, they should show different enzymic and other biochemical properties. It does not follow that if the same properties change in the opposite or in the same direction in neurons and neuroglia that one of them is nutritive, supporting, or satellite in relation to the others. Indeed, although much has been written about the neuron-glia relationship (Hertz, 1965; Hyden, 1967; Johnston and Roots, 1972; Schoffeniels, Franck, Tower and Hertz, 1978; Pevzner, 1979; Giacobini, Vernadakis and Shahar, 1980), it would be difficult, within the context of neurobiology, to define these terms expressing the concept that the neuroglia were somehow 'secondary' to neurons.

CELLULAR IDENTIFICATION – GENERAL COMMENTS




FIGURE 23 Penfield's (1928) drawings of OL-1 oligodendroglia, AS-1 fibrous astrocyte, OL-2 satellite oligodendroglia, AS-2, fibrous astrocyte. This figure is reproduced from Penfield, W. (1928) Neuroglia and microglia: the interstitial tissue of the central nervous system. In Special Cytology, edited by Cowdry, E.V. New York: Hoeber, page 1032.

Cells identified as neuroglia in tissue culture generally do not fire (Hild and Tasaki, 1962; Krnjevic and Schwartz, 1967; Nelson, 1973; Hosli, Andres and Hosli, 1980). Although it is generally believed that neuroglia do not fire, I think that insufficient determined experiments have yet been devoted to posing the questions with an affirmative expectation. The slow potential changes due to the application of aspartate and glutamate are similar in neurons and astrocytes (Hosli, Andres and Hosli, 1979). If one were to show clearly that astrocytes and oligodendrocytes, which had been precisely identified by independent and dependable criteria – under precisely the same conditions as neurons – did not fire, one could use this criterion to distinguish the neurons from the two types of neuroglia. It may be that the necessary exhaustive examinations will be carried out in the future, and electrical excitability will then become a useful criterion.

Microglia and reactive astrocytes are generally considered to be phagocytic. Microglia cannot often be identified with certainty in tissue culture, and it is not at all clear from the literature whether protoplasmic and fibrillary astrocytes are considered to be the same cells as reactive astrocytes, or transformed cells or different cells.

SPECIFIC CELL MARKERS (TABLE 10)

In claiming that a particular marker is 'specific' for a cell type, one would not be

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

satisfied if it were present in preparations of other cell types to any significant degree, since its presence at all indicates that those cells can also synthesize the marker. In some conditions they may synthesize more or less, but the ability to synthesize a naturally occurring marker must derive from the genotype. Therefore, even an apparent concentration in one cell type, in a particular preparation of ten times the concentration in another, does not merit the characterization of a substance as a specific marker, especially in view of the variability of marker concentrations between whole homogenates, synaptosomes, organ cultures, tissue cultures, cell clones, tumours, etc. of the same parent tissue origin. Pevzner (1979, pages 51-63) shows a table of 51 enzymes in 186 preparations, none of which show the enzymes too low for their activities to be measured (Table 7). Therefore, none of these enzymes can be considered specific to a cell type.

Whereas the proteins, transmitters and enzymes occur naturally in or around the cells, the toxins and the antibodies do not. These substances are usually detected using histochemical or immunocytochemical techniques. The use of such techniques - as has been stressed previously - carries the implicit assumption that none of the steps of the procedures used affects the optical properties of the cells substantially. This should have been demonstrated for each step in each procedure for every preparation in the range of conditions under which it is being studied. Markers are sometimes demonstrated in tissue after homogenization - a procedure whose users must also believe that it does not affect the apparent concentration, or distribution, of the marker.

Of the markers, the enzymes and proteins are located within the cells and the toxins are outside the cells. The transmitters may be present inside or outside the cells, and the antibodies usually gain access to the inside of the cells after they have been 'permeabilized'.

It would be fair to say that all the microscopic techniques for indicating markers are qualitative rather than quantitative. They depend upon the following variables: each of the reagents used in the procedures; the concentrations of these reagents; the time for which the tissue has been exposed to each of them; the degree of 'permeabilization' of the cells; the penetration of the substrate or toxin; self-absorption by the specimen; the specificity of the reaction used; the system of microscopy and illumination employed; interference of the reaction demonstrating the marker by substances present in the tissue other than the marker. The effects of most of these variables can be minimized by using suitable 'control' preparations, but the empirical nature of most of the procedures - in addition to one's ignorance about the full impact of these variables on the measurement system - will probably ensure that these techniques will remain largely qualitative for quite a long time to come. Perhaps we will have to await a new epoch, when research workers will have a better stomach for 'control' experiments than they do at present.

APPEARANCES IN TISSUE CULTURE (TABLE 11)

Observations on tissue culture have the advantages that one is observing living whole cells. The cells are not dehydrated, shrunk or sectioned, and virtually all the geometrical problems are obviated. One can compare the relative dimensions of the somas and nuclei and the lengths of the dendrites and glial fibres. In the literature

CELLULAR IDENTIFICATION – GENERAL COMMENTS

there are some clear descriptions of the different kinds of cells in culture, but I cannot find any statistical studies comparing the measurable dimensions of the latter parameters between the different kinds of cells in culture. In tissue cultures the shapes of cells depend upon the following factors: the location in the parent tissue; the age of the parent tissue; the age of the culture; the nature of the culture medium; the frequency of the changes of medium; the number of passages of the culture. The culture may originate from: a whole embryonic organ; a whole adult organ; dissociated embryonic cells; dissociated adult cells; a natural tumour; an experimentally induced tumour. Tumours may be slowly or rapidly growing.

The real difficulty of studies of cell structure in tissue culture is the similarity of appearance of most cells in culture at some time or other during development. I think that tissue culturalists with many years' experience might have great difficulty in distinguishing, for example, cultures of fibroblasts at some stages from cultures of neurons, astrocytes or oligodendrocytes or tumours from them, purely on their appearances alone. It is very likely that they would immediately request knowledge of their tissue origin and clinical history before pronouncing on the identity of a whole culture and the cells in it. Most cells in culture – other than endothelial cells – appear 'neuron'-shaped, and have long processes which grow out in all directions. The processes and the cell bodies sometimes have a thin spread of granules on them. They show intracellular movements.

THE CURRENT VIEW OF THE DIFFERENT KINDS OF CELLS AS DESCRIBED IN THE LITERATURE

With these generalizations and reservations in mind, we may now examine the authoritative literature for descriptions of the different kinds of cells by light and electron microscopy. The intention of this section is to try to find out by what criteria histologists, electron microscopists, histochemists, neuropathologists, immunocytochemists, physiologists and tissue culturalists, distinguish the different cell types, and how real these distinctions are.

In Tables 8-11, one often sees empty columns, in which comparative information is absent from a particular publication, presumably because that information was not investigated in that series of experiments. An occasional space probably has no significance, but I would venture to suggest that sometimes there may well be some good, albeit esoteric, reason for frequent absence of data under a particular heading. This would have been regarded as 'singular' by a celebrated psychobiologist studying an aphasic dog, and would certainly have generated interpretable clues (Holmes, 1905). Clearly, one must distinguish between absence of information and an unequivocal statement denying the possession of a property. The English term, 'fail to show' can either mean that a research worker has tried but failed so far to demonstrate a property he believes to be possessed by his system, or that he has not yet investigated that property by accident or intention. One is entitled to seek an explanation of why such crucial information has not been recorded.

4

Specific Identification

LIGHT MICROSCOPY OF HISTOLOGICAL SECTIONS (TABLE 8)

The literature indicates that neurons may be spherical, elongated, pear-shaped, spindle-shaped, pyramidal, fusiform, conical, unipolar, bipolar, multipolar, polygonal, stellate, irregular or varied in shape; their maximum diameters in stained sections appear to be between 4 μm and 435 μm . The astrocytes are rarely described, but one description is implied in their name, star-shaped; they are the largest glial cells with diameters of 20 μm to 35 μm . Oligodendrocytes are round or oval. They are generally described as 'small' but the only measurement of their diameters is 10 μm to 20 μm . Microglia in normal tissue are, stellate, elongate, spherical, oval, bulb-like, bipolar, tripolar, multipolar, amoeboid, pleiomorphic; they are small or very small, or cannot be seen, or do not exist.

The number of processes - given the name dendrites - arising from neurons is variously put at being from one to thousands, in addition to one thicker axon. They taper, twist, ramify or bifurcate for a few to hundreds of μm , except for the axons which may reach tens of centimetres in length. Astrocytes also have many processes, which are seen to branch, and end frequently on capillaries or blood vessels; the latter connections are often regarded as specific criteria for identifying astrocytes. Oligodendrocytes are generally described as having small, delicate, few processes, or occasionally none at all. Microglia have few processes which branch two or three times; the processes may be straight or tortuous.

By light microscopy the nuclei of neurons appear large, spherical or ovoid and diffuse chromatin is spread within them; they contain conspicuous nucleoli, which are usually centrally placed. The nuclei of astrocytes are 5 μm to 10 μm in diameter, smaller than those of neurons and larger than those of oligodendrocytes. Oligodendrocytic nuclei are often described as eccentric, and often stain darkly. The nuclei of microglia appear spherical, oval or elongate, and stain darkly; nucleoli are rarely reported in microglial nuclei, and most authors seem to regard them as absent.

Neuronal cytoplasm contains Nissl granules, which do not appear in the cytoplasm of other cells. Many authors see a Golgi apparatus, mitochondria and cytoplasmic fibrils; lipofuscin granules may be present, and centrosomes may sometimes be identified. The cytoplasm of astrocytes is not so frequently described. It is sometimes said to be paler and Nissl granules are not reported in it, but Golgi apparatuses,

SPECIFIC IDENTIFICATION

mitochondria and centrosomes can be seen. Oligodendrocytes either have a thin layer of cytoplasm with no fibrils, or they are said to contain Golgi apparatuses, fibrils, lipofuscin and centrosomes. Microglia either have scanty or barely distinguishable cytoplasm. In some publications they contain dense cytoplasm with granules and mitochondria but no Golgi apparatuses or Nissl granules.

APPEARANCES BY ELECTRON MICROSCOPY (TABLE 9)

From the tables one can see that few publications describe the overall appearances of cells by electron microscopy, presumably for the following reasons: firstly, it is believed that the appearances have already been characterized by light microscopy; secondly, the section thicknesses are such that many serial sections would be necessary to describe the whole outline of a single cell; thirdly, one cannot describe the shapes accurately because they are distorted during preparation for electron microscopy; fourthly, the particular authors were not interested in the shapes of the cell bodies.

The neurons appear round, pear-shaped, multipolar, polygonal, angular, irregular or smooth (Table 9). Descriptions of the cell bodies of astrocytes are rare, but they deal mainly with the processes. The processes end on vascular capillaries; astrocytes have less processes than neurons do; they are branched and delicate. There are also few descriptions of oligodendrocytes; they are said to have few to numerous processes, often thin and branching. Microglia are small, may be indented, and have several processes.

The cytoplasm of neurons contains granules believed to be lipofuscin. The cytoplasm of neurons and all types of neuroglia contain Nissl bodies, endoplasmic reticulum, Golgi bodies, mitochondria with cristae, ribosomes, polyribosomes, granules and varying proportions of filaments, tubules and fibrils. The astrocytes have paler cytoplasm than the neurons, while the oligodendrocytes have denser cytoplasm; the microglia have been described as having no cytoplasm, a little cytoplasm or dense cytoplasm (Table 9).

The nuclei of neurons appear 'large'. They may be spherical, oval, lobulated or indented. They may be centric or eccentric. The nuclear membranes appear to be trilaminar, and to be punctuated by nuclear pores or nuclear pore apparatuses. The nuclei contain granules or filaments of chromatin. The nuclei of astrocytes appear as having various shapes: irregular, ovoid, indented, or reniform. They may be centric or eccentric, and appear large or 'prominent'. They also have nuclear pores or nuclear pore apparatuses, and a trilaminar membrane. The nucleoplasm contains masses or granules believed to consist of chromatin.

The nuclei of oligodendrocytes are round or oval, and appear dense. They also have a trilaminar membrane with nuclear pores, and they contain clumps of chromatin. The nuclei of microglia are also oval, elongate, angular, variable or irregular; they are usually dense or light. They contain clumps of chromatin. There seems to be little information on whether or not they have nuclear pores.

The nucleoli of neurons are 'large', centric or eccentric, but have no limiting membranes, visible by electron microscopy. They have a dense granular or filamentous mass, the nucleolonema. Occasionally there is more than one nucleolus in a neuron. The nucleoli of astrocytes are described as absent, occasional, constant or prominent; their shapes have not been examined in detail.

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

In respect of nucleoli, there is unfortunately a lack of information about oligodendrocytes, as there is with microglia. One cannot tell from the literature whether the nucleoli cannot be seen, or cannot be recognized, or do not exist or have not been examined specifically and systematically, or have not been examined at all.

The references given in these tables were not selected for their diversity of views, but represent all the authorities available to the author. Under these circumstances, careful perusal of the original references, or Tables 8 and 9, or my summary of them, leads to the following inescapable conclusion. In the literature the descriptions of the different kinds of cells in the nervous system by light or electron microscopy either overlap so considerably, or are so evidently lacking in comparative data, that one cannot conclude with confidence that the different kinds of cells in the nervous system can be distinguished from each other by light or electron microscopy.

APPEARANCES IN TISSUE CULTURE (TABLE 11)

Prima facie, one would hope that descriptions of cells in tissue culture would be similar to those of cells identified as particular types in histological sections made from adult tissue. Neurons in culture may have cell bodies which are spherical, bipolar or multipolar. They may be relatively large, like Purkinje cells, or small, like dorsal root ganglion cells. They have fine, branching, filamentous processes; filaments are seen in the cell bodies of some of them (Table 11).

Astrocytes have stellate cell bodies with wide undulating processes at their origins, or with multiple branching. Some astrocytes are often described as 'fibroblastic' in appearance. The cell bodies may be contractile, and they may contain more than one nucleus.

Oligodendrocytes may be difficult to distinguish from astrocytes in culture. They appear globose and often have dense nuclei. They also have extensively branching processes, sometimes ending in membranous expansions, but usually fewer in number than the astrocytes possess. The appearance of microglia in culture does not seem to have been described, although motile phagocytic cells are often seen, especially by time-lapse photography.

One may conclude from these descriptions in the literature that neurons, astrocytes and oligodendrocytes cannot be distinguished from each other in culture by their appearances alone, since they overlap so obviously. Identification usually requires knowledge of their tissue origin or pathology, their staining properties and their possession of markers believed to be characteristic for their cell type.

SPECIFIC STAINING

Many colour dyes used in histology, such as methylene blue, haematoxylin, eosin and cresyl violet, are nowadays considered non-specific. The metal stains were developed for biological purposes from their use in photographic techniques, and they were nearly all variations of reactions involving reduction of silver salts. Indeed, before the modern sophisticated mixtures were concocted for colour photography, there were already in commercial use complete systems of colour photography based on silver salts of different hues.

One can list the increasing intensities of colour with which cells in the nervous

SPECIFIC IDENTIFICATION

Table 11 The appearance of allegedly different cellular elements in tissue cultures as seen and described by the authors by phase-contrast microscopy. Lack of information under any heading indicates that it was not supplied in that publication

Neurons	Astrocytes	Oligodendrocytes	Microglia	Author/tissue
Purkinje cells identified by large size and characteristic structure in living preparations, their affinity for silver salts, their content of neurofibrils and basophilic substance. Also present large numbers of small granule cells with delicate processes which could be confused with oligodendrocytes using phase optics and methylene blue. Granules have a clear nucleus and a prominent nucleolus. Well-developed non-membranous processes	Processes form inter-lacing networks, are dense and characterised by presence of undulating membranes; processes have a broad cone shaped base	Similar size cells to granule cells but the perinuclear zone often obscured; processes generally arise directly as a narrow process from the perinuclear cytoplasm; often appeared to glow under phase contrast microscopy, due to their globose form and their rich granular content which cast a halo around the perinuclear cytoplasm		Pomerat and Costero (1956) cat - cerebellum
Identified pyramidal cells, giant pyramidal cells, giant polymorphous neurons and bipolar spindle neurons Cell bodies very from 30 to 300 µm in length; some binucleate and polynucleated neurons; generally one nucleolus present, but 2 or 3 may appear and 1 or 2 nucleolar satellites may be present; neuro-fibrils present in cell bodies as well as in neuronal processes				Geiger (1957) human, rabbit and monkey - cerebral cortex
Cell bodies are small and spherical and show marked refractivity, processes are fine, short and anastomose richly; difficult to distinguish from astrocytes in some cases	Stellate or angulated cells with processes of very great length; processes are slender, arise fairly sharply from the perikaryon; they branch and anastomose freely, appear sticky, elastic, lateralized rather polymorphic nuclei, many are binucleated, sometimes nuclei show amitotic features	Cannot distinguish between primitive oligodendrocytes and astrocytes in culture		Lumsden (1958) human - brain
	In the early stages of cultivation astrocytes from different parts of the brain form a three dimensional network of various densities; the cells are connected by their processes; characteristic shaking movement of the cell bodies; in a thin zone of outgrowth astrocytes present a typical network of stellate processes but at the periphery there is a gradual transformation so more similar to ependyma cells	Small cells with globose, vase-shaped or rounded perinuclear zones rich in highly refractile granules and droplets which tend to obscure the nucleus. A small number of processes arise abruptly (i.e. no cone-shaped base) from the perinuclear cytoplasm		Pomerat (1958) cat - brain and spinal cord
Neurons extended long, delicate processes that form inter-connecting bundles; heterogeneous in size and shape	Fibroblastic	Processes which are usually long extensively branched, dark and thorny; some have small, dark nuclei and dark cytoplasm, some are larger and lighter; processes often end in large membranous expansions	Macrophages - vacuolated cytoplasm which frequently contain debris; occasionally extend short processes	Raff, Fields, Hakamori, Mirsky, Pruss and Winter (1979) rat - optic nerve, cerebellum, corpus callosum and cerebral cortex
10% of cells in dorsal root ganglia culture have typical neuronal morphology, with large refractile cell bodies and long, fine processes	GFAP+ (glial fibrillary acid protein) cells - some large with large vesicular nuclei, others smaller and darker with multiple branching processes	GC+ (galactocerebroside) cells are usually small, with dark nuclei and cytoplasm, 50% of these have multiple, sometimes branching processes, others have rudimentary processes or none at all		Kennedy, Lisak and Raff (1980) human fetuses - optic nerve, spinal cord, dorsal root ganglion
Small and bipolar, or large and or multipolar	Most are flat, fibroblastic - like cells with filament bundles extending into 1 or 2 long processes, some are smaller and darker (GFAP+)	Dark nuclei and cytoplasm; and usually have branched processes (CG+)		Abney, Bartlett and Raff (1981) rat - brain

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

system are stained: blue-violet coloured solutions precipitate as black granules; grey also precipitates as black granules; red-ruby-purple solutions precipitate as brown or black granules. Examination of Table 10 shows that virtually all 'specific' stains eventually show up on cell bodies or fibres as black or brown granules of silver salts or silver metal. One can scan the vertical columns of Table 10, and one cannot fail to see that there is no single staining procedure which all, or even most, authors would agree to be specific for a type of cell or fibre in the nervous system. Of course, individual authors do claim or believe in the specificity of particular procedures indicated in the tables. Part of the problem here is the fundamental one that it is difficult to judge the relative value of different observations by different authors using basically - but not exactly - similar procedures and reagents, when they have often studied cell types in different mammalian species, of different ages, in different nuclei. (Please see also Appendix 1).

Here again, we must be humbled by the conscience of doubt, as well as our unwillingness to adopt an agnostic position, into concluding that staining procedures do not permit us to identify unequivocally the four major types of cell in the nervous system or their processes.

MARKERS (TABLE 12)

'Markers' may be proteins, enzymes, amino acids, transmitters or antibodies. Obviously, the concept of a marker implies that it is thought to be relatively or completely specific for the cell type it is believed to mark. In the list of markers (Table 12) many of the authors have not stated whether or not they have examined the reactions of cell types other than those they believe to be shown up by the markers; in the table these are indicated by the term 'not reported', whereas others, marked 'not labelled', mean that the authors have shown that the markers were not present in the particular cell types. Unfortunately, even the clear demonstration of absence of a marker cannot be used as a certain criterion for the identification of a particular cell type, since frequently the absence of the reaction is itself the only criterion for the identification of the type of cell itself. Thus, it would be a circular argument to define the specificity of a marker for a particular type of cell by the belief that the particular cell is identified by the specificity of the marker.

Many markers, such as tetanus toxin, glial fibrillary acid protein, S-100 or glutamate synthetase, were initially regarded as specific markers for particular cell types until their presence in, or affinity for, other cell types was examined. However, when we take a bird's-eye view of a very large number of markers nestling in the literature and the specificities claimed by the different authors, we come to the inescapable if disappointing conclusion that there is no marker upon whose specificity all authors would agree. (Please see also Appendix 2).

Most of the markers have been tested in tissue cultures, and their specificity may change quantitatively or qualitatively during the life of the culture, and may also be different in the host tissue than in the culture.

One could possibly differentiate the cell types by the intensity with which they are marked, but the concept of specificity rather implies an 'all-or-none' phenomenon, especially in relation to genetic origin or immunological reactions.

SPECIFIC IDENTIFICATION

CONCLUSIONS FROM THE LITERATURE

The evidence in the literature indicates massive and sincere differences of opinion among responsible neurobiologists about the identification of what are believed to be different types of cell, in relation to the following criteria: appearances in adult tissues by light and electron microscopy; light microscopy of cell types in tissue culture; staining procedures believed to be specific; markers generally regarded as specific. In addition, many authors have made the point that they themselves cannot distinguish between the cell types, especially of the neuroglia, or have used the term neuroglia, which implies obviously that they cannot distinguish between the different types of cells described by this term.

We will attempt to draw a conclusion about the reason for all of these difficulties (please see page 205).

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

Table 12 Enzymes, proteins, antibodies, transmitters, and a toxin, which are widely used as markers for different neural elements in tissue culture or immunochemistry. There are no markers for microglia

Marker	Neurones	Astrocytes	Oligodendrocytes	Author
Enzymes				
Carbonic anhydrase	Carbonic anhydrase*	Carbonic anhydrase*	Carbonic anhydrase*	Giacobini (1961)
Glutamate decarboxylase	Glutamate decarboxylase	Glutamate decarboxylase	Glutamate decarboxylase	Roberts (1962)
Gamma aminobutyric acid transaminase	Gamma aminobutyric acid transaminase	Gamma aminobutyric acid transaminase	Gamma aminobutyric acid transaminase	
Beta-galactosidase	Beta-galactosidase*	Beta-galactosidase*	Beta-galactosidase*	Hirsch (1972)
Ceramide glucosyltransferase	Ceramide glucosyltransferase	Ceramide glucosyltransferase	Ceramide glucosyltransferase	Radin, Brenkert, Arora, Sellinger and Flangas (1972)
Beta-galactosidase	Beta-galactosidase	Beta-galactosidase	Beta-galactosidase	Raghavan, Rhoads and Kanfer (1972)
Alpha-glycerophosphate dehydrogenase	Alpha-glycerophosphate dehydrogenase	Alpha-glycerophosphate dehydrogenase	Alpha-glycerophosphate dehydrogenase	Sellinger and Santiago (1972)
Beta-galactosidase	Beta-galactosidase*	Beta-galactosidase*	Beta-galactosidase*	Sinha and Rose (1972)
Carbonic anhydrase	Carbonic anhydrase*	Carbonic anhydrase*	Carbonic anhydrase*	Murai (1973)
Carbonic anhydrase	Carbonic anhydrase*	Carbonic anhydrase*	Carbonic anhydrase*	Nagata, Mikoshiba and Tsukada (1974)
Carbonic anhydrase	Carbonic anhydrase*	Carbonic anhydrase*	Carbonic anhydrase*	Hamberger and Sellstrom (1975)
Not reported	Not reported	2',3'-cyclic nucleotide 3'-phosphohydrolase		Benda (1978)
Not reported	Not reported	Glycerol-phosphate-3-dehydrogenase		
Not reported	Glutamate dehydrogenase	Not reported		Berl, Nicklas and Clarke (1978)
L-glutamate decarboxylase	L-glutamate decarboxylase	L-glutamate decarboxylase		Schousboe (1978)
Not reported	Glutamate synthetase	Glutamate synthetase		Stahl, Spence, Coates and Broderson (1978)
Not reported	Glutamine synthetase	Glutamine synthetase		Tower (1978)
Glycerol-3-P-dehydrogenase	Glycerol-3-P-dehydrogenase	Glycerol-3-P-dehydrogenase		Tower (1978)
Carbonic anhydrase	Carbonic anhydrase	Carbonic anhydrase*		Varon (1978)
Not reported	Glycerol-phosphate dehydrogenase	Glycerol-phosphate dehydrogenase		
2',3'-cyclic nucleotide 3'-phosphohydrolase	2',3'-cyclic nucleotide 3'-phosphohydrolase	2',3'-cyclic nucleotide 3'-phosphohydrolase		
Not reported	Glutamine synthetase	Glutamine synthetase		
Not reported	Not reported	Glutamine synthetase		
Not reported	Not reported	Galactosyltransferase		
Not reported	3-hydroxy,3-methyl glutaryl co enzyme A reductase	3-hydroxy,3-methyl glutaryl co enzyme A reductase		
Not reported	Glutamine synthetase	Glutamine synthetase		Norenberg and Martinez-Hernandez (1979)
Pyruvate phosphokinase	Pyruvate phosphokinase	Pyruvate phosphokinase		Pevzner (1979)
NAD pyrophosphorylase	NAD pyrophosphorylase	NAD pyrophosphorylase		
Glutaminase	Glutaminase	Not reported		Schousboe, Nissen, Bock, Sapirstein, Juurlink and Hertz (1980)
Not reported	Carbonic anhydrase	Carbonic anhydrase		
Glutamate decarboxylase	Not labelled	Not reported		
Not reported	Glutamine synthetase	Not reported		
Not reported	Not reported	Carbonic anhydrase		Sensenbrenner, Labourdette Delaunoy, Pettman, Devilliers, Moonen and Bock (1980)
Not reported	Glutamine synthetase	Not reported		Vernadakis, Parker and Norenberg (1980)
Not reported	Not reported	2',3'-cyclic nucleotide 3'-phosphohydrolase		
Not reported	Not reported	Glycerol phosphate dehydrogenase		
Not reported	Not reported	Carbonic anhydrase isoenzyme II		Ghandour, Derer, Labourdette, Delaunoy and Langley (1981)
Not reported	Not reported	2',3'-cyclic AMP phosphohydrolase		
Not reported	Butyrylcholinesterase	Not reported		
Not reported	Glutamine synthetase	Not reported		White, Dutton and Norenberg (1981)

SPECIFIC IDENTIFICATION

Table 12 continued

Marker	Neurones	Astrocytes	Oligodendrocytes	Author
Enzymes	Not reported	Not reported	2',3'-cyclic nucleotide-3'-phosphohydrolase	Hansson (1982)
Proteins	S-100 Not reported	S-100* Glial fibrillary acid	S-100* Not reported	Hyden and McEwen (1966) Eng, Vanderhaeghen, Bignami and Gerstl (1971)
	14-3-2	14-3-2	14-3-2	Packman, Blomstrand and Hamberger (1971)
	Not reported	Glial fibrillary acidic	Not reported	Bignami, Eng, Dahl and Uyeda (1972)
	S-100	S-100*	S-100*	Moore (1972)
	14-3-2	14-3-2	14-3-2	
	S-100	S-100*	S-100*	Satake (1972)
	14-3-2	14-3-2	14-3-2	
	S-100	S-100*	S-100*	Uyeda, Eng and Bignami (1972)
	Glial fibrillary acidic	Glial fibrillary acidic*	Glial fibrillary acidic*	Jacque, Jørgensen, Baumann and Bock (1981)
	Not reported	S-100*	S-100*	
	14-3-2	Not reported	Not reported	
	Not reported	Glial fibrillary acidic	Not reported	Hamberger and Sellstrom (1975)
	S-100	S-100*	S-100*	Hertz (1977)
	S-100	Not reported	Not reported	
	14-3-2	Not reported	Not reported	Benda (1978)
	Not reported	Glial fibrillary acidic	Not reported	
	S-100	S-100	S-100	Fedoroff (1978)
	Not reported	Glial fibrillary acidic	Not reported	Gheuens, Lowenthal, Karcher and Noppe (1978)
	Not reported	Glial fibrillary acidic	Not reported	Hertz, Schousboe, Boehler, Mukerji and Fedoroff (1978)
	Not reported	Glial fibrillary acidic	Not reported	Jacobson (1978)
	14-3-2	Not reported	Not reported	
	Not reported	Glial fibrillary acidic	Not reported	Jacque and Baumann (1978)
	Not reported	Not reported	Wolfgren W1 and W2	Raff, Mirsky, Fields, Lisak, Dorfman, Silberberg, Gregson Leibowitz and Kennedy (1978)
	Not reported	Glial fibrillary acidic	Not labelled	Schousboe (1978a) Varon (1978)
	Not reported	Glial fibrillary acidic	Not reported	Raff, Fields, Hakamori, Mirsky, Pruss and Winter (1979)
	S-100	S-100		Balazs, Regan, Meier, Woodhams, Wilkin, Patel and Gordon (198?)
	Not reported	Glial fibrillary acidic	Not reported	Bignami, Kozak and Dahl (1980)
	Not reported	Glial fibrillary acidic	Not reported	Currie (1980)
	Not labelled	Glial fibrillary acidic	Not labelled	Kennedy, Lisak and Raff (1980)
	Not reported	Glial fibrillary acidic	Not reported	Schousboe, Nissen, Bock Sapirstein, Juurlink and Hertz (1980)
	Not reported	Glial fibrillary acidic	Glial fibrillary acidic	Schlörch-Rathgeb and Monard (1980)
	Not reported	Glial fibrillary acidic	Not reported	Sensenbrenner, Labourdette, Delaunoy, Pettman, Devilliers, Moonen and Bock (1980)
	Not labelled	Glial fibrillary acidic	Not labelled	Abney, Bartlett and Raff (1981)

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

Table 12 continued

Marker	Neurones	Astrocytes	Oligodendrocytes	Author
Proteins	Not labelled	Glial fibrillary acidic	Not labelled	Bartlett, Noble, Pruss, Raff, Rattray and Williams (1981)
	Not reported	Glial fibrillary acidic	Glial fibrillary	Cohen and Selvendran (1981)
	Not reported	Glial fibrillary acidic	Not reported	Currie and Kelly (1981)
	Not reported	Glial fibrillary acidic	Not reported	Ghandour, Derer, Labourdette, Delaunoy and Langley (1981)
14-3-2		Not reported	Not reported	Jacque, Jørgensen, Baumann and Bock (1981)
	Not reported	Glial fibrillary acidic	Not reported	White, Dutton and Norenberg (1981)
	Not reported	Glial fibrillary acidic	Not reported	Hansson (1982)
14-3-2		Not reported	Not reported	
Not reported		Glial fibrillary acidic	Not reported	
Not reported		Alpha-albumin	Not reported	
Not reported		Not reported	Myelin basic protein	
Not reported		S-100	S-100*	
Not reported		Glial fibrillary acidic	Not reported	Meier, Regan, Balazs and Wilkin (1982)
Anti-bodies	D1, D2 and D3 antigens	Not reported	Not reported	Jacque, Jørgensen and Bock (1979)
	Not reported	NS1 antigen	NS1 antigen	Benda (1978)
	Not reported	G1 and G2 antigens	G1 and G2 antigens	
	N1 and N2 antigens	Not labelled	Not labelled	Jacque and Baumann (1978)
	D1, D2 and D3 antigens	Not reported	Not reported	
	Not reported	NS1 antigen*	NS1 antigen*	Varon (1978)
	Not reported	Anti corpus callosum sera*	Anti corpus callosum sera*	
	Not reported	G1 and G2 antigens*	G1 and G2 antigens*	
	Not reported	Not reported	Anti oligodendrocyte serum*	
	Neural filament antisera	Not labelled	Not labelled	Bignami, Kozak and Dahl (1980)
	Not labelled	Neural-antigen 2 (Ran 2)	Not labelled	Abney, Bartlett and Raff (1981)
	Not labelled	Neural-antigen 2 (Ran 2)	Not labelled	Bartlett, Noble, Pruss, Raff, Rattray and Williams (1981)
	D1, D2 and D3 antigens	Not reported	Not reported	Jacque, Jørgensen, Baumann and Bock (1981)
	D5 antigen	D5 antigen	D5 antigen	
	D1, D2 and D3 antigens	Not reported	Not reported	Meier, Regan, Balazs and Wilkin (1982)
Transmitters	Gamma aminobutyric acid	Gamma aminobutyric acid	Gamma aminobutyric acid	Roberts (1962)
	Dopamine	Not reported	Not reported	Hillarp, Fuxe and Dahlstrom (1966)
	Noradrenaline	Not reported	Not reported	
	5-Hydroxytryptamine	Not reported	Not reported	
	Gamma aminobutyric acid	Not reported	Not reported	Obata (1972)
	Gamma aminobutyric acid	Not labelled	Not labelled	Schousboe, Svenneby and Hertz (1977)
	Gamma aminobutyric acid	Gamma aminobutyric acid	Gamma aminobutyric acid	Kelly and Dick (1978)
	Gamma aminobutyric acid	Gamma aminobutyric acid*	Gamma aminobutyric acid*	Schousboe (1978a)
	Not reported	Acetylcholine*	Acetylcholine	
	Not reported	Dopamine*	Dopamine*	Varon (1978)
	Not reported	Adenosine*	Adenosine*	
	Not reported	Nor-epinephrine*	Nor-epinephrine*	
	Gamma aminobutyric acid	Gamma aminobutyric acid	Not reported	Currie and Kelly (1981)
	Gamma aminobutyric acid	Gamma aminobutyric acid	Gamma aminobutyric acid	Wilkin, Csillag, Balazs, Kingsbury, Wilson and Johnson (1981)
	glycine	glycine	glycine	
Toxin	Tetanus	Not labelled	Not labelled	Dimpfel, Neale and Haberman (1975)
	Tetanus	Tetanus	Tetanus	Mirsky, Wendon, Black, Stolkin and Bray (1978)

SPECIFIC IDENTIFICATION

Table 12 continued

Marker	Neurones	Astrocytes	Oligodendrocytes	Author
Toxin	Tetanus	Not reported	Not reported	Raff, Mirsky, Fields, Lisak, Dorfman, Silberberg, Gregson, Leibowitz and Kennedy (1978)
	Tetanus	Not labelled	Not labelled	Raff, Fields, Hakamori, Mirsky, Pruss and Winter (1979)
	Tetanus	Not reported	Not reported	Balazs, Regan, Meier Woodhams, Wilkin, Patel and Gordon (1980)
	Tetanus	Not labelled	Not labelled	Currie (1980)
	Tetanus	Not labelled	Not labelled	Kennedy, Lisak and Raff (1980)
	Tetanus	Not labelled	Not labelled	Abney, Bartlett and Raff (1981)
	Tetanus	Not labelled	Not labelled	Bartlett, Noble, Pruss, Raff, Rattray and Williams (1981)
	Tetanus	Not reported	Not reported	Cohen and Selvendran (1981)
	Tetanus	Not reported	Not reported	Currie and Kelly (1981)
	Tetanus	Not reported	Not reported	Meier, Regan, Balazs and Wilkin (1982)
Lipid	Not reported	Not reported	Galactocerebroside	Jacque and Baumann (1978)
	Not reported	Not labelled	Galactocerebroside	Raff, Mirsky, Fields, Lisak, Dorfman, Silberberg, Gregson, Leibowitz and Kennedy (1978)
	Not labelled	Not labelled	Galactocerebroside	Lisak, Abramsky, Dorfman, George, Manning, Pleasure, Saida and Silberberg (1979)
	Not labelled	Not labelled	Galactocerebroside	Currie (1980)
	Not labelled	Not labelled	Galactocerebroside	Kennedy, Lisak and Raff (1980)
	Not labelled	Not labelled	Galactocerebroside	Abney, Bartlett and Raff (1981)
	Not labelled	Not labelled	Galactocerebroside	Bartlett, Noble, Pruss, Williams (1981)

This table was compiled from the references indicated, and is not claimed to be comprehensive. 'Not reported' means that the authors have not stated whether or not they examined whether the markers were present in cells other than those indicated. 'Not labelled' means that the authors have shown that the particular marker was not present in the cells indicated. Extremely few authors state by what criteria they identify the particular cell types. An asterisk (*) means that the authors use the generic term 'neuroglia' without indicating a particular cell type.

5

Experimental Studies

HISTOLOGY

Hyden (1959) dissected out single neurons and the pieces of tissue which are adjacent to them (Figure 29) (Hyden, 1959; Hyden and Pigon, 1960; Hyden 1961). They usually picked out clumps of approximately the same volume as the Deiters' neurons they were studying, in order to facilitate comparison of the biochemical properties of the neurons and neuroglia. In view of the paucity of histological studies, we took out pieces of tissue adjacent to neurons, and stained them with the following routine procedures: haematoxylin and eosin; Weil and Davenport (1933); Mallory's (1900) phosphotungstic acid haematoxylin; Patay's (1934) triple stain; chromosome red; Marsland, Glees and Erikson's (1954) stain; Gallyas's (1963) stain (Hillman, Deutsch, Allen and Sartory, 1977). The glial clumps were sandwiched between pieces of liver. With the exception of some areas stained according to Marsland, Glees and Erikson, all the other staining procedures showed: the presence of a small number of star-shaped neurons; the presence of a much larger number of what appeared to be 'naked' nuclei; the tissue between these structures appeared largely unstained by any of the procedures; membranes were not detected by these procedures between the neurons and the naked nuclei, although Weil and Davenport's; Mallory's; Marsland, Glees and Erikson's; and Gallyas's stains are believed to show up neuroglia (Table 13). However, the most surprising, clear and repeatable observation was that with every one of these procedures the membranes between the liver cells could be detected quite clearly (Hillman, Deutsch, Allen and Sartory, 1977). We then concluded that there were no cell membranes in the glial clumps between the neurons - which, of course, have their own membranes - and the 'naked' nuclei. This meant that most of the tissue in neuroglial clumps was either part of a syncytium, or was extracellular. The 'naked' nuclei either represented the nuclei of such a syncytium, or they were cells with little or no cytoplasm. Indeed, the spherical nuclei which are sometimes described as reactive astrocytes, sometimes as oligodendrocytes and sometimes as microglia (Table 8) are usually characterized as having little or no cytoplasm, or having cytoplasm which does not stain with the particular procedures used.

EXPERIMENTAL STUDIES

TABLE 13 The staining procedures used in present experiments and the structures or compounds they are believed to show. These staining procedures can be found in Gray (1958), Gallyas (1963), Romeis (1968) and Ralis, Beasley and Ralis (1973)

Staining procedures	Used to show	Colour
Cajal	Astrocytes	Dark ruby to dark purple
Cresyl violet	Nissl substance	Purple
	Glial cells	Lilac
Gallyas	Nuclei and neuroglial processes	Black or grey-black
Golgi-Cox	Nerve cells and fibres	Black
	Neuroglia	Sometimes stained blue
Haematoxylin and eosin	All cells	Nuclei - blue black Other components - pink
	Glial fibres	Blue
Marsland, Glees and Erikson	Nerve fibres	Dark brown to black
Osmium tetroxide	Proteins and lipids	Black
Palmgren (1948)	Nerve fibres	Dark brown to black
Patay	Nuclei	Red
Weigert-Pal	Myelin sheaths	Dark blue to black
Weil and Davenport	Microglia	Black
	Oligodendroglia	Black
Phosphotungstic acid haematoxylin	Astrocytes	Blue




FIGURE 24 The mitochondria can be well seen in the frontal cortex of rat. The tissue was fixed and embedded in Carbowax (but not stained) and viewed by phase-contrast microscopy (Hillman and Sjuve, 1980). The bar is 5 μm

It was relatively easy to distinguish between the 'naked' nucleus as being surrounded by a syncytium or a curious kind of cell with little cytoplasm. We dissected out neurons and neuroglial clumps, and viewed them both by phase-contrast microscopy, and also after staining them with Janus Green-B neutral red. In all cases, the neuroglial clumps - like the neurons - were absolutely replete with mitochondria (Figure 24) in those parts of the brain (Hillman and Deutsch, 1978b). Therefore, the neuroglial material must be intracellular. This finding, together with the absence of membranes between the neurons and naked nuclei, led to the inescapable conclusion that the neuroglial clumps were, indeed, part of a syncytium.

Subsequently we did a further series of experiments, each on at least 10 pieces of the same tissues under each of the same conditions from 12 rabbits and 12 rats. The following additional experiments were carried out: firstly, whenever central nervous tissue was stained, pieces of kidney and liver from the same animals were taken out and stained simultaneously and with the same procedures; secondly, instead of examining tissue only from the medulla, we then studied pieces from the frontal lobes; the parietal lobes; the occipital lobes; the cerebellum; the thoracic spinal cord and the lumbar spinal cord; thirdly, instead of picking out neuroglial clumps under microscopic vision we placed the stainless steel wire at random points just below the surface of transverse 1 mm sections of the latter parts of the nervous tissue, and picked out pieces of tissue of 500-1000 μm in diameter. Thus, we were examining representative parts of the whole central nervous system, rather than tissue defined by its proximity to neurons or capillaries.

Once again, all these tissues showed the occasional neuron and the frequent naked nuclei in the ubiquitous syncytium (Figures 25-27). With all the staining systems - of which haematoxylin and eosin, cresyl violet and Weil and Davenport are shown - and Mallory's phosphotungstic acid haematoxylin also tested in this series, the membranes

EXPERIMENTAL STUDIES

around the kidney and liver cells could clearly be seen. In addition to staining the central nervous system with the latter stains it has also been examined after staining with all the nine other procedures in Table 13. In all cases no membranes could be seen between the neurons and the naked nuclei.

EXAMINATION OF UNSTAINED TISSUES


Different staining procedures distort the elements of a tissue to different extents. It may not be appreciated that the haematoxylin and eosin, the most popular procedure in use, involves 11 reagents in 14 steps. So we devised a simplified procedure, which ensures that all the tissues to be compared have been subjected to the same reagents and manipulations. This technique, involving five reagents in 10 steps, consisted of embedding tissues in paraffin wax, cutting sections 6 μm thick, mounting them in DPX, and examining them under phase-contrast microscopy (Hillman and Sjuve, 1980). The procedure shows up membranes clearly, as the embedding paraffin wax exaggerates phase differences to the detriment of colour differences, with which histologists and pathologists are much more familiar.

Once again, one saw only two elements, the apparently larger cells with processes, and the naked nuclei. No continuous membranes could be seen between the neurons and the naked nuclei (Figure 28). In addition to embedding the same seven tissues in paraffin wax, we also stained samples of each of them with 2% osmium tetroxide and embedded them in Epon. 'Thick' sections (2 μm) were cut for examination by phase-contrast microscopy, and 'thin' sections (80 nm) for electron microscopy. Once again both of these techniques revealed the same general features (Figure 29): (1) the 'large' approximately spherical cell with cytoplasm and processes, which are generally agreed to be neurons, and whose relative sparsity is known to every jobbing electron microscopist; (2) the spherical nuclei.

When 'naked nuclei' appear by light or electron microscopy to be attached by a 'foot' to a capillary, they are designated astrocytes. The problem with this identification is that, if one cannot see astrocytic 'feet' in a field, unless one examines serial sections, one cannot know whether the 'feet' cannot be seen because they are not in the relatively shallow plane of the section, or because they are not present. Of course, sections for electron microscopy are one-twentieth to one-fifteen-hundredth of the thickness of those used for light microscopy, so that the appearances of any processes would be much rarer on electron micrographs.

CELLS WITH APPARENTLY THIN FILMS OF CYTOPLASM

The cells apparently with little or no cytoplasm were examined carefully by bright-field illumination using approximately $\times 960$ magnification in preparations of rat and rabbit frontal, parietal, occipital, cerebellar, medullary sections, and thoracic and lumbar spinal cord sections, stained by all the staining procedures indicated (Table 13). It rapidly became obvious that the nuclei which appeared to be surrounded by thin rims of cytoplasm were simply out of focus. In any single section one can see a number of nuclei, and since the focal plane at this magnification is only about 1 μm deep, most nuclei are not clearly in focus at any particular setting of the fine adjustment.



FIGURES 25-27 Sections of rat kidney, liver and frontal cortex were stained simultaneously either with haematoxylin and eosin (Figure 25), or with cresyl violet (Figure 26), or with Weil and Davenport (Figure 27), and viewed by phase-contrast microscopy. These three staining procedures (shown) as well as phosphotungstic acid (not shown), all gave the clear appearance of a membrane around the renal tubule and hepatic cells, but none at all between the neurons and naked nuclei in the cortical cells (lower micrographs). It was thus concluded that no such membrane occurs in the latter situation. The bars in figures 25-27 are 50 μm

EXPERIMENTAL STUDIES




FIGURE 26

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM




FIGURE 27




FIGURE 28 Frontal lobe embedded in paraffin wax, not stained, but viewed by phase-contrast microscopy. The bar is 50 μm

The cells out of focus all seem to be encircled by a fine rim of cytoplasm. If one refocuses on one of the latter nuclei, others go out of focus, and they then appear to be surrounded by this much-described thin layer of cytoplasm.

The widespread differences of opinions among distinguished histologists about the specificity of particular staining procedures led us to pose the question, How sure are we that different procedures do indeed show up different cells? We examined this in the following way.

SERIAL SECTIONS

Serial sections were made of the five different brain areas and the two regions of the spinal cord, and each section in a series was stained with a different procedure in the order: haematoxylin and eosin; Patay's triple stain; Weil and Davenport; Mallory's phosphotungstic acid haematoxylin; Gallyas's; Marsland, Glees and Erikson. In these time-consuming experiments we were greatly aided by Mr James Kirby of the St Peter and St Paul's Hospital, London, to whom we wish to record our considerable indebtedness.

In some experiments we picked out an identifiable group of cells which appeared to be typical neurons, and followed them for two or three sections through the sequence of staining procedures. Obviously, since the sections were cut 6 μm thick, it was not easy to identify particular cell bodies, which appeared to have maximum diameters of




FIGURE 29 Sections of a (left) frontal, b (middle) parietal and c (right) occipital cortex of rat were made. They were fixed in glutaraldehyde, stained with osmium, embedded in Araldite, cut 2 μm thick and viewed by phase-contrast microscopy. Only two kinds of cells can be seen: the darkly staining neurons with processes, and the lightly staining naked nuclei usually surrounded by a shrinkage artifact. The bar is 50 μm

EXPERIMENTAL STUDIES

40-60 μm , through more than three to six sections. However, we could follow the same cells - when their positions formed a pattern - relatively easily in photographs of the same areas, and it became abundantly clear that the different procedures generally stained the same populations and not different ones (Figure 30). This could be demonstrated more precisely by counting what proportion of the cells with cytoplasm (other than single nuclei), which could be seen in one section could be identified in any subsequent one of a series. Examination of six serial sections of each of the same parts of the six areas under study showed that about 50% of the same cells could be followed from one section to the next (Table 14). Of course, if a staining procedure were specific for one cell type, only a small proportion of the cells in one serial section should show up in the next one in the series. More remarkably, approximately the same proportion of cells stained by one procedure appeared to be stained by a quite different procedure in the adjacent serial sections. In addition to the remarkable degree of overlap of elements stained, there was another very clear finding in every single section. That was that, with all stains, the bulk of the tissue was only lightly stained by any procedure. This observation was made not only in respect of the six stains mentioned above, but also for all the stains listed in Table 13.

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

TABLE 14 Serial sections were cut from rat frontal cortex 6 μm thick and stained with different stains. The same cells were identified by their positions from one section to the next serial section; the percentage of those seen in the second serial section originating from the first section is given in the third column of figures. The mean percentage for all stains and sections was 55%

	Number of cells stained	Number of cells identified in the first section seen in the second section	Percentage
<u>1st series</u>			
Gallyas: Weil and Davenport	50	24	48
Weil and Davenport: Patay	42	16	38
Patay: Haematoxylin and eosin	35	17	49
<u>2nd series</u>			
Gallyas: Weil and Davenport	84	40	48
Weil and Davenport: Patay	49	28	57
Patay: Haematoxylin and eosin	41	26	63
<u>3rd series</u>			
Gallyas: Weil and Davenport	55	19	34
Weil and Davenport: Patay	33	21	64
Patay: Haematoxylin and eosin	40	22	55
<u>4th series</u>			
Haematoxylin and eosin: Patay	71	30	43
Patay: Weil and Davenport	65	28	43
Weil and Davenport: Phosphotungstic acid haematoxylin	43	28	65
Phosphotungstic acid haematoxylin: Gallyas	44	19	43
<u>5th series</u>			
Haematoxylin and eosin: Patay	46	21	46
Patay: Weil and Davenport	39	19	49
Weil and Davenport: Phosphotungstic acid haematoxylin	37	18	49
Phosphotungstic acid haematoxylin: Gallyas	39	26	67

EXPERIMENTAL STUDIES




FIGURE 30 For legend please see page 104




FIGURE 30 Serial sections of the septal region of rat were cut and stained with (page 3, upper) haematoxylin and eosin; (middle) Patay's stain; (lower) Weil and Davenport; (page 104 upper), phosphotungstic acid haematoxylin (Mallory) and (lower) Gallyas's stain. The cells in each section which appear to be identifiable in the next section are marked by circles on the subsequent pictures. Careful examination reveals that a substantial proportion of the cells stained by one procedure are also stained by the others, that is, the stains (several of which are neuroglial, please see Table 10), are not specific. The apparent shapes, dimensions and positions of the cells obviously change because one is cutting them at different places, because the different staining procedures and the microtomes distort and displace the cells to different extents. Therefore, the identity of cells in successive serial sections is based on their relative position rather than their shapes. The bar is 50 µm

QUANTIMETRY

We quantitated the proportion of the sections of each tissue which were stained by procedures which some authors believed to be specific. If each procedure were specific for neurons, astrocytes, oligodendrocytes and microglia, then the addition of the areas should together add up to approximately 100%. If they were not specific they might add up to more than 100%, and any significant degree of non-specificity would exaggerate the total proportion of the volume of the tissue stained by all the procedures.

We employed the stains classically believed to show: neural cells and their processes (Marsland, Glees and Erikson, 1954); astrocytes, phosphotungstic acid haematoxylin (Mallory, 1900); oligodendrocytes, glia and microglia (Weil and Davenport, 1933). We also used known non-specific procedures, such as haematoxylin and eosin, Patay's

EXPERIMENTAL STUDIES

triple stain and Gallyas's stain (please see Table 13).

The procedure was as follows: the brains of 24 rats were fixed with 10% formalin and embedded in paraffin wax. Over 400 sections were cut across the whole brain and stained with the above procedures. Of these sections, four fields across each of the frontal, parietal, and occipital lobes and of the medulla were taken at random; each of these fields had been stained with one of the six procedures listed in the preceding paragraphs. The fields were examined with bright-field illumination giving a magnification on the screen of $\times 450$.

In each slide, twenty different fields using $\times 16$ objective were scanned under a Quantimet 720 (Imanco Ltd), which has 500 000 Pix cells between the guard frames, and it scans 10 times per second. The microscope was focused and the intensity between 0 and 63 (black to white) was selected for each field to show with maximum contrast the structures which the particular procedure is believed to stain. This adjustment of setting for each field - although time-consuming - corrected for differences of intensity of staining between different areas of a particular section, and also between the intensity and colour seen with different staining procedures. The number of Pix cells scanned and the number of elements counted appeared continuously at the edge of the screen. Each of these two numbers was read five times for a particular field and the mean values for each of the five readings were recorded. The areas stained were expressed as the number of Pix cells divided by 500 000 multiplied by 100; the mean and standard deviations were calculated for a particular staining procedure for all the sections of the particular area under study. Typical measurements are given in Appendix 3. The whole procedure was repeated using different slides but the same procedures, and no significant differences were found between the two sets of measurements. Therefore, the readings of the two sets were combined, and the values given represent eight slides, each counted in 20 different areas, for each of six different staining procedures, taken at random from over 400 slides originating from 24 different rat brains. In addition, slides of thoracic spinal cord, stained with haematoxylin and eosin only, were examined in a similar way.

The results are given in Tables 15 and 16. They show that

1. No single staining procedure, 'specific' or 'non-specific', shows up more than about 10% of the area of any section, or, put another way, most of the central nervous system is not stained by any of the procedures used.
2. If one adds up the areas stained by Marsland, Glees and Erikson (neural cells and processes); Mallory's phosphotungstic acid haematoxylin (astrocytes) and Weil and Davenport (oligodendrocytes and microglia), the total areas stained would only be between 14% and 20% of the total area of the tissue - that is, staining of all the elements together which are believed to be specific does not result in colouration of most of the tissue in the four different areas.
3. Even if one adds up the areas stained by the three 'specific' and the three 'non-specific' stains - whose overlap would much exaggerate the total areas stained - it still does not show up more than 33-41% of the total tissue.
4. Only 2.6-4.2% of the spinal cord was stained, which compares with means of 5.5-7.6% of the brain areas.

TABLE 15. Proportion, % ($m \pm s.d.$) of areas of coronal sections of rat brain (and spinal cord) stained by the procedures indicated ($n = 40$ each).
 The stained sections were scanned by a Quantimet particle counter, using a Leitz microscope, bright-field and a total magnification on the screen of $\times 450$. Please see also text (page 105) and legend to Appendix 3. Please note that the first three procedures are widely believed to be specific for the elements indicated in parentheses.

	Frontal	Parietal	Occipital	Medulla
Marsland, Glees and Eriksson (neural cells and processes)	2.7 ± 1.0	5.2 ± 2.0	7.7 ± 3.3	5.1 ± 1.9
Mallory's phosphotungstic acid haematoxylin (astrocytes)	6.2 ± 2.0	3.6 ± 1.5	8.1 ± 2.1	6.6 ± 2.1
Weil-Davenport (oligodendroglia and microglia)	5.7 ± 1.8	5.2 ± 2.0	4.7 ± 1.9	4.7 ± 1.9
Haematoxylin and eosin (all cells)	5.5 ± 2.0	7.6 ± 2.1	5.9 ± 1.9	6.9 ± 2.6
Patay's triple stain (nuclei)	7.3 ± 2.0	5.1 ± 2.1	5.7 ± 2.6	7.2 ± 2.1
Gallyas (nuclei and neuroglial processes)	6.6 ± 1.7	6.6 ± 2.2	9.0 ± 2.3	9.0 ± 4.6
(Haematoxylin and eosin - spinal cord = 3.4 ± 1.3)				

EXPERIMENTAL STUDIES

TABLE 16 Mean percentages of stained areas as a proportion of total areas of 20 fields, each of two different brains

	Frontal	Parietal	Occipital	Fourth ventricle
Marsland, Glees and Erikson (neural cells and neural processes)	2.7	5.2	7.7	5.1
Mallory's phosphotungstic acid haematoxylin (astrocytes)	6.2	3.6	8.1	6.6
Weil-Davenport (oligodendroglia and microglia)	5.7	5.2	4.7	4.7
'Total mean area' assuming specificity of each stain	14.6	14.0	20.5	16.4

Please see also legend to TABLE 15

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

5. We then examined over 100 sections of brain, and thoracic and lumbar spinal cord of human beings, cats, rats, rabbits and guinea pigs, stained by all the procedures indicated in Table 13, using bright-field illumination and a magnification of x400 (but not the Quantimet).

In every section one sees that only a small proportion of the tissue is stained by any procedure believed to be 'non-specific', or 'specific' for particular elements. Only a small proportion of areas, such as the medulla or the anterior horn, which one is conditioned to believe contain a high concentration of nuclei or neurons, take up a significant quantity of stain. One's interest in particular nuclei conditions one to the expectation that most of that area is stained, whereas, in fact, only a small proportion of the area has been coloured. This phenomenon is analogous to that called 'closure' by psychologists.

ELECTRON MICROSCOPICAL STUDIES

We next looked at electron micrographs. They also show only two kinds of cells, those with processes and the 'naked' nuclei (Figures 31-33). In those with processes the cell membrane can be detected all the way around a substantial volume of cytoplasm. On the other hand the 'naked' nuclei exhibit clear membranes round the nuclei themselves, but evident and striking absence of membranes round an area which would be considered as their cytoplasm. However, all the space between the 'naked' nuclei and the cells with processes is replete with mitochondria.

Thus in unstained sections, sections stained by a variety of histological procedures, and in electron micrographs, the agreement of the findings was excellent. We had cells with processes and 'naked' nuclei, in a ubiquitous volume containing mitochondria.

After considering, in detail, the gulf of uncertainty about the 'specificity' of appearances, staining and markers of different kinds of cells which have been summarized (pages 63-91, Tables 8-11 and Figures 34, 35 and 36), we then sought out our more trusty instrument, 'Occam's Razor'. We concluded that structurally we could classify all cells in the central nervous system into two types - those with processes, and naked nuclei. Those with processes are probably all neurons, and the naked nuclei are the mobile nuclei of the syncytium. These conclusions will be examined in more detail below (please see pages 119-123).

However, we then tested the question directly by looking again at our own 500 or so slides stained by the procedures in Table 13. We could classify all cells seen into one of the three appearances, viz. cells with processes, naked nuclei, and occasionally larger, rounder, paler-staining cell bodies (Figure 36), which we believe to be neurons which have not shrunk or been stained as much as others. In re-examining our histological, including serial, sections, we always bore in mind the many appearances which sections through a polymorphous object could give (Figure 21).

DORSAL ROOT GANGLIA (FIGURE 37)

These were teased out using mounted stainless steel needles, and their capsules torn open. Many oval cells with remarkably few processes came out (Figure 18). Staining with haematoxylin and eosin showed the same types of cells, presumably neurons

EXPERIMENTAL STUDIES




FIGURE 31 An electron micrograph of the parietal cortex of rat showing the relative sparsity of cells, surrounded by other material. Please compare with Figures 32 and 33. The bar is 5 μm




FIGURE 32 Electron micrograph of a neuron in the parietal cortex of rat. The nucleolonema can be seen in the nucleolus. The cell membrane can be detected sometimes as a clear line and sometimes by the change in texture of the cytoplasm on both sides of it. The bar is 5 μ m

EXPERIMENTAL STUDIES




FIGURE 33 Electron micrograph of a neuron and 'satellite' cell in the parietal cortex of rat. The bar is 2 μm




FIGURE 34 Human spinal cord showing ventral motoneurons stained with, (upper left) cresyl violet, (upper right) Palmgren's stain, (lower left) haematoxylin and eosin, and (lower right) phosphotungstic acid haematoxylin. The neurons show up with all these four procedures. The bar is 50 μm

EXPERIMENTAL STUDIES




FIGURE 35 Purkinje cells, of human cerebellum stained with (upper left) cresyl violet, (upper right) haematoxylin and eosin, (lower left) cat cerebellum with Palmgren's stain, (lower right) human cerebellum stained with phosphotungstic acid haematoxylin.
Please note the dendrites of the basket cells arborizing around the Purkinje cells with the Palmgren stain. These are the only dendrites in the mammalian nervous system which appear close enough to cell bodies to permit synapses to be present, but synaptic knobs or expansions cannot be seen even under high-power magnification with oil-immersion objectives. This figure should be compared with views of the cerebral cortex and retina (Figures 50-52), where there are no such intimate relationships. The bar is 50 μm




FIGURE 36 The frontal cortex of rat was stained with (upper left) haematoxylin and eosin, (upper right) phosphotungstic acid haematoxylin, (lower left) Marsland, Glees and Erikson, (lower right) Patay's stain. In all cases, only neurons N, naked nuclei nn and paler, more spherical cells P, could be seen. All cells in all histological sections can be classified as belonging to these three groups, or ependymal cells. The bar is 50 μ m

EXPERIMENTAL STUDIES

(Figure 37). There are a number of small circles between these neurons, which are so small that it is difficult to say whether one is looking at granules or transverse sections of fibres. The neurons stain more or less intensely, which may reflect inhomogeneity of the population or uneven uptake of stain, but there is no clear evidence from light microscopy of a second population, which could be considered as neuroglial cells.

WHITE MATTER

In the white matter we must distinguish between two kinds of fibres. Those in the cerebral commissures or in the cerebellum, for example, are very fine; they have diameters of less than 1 μm , so that their lumens cannot be seen by light microscopy (Figures 38 and 39). The axons of the fibre tracts in the white matter of the spinal cord have greater dimensions - 2·5 μm in stained sections (Figure 40) - much less than those of myelinated peripheral nerves (Figure 41). Nodes of Ranvier and Schmidt-Lantermann clefts are not found in white matter.

Many further studies of the histology of the white matter of the rat, rabbit and human cerebellum and spinal cord were carried out using the routine staining procedures indicated in the legends of Figures 38, 39 and 40. It was noted that there appeared to be very few fibres in the white matter.

AXONS

Eight rats and twelve rabbits were anaesthetized with ether, and their carotid arteries cut. The brachial, femoral and sciatic nerves were excised using fine iris scissors, and the nerves were teased with mounted needles and examined by phase-contrast microscopy. Other nerves were embedded in paraffin wax and transverse sections 6 μm cut and examined under phase-contrast microscopy.

The traditional view of the unfixed preparation was seen - an axon surrounded by a myelin sheath, which was interrupted by nodes of Ranvier. Both the axoplasm and the cytoplasm of the sheath appear clear and translucent (Figure 19). There was no evidence in the cytoplasm of the Schwann cell of any solid structure or lamellae visible within the resolution of the microscope. On transverse section the stained myelin sheaths appeared as uniformly black circles (Figure 41).




FIGURE 37 Transverse section of human dorsal root ganglion stained with haematoxylin and eosin. Note the large neurons, some of them staining much darker than others. Granules are seen between the neurons. The bar is 20 µm




FIGURE 38 Anterior white commissure of human thoracic spinal cord stained with Palmgren's stain, showing a commissure of fibres (left) and syncytial fibres (right). Please see also legend to Figure 39. The bar is 50 µm




FIGURE 39 White matter of the human cerebellum stained with Palmgren's stain. Please note how the fibres in the plane of the field in the lower half swing up towards a transverse orientation at the upper half of the field. The bar is 50 µm




FIGURE 40 Transverse section of white matter fibres of human thoracic spinal cord, stained with Weigert-Pal's stain. The bar is 20 µm

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM




FIGURE 41 Transverse section of part of the sciatic nerve of a rat, embedded in paraffin, not stained, but viewed by dark-field illumination. The bar is 20 μm . Please note the myelin sheaths, and compare with Figures 19 and 40

6

Critique of these Experiments

The demonstration by well-known neurological staining procedures of membranes around kidney and liver cells, but not between nuclei and cells with processes (which we believe to be neurons), could be explained by supposing that the membranes of neuroglia are not visible by light microscopy, but this view is not tenable, since the neuroglia were originally characterized by light microscopy. Alternatively, someone might like to assert that some procedure other than those listed in Table 13 might show up these membranes.

Another possibility is that all cells with processes, which we are now calling neurons, are not just neurons, but also include astrocytes, oligodendrocytes and possibly microglia - in accordance with the current view. We believe that we have brought sufficient evidence to undermine fundamentally the view that the neuroglial cell types can be differentiated with certainty by any - and therefore all - the criteria normally employed; that permits us to proffer the simplifying suggestion that all cells with processes are neurons. This is a working hypothesis, and its implications for neurobiology will be discussed later (pages 205-241).

What then of the large volume of biochemical findings quoted in Table 12 from the groups of Raff in Britain, Hertz in Canada, Eng and Varon in the USA, Hamberger in Sweden, Pevzner in the USSR, Schousboe in Denmark, and many others? The accumulation in one table of these careful experiments carried out in prestigious laboratories all over the world points quite clearly to a Tower of Babel populated by biochemists.

Nonetheless, despite all these reservations, we are bound to accept that significant differences in quantities of markers have been found. Do these mean that the cell types are different? Not necessarily. The cell types have been separated by different procedures involving varying degrees and techniques of homogenization, filtration, and culture in different media. It is therefore not at all surprising that biochemical analysis of the cultures after their hazardous and wayward voyages should find them to be biochemically different. Only if they had all been separated by the same procedures and then illustrated profound biochemical or immunological differences, could one suppose that they were different cell types.

The overall uncertainty about the identity of cells based on the lack of specificity of the markers does not mean, either, that there are no biochemical differences between

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

living neurons and neuroglia *in situ*, or that both the neurons and the neuroglia are homogeneous in their biochemical properties. We would like to stress the distinction between, on the one hand, a dissatisfaction about the adequacy of the evidence identifying the cell types currently characterized by the alleged markers, and on the other, the probability that the population of cells in the living central nervous system is chemically heterogeneous. We regard the latter possibility as being completely open.

We have shown directly that most of the same cells, such as ventral horn motor neurons and Purkinje cells of cerebellum, stain with most of the staining procedures. When we looked at the multiplicity of views about what each procedure does show up (Table 10), we were induced to go back to the originators of classical methods, and ask what they said about the specificity of their own methods. The answer is as follows: the authors either (1) used the generic term 'neural' tissue; (2) did not assert that their procedures were specific; or (3) did not mention control tests on cells other than those in which they were interested. The simple fact is that none of the original authors claimed that their staining procedures were specific. Later authors assumed them to be so.

We have asserted that no continuous membranes could be seen between whole neurons and naked nuclei, either by light or by electron microscopy. We have considered how membranes are detected (Appendix 4). It is widely held that membranes cannot be seen unless either they are normal to the section (Horne and Harris, 1981), or they are not normal to the section (Ham, 1974). The latter two explanations are incompatible. Elsewhere we have shown how the expected incidence of the membrane being seen clearly normal to the section can be calculated, and that the actual incidence with which it is seen thus is much higher (Hillman and Sartory, 1980, page 40). Yet these calculations become unnecessary when one sees that the nuclear, mitochondrial and cell membranes can be detected along most of their courses by electron microscopy; therefore any cell membrane between the neurons and naked nuclei should also be seen as frequently. Liver and kidney cells were surrounded by such membranes. A categorical statement that we have never been able to detect such a membrane in the central nervous system in stained preparations by bright-field illumination, in unstained preparations by phase contrast microscopy, or by electron microscopy, should be taken as an invitation to neurobiologists to produce any evidence to the contrary - if indeed, such evidence exists. In particular we would ask them to show us these membranes in any species or by any histological or electron microscopic technique other than those which we have used.

It might reasonably be asked why we used the neural stains, such as Weil and Davenport's, Mallory's phosphotungstic acid haematoxylin and Marsland, Glees and Erikson's, when we have shown them to be non-specific. When we first used them we shared the widespread belief that they were, indeed, specific for particular elements. We then tested them, and found them to stain the same elements - the cells with processes and the naked nuclei - and sometimes the cell processes. We then searched the literature, and found out what a diversity of opinion there was about which element, and how specifically, it was stained (Table 10). Bearing in mind that the non-specificity would result in the addition of the stained areas being an underestimate of the elements

CRITIQUE OF THESE EXPERIMENTS

present, the staining could still properly be used to examine the proportion of the area occupied by cellular elements. Light microscopic histology is still the only way of quantifying the cellular elements in a significant volume of tissue.

An important potential error in attempting to assert the relevance of these measurements to the tissue in the intact living animal could arise if the cells with processes and the naked nuclei were to shrink significantly differently than the syncytium surrounding them, while they were being prepared for histology. If the former elements shrank less than the syncytium, this would result in the measured proportion they occupied being exaggerated by quantimetry and if they shrank more, it would result in the proportion being understated. Measurements of shrinkage in cerebral slices - representing the whole tissue - and in isolated neurons, showed that the differences in shrinkage were not very large (Hillman and Deutsch, 1978, and sources quoted there).

En passant, one may wonder why many neural staining procedures seem to show up cell bodies, but not - according to the traditional view - most of the cell's processes. The general exception to this proposition is that of the silver salts. We do not believe this to be the case; that is, we do not believe that fixation, staining or dehydration can cause any elements which are present not to show up at all, even though, obviously, some procedures do increase the visibility or contrast of some elements more than others. For example, the failure to see tracts of white matter by electron microscopy is often explained as being due to poor fixation. Such an explanation - which is obviously quite feasible - should not be accepted unless it had been demonstrated; this could be done quite easily.

One may ask why the mitochondria do not appear in a normal haematoxylin and eosin preparation of brain examined under high-power magnification. The simplest explanation is that, since they are moving in a liquid cytoplasm in life (Hillman and Sartory, 1980, page 57), they diffuse out during the rehydration of the sections before staining. Certainly they are seen in abundance in unfixed pieces of central nervous tissue (Figure 42). Another possible explanation for the disappearance of some organelles or fibres is that the vast changes in water content, osmotic pressure, permeability, fragility, pH and pressure occurring during preparation, could cause them to lyse, and then become unrecognizable.

The possibility that in the pieces of brain we were seeing bacteria rather than mitochondria can be discounted, since the preparations were always examined in freshly made up solutions, and the cerebral tissue always came from animals which had been newly killed. There would not be time for the multiplication of bacteria in such an abundance as we have seen.

The neuroglial 'clumps' of Hyden or 'flakes' of Hamberger were defined by their proximity to neurons, and their identification as neuroglial cells depended upon the belief that the neuroglial material was composed of cells. Hertz (1965, 1966) originally used this approach, as did the whole school of Hyden (please see Hyden, 1967), including ourselves (Hillman, Hussain and Sartory, 1973; Hillman, Deutsch, Allen and Sartory, 1977). After the latter experiments, when we realized the ubiquity of the neuroglial material, we then realized that the terms neuroglial 'clumps' and 'flakes' both meant material adjacent to the neurons; that is, all the neural substance other than the neurons and nuclei. We therefore, decided to examine this in a more general

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

way, and take out small pieces of brain and spinal cord at random, to find out if the neuroglial clumps and flakes were, indeed, typical central nervous tissue. This turned out to be the case (page 94).

The detection of large numbers of mitochondria between the neurons and the naked nuclei, as well as within the neurons themselves, indicated that the tissues were intracellular. We detected the mitochondria because we examined the tissue both unstained by phase-contrast microscopy and after staining with Janus green (Hillman and Deutsch, 1978b). Of course, mitochondria are located intracellularly.




FIGURE 42 Mitochondria seen in (left) neuroglial clump, (right) neuron below with neuroglial clump above by phase contrast microscopy; nn is neuron, ng is neuroglia, m is mitochondrion. The bar is 10 μm

CRITIQUE OF THESE EXPERIMENTS

The hypothesis that the third kind of cells - the swollen relatively unstained cells - were neurons which had not shrunk due to dehydration, or had not been as intensively stained as other neurons, fits in with the frequent observation that in an identified population there is a wide variation in the degree of staining of individual cells. This is seen, for example, in the ventral horn cells in the spinal cord and in the Purkinje cells in the cerebellum (Figures 34 and 35).

The wide variation of the appearance of sections of an egg would suggest that sections of neurons could also show a rich range of appearances. A clear space around stained neurons has been recognized as an artifact for many years, and its incidence is widely believed to be diminished by perfusion fixation for light and electron microscopy. The effects of shear during section are not so widely recognized, but they can be appreciated rapidly by comparing the rather elegant appearance of the unfixed unsectioned neuron (Figures 16, upper, and 18) with the more distorted histological appearances of most neurons stained but not sectioned (Figures 16, lower, and 44). I have also noticed it in some unpublished observations. In the early 1970s I attempted to cut sections for electron microscopy of single neurons embedded in epoxy resins. The shrunken torn fragment could hardly be recognized as originating from the whole neuron.

The problems of detecting the presence of, and measuring the thickness of, cell membranes by light and electron microscopy are discussed in Appendix 4.

The fact that different staining systems produce apparently different appearances of cells, even of the same cells, is not at all surprising. One is using an array of fixatives, embedding media, stains, dehydrating agents, differentiating agents, clearing agents and mounting media. These vary in chemistry - aldehydes, heavy metal salts, organic solvents, oils, waxes and epoxy resins. The tissues are subjected to these reagents each with the following conditions varied: thickness of section, length of exposure to the reagent, concentration of the reagent, temperature at which it is applied, sharpness of knife, water solubility of reagent, stability of reagent, subjective assessment and skill of the histologist, the kind of microscopy used and the skill of the microscopist. The likelihood that variation in any of these parameters will affect the appearance and therefore the identification of the cells or their organelles is strongly implied by the necessity to reproduce the conditions of a particular staining procedure prescribed by its author in order to achieve the same appearances. Therefore, the reproducibility of the appearance of a particular structure using a very precisely defined series of conditions cannot be regarded as evidence either that, firstly, that is the original shape of the structure, or secondly, that different shapes resulting from different staining procedures arise from different structures (see also Appendix 5).

If one conceives that each cell in a population may stain and shrink to a different extent during the same histological procedures, one could suppose that a population would show a range of 'plumpness' to 'spideriness', and slight to intense staining. This could depend upon the degree of stress of the animal during killing, the metabolic state of the cells in the tissue at the time of fixation, the depth within the tissue specimen, the size of the specimen, the duration of application of each of the reagents, the distance between the microtome and the cell body in the section, and the proximity

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

of the cell to the surface to which the reagents are added or to the microscope slide. All these variables could also affect the staining of whole cells and each of their individual organelles.




FIGURE 43 Rabbit neurons isolated by hand dissection, stained with, (upper left) Marsland, Glees and Erikson, (upper right) Gallyas's stain, (lower left) Mallory's phosphotungstic acid haematoxylin, (lower right) Weil and Davenport. The bar is 20 μm

In view of the protean appearance of sections of cells, it might be appropriate to ask the following question. How can one be sure that the appearance of 'naked' nuclei in sections is not also due to neurons - that is, that there are not two populations of cells, but only one? Is it not possible that the circles are transverse sections of neurons near the bases of processes. I believe that this is unlikely, because the naked nuclei stain the same colours as the nuclei in the neurons in the same sections by virtually all staining procedures (Figures 25, 26 and 27). Also, one would expect to see an occasional thorn sticking out of the circular structure, if it was indeed a section near the origin of a process of a neuron. I do not know of a report of any such appearance having been seen on the nuclei of classical neuroglial cells.

The naked nuclei seem to be significantly different from the nuclei of the neurons, in that the former do not appear ever to have nucleoli.

When the sections were stained and scanned with the Quantimet, 40 areas were examined in each particular coronal section. This examination, therefore, was a measure of the areas in the whole section and tissue, and not only in the cerebral cortex, or the cranial or spinal cord nuclei. While our specific studies were done on

CRITIQUE OF THESE EXPERIMENTS

rats and rabbits, we have also examined tissues from the other species mentioned, and stained by the other procedures listed (Table 13). We have never seen any stain which colours equally most of the area or volume of any nervous tissue. We would request any neurohistologist who has found any procedure which is claimed to be specific for any cell type in any species to scan it with a particle counter, or to send it to us to examine ourselves. Naturally, we would be particularly interested to see histological sections or light micrographs of any tissue in which most of the tissue has been stained. Nevertheless, we have shown that the usual assumption that the 'specific' neurohistological stains would together colour the totality of the tissue, and that, therefore, the neurons and neuroglial cells together make up most of the nervous system, is simply not tenable, at least for the rat and rabbit central nervous system.

7

The Fine Structure of the Nervous System

THE NEURON

The structure of the generalized cell has previously been reviewed (Hillman and Sartory, 1980a), and I would refer the reader to this publication for detailed evidence that the following structures could not exist in living cells: Nissl body, endoplasmic reticulum, ribosomes, lysosomes, Golgi apparatuses, cristae of mitochondria, nuclear pores, neurofibrils, neurofilaments, neurotubules, microtrabeculae, intermediate filaments and actomyosin filaments (other than in muscle) (Table 17). The evidence brought there may be summarized briefly. Living cells show intracellular movements which are visible under relatively low-power light microscopy ($\times 200-1000$) (Figure 44). Cytoplasmic movements are incompatible with the existence of a fine cytoskeleton, which requires 10-100 times greater magnification to see, and which is supposed to permeate the cytoplasm and to be attached to the cell membrane and to the nuclear membrane. Intracellular viscosity has been measured in a large number of living cells and has generally been lower than that of glycerol. Observations on many living cells indicate that the cytoplasm is a liquid. In addition to these biological grounds for doubting the existence of many of these apparent structures, most of them appear in the plane of the picture far too frequently than simple geometry would dictate; they are two-dimensional and could not exist in three dimensions. No three-dimensional transparent scale models in which intracellular movement might appear possible have ever been made.

On geometrical grounds it has also been shown that the trilaminar appearance of the cell membrane, the nuclear membrane, the mitochondrial membrane, and the neuromuscular junction, is an artifact, although the membranes themselves are not. The synapses will be dealt with below (please see pages 135-156).

In response to our assertion that the 'cytoskeleton' is an artifact of preparation for electron microscopy, it has been alleged that it can be seen by light microscopy. The filaments are fibrils and are believed to have diameters between 10 nm and 15 nm (Lazarides, 1980; Shelanski, Liem, Letterier and Keith, 1981). Until recently it has generally been accepted that the maximum resolution of the light microscope using a monochromatic source under optimal conditions is 200-250 nm. Therefore one could only see structures 10-15 nm in diameter by light microscopy, if they occurred in bundles whose total diameter would be sufficient to be resolved. However, when one looks at

THE FINE STRUCTURE OF THE NERVOUS SYSTEM

the elements of the cytoskeleton by electron microscopy, they do not appear to be associated in close bundles (Table 18). It is obviously inconceivable that a structure which appears by light microscopy to have a minimum diameter of 200 nm would shrink to only 10 nm, bearing in mind that the preparations are observed after dehydration both in light and electron microscopy.

TABLE 17 Artifacts in nervous tissues and suggestions about their most likely origin during preparation

Artifactual structures	How they were caused
Trilaminar <u>appearance</u> of membranes	Groove caused by evaporation and explosion of tissue under thin metal deposit (please see Figure 45)
Golgi apparatus	Precipitate of cytoplasm with heavy metal
Endoplasmic reticulum	
Lysosomes	
Cytoskeleton	
neurofibrils	
neurofilaments	
actin filaments	
neurotubules	
microtrabeculae	
Myelin lamellae	Dehydration and precipitation of cytoplasm with heavy metal
Anatomical synapses (light microscopy)	Dehydration and precipitation of contents of myelin sheaths
(electron microscopy)	
Structures in synaptic cleft	Metal deposits on regions of cell membranes
Exo- and endocytotic vesicles	Dehydration and precipitation
Nuclear pores	Granules adjacent to membranes around which metal has deposited

Recently, Allen and his colleagues have introduced a video-enhanced contrast system, with which they claim to see by light microscopy structures as small as 5 nm in diameter moving along nerve axons (Allen, Travis, Allen and Ilmaz, 1981; Allen, Metuzals, Tasaki, Brady and Gilbert, 1982; Brady, Lasek and Allen, 1982; Allen and Allen, 1982).

If one cuts sections of any fibrous tissue at random, the majority of fibres are very unlikely to be found well orientated to the microtome either longitudinally or transversely. Why are the fibres, filaments, fibrils and trabeculae - which are believed to be orientated randomly in tissues other than muscle - so shy of exhibiting their oblique profiles? One would not say that they are never seen obliquely, but they do appear in nerve cells far more frequently in longitudinal than in transverse section, and they appear obliquely hardly at all. Moreover, attempts to adduce biochemical or




FIGURE 44a Intracellular movements in a rabbit Deiters' neuron. Changes in the appearance of the nucleolus can be seen clearly. The neuron was isolated by the technique of Hyden (1959) and placed in the '199' medium of Morgan, Morton and Parker (1950) in a parallel-walled chamber (Sartory, Fasham and Hillman, 1971). It was examined under phase-contrast microscopy, a, 28 hours after the animal was killed, b, 10 minutes later, c, 20 minutes later. The bar is 5 μm




FIGURE 44b




FIGURE 44c

THE FINE STRUCTURE OF THE NERVOUS SYSTEM

TABLE 18 Publications of mammalian structures showing intracellular trabeculae, filaments and tubules in the nervous system.

Structures	(m)	Animal/tissue	Authors
Neurofilaments		Mammalian cerebral cortex	Rhodin(1963)
Neurofilaments (9-10)		Rat optic nerve	Peters and Vaughn(1967)
Microtubules (23-26)			
Gliofibrils		Rabbit neocortex	King(1968)
Microtubules			
Neurofilaments		Rat cerebral cortex	Peters, Proskauer and Kaiserman-Abramof(1968)
Microtubules (22)			
Neurofilaments (10)		Rat anterior horn cell	Wuerker and Palay(1969)
Microtubules (27)			
Neurofilaments		Cat somatic sensory cortex	Jones and Powell(1970)
Microtubules			
Filaments (9)		Rabbit spinal cord	Raine and Wisniewski(1970)
Microtubules (25)			
Neurofilaments		Rat brain, parietal lobe	Tani and Ametani(1970)
Microtubules (22-27)			
Microfilaments (4-6)		Rat dorsal root ganglion	Yamada, Spooner and Wessels(1971)
Neurofilaments (9-10)			
Microtubules (24-28)			
Neurotubules (24-28)			
Neurofilaments (6-10)		Rabbit nodose ganglion	Johnston and Roots(1972)
Neurofilaments		Rat cerebellar cortex	Palay and Chan-Palay(1972)
Microtubules			
Neurofilaments (10)		Rat cerebellar cortex	Wuerker and Kirkpatrick(1972)
Microtubules (25)			
Neurofilaments		Monkey visual cortex	Palay, Billings-Gagliardi and Chan-Palay(1974)
Microtubules			
Microtrabeculae (3-6)		Human fibroblast	Wolosewick and Porter(1979)
Microfilaments (6-10)			
Microtubules			
Neurofilaments (7)		Human unmyelinated nerves	Han and Holmstedt(1981)
Microtubules (24)			
Neurotubules (24)			
Microtrabeculae (3-12)		Rat cerebral cortex	Cohen, Wolosewick, Becker and Pappas(1984)

In all of these publications the structures in the illustrations shown appear single on the electron micrographs and not in bundles. Their diameters, as noted by the authors, are also indicated. It should be noted that as single strands their diameters are at least one order smaller than could be seen by the light microscope. These publications were not selected because the structures were not in bundles, and I have not been able to find any electron micrographs in the literature on mammals showing strands sufficiently close together that a bundle of them would be seen with the resolution of the light microscope

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

biophysical evidence to explain the geometrical paradox cannot be successful, since they are epistemologically of quite different dimensions.

If it is accepted that the 'trilaminar' appearance of membranes is artifactual - and I stress that the detailed evidence has been published elsewhere (Hillman and Sartory, 1980) - one is edged along to the conclusion that the cell membrane, the nuclear membrane and the mitochondrial membranes are all of single thickness. The difficulties both of detection and of measurement of membrane thickness are discussed in Appendices 3 and 4.

The 'trilaminar' appearance could be due to the two layers of metal deposited on either side of the water and lipid extracted membranes. Alternatively, it could be the groove resulting from the explosion of what was left of the cell membrane through the layer of heavy metal salt above it. This would occur when the highly energetic electron beam, passing through the atmosphere at a very low pressure in the electron microscope, hits the membrane beneath the heavy metal salt and vaporizes it. The resultant explosion would blow out a groove (Figure 45). When the thickness of the trilaminar membrane is measured in a dehydrated specimen, it must be underestimated.

The well-known Davson-Danielli hypothesis about the structure of the membrane dominated the textbooks between the 1930s and the early 1970s. It proposed that the membrane consisted of a lipid molecule sandwiched by two protein molecules (Figure 12). A critical study of the evidence for the hypothesis would be a valuable addition to the literature, but it is outside the main theme of this monograph. However, there are a number of anomalies about belief in this particular model, which will be dealt with here. The Robertson 'unit' membrane is considered by some to be the structural expression of the Davson-Danielli model, and by others to be independent here in terms of the former idea.

A 'microsomal' fraction believed to contain the cell membrane and the endoplasmic reticulum, has been separated by subcellular fractionation, and the analysis of its composition is generally believed to yield data about all the 'membranes' of the 'cytoskeleton' or 'endoskeleton'. Thus, the belief has gradually gained credence that all the 'membranes' are chemically similar, and - as a corollary - that unless a structure has been shown to have the particular characteristic chemical constitution, it is probably not a membrane.

Both the Davson and Danielli membrane and the 'unit' membrane are agreed to be 6-12 nm thick throughout the literature. With transmission electron microscopy the membrane appears quite smooth, although the use of freezing techniques frequently shows it to have little knobs on it. Such knobs are believed to be proteins, enzymes or receptors of a large and increasing variety (please see Table 29).

The fluid mosaic hypothesis of membrane structure (Singer and Nicolson, 1972) is now widely accepted, as if it were a finding (Semenza and Carafoli, 1977; Repke, 1980; Kates and Kuksis, 1980). It originated from some thermodynamic considerations and studies on the distribution of antigens applied to red cell membranes and detected by freeze etching. 'A prediction of the fluid mosaic model is that the two-dimensional long-range distribution of any integral protein in the plane of the membrane is essentially random' (Figure 13) (Singer and Nicolson, 1972). The following questions immediately arise: What does 'model' mean in this context? Is this what they believe the




FIGURE 45 Suggestion about the mechanism by which a membrane appears as two lines by electron microscopy. Upper row side views, lower row plan views, of a membrane (coarse mottling), embedded in a medium (fine mottling). Left shows an embedded membrane; middle shows the deposit of heavy metal (black) on the surface of the membrane. The electron bombardment, heat and high vacuum causes the tissue to vaporize and explode through the metal deposit, leaving the two metal tracks at the edge of the previous location of the membrane, right

structure to be? What does a 'two-dimensional orientated viscous fluid' mean? Does it mean a thin layer? How can a fluid be two-dimensional? Do their experiments tell us any more than that the cell membrane contains water, and there is water on both sides of it? If one took a coverslip with water on it and placed a drop of protein on one corner, would it not diffuse across its surface? What would that indicate about the properties of the glass? If one took two adherent wet pieces of paper and injected antibody between them, would that not also spread 'two-dimensionally' as well as through the paper? How can one study a 'fluid mosaic' using freeze etching which freezes the tissue to below its eutectic point, until it becomes solid?

I think one should state unequivocally that the evidence upon which the theory is based can be accounted for by stating that living membranes contain water, and that diffusion occurs in aqueous media.

The idea that cell membranes might be fluid or liquid, implies the so-far unanswered question, 'How do the cells have sufficient tensile strength to contain the cytoplasm?' This is especially pertinent for myelinated nerves, muscle cells and epidermal cells, which may be exposed to considerable pressures. How could one pick out a neuron by hand dissection? How could Cummins and Hyden (1962) cut open neurons and pick up their membranes?

The real difficulty about both the Davson and Danielli and the Singer-Nicholson hypotheses is that neither of them can be demonstrated clearly to be true or untrue. Once again, we have evidence of relatively simple hypotheses, in proof of which cornucopias of complex experiments have been concocted, but which are not amenable to direct proof or falsification. The facility with which monolayers of lipids can be made does not and cannot prove that this is what happens in cell membranes, although many membranologists seem to believe it does.

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

The presence of water in a solid membrane should be no more surprising than water in an orange skin, which does not mean that the skin is fluid.

It would seem that much of the difficulty about any discussion about the solidity or fluidity of the cell membranes arises from two different uses of the word fluid. For example, glass has the mechanical properties of a solid, but physicochemically it is a fluid. If Singer or Nicholson were to draw a plan of a house, they would no doubt draw the windows as solid structures, even although they certainly know that glass is a fluid molecularly. The positions should be clarified. We believe that mechanically all the real cell membranes are solid structures, probably containing free water. Our view is mainly based on the ability with which whole neurons can be separated, and the possibility of cutting them open (please see page 231). We have no accurate knowledge of the precise chemistry of the membranes (Hillman, 1972). We are not at all clear whether Singer and Nicholson make the distinction between the mechanical and physicochemical meanings of the word 'fluid', and, if so, whether they are claiming that the cell membranes are fluid or liquid both mechanically and physicochemically. However, our main contention must be stated categorically, all the cell membranes in life are mechanically solid. Definitions of fluid, liquid or solid can be found in standard textbooks of mechanics or physical chemistry.

In relation to the membranes of the 'cytoskeleton' it is pertinent to ask why, in electron micrographs, the endoplasmic reticulum appears in sections as parallel lines, but their supposed subcellular offspring appear as rings. The electron microscopists and the biochemists both vehemently maintain that the two structures are identical. They assert that the reticulum 'rounds off' during separation. Yet it is simpler to suggest that the parallel lines and the circles represent different structures. In this sceptical age it is extraordinary that the identity of two such different shapes should be accepted as representing the same object on the basis of a verbal explanation, in the absence of demonstration of how the one shape could change into the other. If it did, it would imply the following train of events: pieces of two parallel lines representing a net or two layers of flattened vesicles would join together at two ends of a length of reticulum simultaneously; such two parallel strands or plates would expand to form a complete circle but would not burst; the ends beyond the join would disappear; the rings representing the thin sections would orientate themselves almost completely in the plane of the electron micrograph.

It should be noted that: (a) the microsomes are not seen joined at both ends on a sort of knot; (b) no intermediate shapes between the parallel lines and the almost perfect circles are seen; (c) segments like an arc of a circle, or a piece of orange skin, would be expected if the fraction contained pieces of reticulum not in the plane of the section; (d) unstuck pieces are not seen equatorially on microsomal circles. These vagaries of geometry are so striking that they shake one's faith in the identity of the endoplasmic reticulum and the microsomes, an identity lacking evidence.

Synapses and synaptosomal fractions are beset by similar structural difficulties. The electron microscopist's synapse is supposed to be a knob connected by a dendrite. When the knobs are separated in the sub-cellular fractions of nervous tissue (Pappas and Purpura, 1972; Cottrell and Usherwood, 1977, many papers) it is generally believed

THE FINE STRUCTURE OF THE NERVOUS SYSTEM

that the resultant hole in the synapse closes up, so that the synaptosome looks like a perfect circle.

The evidence adduced here that the Davson-Danielli, the Robertson, and the Singer-Nicholson hypotheses, have serious and unacceptable inconsistencies (pages 251-254) should not be taken as denying that cells, nuclei or mitochondria have membranes. If we must be rigorous in our thinking, we can only conclude the following points about the cell membranes:

1. The latter membranes exist in life, although other 'cyto-membranes' do not (Table 17), (please see also Appendix 4 on detection of membranes).
2. The membranes are insoluble, and probably contain substantial amounts of proteins and lipids. The measurements made on subcellular fractions must underestimate the contents in the cells of all soluble compounds including proteins and fatty acids. These must be lost differentially from different fractions, since all procedures involve the additions of a large volume of water and other chemically active reagents. The precise quantity of materials lost from each location in the cell depends on: the chemistry of all the reagents; the time for which they are applied; the physical procedures used; the rate and total quantity of energy dissipated during the procedures; the affinity of each natural component of the tissue for each other and for all the reagents used.
3. Structurally the membranes are each one layer thick. At a molecular level they may be chemically heterogeneous, but it is likely that their dimensions dictate some sort of orientation to them. Semi-permeability, active transport and excitability are all phenomena which can be more easily visualised if the molecules in the membranes are orientated, but one must still be cautious about interpreting the orientation of molecules from studies of dehydrated tissues.
4. The thicknesses of the membranes in life are unknown, but are certainly more than the 6-8 nm seen in dehydrated specimens by the electron microscopist. Since they appear to be of uniform thickness at high magnification in electron micrographs, it is likely that this reflects a uniformity of thickness in the original living material upon which the heavy metal was deposited.
5. The nuclear membranes are not punctured by pores (for evidence, please see Hillman and Sartory, 1980b, pages 67-75).
6. In addition to shrinkage of the width of the membranes resulting from dehydration, it must also shrink in the longer axis. Therefore, in life the membranes around the cell, the nuclei and the mitochondria must contain greater volumes than are seen by histology or electron microscopy. They must be more convex in life than in histological preparations.

THE SYNAPSE

The emergence of the synapse and the neuromuscular junction first as physiological, and then as anatomical, entities has been discussed briefly (pages 15-17). The time has now come to examine the current evidence for their appearances. There are several references in the literature to the suggestion that synapses are difficult to find in the brain (Glees, 1944; Wyckoff and Young, 1954, 1956; Boycott, Gray and Guillery, 1960).

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

When one looks at the mammalian literature, most of the synapses seen by light microscopy are found in the spinal cord or in the cerebellum, or are descriptions of neuromuscular junctions, or are diagrams. We show here a fine drawing from Held (1897) (Figure 9) who described 'end-fusse', and a more recent light micrography of a silver preparation from Wyckoff and Young (1956) as Figure 46. Small silver granules can be seen on the surface of the neuron soma. Careful examination of the latter figure reveals two important features which are characteristic of all light micrographs: firstly, that granules are seen both on the somas, and also in the background between the somas; secondly, relatively few of the granules have pre-terminal fibrils which can be seen to be connected to other somas. In Table 19 we list 18 publications showing synapses in the illustrations of which granules of approximately the same sizes appear clearly outside the neuron somas. Few of the authors comment on this, although occasionally it has been explained by the suggestion that those particular granules are on the surfaces of somas which are out of focus. The short connections of the synaptic knobs are recognized in the diagram which De Robertis (1959) has put in a review of the subject, and has kindly permitted us to reproduce as Figure 47.

In respect of the granules or knobs being located away from the somas or on somas which are out of focus, one has, of course, no way - other than their appearances - of identifying these knobs as synapses in these particular preparations. Perhaps, in the future, an author who claims that the granules or synapses are on surfaces of somas out of focus will add an adjacent photomicrograph showing the surface of the somas in focus, with the granules neatly situated on them.

As regards the second feature - the shortness of the connections to these knobs - one has a further difficulty. The fact that one can see fibrils arising from the knobs in some instances (Figures 46-48) means that their diameters are wide enough to be within the resolution of the microscope used. As one can see the dendrites arising from the cell bodies on the same illustrations, one must ask why one cannot see the fibrils from the alleged synapses wandering for a good distance across the microscopic field. When one examines cerebral commissures or cerebellar white matter, stained, for example, with Marsland, Glees and Erikson's or Palmgren's stain (Figures 38 and 39), one sees many dendrites, and fine fibres all of which are sufficiently thick to be seen within the depth of 4-10 μm sections; this is evidence that one has sufficient magnification to see fibres connected to the synaptic knobs, if they were present.

The expected appearances of these connections on plan view and on side view have been indicated on a diagram (Figure 49). The latter are simply not seen. With the exception of the fibres from the basket cells surrounding the cerebellar Purkinje cells (Figure 35), I have been unable to find anywhere in the mammalian literature any light micrographs showing even fine fibres apparently investing somas; however, even on Purkinje cell bodies, knobs are not generally seen, and the encirclement of fibres does not necessarily mean that these fibres contact granules on the soma surface.




FIGURE 46 'End-feet' (synapses) on the neurons of the ventral horn of the thoracolumbar region of the cat. The 'end-feet' were stained with silver stains. Please note the following features: the diameters of the end-feet; the rarity, except on a few endfeet in 11, of preterminal fibres attached; the frequent presence of silver-stained granules and fibres, in regions not obviously on the neuron surface. This figure is reproduced from Wyckoff, R.W. and Young, J.Z. (1956) The motoneurone surface. Proc. Roy. Soc. B. Lond., 144, 440-450, by kind permission of Professor Young and the Royal Society

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

TABLE 19 Illustrations by light microscopy showing some knobs of similar size to synapses outside the cell body

Tissue	Author
Mammalian spinal cord	Phalen and Davenport (1937)
Cat ventral ganglion of acoustic nerve	Lorente de No (1938)
Cat ventral horn cell	Barr (1939)
Cat cervical sympathetic ganglion	Gibson (1940)
Cat spinal cord	Barnard (1940)
Human cerebral cortex	Meyer and Meyer (1945)
Mammalian nervous system	Weber (1955)
Cat visual cortex	Sholl (1956)
Cat spinal cord	Wyckoff and Young (1956)
Rat superior cervical ganglion	Couteaux (1958)
Kitten spinal cord	Madge and Scheibel (1958)
Dog spinal cord	Gelfan (1964)
Cat spinal cord	Sprague and Hongchien (1964)
Kitten spinal cord	Szentagothai (1964)
Cat reticular formation	Brodal (1969)
Cat medulla oblongata	Windle (1969)
Rat reticular formation	Curtis, Jacobson and Marcus (1972)
Cat ventral horn cells }	
Rat anterior horn cell	Leeson and Leeson (1981)

The terms used are those of the authors

THE FINE STRUCTURE OF THE NERVOUS SYSTEM




FIGURE 47 A diagram of the current view of the attachment of synapses to the neuron soma, from de Robertis (1959). Please note that the axon and dendrites are shown travelling a long way away from the soma in A, but neither in A nor B do the synaptic connections reach beyond a small distance from the soma. This diagram is reproduced by kind permission of the author and publishers from 8, 65 (Academic Press). This diagram should be compared with Figures 49 and 53, showing the expected view of synapses




FIGURE 48 Synapses on neurons of A, rat reticular formation, B, ventral horn of cat, stained with silver stain. In the reticular section many stained granules are not obviously associated with neurons. In the ventral horn neuron many fibres can be seen, but they are rarely attached to the synapses. These illustrations are reproduced from Curtis, B.A., Jacobson, S. and Marcus, E.M. (1972) Introduction to Neurosciences, Philadelphia; Saunders, page 54, by kind permission of Dr Jacobson and the W.B. Saunders Company. The bar is 10 μm




FIGURE 49 Expected views of synapses seen by light microscopy on the soma of a neuron. Please note the synaptic connections ('telodendroglia') spreading to the edge of the field. This should be compared with the previous Figures 46-48

In a Golgi preparation of brain, 150 µm thick, one can focus on the dendritic tree at various planes in the tissue; the dendrites form an aesthetic arborization (Figure 50). As one follows any particular dendrite under the microscope, especially using a x90 or x100 oil-immersion objective, one can see that there are no connections between the dendrites and axons from a particular soma to any other soma, just as the branches of trees in a closely planted grove do not make connections with each other. However, they appear to do so on a normal two-dimensional photograph (Figure 51).

In cats, Wassle (1982) demonstrated the dendritic trees of ganglion cells in whole mounts of retina after retrograde labelling with horseradish peroxidase, and has kindly provided us with Figure 52. Although occasional dendrites from one soma overlap one another, and fluorescent granules are seen on many dendrites, it is difficult to suggest that there are any connections between the cells or their processes.

Boycott, Gray and Guillory (1960) explained the failure to see synapses frequently in the neo-cortex by their not being stained (see also Gray, 1966).

En passant, it should be noted in the illustration from Wyckoff and Young (1956) (Figure 46) that the synaptic knobs appear empty on light micrographs, while the electron microscopists fill them with a treasury of synaptic vesicles.

The histologists searched for the synapse, while the physiologists continued to describe the properties of reflexes, of excitation and inhibition, in the belief that they were all located in the anatomical structures which were still rather elusive. They were both ready to accept that the location by the electron microscopists of the synapses not only between neurons, somas and fibrils, but also between virtually every part of the neuron including the axon and many neuroglia (please see page 17). They did not appreciate the serious three-dimensional difficulties in interpreting the micrographs,




FIGURE 50 The original legend reads 'A section showing a number of the smaller pyramidal neurons in the outer part of the visual cortex of a cat stained by the Golgi-Cox method. Technique: mercuric chloride-dichromate. For details see Sholl (1953). Section 150 μm thick'. Apparent contacts between the processes can be seen to be extremely rare even in this photograph. This figure is reproduced from Sholl, D.A. (1956) *The Organisation of the Cerebral Cortex*. London: Methuen, page 14, by kind permission of the publishers




FIGURE 51 Dendritic arborizations in the human cerebral cortex: left, neonatal, right, after 24 months. These illustrations are reproduced from Conel, J.L. (1939, 1959) *The Postnatal Development of the Human Cerebral Cortex*, Vols. I, VI. Cambridge, Mass: Harvard University Press, by kind permission of the publishers and the President and Fellows of Harvard College. It is difficult to imagine that the bodies of neuroglial cells could fit in between these dense arborizations.




FIGURE 52 Micrograph from a whole mount of the cat retina after retrograde labelling of the ganglion cells with HRP injected into the ipsilateral lateral geniculate nucleus (scale 100 µm, eccentricity 7 mm, temporal retina). This figure is from Wassle, H. (1982) Morphological types and central projections of ganglion cells in the cat retina. In Progress in Retinal Research, 1, eds Osborne, N.N. and Chader, G.J. Oxford: Pergamon, page 143; it is reproduced by kind permission of the author and publishers. Please note that there is little evidence of contacts between dendrites

nor the probably irreconcilable discrepancies between the views of the light and electron microscopists of what are believed to be the same structures.

The geometrical problems are summarized in Figure 53; both kinds of microscopists have different problems. By light microscopy the vast predominance of views could be represented by a section through an egg lying on its side with its long axis parallel to the surface upon which it was lying. One does not see sections in all the other orientations in light micrographs. Obviously, if the synaptic knobs are distributed all




FIGURE 53 Drawings of possible appearances of synapses by light microscopy (top line, a,b,c,d) and electron microscopy (lower lines, A,B,C,D,E); the same letters should refer to the expected views of the same shaped synapses at different magnifications and orientations. The black lines and shading indicate pre- and post-synaptic thickenings; the grey shading indicates the cytoplasm of the synapses and the neurons. These diagrams are based on the assumption that the synapses are randomly distributed over the surface of the neurons, as shown, for example in the diagram of Barr, 1939. The appearances of the synapses by light microscopy should be completely compatible with their appearances by electron microscopy. Geometry dictates unequivocally that all the appearances of the electron microscopic images indicated by a particular capital letter should be seen, and that they should occur in approximately equal frequencies. Examination of the considerable literature on the structure of synapses fails to show a significant incidence of any of the images of the third line down (with oval pre- or post-synaptic membranes), nor of the arcuate sections of the synapses as seen at the bottom extreme left and extreme right. Light microscopic images corresponding to the 'double' synapse seen on electron microscopy (E) do not seem to have been reported either (please see references in Tables 21-26). It should also be noted that the common views by light microscopy a,b,c, do not correspond to the common view by electron microscopy, C, in the right-hand column

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

over the neuron surface, any section should go randomly through single synapses in a complete range of orientations and sizes. For example, the pear shape should not appear more frequently than the circular shape, which would be at right angles to the pear shape. We have tilted the model and drawn the resultant shapes which would be expected (Figure 53). In these diagrams the top line represents the views seen by light microscopy, with the exception of the c in the middle of the diagram.

One would expect to see the same structures on electron micrographs at higher magnifications than they are seen in light micrographs. These are represented by capital letters on Figure 53. Simple geometry again tells us that for an object which is randomly orientated it should be seen equally frequently in any two planes at right angles to each other. When one takes the top line of expected electron microscopic images A, B, C and D, one can then tilt them at 45° away from or towards the plane of the picture, and look at the resultant expected images depicted in the next line down using the same capital letters. One then obtains a series of images in our diagrams, which are simply not seen on electron micrographs. Put another way, one rarely if ever sees a synapse whose thickening obliterates the synaptic cleft and which bridges as an oval shape the pre- and post-synaptic region. If one then tilts the synapses a further 45° to examine their expected appearances at right angles to their long axes, one would expect to see the appearance of a black circle due to the synaptic thickenings not apparently connected to the membranes of either of the cells. Again, this appearance should occur as often as the right-hand top electron micrographic diagram C.

A further generalization may be made. One hardly ever sees the preterminal fibre of the synapse on electron microscopy. The top right-hand image C appears to be the commonest picture. Why is the connection of the synapse to the parent soma seen so rarely on electron microscopy, when - although too short for distant connections on light microscopy - it is at least seen sometimes (Figure 46). Fortunately, the neuromuscular junction represents a control tissue, in which the preterminal fibre is always seen by electron and by light microscopy. As far as the classification of synapses into Types I and II (Gray, 1959), represented in Figure 53 by E, it should be pointed out that that appearance does not seem to have any counterpart in light microscopy; perhaps the magnification is not high enough to show it. However, neither Gray himself nor those who accept his classification appear to have realized that a section through a Type II synapse E at right angles to, and in the long axis of, the one shown would give the typical appearance of a type I synapse; thus, there is no necessity to postulate the two types.

Hyden and his school (Hyden, 1961, 1967a) have published a large number of photographs of isolated unfixed neurons, similar to our micrographs (Figures 16 and 44). At the magnifications used one should be able to see synaptic knobs, if they exist, but they are difficult to recognize. This could be because (1) they have all been pulled off during the isolation by hand dissection; (2) the granular appearance of the neuron surface under phase-contrast microscopy is due to many small synapses; (3) the synapses in unfixed tissue have the same refractive index as the media in which the neurons are prepared (usually 0.25 mol/l sucrose); (4) synaptic knobs do not occur on neurons. Roots and Johnston (1965) found rather few synapses on isolated

THE FINE STRUCTURE OF THE NERVOUS SYSTEM

neurons using electron microscopy and they also believed that the membranes of the neurons are largely stripped off during preparation. The latter explanation seems unlikely to us, since isolated neurons after isolation possess resting potentials normally believed to be associated with a membrane (Hillman and Hyden, 1965); these potential differences could be changed reversibly by altering the oxygen, potassium ions or sodium ions in the incubating media. Many of those who isolated unfixed neurons believe that the granular appearance is due to synapses.

We believe that the granular appearance is due to intracellular structures for the following reason. One can draw diagrams of the appearance of granules on the surface of the soma (that is, synapses), and compare the appearances with those of granules in the cytoplasm (that is, mitochondria) (Figure 54). When one examines cells by low-power light microscopy with a considerable depth of focus, the number of granules seen over the nuclei would be more if the granules were synapses than if they were mitochondria. Neurons which have synaptic physiology in tissue culture have been illustrated by several authors (Table 20). Invariably fewer granules appear over the nuclei than over the cytoplasm at these relatively low magnifications (Figure 55). (Of course, when one focuses on the nuclei with high-power light microscopy when the depth of focus is only 1-2 μm , one cannot distinguish whether granules are in the cytoplasm or on the surface of the cell.) The fewer granules over the nucleus than over the cytoplasm under low-power microscopy must be interpreted as meaning that the granules are intracellular - that is, mitochondria and not synapses.

One must readily admit the difficulty of divining the total specifications prescribed by the original designer of anatomical structures, but it is interesting to speculate on the reason for which the precise area through which the transmitters are believed to diffuse and attach - the pre- and post-synaptic regions - should have been so armorially thickened. It is strange that these should be chosen to be sites of the entry and exit of large insoluble molecules or granules - processes known as endocytosis and exocytosis.




FIGURE 54 If one sees granules in a cell under the light microscope, one can determine if they are on the membrane of the cell (left-hand figures) or in the cytoplasm (right-hand figures); that is, whether they are synapses or mitochondria. The cell is viewed in transverse section (upper diagrams) and then from above (lower diagrams). If the granules are on the surface they will appear in only slightly less density over the nucleus than over the cytoplasm (lower left). If they are in the cytoplasm but not in the nucleus (upper right) they will appear from above in much lower density over the nucleus than over the cytoplasm (lower right). In cells examined under low-power magnification when the depth of focus is several micrometres, one can see, nevertheless, that fewer granules are visible over the nucleus than over the rest of the cytoplasm (Table 20), suggesting that the granules are intracytoplasmic, that is, they are mitochondria.

TABLE 20 Neurons in tissue culture in which the nucleus appears clearer than the cytoplasm, which indicates that the granular appearance is more likely to be due to mitochondria than to synapses

Human cerebral and cerebellar cortex	Costero and Pomerat (1951)
Human cortex	Geiger (1957)
Rat and cat cerebellum	Hild and Tasaki (1962) *
Rabbit cerebral hemispheres and cerebellum	Varon, Raiborn, Seto and Pomerat (1963)
Chick and rat dorsal ganglion	Pomerat, Hendleman, Raiborn and Massey (1967)
Mouse spinal cord and dorsal root ganglion	Nelson (1973) *
Mouse dorsal root ganglion	Shahar, Grunfeld, Spiegelstein and Monzain (1975)
Mouse spinal cord	Fischbach and Nelson (1977) *

* In these publications, neurons have been shown to fire. Please see Figure 54.




FIGURE 55 Purkinje cells of cat cerebellum in (left) a 25-day-old culture; (right) a 42-day-old culture. The cell on the left is being penetrated by an intracellular electrode, which records an action potential. This photograph is reproduced from Hild, W. and Tasaki, I. (1962) Morphological and physiological properties of neurons and glial properties of neurons and glial cells in tissue culture. *J. Neurophys.*, 25, 277-304, by kind permission of the authors, the editors of the *Journal of Neurophysiology*, and the American Physiological Society. The cell on the right is reproduced from Pomerat, C.M. and Costero, I. (1956) Tissue culture of cat cerebellum. *Am. J. Anat.*, 99, 211-247, by kind permission of the editors of the *American Journal of Anatomy*, and Alan Liss, Inc. Please note that in both the nucleoplasm appears less granular than does the cytoplasm

Practically every part of the neuron has been claimed to be seen on electron microscopy to synapse with other neurons, and, occasionally with neuroglial cells (please see references on pages 15-17). This raises a number of questions. Why do these synapses not appear as knobs on light microscopy? How widespread is their distribution? Does an electron microscopist examining the mammalian central nervous system see thickenings on the surface of cells which he had previously thought of as neuroglia, then proceed to reclassify the cells as neurons? Have systematic attempts been made to search for them and examine their incidence widely in relation to mammalian neuroglia? Are there any other certain and unequivocal criteria which a microscopist can use to identify a synapse?

It is generally held that the careful histological studies of Ramon y Cajal - summarized in 1954, please see page 13 in which he pointed out that neurons did not have contact with each other, support the existence of synaptic clefts. The apparent absence of continuity of neurons has been used as one element of evidence that synaptic transmission is chemical. What does one mean by 'contact'? In an electric plug,

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

a wire is screwed tightly on to a metal contact. Most authors see the synaptic cleft 20-30 nm wide (Table 21), which is probably much narrower than the gap in parts of a fully functional electrical contact. In a synapse, the pre- and post-synaptic sides would both be in a tissue fluid which is highly conductive.

Gray (1959) classified cortical synapses into two types, based on the width of the synaptic clefts, the depth of the post-synaptic thickenings, and whether the synapses appeared to be axodendritic or axosomatic. Subsequently, he found that the same classification could be extended to cerebellar synapses (Gray, 1961). Van der Loos (1965) doubted the distinctness of the two types, and Karlsson (1966) did not accept that the classification was applicable to other parts of the central nervous system.

TABLE 21 The width of the synaptic cleft in mammalian central nervous system measured by electron microscopy

Tissue	Width (nm)	Author
Rat abducens nucleus	20	Palay (1958b)
Rat abducens nucleus	20	Hyden (1961)
Rat dorsal cochlear nucleus	20	Eccles (1964, page 17)
Monkey lateral geniculate nucleus	20	Gray and Guillery (1966)
Cat subfornical organ	20	Akert, Pfenninger, Sandri and Moor (1972)
Rat cerebral cortex	30-40	Curtis, Jacobson and Marcus (1972, page 52)
Rat prepyriform cortex	20-30	Johnston and Roots (1972, page 35)
Rat cerebral cortex	20	Jones and Brearley (1972a)
Cat oculomotor nucleus	20	Pappas and Waxman (1972)
Rat ventral horn of spinal cord	20	Bloom and Fawcett (1975, page 37)
Neurons from various species	20-30	Peters, Palay and Webster (1976, page 135)
Neurons from various species	15-25	Brodal (1981, page 11)

The identification and therefore confirmation by electron microscopy of the existence of synapses, which had been previously seen and described by light microscopy, was generally regarded as having settled the problems of the elusiveness and shapes of the synapses. There are, however, further disquieting discrepancies. Comparison of the numbers of synapses per soma by light microscopy (Table 22) with the numbers by electron microscopy (Table 23), shows that - with a few exceptions - the former technique reveals incidences in the 10s and 100s, while the latter technique reveals 1000s to 10 000s. One may assume that when light and electron microscopists measure synaptic diameters, they publish values of average diameters, or their photographs show 'typical' synapses. One has to be a bit careful in comparing values for light micrographs, which are normally given by the authors (Table 24) with measurements from electron micrographs (Table 25). The electron microscopists have generally avoided giving values for the diameters of their synapses, so we have had to measure them off their figures. The precision of values given by authors of light micrographs is low because the synapses are too small to measure them accurately,

THE FINE STRUCTURE OF THE NERVOUS SYSTEM

so they are usually estimates; in electron micrographs the precision is diminished by the accuracy with which measurements can be made from published micrographs. With these reservations firmly in mind one can compare the diameters of synapses measured by light microscopy (Table 24) with those measured by electron microscopy (Table 25). Although the range of the former is between 500 and 15 000 nm in diameter, and the latter is from 250 to 2500 nm, in general the overlap between the two sets of dimensions is so poor as to cast doubt as to whether the two sets of measurements were both of the same objects. The differences could not be accounted for by supposing that the shrinkage during silver staining for light microscopy would be less than that during osmic acid staining for electron microscopy, bearing in mind that shrinkage resulting from dehydration with ethanol is common to both techniques; also a linear dimension which is double another encloses a volume eight times that of the other.

We may summarize the conclusions about the synapses. Synapses seen by electron microscopy do not respect Euclidean geometry. They are seen in larger numbers, but in smaller dimensions, by electron than by light microscopy. Although some authors have shown the appearance by light microscopy of some areas where dendrites from different cells appear to overlap (Van der Loos, 1960, 1974; Van der Loos and Glaser, 1972). I have been unable to find in the literature micrographs showing a parent soma connected to a dendrite whose preterminal fibre is attached to a synapse. At a full meeting of the Physiological Society at University College, London, on March 28th, 1985, I invited colleagues to send me micrographs or references to literature showing this.

The synaptic thickenings of the electron microscopists may be due to the heavy metal stains attaching themselves to apparent rings believed to be parts of the membrane for which they have a greater affinity than other parts, or they may be due to random deposits of the heavy metal on such rings. There is no way of distinguishing between these two possibilities. Furthermore, even if the heavy metal does have a greater affinity for 'specific' sites on the cell membrane, that does not mean that these represent the sites of the physiological activities. Even if the measurements of the dimensions of the synapses were accurate to within one order of what they appear to be, the widths of the intracellular electrodes with the finest tips would be far too large to be of use in locating the site of a single synapse. Since one cannot specify the sites of physiological events of transmission, it follows that findings about physiological events do not necessarily support or deny any hypotheses about those events. (Please see also Hillman, 1985.)

The discrepancies between the light and microscopic views of synapses and their apparent absence in some sites does not necessarily mean that they do not exist, but they certainly raise some difficult questions. However, our discussion of the lack of cogent evidence for long connections between the synaptic endings and other dendrites presents a much more serious problem, especially for the physiologists. This is dealt with on page 197.

I would like to stress that I am not denying the existence of neuromuscular junctions or pre-terminal fibres to them. However, evidence from them may not be relevant to synapses, without demonstration of identity of the particular property in both tissues.

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

TABLE 22 Number of synapses per neuron soma reported by the authors or counted from their publications, as detected by light microscopy

Neuron	Synapses per cell	Stated (s) or counted (c)	Authors
Guinea-pig trapezoid nucleus	54 48 46 36 26	c) c) c) c) c)	Barker (1899)
Mammalian spinal cord	0-10	s	Phalen and Davenport (1937)
Cat motor neurons	14 38 38 22 30	c) c) c) c) c)	Lorente de Nò (1938)
Cat ventral horn cell	1250	s	Barr (1939)
Cat anterior horn cell	192 388 156 130 416 68 106 148 114 234 150 70	c) c)	Barnard (1940)
Cat cervical sympathetic ganglion	Up to 13	s	Gibson (1940)
Monkey lateral geniculate body	1	s	Glees and Le Gros Clark (1941)
	1	c	
Human dorsal sensory and internuncial	97	s))	Minckler (1942)
Human intermediate sensory and internuncial	152	s))	
Human ventral sensory and internuncial	116	s))	
Human nucleus dorsalis - thalamus	348	s))	
Human intermediolateral - thalamus	122	s))	
Human posteromedian - thalamus	362	s))	
Human anteromedian - thalamus	675	s))	
Human anterolateral - thalamus	1624	s))	
Human posterolateral - thalamus	3990	s))	
Human cerebral cortex	16 24	c) c)	Meyer and Meyer (1945)
Cat ventral horn of spinal cord	2000	s	Wyckoff and Young (1956)
Rat superior cervical ganglion	12 14	c) c)	Couteaux (1958)
Dog lumbosacral cord neuron	800	s	Gelfan (1964)
Dog motor neuron	6000-10 000	s	
Cat spinal cord	84	c	Sprague and Hongchien (1964)
Rat cerebral cortex	10 000	s	Johnston and Roots (1972)

The terms used in Tables 22-25 are those used by the Authors

THE FINE STRUCTURE OF THE NERVOUS SYSTEM

TABLE 23 Number of synapses per neuron soma reported by the authors as detected by electron microscopy

Neuron	Synapses	Authors
Rat spinal cord	2000	De Robertis (1959)
Mammalian	Many hundreds	Eccles (1964)
Rabbit Deiters cell	10 000	Hyden (1964)
Rat lateral geniculate nucleus	133 83	Karlsson (1966)
Guinea pig cerebral cortex	800	Blackstad (1967)
Mouse visual cortex	7000) Cragg (1967)
Mouse motor cortex	13 000)
Monkey visual cortex	5600)
Monkey motor cortex	60 000)
Vertebrates	10 000	Johnston and Roots (1972)
Cat cerebral cortex	870	Kaisman-Abramof and Peters (1972)
Cat spinal cord	Several thousands	Conradi (1972)
Mammalian spinal motor neuron	500) Bullock, Orkand and Grinnell
Mammalian cortical pyramidal cell	30 000	(1977)
Mammalian cerebellar Purkinje cell	Up to 80 000)

TABLE 24 Diameters of synapses seen by light microscopy in the literature. The values were stated by the authors

Tissue	Width (nm)	Authors
Mammalian nervous system	500-7000	Ramon y Cajal (1909)
Monkey spinal cord	950 x 800-3330 x 2600	Phalen and Davenport (1937)
Cat spinal cord	500 x 100-4000 x 5000 (mean 1500 x 2000)	Barr (1939)
Cat spinal cord	7000 (mean 2000-3000)	Gibson (1940)
Cat ventral horn motor neuron	1000-5000	Bodian (1952)
Cat cerebral cortex	2500	Entin (1954)
Cat spinal cord	15000 x 8000	Szentagothai-Schimert and Albert (1955)
Rat spinal cord	1000-5000 (mean 2000)	Wyckoff and Young (1956)
Human cerebrum	800-2500	Smythies, Gibson and Purkis (1957)
Rat visual cortex	2000-4000	Boycott, Gray and Guillery (1960)
Cat spinal cord	1000-4000 few 15000	Gray and Guillery (1966)
Cat reticular formation	500-3000	Brodal (1969)
Rat cerebellum	1500-6000	Valdivia (1971)
Cat motor neurons	1000-3000) Curtis, Jacobson and Marcus (1972)
Cat granule and stellate cells in cerebellum	500)

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

TABLE 25 Diameters of synapses as measured from illustrations of electron micrographs in the literature. The accuracy of measurements is estimated to be about 50 nm from the printed micrographs

Tissue	Width (nm)	Authors
Rabbit ventral horn cell	1200	Wyckoff and Young (1956)
Cat upper lumbar spinal cord	600-1000	Gray (1959, 1961)
Rat hippocampus	500	Blackstad (1967)
Ox brain	500	Roots and Johnston (1964)
Rat prepyriform nucleus	800	Gray and Guillory (1966)
Rat medial mammillary nucleus	800	Heimer and Ekholm (1967)
Rat cerebellum	1500	Lumsden (1968)
Cat lateral vestibular nucleus	1300	Brodal (1981, page 10)
Cat subfornical organ	500-800)	Akert, Pfenniger, Sandri,
Monkey spinal cord	500)	and Moor (1972)
Rat cerebellar cortex	500-2000	Bloom (1972)
Rat prepyriform cortex	1300	Johnston and Roots (1972, page 38)
Rat cerebral cortex	1200)	
Cat oculomotor nucleus	1000-1500)	Pappas and Waxman (1972)
Cat sensorimotor cortex	650-1000)	
Cat hippocampus	1200)	
Rat median septal nucleus	500-2500	Raisman and Matthews (1972)
Rat olfactory glomerulus	600-1000	Reese and Shepherd (1972)
Monkey lateral geniculate body	1500	Glees and Hasan (1973)
Rat ventral cochlear nucleus	250-600)	
Rat cerebral cortex	250)	
Rat spinal cord	250-1000)	Peters, Palay and Webster (1976, pages 118-180)
Rat auditory cortex	500-750)	
Rat visual cortex	400-900)	
Monkey visual cortex	600	Somogyi and Cowey (1981)

The synapses are believed to contain 'vesicles' (Gray and Whittaker, 1962), which are 10-70 nm in diameter (Table 26). It is very likely that the vesicles are artifacts for the following reasons. Inspection of the references in Tables 23, 25 and 26 shows that in any individual preparation vesicles appear far too frequently of uniform size. If they were uniform in diameter in the living tissue, they should be seen in sections in a normal distribution of diameters. Such studies which have been done show certain diameters to predominate; very few small contours of vesicles are seen (please examine illustrations in Birks, Huxley and Katz, 1960; Pappas and Waxman, 1972; Gray, Gordon-Weeks and Burgoyne, 1982). Secondly, the number of vesicles seen depends upon the fixative used to stain them, and the concentration of magnesium ions in the preparation medium (please see page 229). Thirdly, much of the evidence which is believed to be compatible with the transmitter hypothesis, but in our opinion, does not necessarily bear upon it, has been reviewed in this monograph (pages 226-230), and has been shown to contain a number of important but unlikely assumptions. Fourthly, there are believed to be about 50 neurotransmitters (Table 27). One must ask if each

THE FINE STRUCTURE OF THE NERVOUS SYSTEM

TABLE 26 Diameters of synaptic vesicles seen by electron microscopy in the literature.
The values were stated by the authors

Tissue	Width (nm)	Authors
Guinea-pig forebrain	50	Gray and Whittaker (1960)
Rat abducens nucleus	20-65	Hyden (1961)
Rat cerebral cortex	20-70 (mean 40)	De Robertis (1963)
Mammalian central nervous system	30-60	Eccles (1964)
Rat cerebellar cortex	15-60	Palay (1964)
Rat prepyriform cortex	30-60	Gray and Guillory (1966)
Cat dorsal nucleus of lateral geniculate body	30-50	Peters (1968)
Rat ventral horn of spinal cord	20-65	Bloom and Fawcett (1969)
Cat reticular formation	30-60	Brodal (1969)
Rat cerebellum	40 50-60 x 25	Valdivia (1971)
Kitten subfornical organ	50	Akert, Pfenniger, Sandri and Moor (1972)
Rat cerebral cortex	10-20	Johnston and Roots (1972)
Cat oculomotor nucleus	40-60	Pappas and Waxman (1972)
Various mammalian neurons	40-50	Jones (1975)

TABLE 27 Substances which have been proposed to be involved in the communication between neurones (partly from Barchas, Akil, Elliott, Holman and Watson (1978) and partly from Osborne (1981)) Communication between neurons. Neurochem International, 3, 3-16.

Adrenalin	Oestrogens
Dopamine	Testosterone
Noradrenalin	Thyroid hormone
Tyramine	Bombesin
Octopamine	Cholecystokinin (CCK)
Phenylethylamine	β -Endorphin
Phenylethanolamine	Gastrin
Dimethoxyphenylethylamine(DMPEA)	Neurotensin
Tetrahydroisoquinolines	Proctolin
Serotonin (5-hydroxytryptamine)	Prolactin
Melatonin	Oxytocin
Tryptamine	Substance P
Dimethyltryptamine (DMT)	Somatostatin
5-Methoxytryptamine	Angiotensin
5-Methoxydimethyltryptamine	Luteinizing hormone releasing hormone (LHRH)
5-Hydroxydimethyltryptamine (bufotenin)	Vasopressin
Tryptolines	Vasoactive intestinal polypeptide (VIP)
ATP	Adrenocorticotrophic hormone (ACTH)
Acetylcholine	Thyrotropic releasing hormone (TRH)
Carnosine	Sleep factor delta
Histamine	
Gamma - aminobutyric acid (GABA)	
Gamma - hydroxybutyric acid (GHB)	
Glycine	
Taurine	
Purine	
Aspartate	
Glutamate	
Prostaglandins	
Corticosteroids	

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

of them has its own vesicles. Presumably, the general view is that those neurons excited or inhibited by a particular putative neurotransmitter have in their synapses vesicles containing 'packets' of their own transmitter. The transmitter hypothesis has only been worked out in detail for acetylcholine and some catecholamines, (see pages 226-230).

DENDRITES

Each neuron has up to about 20 primary dendrites and their number increases with old age (Golgi, 1879; Conel, 1939-1963; Sholl, 1956; Nauta and Ebbeson, 1970). It should be noted that there are relatively few primary dendrites and that they divide into many secondary ones (Figures 50 and 51). The dendritic tree appears like a sapling in winter; that is to say, each neuron has probably one main axon and three to six larger primary dendrites, each giving out few branches.

In our view the dendrites radiate out into the syncytium and make no contact whatsoever with other dendrites, synapses or cell somas. This observation was first made by Ramon y Cajal (page 12). The morphology of the dendrites is probably similar to that of the free nerve endings found in the cornea, skin, blood vessels and viscera (Boeke, 1932; Kuntz and Hamilton, 1938; Tower, 1940; Woppard, Weddell and Harpman, 1940).

The ideas that the axons can be distinguished from the dendrites by any criteria other than their size, or that each neuron has only one or two axons, are both assumptions.

The surfaces of the dendrites in tissue culture appear much smoother than they do in histological sections, probably because they shrink and their surfaces become granular during preparation; heavy metal stains and intracellular components precipitate during dehydration.

If, as is proposed here (page 92), the cells generally described as astrocytes and oligodendrocytes are neurons, it follows that all neuroglial processes and fibres are dendrites.

NEUROMUSCULAR JUNCTIONS

In mammals neuromuscular junctions can be seen unfixed, or stained by light microscopy, and by electron microscopy (for reviews, see Cole, 1955, 1957; Couteaux, 1960; Zacks, 1964; Coers, 1967; Cold Spring Harbor Symposium, 1975; Gauthier, 1976); they are sometimes visualized by the presence of cholinesterases (Koelle and Friend-enwald, 1949; Karnovsky and Roots, 1964). By light microscopy they appear as tubular strands ending in muscle (Figure 56), or they may sport knobs at the ends of them (see, for example, McMahan, Spitzer and Pepeu, 1972). Sometimes there appears to be one junction for a group of muscle fibres (Figure 56), sometimes one for each muscle fibre, and sometimes several for each muscle fibre (Vrbova, Gordon and Jones, 1978); even in the latter case, a single neuromuscular junction services many sarcomeres. This means that excitation spreads across the sarcomeres, a long way from the end-plate, visible by light or electron microscopy.

By electron microscopy one usually sees between the axon and the muscle invaginations

THE FINE STRUCTURE OF THE NERVOUS SYSTEM

or junctional folds; these may branch and their appearances are highly variable (*op. cit.*). The axonal end contains a large number of vesicles, which should be considered in the same light as synaptic vesicles, and are probably artifacts (page 154). The junctional folds themselves have a dangerous proclivity to lie nearly always in the plane of the electron micrographs. Their dimensions are such that they or the space believed to be occupied by them should be visible by light microscopy. Perhaps micrographs showing these have not yet reached the literature, but I would be grateful if anyone would send me a light micrograph showing them, or draw my attention to a publication in which they feature.

CYTOPLASM OF NEURONS

In life, the cytoplasm is a liquid of low viscosity permitting diffusion, Brownian movement and the movement of mitochondria; pinocytosis may also occur. The mitochondria are filamentous structures (Figures 24 and 42) with single-layered membranes and they contain a liquid mitochondrioplasm. They are the only identifiable structures in the cytoplasm, although other granules - which are difficult to classify, name or analyse chemically - are sometimes also present there (Figure 44, pages 128-130). In tissue cultures, the cytoplasm is of low refractive index and is translucent. In living cells there are no Golgi apparatuses, lysosomes, fibrils, filaments, trabeculae, reticula or ribosomes (Table 17).

The intracellular movements would be accelerated by heat released as a consequence of metabolism, by pressure changes in the tissue due to changes in posture and muscle action, and by changes in the chemistry of the extracellular fluid, cytoplasm, nucleoplasm, mitochondrioplasm or pars amorpha; the chemical constitution of fluid compartments would change continuously as a consequence of transport, secretion and changes in permeability of the membranes between the relevant compartments.




FIGURE 56 Neuromuscular junction of human being stained with Schofield's modification of Bielschowsky's silver stain. The bar is 100 µm. This figure is reproduced from Ralis, H.M., Beasley, R.A. and Ralis, Z.A. (1973) Techniques in Neurohistology (London: Butterworth), page 141, by kind permission of the authors and publishers. Please note the asymmetry of the entry of the nerve fibres and the rather few neuromuscular junctions

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

The questions have been raised as to what is the site of protein synthesis, if the ribosomes do not exist? Where do acid hydrolases occur, if not in the lysosomes? The answers to these two questions are the same. These activities occur in life in solution or in suspension in the cytoplasm. Where do the enzymes for oxidative phosphorylation sit, if not in lines along the cristae? The answer is in the mitochondrioplasm. Moreover, posing these raises several other pertinent questions. What is the site of protein synthesis, when ribosomes cannot be seen? Are 'free ribosomes' the name given to any granules, which must be presumed to be the sites of protein synthesis whenever ribosomes cannot be seen lining the endoplasmic reticulum? Similarly, where do acid hydrolases occur in those cells such as neurons which do not seem to contain lysosomes? Is there any evidence that the enzymes of oxidative phosphorylation do line the cristae, other than that they provide greater surface area (greater than there would be without them)? Does not the concept of 'pools' imply the existence of biochemical compartments which are not physical compartments? Why is the intracellular viscosity so low if the cytoplasm is full of 'endoskeleton' and their dependent granules? Finally, if prokaryotes can pack all their metabolism and genes in one compartment, why cannot eukaryotes do likewise?

It is universally agreed among all biologists that living cells contain hundreds to thousands of enzymes or enzyme activities. The subcellular biochemists believe in the existence of the following 13 compartments or phases: extracellular space; intramembrane space; cytoplasm; cisternae; lysosomes; mitochondrial intramembrane space; intramitochondrial space; intramembrane nuclear space; Golgi body space; ribosomes; nucleoplasm; nucleolonema; pars amorpha, of which we accept the existence of the six compartments or particles underlined. It is clear that our housing problem for the thousands of enzymes is serious, but the same problem is shared to a lesser extent by those who would multiply the number of compartments by 13/6, i.e. 2.2. Neither of us has anything like the problem of the prokaryotes, who have to pack all their biochemistry into one compartment. The biochemists cannot solve this problem because so far they have concentrated on finding out where the enzymes end up in the subcellular fractions *in vitro*, and not where they are located undisturbed *in vivo* (Hillman, 1972).

THE NUCLEAR MEMBRANE OF THE NEURONS

The nucleus is spherical or oval. It is surrounded by an imperforate membrane which is one layer thick. The chemical composition of the nuclear membrane is unknown, since it cannot be isolated and identified unequivocally in a fraction which has been demonstrated to contain no significant concentration of other cellular elements, or to have lost major components during preparation. Nevertheless, it is widely believed to have the same chemical composition as that proposed for the cell membrane by Davson and Danielli or Singer and Nicholson.

NUCLEAR PORES

Nuclear pores can be seen by electron microscopy (see, for example, Peters, Palay and Webster, 1976, page 55). We have concluded that they are artifacts and suggested

THE FINE STRUCTURE OF THE NERVOUS SYSTEM

a number of ways in which they may arise (Hillman and Sartory, 1980, pages 68-75). At present, the best explanation for the appearance on electron micrographs of granules passing across the pores is as follows. Any particle which happens to be present on the nuclear membrane, either during the deposit of the heavy metal stain, or after microtomy, or at the time when the electrons bombard the membrane in the electron microscope, will either prevent the heavy metal stain being deposited on the nuclear membrane, or prevent access of the electrons to that part of the membrane. This will create the appearance of a pore in the membrane, especially if the particle explodes. The same explanation seems likely for apparent endocytosis and exocytosis at synapses.

It has also been pointed out elsewhere that the connections of the nuclear membrane to the endoplasmic reticulum are seen frequently in diagrams (Hillman and Sartory, 1980, pages 102-103), but rarely in electron micrographs.

NUCLEOPLASM

In tissue culture the nucleoplasm appears clear and translucent (Table 20) and contains the nucleolus. In stained tissues it appears as a granular uniform precipitate.

NUCLEOLI OF NEURONS

The nucleoli of unfixed neurons of rat, rabbit and guinea pig, can be seen by phase-contrast microscopy to be surrounded by a relatively thick membrane (Figures 3, 18 and 44) (Hussain, Hillman and Sartory, 1974). If cells are stained, the membrane can no longer be seen, because the contents of the nucleoli precipitate (Hillman, Hussain and Sartory, 1976). Nothing is known about the chemistry of the membrane, except that it can only be seen in unfixed cells if the neurons are isolated in saline, since sucrose also precipitates the contents of the nucleoli (*op. cit.*).

By light microscopy one can also see the nucleolonema, which is in a state of movement in unfixed cells (Figure 44), but stops after the addition of phenol (Sartory, Fasham and Hillman, 1971). Some of the shapes of the nucleolonema present in the nucleoli resemble rosettes, stars or umbrellas, as they move, change, and go in and out of focus continuously (Figure 57). Very little attention has been given in the literature to the nucleolonema, its movement and its structure, probably because neurons, even in cultures, are usually examined in fixed tissues, in which this movement does not occur. Furthermore, by electron microscopy, the nucleolus looks like a granule with little structure (Figures 32 and 33).

ASTROCYTES, OLIGODENDROCYTES AND MICROGLIA

A few classical views of these kinds of cells are shown (Figures 58-62), and the difficulties of the classification have been illustrated in Tables 2-12 and Appendices 1 and 2.

All light and electron microscopists have accepted the view of Golgi (1885) that astrocytes are characterized by their attachment to blood vessels. Del Rio Hortega (1928) was also of the view that they were all attached to blood vessels or pia mater. However, Penfield (1932, page 424) says that this is 'a sweeping statement to which we cannot at present subscribe'. Nevertheless, the view is widely held at present,




FIGURE 57 The fine structure of the nucleoli of rabbit neurons, from the Deiters' nucleus, as seen by phase-contrast microscopy. Various shapes could be detected in the nucleolonema which was changing continuously. The neurons were isolated in 0.25 mol/l sucrose, which does not permit one to see the nucleolar membrane clearly. The movements stopped on the addition of phenol. The same structures were also seen in neuronal nucleoli from rat, guinea pig and frog. The bar is 1 μm long. This illustration is from Sartory, P., Fasham, J. and Hillman, H. (1971) Microscopical observations of the nucleoli in unfixed rabbit Deiters' neurons. *Microscopy*, 32, 92-100. These drawings are reproduced by kind permission of the Quekett Microscopical Club




FIGURE 58 Different cells in the cat cortex stained by Rio del Hortega's modification of the Golgi method, x160: 1 neuron; 2 fibrous astrocyte; 3 protoplasmic astrocyte; 4 above, protoplasmic astrocyte, below, oligodendrocyte; 5 oligodendrocyte; 6 transitional oligodendrocyte; 7,8,9, microglial cells. These identifications are due to the author. This illustration is from Ramon-Moliner, E. (1968) The morphology of dendrites. In The Structure and Function of Nervous Tissue, ed. Bourne, C.H., Vol. 1, (New York: Academic Press), pages 205-267, and is reproduced by kind permission of the author and Academic Press. We regard 1-6 as being neurons (page 189)




FIGURE 59 A human astrocyte from the cerebral cortex stained with Cajal's gold sublimate. The slide was kindly supplied by Dr C. Treip of Cambridge University. The large astrocyte is probably of the reactive type. The bar is 50 μm




FIGURE 60 Cells which are identified by arrowheads as oligodendroglia in white matter. The original legend reads: 'They are not always readily distinguishable from astrocytes'. This section is stained with haematoxylin and eosin, and the bar is 10 μm long. From Lewis, A.J. (1976) Mechanisms of Neurological Disease (Boston: Little, Brown and Co.), page 44, by kind permission of the author and publishers




FIGURE 61 Electron micrograph of a cell from rat lateral geniculate body identified by Professor Paul Glees, who kindly provided it; the cell is believed to be an oligodendrocyte because of its elongate nucleus, dark cytoplasm and clear mitochondria. Please note that the cell membrane is visible all round the cytoplasm. The bar is 10 μm . We would classify this as a neuron, as it has a discernible membrane




FIGURE 62 A microglial cell stained with Del Rio Hortega's stain. At this scale the diameter of the circle would be approximately 170 µm. Please note that the cell in the centre which the authors call a microglial cell has processes, and there are other spherical bodies, probably neuroglial nuclei without such processes, for example, between 1 o'clock and 2 o'clock. From Bouman, L. and Bok, S.T. (1932) Histopathology of The Central Nervous System (Utrecht: Oosthoek's Publishing Comp.), plate 2. We would classify this as a neuron.

both by light and by electron microscopists.

Golgi staining procedures require thick sections, sometimes as much as 150 µm (Sholl, 1953), so that one should be able to see a great deal more of the three-dimensional structure of cells than is possible with the usual thinner sections. The earlier electron microscopists cut 'thick' sections of 0.5-1 µm, and modern practice is to cut them 10-50 nm thick. Even the use of the thickest of these sections creates difficulties in identifying astrocytes by their 'feet'.

A micrograph is taken of a fibre ending on a capillary and its cell of origin is then dubbed an astrocyte. If such a foot is not seen, but it is believed that that particular cell is an astrocyte, other morphological criteria are used to identify it, or it is supposed that the particular foot would be seen in another serial section of the same cell.

As far as I am aware, no-one examining sections, say 100-150 µm thick by a Golgi-Cox procedure, has shown that processes of the cells which they identify as astrocytes are attached to cerebral capillaries and blood vessels at a greater incidence than are processes originating from neurons or oligodendrocytes. The alleged high incidence of relationship between astrocytic 'feet' and capillaries should not be accepted without comprehensive calculations being made, and several positive and statistically significant studies being carried out at several different neurobiology research centres. One would also have to show that capillaries are associated with astrocytic feet with a significantly greater incidence than they are associated with oligodendrocytes, microglia and neurons.

The other difficulty is that both the neurons and neuroglia with their processes - as well as the capillaries - are widely distributed in three dimensions in the central nervous system, so that it is very likely that many of them would appear to be in contact at low magnification, when virtually the whole thickness of the section is in focus.

THE FINE STRUCTURE OF THE NERVOUS SYSTEM

The astrocytic 'feet' appear to electron microscopists as spaces between cells (Figure 63). The space stains lightly and contains relatively few particles. Close examination reveals that this 'foot' could not be an extension of a cell. It simply has no membrane around it. If one looks at this region on an electron micrograph at relatively low power electron microscopy, one sees the membrane of the cells adjacent to the foot as a single line. If one views it at higher power these cells are seen to have membranes appearing as the 'unit' membrane of Robertson. Unfortunately, the astrocytic foot itself completely lacks membrane appearing either as one or as two lines around it. It seems to 'share' the membrane of the adjacent cells. I would like to invite any electron microscopist to send me micrographs showing the expected adjacent membranes represented in our diagram (Figure 64). It has been argued that the membranes are only seen when they are almost normal to the plane of section, but this would not explain the absence of a membrane visible around the 'foot', when it is nearly always clearly present along most of its route around the adjacent cells, as well as, of course, around the nuclei and the mitochondria. The simplest explanation for the appearance of the 'foot' by electron microscopy, the sparsity of its contents, and its lack of personal membrane, is that it is due to shrinkage of two adjacent neurons, at the end of which there is a space containing a naked nucleus.

Astrocytes and oligodendrocytes are difficult to demonstrate by light microscopy in normal tissues, and some authors deny the existence of microglia. Clear light micrographs of astrocytes and oligodendrocytes - as opposed to diagrams - are hard to come across in the literature.

Usually when a nucleus is seen near an axon, the adjacent cell is called an oligodendrocyte. The rationale behind this is not at all clear. One may surmise that it is because most histologists regard any nucleus with little cytoplasm evident around it as an oligodendrocyte. In this monograph it is suggested that the nuclei of Schwann cells are the nuclei of adjacent connective tissue (Figure 65). In this location we would regard these oligodendrocytes as naked nuclei, and not neurons.

A few 'classical' views of these kinds of cells are shown (Figures 58-62), but a serious effort needs to be made to resolve the contradictions and uncertainties about the different kinds of allegedly different neuroglial cells exhibited in Tables 2-12 and Appendices 1 and 2. I would suggest that all the following cells with processes are neurons: astrocytes other than the reactive ones; oligodendrocytes and microglia, when they are described as having processes. Cells which have no processes and no cell membranes hard by are naked nuclei; reactive astrocytes; microglia; 'satellite' cells; oligodendrocytes, when they are described as having little cytoplasm. The latter group are probably identical with the original cells of Deiters' (1865), but they should not be confused with the neurons of the lateral vestibular nucleus of Deiters', which have been so extensively studied by Hyden and his school (Hyden, 1961, 1967).

THE SYNCYTIUM

This pervasive tissue consists of a fluid probably of similar composition to that of the cerebrospinal fluid, containing naked nuclei, mitochondria and fibres. It would be of

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM




FIGURE 63 Electron micrograph of astrocytic foot from adult rat cerebral cortex. The original text reads: 'At the bottom of the field is the nucleus (Nuc) of a protoplasmic astrocyte. This shows condensations of karyoplasm beneath the nuclear envelope. Arising from the top of the cell body is a thick process that contains a bundle of filaments (f). Other organelles present in the cytoplasm include mitochondria (mit), microtubules (m), free ribosomes (r), rather short cisternae of the granular endoplasmic reticulum (ER), and two lysosomes (Ly). The outline of this astrocyte is very irregular and is partially picked out with triangles, which show that its contours follow those of the dendrites and axons in the surrounding neuropil. Sheet-like processes (asterisks) from neuroglial cells such as this one pervade the neuropil and are recognised by their irregular contours.' This figure is reproduced from Peters, A., Palay, S.L. and Webster, H. de F. (1976) *The Fine Structure of the Nervous System* (Philadelphia: W.B. Saunders Co.), page 237, by kind permission of the authors and publisher




FIGURE 64 Diagram of an astrocytic 'foot' between four neurons, abutting on to a capillary as visualized by electron microscopy. The membrane around the neurons and capillary cells can be seen, but where is the membrane of the astrocyte itself(?) of which the foot is an outgrowth? The astrocytic foot should always be seen as having a membrane of its own, as well as those of the neurons between which it protrudes. In electron micrographs the membranes from what are believed to be the neurons can be seen clearly along most of their courses, so that the suggestions that the absence of the astrocytic membrane can be explained by the orientation of section or by the lack of staining of oblique sections are not satisfactory for the usual absence of the membrane of the astrocytic foot.




FIGURE 65 A whole mount of a small nerve in a thin spread of rat mesentery, showing the longitudinally orientated connective cells of the perineurium, and the nuclei of the Schwann cells in the interior of the nerve. May-Grunwald-Giemsa stain. The bar is 50 µm. This figure is reproduced from Bloom, W. and Fawcett, D.W. (1968) *A Textbook of Histology*, 9th edn (Philadelphia: Saunders), page 325, by kind permission of the authors and publisher. Please note the similarity of appearances and dimensions of fibroblast, pericyte and Schwann nuclei.

low viscosity were it not for the latter three components, which have the effect of stiffening it. When neuroglial clumps are examined in 0.25 mol/l sucrose by vertical illumination, they glisten white, and they show autofluorescence (Hillman, Hussain and Sartory, 1973). The syncytium or 'ground substance' is translucent to transmitted light. It should not be confused with the intracellular 'ground substance' of the electron microscopists (see, for example, Wolosewick and Porter, 1979).

The syncytium contains two kinds of fibres with diameters of less than 2-3 µm, whose lumens are not visible even under high-power light microscopy; there are bundles of fibres constituting tracts and commissures. We have reproduced a diagram produced by Bonin in Bailey and Von Bonin (1951), to show the richness of intracortical connections which had already been shown by physiological and histopathological techniques by that date (Figure 66). Of course, there are now known to be many more. There is also a rich network of apparently single fibres going in every direction. These show up well with Palmgren's staining procedure (Figure 38). In the literature, little attention has been paid to the latter fibres, and no-one has speculated about what they do. One can easily imagine that a particular such fibre would be very difficult to trace histologically, and it would be even more difficult to find the similar fibre again in another animal or post mortem human specimen. Nevertheless, it would probably be of value to try to envisage experiments which could find out what these ubiquitous straight single fibres can do.




FIGURE 66 Cortico-cortical connections revealed by physiological neuronography in the chimpanzee (after Bonin). This diagram is reproduced by kind permission of the publishers, from Bailey, P. and von Bonin, G. (1951) *The Isocortex of Man* (Urbana: University of Illinois Press), page 239

physiologically. They do not appear to touch, to connect, or to synapse with, each other.

NAKED NUCLEI

The naked nuclei are ubiquitous in the central nervous system (Figures 25-36). Some can probably be identified as free nuclei in subcellular preparations (Figure 67). These nuclei are denoted as 'naked' here because they are devoid of cell membrane (although not of nuclear membranes); the term 'naked' here does not mean that they do not have cytoplasm, which we have concluded is the whole syncytium. Since one never sees nucleoli in these nuclei, it may be presumed that they do not possess them (Figures 60 and 62). Their nucleoplasm is granular after staining, but it seems very likely that in life it is a uniform fluid, and only becomes granular on dehydration.

In tissue culture active phagocytes can be seen and they are often identified as microglia. They are probably also naked nuclei.

WHITE MATTER

White matter can be seen in the cerebrum, the cerebellum and the spinal cord (Figures 38-40 and 68). The individual axons are of a thickness which permits them to be seen by light microscopy, especially when stained with silver salts. At low-power magnification, bundles of fibres can be seen in longitudinal section (Figure 40). In thick transverse sections of the spinal cord one can focus at high-power magnification through the depth of the section, and see that one is observing fibres which are truly longitudinal and three-dimensional.

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM




FIGURE 67 Light micrographs of free nuclei from rat brain liberated into 0.2 mol/l KCl by ball mill agitation. Nuclei are stained with the counting diluent. 0.1% methylene blue-chloride in buffered 0.2 mol/l KCl. 1, Nuclei of neurons from several neuronal types; note heavily stained prominent central nucleolus and pale chromatin network (except in folded and partly disintegrating nucleus at left centre). 2, Nuclei of glial elements, probably oligodendroglia; note dense, heavily stained chromatin mesh and dark multiple paracentral nucleoli; these nuclei are uniformly oval or round and are about the same size as an erythrocyte; in this medium they stain brilliantly and have an iridescent sheen. 3, Large oval to round pale nuclei with multiple paracentral nucleoli and diffuse chromatin mesh; these presumably derive from astrocytes. 4, Small, crescentic nucleus which may either be of microglial origin or from vascular endothelium; such nuclei are relatively common in cortical samples. 5, Myelin fragment. 6, Erythrocyte. All photo-micrographs on recorded Eastman ortho-press plates at a uniform magnification of $\times 1,450$. This figure is reproduced from Nurnberger, J.I. and Gordon, N.W. (1957) in Ultrastructure and Cellular Chemistry of Neural Tissue, ed. Waelsch, H. (London, Cassell), page 109, by kind permission of the authors and publisher. Please compare these nuclei with those of oligodendrocytes (Figure 60)




FIGURE 68 Drawing of a transverse section of a gorilla spinal cord, with the ventral surface above, and the dorsal surface below. Fibres can be seen in the ventral and dorsal horns by light microscopy, and their diameters are of the order of micrometres. From Waldeyer, H. (1888) Das Gorilla Ruckenmark. Akad. Deutsch. Wissenschaft, Phys. Abh. III, Table 13

It is generally held that, like peripheral nerves, the fibres of the white matter are myelinated, although they have neither nodes of Ranvier nor Schmidt-Lantermann clefts. The only evidence for their similarity is that they both appear as relatively thick concentric circles. It is hard to find electron micrographs in the literature of 'myelin lamellae' in white matter of the spinal cord, while they are always found in peripheral myelinated nerves (page 176).

GANGLION CELLS

Cells from dorsal and sympathetic ganglia are spherical or oval. They are best visualized by phase-contrast microscopy of hand isolated cells (Hillman, 1966;

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

Hillman and Khallawan, 1970). Under these conditions they can be seen to have extraordinarily few axons or dendrites, perhaps only one or two. The possibility that many dendrites are not seen, because they adhere to the cell body by surface tension after isolation, can be discounted, since one does not see such processes waving about when the cell bodies are manipulated in aqueous media - as, for example, cilia are observed *in vitro* on tracheal epithelia or on paramecia. The cytoplasm of the ganglion cells is granular, probably due to intracellular mitochondria. There are no end-feet or synapses apparent by light microscopy on the surfaces of the ganglion cells, despite the physiological evidence that synaptic activity as defined physiologically is a characteristic of some sympathetic ganglion cells.

The population of neurons in dorsal and sympathetic ganglia appears to be homogeneous, when viewed by light microscopy, either unfixed or stained (Figures 17, 18 and 37), although there appear to be approximately two intensities of staining. There is a fluid between the neurons, and the ganglion cells are surrounded by capsules - diamond-shaped in the case of the autonomic ganglia, and spherical in the case of the dorsal root ganglia. The capsules form a considerable proportion of the weight and volume of the whole ganglia.

Ependymal Cells and Endothelial Cells

In the central nervous system the ventricles are lined by columnar or cubical ependymal cells (Figure 69). The capillaries are composed of the typical small narrow flattened endothelial cells lining capillaries elsewhere in the body, and no further description is necessary.




FIGURE 69 Anterior horn of lateral ventricle of rat stained with Holzer's stain showing the cubical to columnar ependymal cells, and syncytial fibres. The bar is 50 μm

MYELIN SHEATH

Much of the understanding of the structure of cell membrane is extrapolated from studies on myelinated nerve fibres into which the Geren model (1954) has been incorporated (Figure 70). Both the lamellar structure of the myelin sheath and the Geren model need careful examination. We will discuss the Schwann cell separately.

1. The myelin lamellae are virtually always seen perfectly transversely or as equally spaced parallel lines (Figure 71). Oblique views are hardly ever seen. If one looks at each of the illustrations in the references listed (Table 28), one can see many axons which have obviously been cut obliquely, since they appear oval instead of circular. Yet in none of these is there a significant incidence of oblique views of the individual lamellae. In general the width of the individual lamella as seen at the maximum diameter of the whole myelinated nerve should bear the same relationship to the width of the lamellae at its minimum diameter, as the whole nerve maximum and minimum diameters bear to each other. Examination of electron micrographs reveals a uniformity of spacing of the lamellae virtually independent of the orientation of the section.

This uniformity of spacing could occur if, for example, the nerve were compressed while it was being sectioned. However, these references were only chosen because in each of their illustrations nerves can be seen in several orientations. It would be inconceivable that compression, either during separation of the nerve or during section, would cause the myelin sheaths to be thinned in a random orientation relative to the source of compression, rather than clearly related to the direction from which it came. Of course, if an illustration contained only one myelinated nerve, compression could apparently cause the same spacing in the longer and shorter axis of the visible face of the section.

2. The lamellae show a constant spacing on longitudinal section. If one were to cut a Swiss roll (jelly roll) longitudinally through its maximum diameter, one would expect the layers to appear equally spaced. If one cut a longitudinal section away from the maximum diameter, one would expect the layers nearer the centre to appear further apart (Hillman and Sartory, 1980, page 37). Therefore, there are two geometrical constraints imposed on the myelin sheath as seen in longitudinal section. If the section has gone through the maximum diameter, firstly, the lamellae should appear equally spaced; secondly, the axon should be clearly visible; and thirdly, the sheath should be seen on both sides of the axon. These three geometrical imperatives rarely speak together.

If, on the other hand, one can not see the axon, it follows that the longitudinal section has not passed through the maximum diameter of the axon, and therefore the lamellae should again show splaying. It has proved impossible to find in the literature, or to coax from any electron microscopist, any micrograph showing the lamellae in any spacing other than equal in longitudinal or transverse section. In summary, one can say that the commonest expectations arising from geometry of longitudinal sections of real tubes are realized rarely by the electron microscopists, but, alas, more commonly by the graphic artists.

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM




FIGURE 70 A diagram of how the Schwann cell is believed to ensheathe the axon according to the Geren model. CP is the cell process, P is the myelin-forming process, M is the mesaxon, D is the major dense line, I is the intraperiod line, t is the external tongue process. This figure is reproduced by kind permission of the author and publishers, from Peters, A. (1968) The morphology of axons of the central nervous system. In *The Structure and Function of Nervous Tissue*, ed. Bourne, G.H., Vol. 1 (New York: Academic Press), page 170

THE FINE STRUCTURE OF THE NERVOUS SYSTEM




FIGURE 71 Electron micrograph of a transverse section of the cochlear nerve of a cat. This figure is reproduced by kind permission of the author, Mugnaini, E., and publishers in Fawcett, D.W. (1981) *The Cell* (Philadelphia: Saunders), page 31

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

TABLE 28 Publications showing myelin at several orientations in a single illustration, but in which the spacing of the lamellae do not appear to differ significantly in the various orientations seen in the individual illustration. Also there is no significant variation in spacing of lamellae either between several illustrations in the same publication, or between publications, although, obviously they would not have been sectioned all at the same orientations

Tissue	Authors
Cat skin nerve	Gasser (1958)
Rat spinal cord	De Robertis, Gerschenfeld and Wald (1960)
Rabbit peripheral nerve	Causey (1962)
Mouse sciatic nerve	Robertson (1962)
Rat sciatic nerve	Schade and Ford (1965)
Rat dorsal root ganglion	Novikoff (1967)
Monkey cerebellum	Glees and Meller (1968)
Rat sciatic nerve	Elfvin (1968)
Rat optic nerve	Peters (1968)
Rat optic nerve	Vaughn and Peters (1968)
Rat optic nerve	Hirano (1969)
Rat medullary pyramid	Curtis, Jacobson and Marcus (1972)
Rat sciatic nerve	Johnston and Roots (1972)
Calf cerebral cortex	Deutsch (1973)
Monkey lateral geniculate body	Glees (1973)
Guinea pig sciatic nerve	Bloom and Fawcett (1975)
Rat spinal cord	Rhodin (1975)
Baboon peripheral nerve	Landon and Hall (1976)
Baboon middle popliteal nerve	Ochoa (1976)
Cat medial gastrocnemius nerve	Aoki and Munoz-Martinez (1981)
Rat optic nerve	Leeson and Leeson (1981)

3. The lamellae appear close together by electron microscopy. Although each individual lamella is below the resolution of the light microscope, together the whole sheath should be thick enough to appear solid on light microscopy. Although the Schwann cell membrane can be seen or detected by light microscopy, the lamellae of which it is supposed to be composed - according to the generally accepted Geren view - cannot be seen in longitudinal or transverse sections of unstained peripheral nerves (Figures 19,41,73).

4. By high-power phase-contrast microscopy, Brownian movement can be seen in the cytoplasm of the Schwann cell. This is well illustrated in the beautiful time-lapse films of Costero and Pomerat (1951).

5. 'Myelin figures' are seen in normal kidney (Figure 72) and more frequently in liver. In a number of toxic conditions it has been shown by electron microscopy that the

THE FINE STRUCTURE OF THE NERVOUS SYSTEM

number of 'myelin figures' increases in mammalian liver (Emmelot and Benedetti, 1960, 1961; Albot and Jezequel, 1962; Herman, Eber and Fitzgerald, 1962; Rouiller and Simon, 1962; Salomon, 1962; Steiner and Baglio, 1963; Mikata and Luse, 1964; Steiner, Mujai and Phillips, 1964; Stenger, 1964; Herdson and Kaltenbach, 1965; Campbell, 1967). It is hardly likely that there are occasional pieces of real myelin (as claimed to be present around peripheral axons) in the liver and kidney; secondly, it seems difficult to believe that toxic agents, such as dimethyl nitrosamine, hepatitis virus, ethionine, thioacetamide, naphthylisocyanate, N-2-fluoronyldiacetamide, carbon tetrachloride or thiohydantoin, would themselves cause myelination. Many toxins like lead, mercury and diphtheria cause demyelination. It is interesting that Emmelot and Benedetti (1961) talked of 'parallel-packed smooth membranes, resembling myelin sheaths'. Others have talked of 'myelin figures' or 'myelin-like figures'.

THE GEREN MODEL

6. The Geren model is illustrated in Figure 70. If the Schwann nucleus did wrap around the axon, as the Geren 'model' proposed, the nucleus would be expected to appear as if it were at the apex of the wrapping cell, i.e. in the middle of the angle subtended by it at the anti-node, rather than the way which it is drawn in diagrams of light and electron micrographs as the occupant of a concentric tube of Schwann cell cytoplasm parallel to the axon.

7. The Geren model shows the unfolding of the double-layered myelin sheath externally, which has been named the external mesaxon, and their parting at the internal mesaxon. The joining or separation of two double layers should, of course, make a four-layered membrane, which should thus be double the thickness on low-angle diffraction of the cell membrane. A four-layered mesaxon is simply not seen. One could postulate a mechanism whereby the two double membranes would 'fuse' together, but this would require very active proteolytic and lipolytic enzymes, which would also have the remarkable properties of knowing which two of the four membranes should be dissolved away, and when to stop before dissolving the other two. This problem would only be slightly simplified if our view that the membrane was of single thickness (page 133) were accepted.

8. Another requirement of the Geren model is that the Schwann cell cytoplasm is continuous between the two double-layered membranes investing the axon (Figure 70). Can any histologist, electron microscopist, or neurobiologist using tissue culture produce a convincing photograph of this?

9. In tissue cultures of myelinating fibres the movements of the swellings along the axon were observed (Figure 73); electron microscopy of myelinated axons showed lamellae around the axons. These findings led to the 'Geren model', which was a hypothesis that the bulges on the axon were the nuclei of Schwann cells, and that they rotated around the axon with their cytoplasm to produce the lamellae; implicitly, this would involve rotation around the axon in a transverse plane, since diagonal rotation would produce lamellae of characteristic but totally different orientation.




FIGURE 72 Electron micrograph of rabbit kidney fixed with glutaraldehyde and osmium tetroxide (2%), and stained with lead citrate and uranyl acetate. Please note the large 'myelin figure' in the centre. The bar is 1 μm

However, to my knowledge no-one has demonstrated the two main findings which would be required by, and would justify, the hypothesis; viz: firstly, by time-lapse photography of tissue cultures one should be able to take successive views of the nucleus rotating radially around the axon, not just appearing in different positions diagonal to each other; secondly, one should be able to make histological and electron microscopic preparations of successive stages of the invagination and rotation of the Schwann nucleus, including views of the membranes before they have fused together. Until these findings have been made positively, the 'Geren model'

THE FINE STRUCTURE OF THE NERVOUS SYSTEM

must remain a hypothesis. I think that we have here a clear example of the trans-substantiation of a much-repeated hypothesis into a hallowed finding.

The explanation for the appearance of the lamellae is that they represent deposits of the intracellular fluid of the narrow tubular sheath around the axon, when it is dried out by subjection to the high vacuum and the bombardment by the electrons in the electron microscope. In longitudinal section the salty solution would dry out as equally spaced lines parallel to the walls of the sheath in which it was enclosed; in transverse section it would deposit as the fluid within a thin vertical tube as parallel concentric circles. This explanation is similar to that given for the appearance of the endo-skeleton in the axon.

It should be noted that although the electron microscopists claim that they are looking at membranes largely consisting of lipids and proteins, the thin sections have probably been prepared with at least one of the following agents, which can extract lipids: ethanol, glutaraldehyde, propylene oxide or xylene (Hopwood, 1969, 1970; Vanha-Pertulla and Grimley, 1970). In addition, water-soluble proteins must be extracted by all aqueous reagents.

The myelin sheath is usually regarded as the insulation of the axon. The staining of the sheath by such reagents as osmium salts and Sudan III makes it likely that the liquid within the sheath contains a high concentration of lipids and/or proteins, which would impart a high resistance to the sheath, and would render the antinodal region more resistant than the region of the node of Ranvier. Thus the simpler concept of the 'myelinated' nerve presented here also fits in with saltatory conduction (Stampfli, 1954).

If the lamellae are an artifact of preparation, they cannot represent a useful model for study of individual cell membranes.

It should be emphasized that, for the reasons presented above, we do not believe that the lamellae occur either in the living peripheral or central nervous systems. The cytoplasm appearing to be present in the Schwann cells is simply the cytoplasm of the connective tissue between the axons.

Thus the myelinated and unmyelinated nerves have a simpler structure than the electron microscopists ascribe to them. The myelinated nerve consists of two concentric tubes containing clear fluids in which intracellular movements occur in life; the tubes are interrupted by nodes of Ranvier and incised partially by Schmidt-Lantermann clefts. The unmyelinated nerves exhibit the elegance of a narrow capillary tube made of the finest bohemian glass.

Furthermore, it must be stressed that the chemistry of extracts of neural tissue does not necessarily provide any direct information about the structure of the cells in the intact living animals.

THE SCHWANN CELL

The Schwann cell represents an interesting problem. Schwann's (1838) original diagram shows what appears to be a nucleus lying on the outside of the axon (Figure 11). Of course, his drawing was made before photomicrographs had become a practical proposition. They had only been developed by J.B. Reade in 1836 (please see Henderson,




FIGURE 73 Living cultures of kitten cerebellum. (Upper) 43-day-old culture; single swelling with maximum thickness of $6.32\text{ }\mu\text{m}$ in nerve fibre of $1.3\text{ }\mu\text{m}$ average thickness. Phase-contrast (lower) 34-day-old culture; multiple irregularities in myelinated fibre. Phase-contrast from Hild (1959), Myelin formation in cultures of mammalian central nervous tissue. In *The Biology of Myelin*, ed. Korey, S.R. (London: Cassell), page 195, reproduced by kind permission of the author and publishers

1978) and demonstrated before an audience by John Dancer in 1840. However, the following authoritative works - including Causey's monograph entirely devoted to the Schwann cell - show diagrams of the Schwann cell, rather than photomicrographs (Ramon y Cajal, 1909; Schmitt, 1958; Causey, 1960; Wechsler, 1970; Peters, Palay and Webster, 1976; Landon, 1976; Sunderland, 1978). This curious lacuna in the literature should be a matter of concern. The diagrams in the works cited generally show the Swiss, or jelly roll, tube, surrounded by an apparently further tube containing a nucleus; the latter is believed to be the nucleus of the Schwann cell. In the rare

THE FINE STRUCTURE OF THE NERVOUS SYSTEM

circumstance of the nuclei being seen by light microscopy, it may appear as:

1. A granule within the myelin sheath (Nageotte, 1932). Nageotte, like some of the previous authors he cited, described it as the 'Schwann cell syncytium'. In a comprehensive review of the literature, Sunderland (1978, page 14) gives the thickness of the myelin sheath as between 0.25 µm and 3 µm in various myelinated nerves. Thus, one would have to suppose that the diameters of the nuclei were of the same order as those values. These would make them hardly visible by light microscopy. In fact, if one views them carefully, they appear more as granules without clear membranes than as nuclei.
2. Another appearance by light microscopy, of which Bloom and Fawcett (1968, page 325) gave a beautiful illustration, is of nuclei lying on the surface of the axons (Figure 65). The cytoplasm around these nuclei is not evidently connecting them to the axon. Also, they are quite indistinguishable in appearance from the nuclei of capillaries or connective tissue.
3. The third appearance is well seen in the tissue culture of Hild (1957, 1959) reproduced here as Figure 73, as well as Pomerat, Hendleman, Raiborn and Massey (1967). These shots are taken from time-lapse motion films, similar to those of Costero and Pomerat (1951). They are particularly important, since they are regarded as being clear evidence for the Geren model, which has already been discussed. However, when one examines these pictures carefully, it becomes extremely doubtful whether they can be described as nuclei at all. Nuclei are characterized both in unfixed and stained cells of all species by having a different refractive index or staining more darkly in cells of all tissues. It seems quite inconceivable that they should appear transparent and structureless as they do in tissue culture. I would suggest that these 'Schwann cell nuclei' seen in culture are early swellings of the axoplasm, which subsequently enlarge and widen to give the classical appearance of the sheath.

Sometimes by electron microscopy large nuclei are seen adjacent to the lamellae, and these are called either Schwann cell nuclei or oligodendrocytes; the latter identification is based on their having similar appearances to those of cells called oligodendrocytes in the central nervous system, since they have scant cytoplasm (references in Table 28). It is noteworthy that the asymmetrical appearance of the nuclei in the Geren model, like the appearance of the mitochondria in the cytoplasm of the Schwann cell, is very rarely seen in electron micrographs.

One must then conclude: (1) that the granules seen sometimes by light microscopy within the antinodes are precipitates due to staining; (2) the larger nuclei on the surface of myelinated or unmyelinated axons by light or electron microscopy are the nuclei of connective tissue or endothelial cells outside the sheath; (3) the Schwann cells in tissue culture are concentric tubes originating from the axons (as was suggested by Hardesty in 1904); (4) the diagrams in the textbooks represent artists' views based upon a hypothesis stretching beyond the discipline of careful observation.

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

The real sheath probably contains high concentrations of lipids and proteins because it stains with osmium salts and Sudan III, although the transparency of the sheath and the Brownian movement within it make its contents liquid. We conclude that the Schwann cell nucleus is the nucleus of the connective tissue adjacent to the axon, and it may well be mobile (Figure 65).

8

Other Evidence Supporting these Conclusions

TISSUE BETWEEN NEURONS

The original observation of Virchow (1846) of a 'nerve glue' was widely accepted at the time. Bevan Lewis (1889) characterized the ground substance between the neurons as 'vastly preponderating over the cell and nuclear elements of the neuroglia'. Brierley (1957) quoted Bevan Lewis (1889), Obersteiner (1890), Taft and Ludlum (1929), Taft (1938) and Hogue (1946) describing it as a fine granular mass, and similar observations were made by Held (1909), Hess (1953) and Windle (1958). Brierley (1957) said that this ground substance was 'the main factor responsible for the blood brain barrier'.

Von Economo and Koskinas (1925) and von Economo alone (1929) stained human cerebral cortex probably with cresyl violet and toluidine blue (*ibid.*, page 24). The latter author (1926) had calculated a grey matter-cell coefficient, which was the ratio of the mass of the grey matter of the cortex to the total cell mass. This turned out to be 20-27:1, giving the cells 4-5% of the weight of the grey matter. He suggested that their ratio could be used as an index of the stage of development of any brain, since higher animals had a higher proportion of cells. The cellular elements were found to be a small proportion of the tissue as we found (Tables 15 and 16). Of course, in both studies the proportion could have been depressed by the staining procedure not showing up all the elements, as Wyckoff and Young (1956) and Nauta and Feirtag (1979) among many others have asserted. Yet it is difficult to allege that some elements have not been stained without careful and comprehensive studies identifying all the cells, using independent 'specific' procedures, which have been demonstrated to show up all of, and only, the elements for which they are claimed to be specific.

Alzheimer (1910) and del Rio-Hortega (1919b) suggested that there were some amoeboid cells in the brain which could move, and they called them microglia. Movement of the cells in cerebral slices was seen directly by Dodgson (1948). He observed sections of mouse brain slightly stained with methylene blue; when he applied light pressure 'sluggish streaming movements', as well as Brownian movements of granules, occurred. His tissue was fresh, and he did not have to bathe it in any medium. Yet the movement could have resulted only from the damage when the slices were cut, or

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

from the pressure, and one cannot exclude the possibility of it being an artifact. He did not pursue these investigations (personal communication, 1978).

Cells migrate during the development of the central nervous system. Cells adjacent to the lumen of the neural tube divide first, and then migrate centrifugally through the tissue of the neural tube; this, therefore, could not be solid, and must be syncytial at that stage. Sauer (1935), like the many electron microscopists who have since denied the existence of a syncytium (please see Jacobson, 1978, page 32) nevertheless show the migration of single neuroepithelial germinal cells up columns of presumed cytoplasm, which are sometimes claimed to be attached to a basal membrane. However, these columnar cells stand like grenadier guardsman in sentry boxes with spaces containing no nuclei between them. Must not these spaces be dubbed as syncytial? It is extremely difficult to conceive of a column of cells fleeing centrifugally away from the lumen of the neural tube if the tube itself was solid with cells; there would have to be either a counterflow of cytoplasm, or radial growth of the whole neural tube at quite an explosive rate.

At a later stage of development, cells in the spinal cord, cerebellum and isocortex have been shown to migrate from the ependyma over a period of a few days (Figure 74). One cannot avoid the conclusion that at this stage, also, they must be migrating through a syncytium. Nevertheless during the whole of this period the cortex appears largely solid by electron microscopy (Voeller, Pappas and Purpura, 1963; Hassler and Stephan, 1966; Aghajanian and Bloom, 1967; Gruner and Zahnd, 1967; Meller, Briepohl and Glees, 1968a,b).

The existence of a syncytium in the brain during development is not, in itself, proof that it persists into adulthood. However, there are two other lines of evidence for this belief. Smart and Leblond (1961), using autoradiography, showed that new neuroglial cells could be formed in the adult brain. Microglia may surround injured brain as a result of proliferation of cells adjacent to the wound (Maxwell and Kruger, 1965; Cammermeyer, 1970; Vaughn, Hinds and Skoff, 1970; Stenwig, 1972; Korr, 1980). Or they may originate from leucocytes in the blood which escape into the region of injury (Konigsmark and Sidman, 1963; Oehmichen, Gruninger, Saebich and Narita, 1973; Matthews, 1974; Oehmichen, 1978). Whatever their origin, microglia or phagocytes - with apparent nuclear diameters of 5-20 μm in the stained preparations, and, therefore, larger *in vivo* - migrate in considerable numbers around the site of injury, inflammation or infection, starting as early as 12 hours after (Del Rio Hortega, 1932, pages 520-521, quotes 29 references). These movements are comparatively rapid, of the same order as occurs in 'axonal flow'. It seems extremely unlikely that the cells or nuclei could move at such a rate if, indeed, the central nervous system was composed of a testudinous block of cells with an extracellular space of between 4% and 25%; this would give linear extracellular spaces of approximately 1-10%. Since the diameter of the nuclei is much larger than 1-10% of the maximum diameter of the neuronal somas, it could not be supposed that microglia or phagocytes could interpolate themselves between what are considered to be the packed neurons and neuroglia. Furthermore, if they were, they should be caught in the act at least occasionally by zealous microphotographers.

OTHER EVIDENCE SUPPORTING THESE CONCLUSIONS




FIGURE 74 Germinal and other cellular zones and layers in the developing cerebral isocortex, spinal cord and cerebellum of mammals. Progressively later stages of development are shown from left to right. CP, Cortical plate; E, ependymal layer; EG, external granule layer; Gr, granule layer; I, intermediate zone; LD, lamina dissecans; M, marginal zone; Mo, molecular layer; P, Purkinje cell layer; S, subventricular zone (also called subependymal zone); V, ventricular germinal zone. Mitosis can sometimes be seen. This figure is reproduced from Jacobson, M. (1978) *Developmental Neurobiology*, 2nd edn, (New York: Plenum Press), page 62, by kind permission of the author and publishers. Note the cell migration.

In a similar fashion, many large molecules like inulin, serum proteins, ferritin and horseradish peroxidase enter the parenchyma after injection from the ventricles, the choroid plexus or the subarachnoid space (Klestadt, 1915; Bowsher, 1957; Lee and Olszewski, 1960; Pappenheimer, Hersey, Jordan and Downer, 1962; Klatzo, Miquel, Ferris, Prockop and Smith, 1964; Rall, 1964; Brightman, 1965a,b; Kitai and Bishop, 1981). It is quite possible that the damage due to the manipulation of the brain or pressure on it adds to the spread of these large molecules, or their micelles, but there is too much evidence of such 'pinocytosis' in the central nervous system, to explain away all the findings describing the passage of these large particles. Obviously, the movement of particles, cells or nuclei implies that the resistance to them is low.

Conel (1939–1963) made a monumental six-volume study of the human cerebral cortex from babies of 1 month to 2 years old, and we have selected drawings from the two extremes of age, which illustrate the following points (Figure 51). With increasing age, there is an increase in (a) the number of neurons, (b) their density, (c) their arborization, and (d) the overlap of these fibres. He did not show any microglia or naked nuclei. Dendrites are generally believed to connect with the hundreds to thousands of synapses which most authors believe inhabit the surfaces of neurons (Tables 22 and 23). In the presence of such a mass of processes radiating

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

from the soma in all directions, it is difficult to imagine how the same space of the neuropile could also contain neuroglial cells of the form shown (Figure 50; see also Conel, *ibid.*; Kitai and Bishop, 1981). The only circumstance in which they could coexist as close neighbours to each other would be if the neuroglia contained holes like an Emmentaler cheese, through which the processes of the somas could pass. I do not think that anyone has described this caseous appearance, nor have I seen it myself in histological sections or in micrographs.

RESISTANCE AND IMPEDANCE OF BRAIN AND SPINAL CORD

Many measurements have been made of the resistance and impedance of cerebrospinal fluid, cerebral cortex, white matter and spinal cord, of rabbits and cats (Crile, Hosmer and Rowland, 1922; Freygang and Landau, 1955; Ranck, 1963a,b; van Harreveld, Murphy and Nobel, 1963; Li, Bak and Parker, 1968). These authors found values in the range 100-1200 Ω cm for either of these parameters. However, the resistance of the membranes of individual excitable cells measured using intracellular microelectrodes was 4-43 M Ω (Coombs, Eccles and Fatt, 1955; Frank and Fuortes, 1956; Lux and Pollen, 1966; Nelson and Lux, 1970; Li, Okujava and Bak, 1971). Neuroglial cells in tissue culture (Hild and Tasaki, 1962) were calculated to have resistance of 2 M Ω when the processes were taken into account (Li, Okujava and Bak, 1971). In reviewing these measurements it is quite clear that the values of resistance and impedance of excitable cells, and also cells identified by the authors as neuroglia, were very high relative to the values for the tissues as a whole, and it was concluded that 'the low impedance of gray matter and of white matter measured in the fibre direction is consonant with the presence of an appreciable extracellular space in central nervous tissue, and can be reconciled only with difficulty with the paucity of extracellular space shown in many investigations of the tissue with the electron microscope' (Van Harreveld, 1966, page 69). Thus, all the findings agree that tissue impedance is low. It is usually assumed that this is due to shunting in the extracellular space, but it also fits in well with the high-resistance and impedance neurons being surrounded by a low-resistance syncytium.

PROPORTION OF BRAIN IN WHICH ABRUPT DIRECT CURRENT POTENTIALS CAN BE RECORDED

When an intracellular electrode penetrates a living tissue the sudden change in direct current voltage which is recorded is normally interpreted as penetration of the membrane of a cell. If the central nervous tissue is then stimulated by electric current, transmitters, hypoxia, or potassium ions, and an action potential results, this is taken to mean that the cell was a neuron; if not, the cell penetrated is believed to be a neuroglial cell.

All living cells exhibit transmembrane potential differences so that a measurement of the proportion of the distance in which an electrode tip, say, less than 1 μm in diameter, records a voltage compared with the total distance which the electrode travels gives one an assessment of the closeness of the packing of the cells in the tissue; the measurements are not precise, because the tissues are elastic and the electrodes may bend. Very often the electrode does not record the maximum voltage

OTHER EVIDENCE SUPPORTING THESE CONCLUSIONS

across the cell membrane, because the larger the diameter of the tip of the electrode relative to that of the soma, the smaller the voltage it records; also, the electrode may become blocked (Adrian, 1956), or it may penetrate the soma or its processes obliquely (Woodbury, Hecht and Christopherson, 1951; Frank and Fuortes, 1955; Haapanen, Kolmodin and Skoglund, 1958). However, if one simply adds up total distances during which penetrations record voltages of any value, one derives a measure of cell density. This proportion was only 22% in cerebral slices *in vitro* (Figure 75; Hillman, Campbell and McIlwain, 1963). Even allowing for a 15-50% increase in volume of cerebral slices after incubation *in vitro*, it is evident that only a minority of the tissue consists of cells with membranes. In muscle, the proportion of distance of travel of the electrode without recording a voltage is much smaller, probably 5-10% (H. Hillman, 1962, unpublished).

VISCOSITY OF THE BRAIN AND SPINAL CORD

Anyone who has operated on, or assisted in an operation on, the brain or spinal cord of a living mammal, has the experience that the living tissues are extremely soft compared with other solid organs, such as the liver or muscle. This, of course, is an anecdotal observation, which one would like to have made more scientifically respectable by measuring such parameters as viscosity, compressibility, impedance to sound, or elasticity. Unfortunately the tissue is heterogeneous, and measurements one would make would be likely to be operational. Measurements on dead tissue, or tissue suffering from ischaemia, acidosis, autolysis, cooling or substrate deprivation would probably not be wholly relevant.

ASTROCYTES AND OLIGODENDROCYTES

Many neurohistologists, including Purkinje, Deiters, Hyden, Giacobini, Pavlin, Hertz and Hamberger, have dissected out single neurons by hand, but none of them have ever claimed to separate single astrocytes or oligodendrocytes, either deliberately or by accident. Hyden and Pigon (1960) reported that the neuroglial clumps mainly consisted of oligodendrocytes, with some astrocytes adjacent to blood vessels.

Hamberger (1963) said that his neuronal glia were 90% oligodendrocytes and 10% astrocytes, and capillary glia were 70% oligodendrocytes and 30% astrocytes, but nevertheless these cell types were never isolated as individual bodies from the neuroglial clumps. Of course, the fact of not having found particular evidence is not as cogent as positive evidence; this is especially true when it is not clear if the research workers mentioned actively sought to isolate astrocytes and oligodendrocytes, and failed, or if they did not try to do so. I, myself, have never had difficulty in isolating the characteristically shaped 'neurons', and I could also point to the naked nucleus in a neuroglial clump. Following Cummins and Hyden (1962), I could dissect out the nucleus from a neuron, but I, also, have never seen any other recognizable shape of a whole cell in a clump.

Throughout the literature on cell types in the nervous system, there are references to 'nerve glue', 'nutritive' cells, 'satellite' cells, and 'supporting' cells. These terms indicate both ignorance about what they do, and an albeit subconscious feeling that they represent a less worthy class of cell in the architectonic pantheon.




FIGURE 75 Four photographs of typical changes in potential differences during penetrations of $310 \mu\text{m}$, from the surface into slices of guinea pig cerebral cortex *in vitro*. At (a) the vertical displacement is 100 mV and the horizontal displacement 0.5 ms . In cerebral slices whose volume increases about 25% on incubation, the tips of the electrodes recorded a voltage for only a mean of 28% of their total excursions. As can be seen, many of the voltages recorded were very low, indicating very few cellular elements with large voltages. This recording is reproduced from Hillman, H., Campbell, W.J. and McIlwain, H. (1963) Membrane potentials in isolated and electrically stimulated mammalian cerebral cortex slices. *J. Neurochem.*, 10, 325-339, by kind permission of Pergamon Press

This may not be a trivial matter, because it preconditions the belief that the total volume of the brain is largely occupied by the 'more important' neurons. This seemingly semantic difficulty is compounded by the fact that 'satellite' cells and microglia being apparently smaller than neurons would be much more difficult to penetrate with an intracellular electrode to study their membrane properties, for example by iontophoresis. In general the electrophysiology of smaller cells is more difficult to study, and so less is known about them in the central nervous system.

9

Summary of Conclusions about Nervous Tissue in *Living Mammals*

1. The central nervous system contains neurons, naked nuclei, fibres, capillaries, ependymal cells and a syncytium, which is the largest component by volume.
2. The syncytium is a viscous jelly, in which naked nuclei move slowly and mitochondria move rapidly. The mechanical, physical, biochemical and pharmacological properties of the central nervous system are dominated by the properties of the syncytial ground substance.
3. All cells with processes are neurons. The only other cells are the naked nuclei, the ependyma and the capillary endothelium.
4. Cells in the central nervous system are not in contact with each other, even when their processes are close in rare locations like the Purkinje and the basket cells of the cerebellum.
5. Neuron somas are almost spherical in shape, and are invested in solid membranes. All the membranes are one layer thick. (Synaptic knobs do not exist in life, although the physiological properties generally attributed to synapses are real. Synaptic vesicles do not exist.) Each neuron has probably only one axon, and usually very few primary dendrites (Figure 76).
6. The cytoplasm of the neurons is a fluid of low viscosity, in which mitochondria and a few other unidentifiable particles move continuously. (The Golgi body, the endoplasmic reticulum, the lysosomes, the tubules, fibrils, filaments and trabeculae do not exist in living cells.)
7. The nuclear membranes of neurons and naked nuclei are imperforate.
8. The mitochondria both in the neurons and in the syncytium contain a clear fluid, the mitochondrioplasm. (The cristae do not exist in life.)
9. The nucleoli of neurons are surrounded by single-layered membranes. They contain a skein-like nucleolonema.
10. Naked nuclei are spherical and contain a uniform translucent liquid nucleoplasm; they do not have nucleoli.
11. Ganglia contain only spherical or oval neurons; synaptic knobs are not seen there.
12. Unmyelinated axons and nerve fibres are tubes. The axoplasm is a clear fluid of low viscosity. (It does not contain an endoplasmic reticulum, neurotubules, neurofibrils, neurofilaments or microtrabeculae.)




FIGURE 76 A diagram of a high-power view of a live neuron. Please note the absence of an endoplasmic reticulum, Golgi bodies, synapses, nuclear pores, endoskeleton or mitochondrial cristae. The cytoplasm, the nucleoplasm and the mitochondrioplasm are of low viscosity and each is uniform. Please compare with Figure 44

SUMMARY OF CONCLUSIONS

13. The myelin sheaths around peripheral axons are concentric tubes containing a viscous fluid. Nodes of Ranvier and Schmidt-Lantermann clefts are spaced along its length. (The Schwann cell does not exist.) The Schwann nucleus is the nucleus of a cell in the adjacent connective tissue. (The myelin lamellae do not exist in life.)
14. The fibres of the tracts in the white matter of the spinal cord are smaller in diameter than those in peripheral myelinated nerves.
15. Transmission is mainly electrical, but the excitability of the cell membranes can be conditioned by the chemical nature of the syncytium, including the presence and interactions of substances believed to be transmitters.
16. The syncytium and the choroid plexus are the blood-brain barrier.
17. The Geren model does not describe the events during the myelination of the axon.
18. Axoplasmic flow is a combination of diffusion, Brownian movement, streaming, laminar flow and convection.

Our views on the cellular structure of the brain are summarized in Figures 77 and 78.




FIGURE 77 Diagram of a low-power light microscopic view of brain adjacent to the ventricle, according to conclusions arrived at here. In addition to the ependymal cells at the upper margin, and two capillaries depicted, the other elements are neurons, naked nuclei and syncytium. Please note: there are no synaptic connections between any of the dendrites or cell bodies, nor are there any synaptic knobs. The processes are much straighter than appear in fixed dehydrated sections. In life, the brain is very soft. This figure should be compared with Figures 50 and 51




FIGURE 78 Diagram of a high-power view of the central nervous system of the mammal as it would be seen in life. A nerve tract traverses the right-hand corner. Please note the mitochondria inside and outside the neurons (page 122); they appear in all orientations. The neuron cell bodies are rounder than they usually appear in dehydrated sections (page 35). The neuronal nuclei contain nucleoli, which are surrounded by a membrane (fig. 18); the nucleolonema is seen in the nucleoli (page 160). In life the mitochondria, the naked nuclei and the nucleolonema are in constant motion. In the syncytium one can see fibrils much finer than the mitochondria (fig. 38). Nucleoli are not seen in the naked nuclei, whose nucleoplasm is uniform

10

Problems Arising from the Latter Conclusions

The first question which arises is, 'Have these views been held before, and, if not, why not?'

During the nineteenth century it was generally held that the central nervous system consisted of ganglion cells (neurons), Deiters' cells (microglia or naked neurons) and neuroglial material (pages 11-13). This view is similar to our own. In the twentieth century the different kinds of cells were identified by their being stained by newly described procedures. The idea then evolved, rather than originated, that if all the different kinds of cells characterized thus could all be stained in the same preparation, the whole tissue - with the exception of a small extracellular space - would be seen to be filled with these named elements. This was most recently stated by Nauta and Feirtag (1979). This belief was based on the acceptance by neuropathologists of the specificities of the staining procedures which their original authors had neither tested nor even alleged. Unless one examines the original papers (Table 10), one would naturally assume that the specificities and non-specificities of each staining procedure would have been rigorously tested. The reasons for the early failure to do so have been discussed briefly, viz: respect for the authority of the early neurohistologists; the fact that the concepts of control experiments and specificity were not considered important by then; the failure to appreciate that different procedures distort cells differently. We would not wish to encourage disrespect for authority, but would urge that experiment should always be more persuasive than the power or pelf of its protagonists. By and large the relevant control experiments are still waiting to be done (page 242), and specificity is better known for its assertion than its demonstration. History may well view even more unkindly the premature derivation of conclusions from these incomplete experiments, than the fact that the experiments themselves have not yet been completed - albeit after more than a busy half-a-century.

In the late 1940s and early 1950s the electron microscopists looked at the central nervous system and found it full of nuclei and mitochondria. They were very good at seeing nuclear and mitochondrial membranes but did not ask themselves why they could see the latter membranes so clearly and along most of their routes, but could not see the cell membrane so well around 'astrocytic feet' or cells they characterized

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

as oligodendrocytes (Figures 61 and 63). Whatever optical or geometrical explanations they could produce for their failure to see the latter as frequently as expected (Ham, 1974; Horne and Harris, 1981; Michell, Finean and Coleman, 1982) were rapidly shown to be irrelevant by the high frequency with which they could see the nuclear and mitochondrial membranes (Figures 31-33, 61 and 79).

The belief that the whole central nervous system consisted of neurons, neuroglial cells and a small extracellular space naturally led to the supposition that the pervasive mitochondria outside neurons seen by electron microscopists were located in the neuroglial cells thought to be present between the neurons. When they could not see the membranes which they expected to surround the neuroglial cells, they looked for geometrical explanations to explain the invisibility.

A number of possible educational and sociological factors which have predisposed to uncritical acceptance of the validity of the findings of electron microscopy have been discussed elsewhere (Hillman and Sartory, 1980b).

A further fundamental point must be made here. Observers use their brains as well as their microscopes. There is a mind in the eye of the observer, as well as an eye in the mind of the microscope. One is conditioned by expectations which have been implanted by education. Two observers see different parts of the same image, attach different degrees of importance to each part, and often interpret observations of the same image differently. For this reason I would strongly advocate the fabrication of three-dimensional models of any alleged structure (page 126).

IDENTIFICATION OF ASTROCYTES

The Golgi stain requires thick sections (up to 150 μm ; Sholl, 1953), so that one should be able to see a great deal more of the three-dimensional structure of sections than is possible with the usual thinner sections; modern histologists cut 4-6 μm , the earlier electron microscopists cut 'thick' sections of 0.5-1 μm , and modern practice is to cut sections 10-100 nm thick. Even the use of the thickest of these sections creates difficulties in identifying astrocytes by their 'feet'. If one cannot see a 'foot' it may be either because the cell is not an astrocyte, or because no 'foot' happens to be present in the plane of section, or because 'feet' do not exist. Thus serial sections have to be examined to identify with certainty a particular cell as an astrocyte by this criterion.

The lack of membrane of the astrocytic foot has been discussed (page 167). The simplest explanation for the appearance of the 'foot' by electron microscopy, the sparsity of its contents, and its lack of personal membrane, is that it is a shrinkage artifact between two neurons, at the end of which there is a lightly staining space containing a naked nucleus. Thus, in our view, the astrocytes are the nuclei of the syncytium, which are identical with microglia.

SYNAPSES AND TRANSMISSION

It has been shown that the vesicle hypothesis of transmission embraces a number of unlikely assumptions and findings which have not yet been demonstrated (pages 226-230). Nevertheless, there is a vast bank of data on the iontophoretic application of transmitters



FIGURE 79 Electron micrograph of layers I (above) and II (below) of cat cerebral cortex. The original legend of Jones and Powell's paper (1970) shows 'the bifurcating (ringed arrows) apical dendrite of a pyramidal neuron passing between two other cells to reach layer I. Note the densely packed neuropil of layer II with many small axon terminals, dendrites and dendritic spines; many of the spines can be seen to be derived from the small dendrites of the layer (arrow)'. We show this micrograph to indicate the appearance, at 2 o'clock and 9 o'clock, of two neurons with their membranes and, at 5 o'clock of one nucleus, with no apparent membrane around it. This electron micrograph is reproduced from Jones, E.G. and Powell, T.P.S. (1970) Electron microscopy of the somatic sensory cortex of the cat. II. The fine structure of Layers I and II. Phil. Trans. Roy. Soc., B, 257, 13-21, plate 17, by kind permission of the authors and the Royal Society.

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

and 'putative' transmitters (Table 27) to cells in the brain and spinal cord presumed to be neurons (Eccles, 1961, 1964; Triggle, 1967; Katz, 1969; Berl, Clarke and Schneider, 1975; Iversen, 1975; Fonnum, 1978; Pepeu and Kuhar, 1980; Bradford, 1982). The injection of each of these substances on the surface of particular cells either excites them, inhibits them or has no effect on them. Such findings have traditionally been interpreted in relation to the existence of synapses and the vesicle hypothesis, but they can also be interpreted simply without the necessity of belief in these structures. The experiments tell us quite clearly that some cells react to some naturally occurring substances in the brain, but not to others. Sometimes they are injected at rather high concentrations, since it is difficult to know precisely the concentrations at different distances away from the micropipettes, and it is also virtually impossible to know the relevant natural tissue concentration *in vivo*. Furthermore, the assumption has to be made that the current used to inject the transmitters does not itself have any significant effect on the permeability or excitability of the membranes of the cells being examined; this assumption has usually been tested.

Despite these reservations the abundant evidence of iontophoretic experiments tells us only that different cell membranes are sensitive at low concentrations to different naturally occurring substances. It does not prove that the loci of these sensitivities are the knobs on the surface of the neurons. Nevertheless, this does pose a considerable question. If the neurons are not connected at synapses, how is an action potential from one cell communicated to another? For the time being I must put forward the following hypothesis: the action potential in one cell body is conducted (in an electrical sense) through the syncytium in all directions, and decrements according to its space constant. The nearest membrane is excited if the current reaching it is sufficiently high. The sensitivity of the membrane of the 'receiving' cell mainly determines if the cell fires. The various neurotransmitters, chelating agents, labilizers, stabilizers, depolarizing agents and toxins, all determine the sensitivity to incoming current of particular cell membranes. It should be noted that it may be difficult to devise experiments which could distinguish unequivocally between the direct effect of a substance on the membrane and a 'modulating' effect on its sensitivity. For example, a neuron may fire because it is depolarized by acetylcholine, made hypoxic by diminishing its oxygen supply, 'labilized' by a low concentration of calcium ions in its immediate environment, or deprived of substrate during dying.

The scarcity of cogent evidence for the existence of structural synapses induces one not to accept their reality in life. I have therefore felt obliged to propose an alternative explanation to the widely held transmitter and vesicle hypotheses, which are partly based on microscopical findings. I believe that I have brought sufficient evidence against the synapses that they should be regarded as artifacts unless new data be produced about them. Logically, of course, if the hypothesis which I have outlined briefly above should prove to be wrong, this would in no way invalidate the evidence that the synapses are artifacts. Rather, it would oblige me to propose another hypothesis for transmission which would also not require the existence of synapses.

PROBLEMS ARISING FROM THE LATTER CONCLUSIONS

It should be made clear that the term 'synapse' is being used here in the sense of a nerve-nerve connection, and not as a neuromuscular junction.

I would propose that excitability in the nervous system spreads across the syncytium by electrical conduction. This suggestion is not difficult to accept since it is evident that when electrical activity is recorded with extracellular electrodes, such as fine metal microelectrodes, or coarser electrodes used in electrocorticography, or in electroencephalography, or in class experiments recording action potentials in sciatic nerves, excitability has passed through the bulk nervous tissue, without necessarily involving synapses. Electrical conduction could provide a starting point for an attempt to synthesize a general theory, involving nerve conduction (Hodgkin, 1951), the transmitter hypothesis (Katz, 1969) and the chemical hypothesis of nerve conduction (Nachmansohn and Neumann, 1975). Such a synthesis would have the effect of making the hypotheses much less incompatible.

In summary, then, having shown difficulties of accepting the existence of synapses, I wish only to initiate discussion to encourage physiologists, biochemists, pharmacologists and biophysicists to re-examine their views on transmission in so far as they are linked in their minds with the current view of the structures they call synapses.

THE NATURE OF THE STRUCTURES DESCRIBED AS SYNAPSES SEEN BY LIGHT AND BY ELECTRON MICROSCOPY

We have shown that the granules seen when examining unfixed neurons (Figure 54) are very likely to be intracellular, and, therefore, they are probably mitochondria (page 148). We have further shown that the granules seen on the surfaces of cell somas in stained sections of spinal cord and brain are not connected with any preterminal fibres, teloglia, telodendroglia, dendrites, fibrils or cell bodies (pages 141-144). Their dimensions and their incidence of occurrence as seen by light microscopy hardly overlap the same parameters seen by electron microscopy (Tables 22-25). The stains used in histology are generally silver salts, which are reduced, react with, and precipitate proteins and lipids. This gives a clue as to the nature of the appearance of granules described as synapses. They are most likely to be silver precipitates on tissue, bearing in mind that the silver salts are reduced and the silver-tissue reaction product is dehydrated subsequently. That this explanation is likely is demonstrated by the following experiment. A smear of egg albumin was stained with Weil and Davenport's stain, and another with haematoxylin and eosin. It is quite clear that all sorts of artifactual appearances similar to granules, fibres and fibrils can result (Figure 10).

The electron microscopic image of the synaptic cleft appears as two parallel lines far too frequently than solid geometry would permit, and is, therefore, a two-dimensional deposit (Figure 53, page 145). Please see references in Tables 21-26.

The most likely explanation for both the pre- and the post-synaptic thickenings is as follows. A certain proportion of granules of osmium salts deposits in the region of the membrane of the neuron. When the chamber of the electron microscope is evacuated and the specimen is subjected to electron bombardment, X-radiation and high

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

temperature, the underlying membranes evaporate beneath the osmium granule and crack it.

However, irrespective of the latter unproven explanation, I would like to stress that the inspired hypothesis of synaptic chemical transmission could never be proved unequivocally with respect to mammalian neurons, since one cannot localize the tip of a relatively large iontophoretic micropipette sufficiently accurately to be certain that the transmitter was actually being applied to a single very small synapse (pages 153-154). It should also be noted that - whereas in autonomic ganglia one can distinguish between the pre- and the post-synaptic fibres in mammals - it is also difficult to know for certain the precise localization of applied drugs in this sense as well. The use of the terms pre- and post-synaptic is often based on pharmacological criteria alone.

'UNIT' MEMBRANES

The 'trilaminar' appearance of various cell membranes can be seen in any electron micrograph and all textbook diagrams, except our own (Hillman and Sartory, 1980). We have shown the trilaminar appearance - as opposed to the cell membrane - to be an artifact, and we would proffer the following explanation for its genesis. Since it is two-dimensional, and the fixatives and heavy metals are added to the tissue before section, it follows that the 'trilaminar' appearance must have arisen after the sections were cut. The focal plane of the electron microscope is very deep, but the image of the membrane is usually only on the surface (*Ibid*, page 41). Therefore, the 'trilaminar' appearance must arise on the surface as a consequence of the exposure of the heavy metal covering the membrane to the vacuum, heat and irradiation. This must cause the membrane beneath the heavy metal to expand, vaporize and probably explode, leaving a groove lined by heavy metal on both sides (Figure 45). Please see also Appendices 4 and 5.

CYTOSKELETON OR ENDOSKELETON

We have already concluded that all the elements described by this term (Table 17) are artifacts - that is, they do not exist as structures in living cells (for detailed discussion, please see Hillman and Sartory, 1980, pages 44-57). In addition to the geometrical objections to its existence in life, we have pointed to the several different kinds of movements within living cells, which would be incompatible with any network or large body within the cytoplasm.

Cytologists have accepted that there is a problem here, and there is a considerable literature on cell motility. They see the endoskeleton by electron and light microscopy (Allen and Kamiya, 1964; Goldman, Pollard and Rosenbaum, 1976a,b,c; Dustin, 1978; Roberts and Hyams, 1979; De Brabander and De May, 1980). The ability to extract actin and myosin from many cells has led to the suggestion that they can cause the movements, as actin and myosin are claimed to do in muscle. This has been regarded as the answer to the question as to how the cytoskeleton would permit intracellular movement. This view is based on the following evidence: firstly, the ability to extract actin and myosin chemically; secondly, the appearance by electron microscopy of largely two-dimensional strands said to be of the expected

PROBLEMS ARISING FROM THE LATTER CONCLUSIONS

size of actin and myosin molecules; thirdly, the antigen reactions which are reported to occur only after cells have been fixed and dehydrated; fourthly, the strands visible by light microscopy (page 131). All of this evidence together does not prove the hypotheses either that the fibres do consist of actin or myosin, nor that if they are of this chemical nature, they do indeed cause intracellular movements. At this point I would like to reiterate my view that we cannot be agnostic on this or other hypotheses (page 4); either there is enough cogent evidence to prove the matter beyond reasonable doubt - or we cannot accept it. The fact that an increasing number of people believe the hypotheses encourages us to examine the evidence carefully, but it is not itself evidence of the validity of the hypotheses. The validity of a viewpoint cannot be measured by the number of its adherents.

Perhaps the hiatuses in the thinking should be spelled out precisely. The apparent similarity of dimensions of some fibrils to those of actin and myosin is quite insufficient proof that these fibrils are of this chemical nature. The demonstration of the presence of markers on fixed, dehydrated and sectioned tissue - often with added fluorochromes - is quite insufficient evidence of true specific localization of antigens unless and until the antigen-antibody reaction has been shown to occur in unfixed, fully hydrated sections. The failure to carry out or to publish the latter experiments so far should only serve as a stimulus for anyone claiming localization by these techniques to proceed with these definitive experiments without delay; they should not continue using quite incomplete experiments distant from natural conditions, as sources for substantive findings. Thus, I would suggest that if these reactions claiming to show localizations cannot be demonstrated in fresh tissue to which no fixatives or alcohols - which might reasonably be expected to alter antigen-antibody reactions - have been added, then we should cease believing in these results, rather than abstain from doing the relevant crucial experiments or accepting hypotheses with their major assumptions unproved.

There are further reasons which make it unlikely that contraction of acto-myosin filaments or of other contractile proteins could cause intracellular movements, like streaming, Brownian movement, pinocytosis, phagocytosis, mitochondrial movements, etc.

Firstly, if, for example, mitochondria were to be moved by acto-myosin filaments, the filaments would have to be attached not only at the mitochondrial end, but also to the cell membrane, or to the nuclear membrane, in the direction of which the mitochondrion was about to move. However, contraction of a filament at the east end of a cell would stretch the opposite filament at the west end of the cell. After a finite contraction the filament at the east end would be at its shortest length and the one at the west end would break. That is to say, a one-step movement could occur, but not a continuous movement, like streaming or pinocytosis, nor any reversible movement.

Furthermore, we would ask for evidence that: the filaments are attached to another firm structure; they are not randomly orientated; they are seen both in the contracted and in the relaxed state; and, of course, that they are composed of actin and myosin.

The weakness of the suggestion that acto-myosin filaments could initiate or

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

accelerate movements of mitochondria, Golgi bodies or lysosomes, can be illustrated by an analogy. If a crowd is trying to move along in a packed closed underground train, under no circumstances can the introduction into the carriage of any persons, animals, machinery or appliances, increase the rate at which the passengers could change places in the compartment. It could only slow down their movement.

It is also noteworthy that during the last century there were many publications on 'colpoids' which were mixtures of salts, oils, granules and coloured reagents, whose microscopic properties imitated those of cells (Herrera, 1869-1942; Butschli, 1894; Hardy, 1899; Bastian, 1913). Suitable mixtures showed streaming, Brownian movement, laminar flow, and coalescence of globules. (Nowadays, the latter would probably be called 'endocytosis'.) Of course, the colpoids possessed no metabolic source of energy, acto-myosin filaments or 'cytoskeleton'.

We have concluded that the whole 'cytoskeleton' is a precipitate of the cytoplasmic constituents, with heavy metals, after having been bombarded by electrons under conditions of very low pressure. Unpalatable as it may be, one must insist that direct evidence of intracellular movement can never be derived from examination of fixed, dehydrated, frozen or dead tissues, or of electron micrographs.

One reason for the difficulty in relating movements to chemical phenomena arises from the use of the word 'movement' to cover: intracellular movements; movements of whole cells such as phagocytosis; putting out pseudopodia; flagellar movement; muscle contraction.

If the ribosomes and lysosomes do not exist in the living cells, where does protein synthesis and acid hydrolysis occur in the living cell? We have shown that the biochemical compartmentation is unknowable by current destructive techniques (Hillman, 1972). Protein synthesis and acid hydrolysis occur in virtually all cells, yet ribosomes and lysosomes are not seen as structures in all tissues. (For example, they do not appear in muscle or kidney cells.) If one admits that either the biochemical activity can be present in cells which do not appear to have the granules at all, or that it may be present in compartments which are neither ribosomes nor lysosomes, one finds oneself in the difficult position of describing 'free ribosomes' or 'free lysosomal activity'. The implication of this is that one is admitting that these particles are not necessary to house the biochemical activities, or the biochemical activities must be in particles one cannot identify. The first of these alternatives admits that the biochemical activities do not require these locations; the second involves one in the difficult logical position of attempting to use supposition as evidence.

Of course, denial of the existence of the cytoskeleton in no way means denial of the existence of the biochemical activities alleged to be in it, although this should not be taken as implying that we accept that the locations of the activities found by subcellular fractionation techniques are true representations of their locations in intact living tissues.

THE INTEGRITY OF THE BRAIN

We have concluded that the brain is composed of neurons with their axons and dendrites within an embracing fluid syncytium. I should like to stress that we regard the neurons

PROBLEMS ARISING FROM THE LATTER CONCLUSIONS

and all their processes to be in fixed positions. The rest of the central nervous system, which is the majority, is a fairly viscous syncytium in which the naked nuclei may move and they may act as phagocytes. Neurosurgeons taking biopsies invariably find that the texture of the living brain and spinal cord is extremely soft.

The question then is raised as to what the central nervous system owes its mechanical integrity. The tissue itself is probably bound together by the axons, the processes, the blood vessels and the capillaries, as concrete is reinforced by steel rods. On a larger scale, the processes are seen as commissures, tracts, and projections. The ventricular system and the central canal is lined by an ependymal epithelium internally (Figures 6 and 77), and by the meninges externally. The pia mater, of course, embraces the central nervous system most intimately. I would suggest that the system has sufficiently strong covering and internal reinforcement to account for its mechanical integrity, without requiring all the cells to adhere to each other.

THE CENTRAL NERVOUS SYSTEM AS A SYNCYTUM

There have been three different concepts of the syncytium (pages 11-13). The reticularists, like Gerlach (1872) and Golgi (1883), believed that all the neurons were connected by a continuous fine meshwork of fibrils. Bauer (1953) had a different view of the syncytium - that the fibrils within the neurons connected through the membrane to dendrites outside in the syncytium (Figure 7). A third view - to which we adhere - is that the neuroglial tissue between the neurons only is a syncytium; this is illustrated in our generalized diagram of the central nervous system (Figure 78).

In denying the existence of connections to synapses, we have implicitly ruled out the first possibility (pages 136-146). Our conclusion that there are no intracellular filaments in the cytoplasm other than mitochondria would require us to deny the possibility of any connections between extracellular or syncytial fibres, on the one hand, and any intracellular fibres, on the other; obviously, we would hold it as extremely unlikely that dendrites are attached to mitochondria, when the latter are moving continuously in life. Furthermore, under the microscope, one simply does not observe such fibres passing through the neuronal membrane. Our explanation for the belief in such an apparently continuous syncytium is that the mitochondria are seen in the syncytium between the neurons, and are also seen in the neuronal cytoplasm; a small indulgence in unconscious extrapolation (Cook, 1983) has joined them together in the minds of some observers.

EVIDENCE OTHER THAN STANDARD TRANSMISSION ELECTRON MICROSCOPY

We have denied the existence in life of a wide range of structures, all - except the Golgi apparatus - originally seen by electron microscopy (Table 17). Horne and Harris (1981) talk about the 'enormous range of structures which have been meticulously investigated by independent measures', and Michell, Finean and Coleman (1982) stress particularly that freezing techniques are productive sources of independent data. A few comments might be apposite here:

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

1. Biochemical evidence cannot be used in support of, or to deny, morphological findings. They are different epistemologically, often different in dimensions by several orders, and their interpretation has usually been based on certain conceptual models of the structure.
2. Low-angle diffraction studies are usually made on partially or completely dehydrated tissues - except for the studies on frog peripheral nerves (Finean, 1954, 1960). Obviously, unless one attempts to examine, calculate and correct for, the distortions and shrinkages due to the reagents, the dehydrations during preparation, and the effects of sectioning and the electron bombardment, conclusions about shapes and sizes of organelles or the distances between them must be taken magno cum grano salis. Nevertheless - as we have stressed frequently - the desire to derive meaningful conclusions obliges one to attempt to quantitate the latter changes in every tissue whose microscopic structure one wishes to elucidate. Furthermore, it is not good enough to pay lip service to the known or suspected changes during tissue preparation, without using them to correct all one's measurements and calculations (Hillman, 1983).
3. Freezing procedures used with electron microscopy are said to provide independent evidence for beliefs in some of the structures seen by standard transmission electron microscopy. This view is not tenable because:
 - (a) Like biochemical experiments, the results of experiments involving freezing are interpreted in relation to beliefs derived from the earlier techniques.
 - (b) Freezing is only used to fix tissues, which are subsequently stained with heavy metals, dehydrated and subjected to vacuum and electron bombardment, just as it is with other techniques of electron microscopy (Stolinski and Breathnach, 1975; Rash and Hudson, 1979; Sleytr and Robards, 1982). Since we have concluded that the artifacts of electron microscopy originated after fixation of the tissue, in our view, evidence derived by the application of freezing techniques could only be regarded as independent of that of standard electron microscopy, if the two techniques did not share the same steps during which the artifacts may arise.
 - (c) Nowadays, it is considered that tissue should be frozen as rapidly as possible to minimize the size of ice crystals. Unfortunately, the greater the rapidity of cooling, the less time the bubbles of gas coming out of solution at the lower temperatures have to escape from the tissue.
 - (d) Although it is widely believed that vitrification occurs when cooling is very rapid, specific investigations of the simple systems of water and dilute aqueous solutions have demonstrated how difficult it is to achieve (Bruggeller and Mayer, 1980). The belief implied here is that if vitrification occurs, the tissue is not dehydrated or deformed. The technique called 'critical-point drying' is widely used in the belief that it preserves the structure of tissues. The term is misleading, since at the critical point, the solid, the liquid and the vapour phases are in equilibrium, and no drying would occur. Drying takes place only when one moves away from the critical point by raising the temperature or lowering the pressure. It is intended that sublimation of the

PROBLEMS ARISING FROM THE LATTER CONCLUSIONS

ice to vapour will avoid the surface tension and electrostatic effects in the tissue, which may distort the cellular structure. However, I am not aware of any studies of specific cells before and after the application of freezing techniques, showing that shrinkage and deformation do not occur. Sleytr and Robards have made very important contributions to the study of artifacts due to freezing techniques (Sleytr and Robards, 1977a,b, 1982). In the latter review they have listed 11 main agents during freeze-fracture, which have been shown to cause 30 different kinds of artifact during various procedures. They continue to use the techniques although they concluded that 'it is possible to assert that early hopes that freeze-fracture and etching techniques might provide artifact-free images because chemical fixation can be replaced by (undamaging) cryofixation have not been fulfilled....'

It is now abundantly clear that each of the factors involved in the formation of the final image can either enhance, or compensate for, each other.'

- (e) Since the images produced by freezing techniques appear as two-dimensional as those by the earlier techniques, those who practise them must produce their own explanations for these geometrical difficulties. They cannot use the shared problems as evidence that the problems do not exist.

GLIOSIS

The reaction of central nervous tissue to infection has a cellular and a fibrillary component. The cell bodies may have little or no cytoplasm and pathologists call them variously, reactive astrocytes, oligodendroglia, microglia, phagocytes or histiocytes. The pathologists assign the names partly on account of their knowledge of the clinical conditions which afflicted the patients (Zulch, 1979). It is generally agreed that the cells can be mobile. They may arise acutely by migration from other parts of the central nervous system, by division locally, or by diapedesis from blood vessels (for excellent modern reviews, see Jacobson, 1978; Oehmichen, 1978; Korr, 1980). We would characterize these cells on the basis of their lack of cytoplasm or obvious processes, as 'naked nuclei'.

The increase of 'protoplasmic' or 'fibrillary' astrocytes, with processes which occur as a consequence of inflammation (see, for example, Glees, 1955; Windle, 1958; Nakai, 1963; Greenfield, 1976; Treip, 1978) raises a difficulty for us, since we have put forward the hypothesis that, in normal central nervous systems, all cells with processes are neurons. If, then, these cells occur in normal tissue, they must be neurons. If they only occur in ill tissue - of which most slides and micrographs appear to be representations - we must ask if neurons can proliferate in these conditions. We cannot answer this question at the moment but would list the kinds of ways fibrils can arise in living tissues, or precipitates imitating them can appear as a result of the staining procedures:

1. Sprouting of 'neurites' has been shown to occur in cultures of sympathetic and sensory ganglia if nerve growth factor is added to them (Levi-Montalcini, Meyer and Hamburger, 1954; Symposium, 1964). In young rats, neurons isolated from the medulla, as well as from sympathetic and sensory ganglia, put out these fine

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

processes, and the addition of nerve growth factor increased the frequency with which they did (Hillman and Sheikh, 1968; Hillman and Khallawan, 1970). These fibrils did not develop into axons or dendrites in our cultures, and such fibrils are seen in cultures of growing cells, irrespective of the tissues from which they originate.

2. Fibrils have been shown to grow out of the proximal end of a transected peripheral axon, when it begins to regrow towards the distal end (Weddell and Glees, 1941; Guth, 1956).
3. If the protoplasmic or fibrillary astrocyte were really a neuron, and if it had been demonstrated rigorously that the number of processes increased due to inflammation, then one could suppose that real new dendrites could grow out of the cell somas in response to pathological conditions.
4. Inspissated pus may appear as fibres after precipitation by drying, and after staining - as egg albumin may do (Figure 10).
5. The fibrils could be a staining artifact arising from a change in local chemistry of the central nervous system, which would make the silver-tissue complexes less soluble, and thus precipitate out in rows. Although it may be difficult in micrographs to distinguish between precipitates and dendrites, no such difficulty presents itself when one looks at the stained histology slide using oil-immersion objectives, and can follow them through several focal planes. Precipitates appear two-dimensional, whereas dendrites appear in three dimensions.
6. 'Non-specific' fibrils are found in many tissues, such as fibrous tissue, mesenteries, ligaments and tendons. There seems to be no *prima facie* reason why they should not occur in the connective tissue of the central nervous system, nor why they should not increase following inflammation, especially if it is chronic.

We should reserve judgement upon which is likely to be the true explanation for the apparent increase in fibrils after inflammation, until it has been shown positively: that the number of fibrils does increase; that astrocytes are not neurons; that the affinity and specificity of the staining procedures has not changed significantly as a consequence of the pathological conditions; that clots containing fibrin or dried inspissated pus can be differentiated histologically from dendrites or glial fibres; that one can distinguish histologically between dendrites and neuroglial fibres.

11

Properties of the Nervous System in the Light of Present Conclusions

CLASSIFICATION OF CELL TYPES, NORMAL AND PATHOLOGICAL

We have characterized all those cells which have membranes and processes projecting from them as neurons; these include cells currently designated as neurons and astrocytes. Cells without membranes around each nucleus and without nucleoli are naked nuclei, which are sometimes called oligodendrocytes, astrocytes or microglia. Capillary cells and ependymal cells are also present in the central nervous system. The simpler classification would solve the major problems of disagreements about their characteristic structures by light microscopy, by electron microscopy, and in tissue culture (Tables 2-11). It would also solve the problem of the multiplicity of views about the specific or recommended staining systems appropriate to each kind of cell (Table 10). It would clearly imply the necessity to carry out control experiments on all staining systems used in pathology and histology, in an attempt to define the real specificity of each staining system, and whether there are morphologically recognizable sub-populations of neurons and neuroglia.

Different populations of cells are, indeed, defined by their immuno-chemical properties. Some neurons show up after complex histochemical procedures intended to indicate the presence of, say, catecholamines. As I have indicated previously (Hillman, 1972, pages 41-53; Hillman, 1983; and in this volume, pages 198-199). I do not think that one can accept that substances - mostly transmitters - are necessarily found in the same locations after fixation, embedding, sectioning and processing as they were in the living tissues. However, since the cells in a particular region after such processing do indicate different chemical and microscopical properties, which are constant (Adams, 1965; Ungerstedt, 1971; Pearse, 1972, 1980; Moore and Bloom, 1978, 1979; Pickel, 1981), one must conclude that there are, indeed, different biochemical sub-populations.

It is curious that, in general, histochemists find that most transmitters are located on neurons. This is probably because they believe that the transmitters nearly always arise in or near neurons, therefore any cells which apparently show the presence of a widely accepted or putative transmitter by that token define themselves as neurons. Alternatively, histochemists may have carried out careful examinations of cells they believe to be neuroglia and concluded that such cells independently identified do not

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

contain the transmitters. However, gamma-aminobutyric acid, adenosine, acetylcholine, and dopamine in rather few systems have been reported in neuroglial cells (Table 12).

Whereas the classification of cell types may be considered an academic exercise in histology, it has an important relevance in neuropathology. There are various classifications of tumours: Bailey and Cushing (1926) listed 14 types of gliomas; Zulch (1965, 1979) listed 30 different brain tumours and other space-occupying lesions; the Unio Internationalis Contra Cancrum in 1958 (see Zulch, 1965, page 37) classified 30 different tumours mainly on the basis of tissue of origin (see also Kernohan and Sayre, 1952). The classifications take into account (1) clinical history of the patient, age of onset of the disease and rapidity of spread of the tumours; (2) whether the tumours arise in the more primitive cells, such as pineoblasts or astroblasts, or the later cells, the pineocytes or astrocytes; (3) whether they are believed to arise from neuroepithelium, mesoderm, ectoderm, the whole embryo or the vasculature; (4) their tissue of origin, such as the medulla, meninges or retina; (5) the cell type from which they are believed to arise, such as the astrocyte, oligodendrocyte or ependymal cell. Most authors are rather uncertain as to whether there are tumours in true adult neurons of the cerebrum, cerebellum or spinal cord. However, there is general agreement that in all tissues, tumours of differentiated cells and tissues are less malignant than those of undifferentiated cells; the degree of malignancy is assessed by the incidence of the latter cells, the degree of anaplasia, the frequency of mitotic figures, the amount of cellularity, the involvement of endothelial cells and the extent of necrosis (Kernohan and Sayre, 1952). I would suggest that the naked nuclei are the cells of origin of all undifferentiated tumours of the central nervous system; when they become pleiomorphic their precise identification becomes difficult, but the failure to see a membrane around the cytoplasm in histological sections of these tumours, and in the illustrations in all the publications cited, is really quite striking. Nishii (1929) characterized as sarcomas the undifferentiated tumours in the brain which contained cells reminiscent of our naked nuclei, but this idea is not generally accepted. However, the high grade of malignancy of undifferentiated cells, their invasiveness, the uncertainty about from which kind of glial cell types particular tumours originate, and the similarity of these undifferentiated nerve cells, all taken together, encourage one to attempt to simplify the cellular classification of tumours into two classes only - those showing malignant cells with fibres or processes, and those showing only naked nuclei. There are two possibilities about the origin of the cells in these tumours. Firstly, the cells with fibres are tumours of neurons and the others are tumours of naked nuclei. A second hypothesis, which I favour slightly more, is that primary tumours of differentiated adult neurons are extremely rare or do not exist at all, and that all tumours arise from naked nuclei; the fibres of the first kind in this case would not be dendrites or axons. The latter explanation seems more likely since (1) there is a general disinclination to identify, in adult animals, tumours originating from neurons in the central nervous system, as opposed to their common occurrence in the peripheral neurons (Russell and Rubinstein, 1977, page 260; Treip, 1978); (2) there do not appear to be reports of mitoses having been seen in neurons of adult mammals; (3) I have rarely seen any cell in a tumour, which had recognizably originated from a neuron,

PROPERTIES OF THE NERVOUS SYSTEM IN THE LIGHT OF PRESENT CONCLUSIONS

and, for example, had the normal complement of processes attached to its soma, but had clear signs of being malignant. Russell and Rubinstein (1977, pages 260-269) devote a section to ganglioneuromas and gangliogliomas, and indicate that such tumours contain considerable glial elements. However, with the exception of one bizarre ganglioglioma with gemistocytic astrocytes (their Figure 202, page 263), most of the neurons themselves have appearances which are not grossly different from those found in normal tissue. Cavanaugh (1970) fixed cerebral tissues rapidly after he made lesions in the brain with needles, and he was able to observe many mitoses in reactive astrocytes around the lesions. Thus, we know that such cells (naked nuclei) may divide in adult animals. This would make them obvious candidates for malignancy. Their location in a syncytium, and their ability to move within it, may explain the relatively rapid spread and unclear boundaries of these malignant tumours of the central nervous system (Figure 80).




FIGURE 80 Human cerebral astrocytoma stained with haematoxylin and eosin. Please note that the nuclei have little or no adherent cytoplasm and no cell membrane. We would classify these as naked nuclei. The bar is 20 μ m

MEMBRANE TRANSPORT

This term is used to mean transport both across a membrane, and by a membrane. Transport, itself, is a portmanteau word, since it encompasses a number of different mechanisms characterized by their individual kinetic characteristics (Bayliss, 1959; Wilbrandt and Rosenberg, 1961; Stein, 1967; Harris, 1972). The close use of Occam's Razor conditions us to recall that the simplest form of transport is diffusion, and that substances distribute themselves in different phases according to their different affinities for all the chemical species in each of the phases (see, for example, Goldstein, Aronow and Kalman, 1968). Therefore, unless one shows that any movement occurs at a significantly different rate than diffusion, one must assume that this simplest mechanism alone is operative. Indeed, in order to stop it, a tissue or reaction mixture would have to be cooled to absolute zero temperature, although it is drastically slowed if the

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

whole system is solidified.

In order to demonstrate that a movement of molecules is due to 'active' transport or is dependent upon metabolism, it is usual to show that subjection of the tissue to hypoxia or poisons, or deprivation of substrate, causes a massive redistribution of the components of the system in the direction of the apparent electrochemical gradients. Unfortunately, we are using crude measures here, since none of these experiments demonstrates that metabolism is directly involved. For example, the effects seen may result from increased lactate or lowered hydrogen ion concentration, and may not be a simple consequence of interference with metabolism. Indeed, it is very difficult to design experiments which can show unequivocally that the rates of particular reactions or movements require specifically biological mechanisms. Even the measurements of temperature dependence or oxygen uptake may show clearly simultaneous changes, whose relationship is not necessarily causative. Perhaps we are peering over the battlements at the joust between the vitalists and the mechanists, whose outcome it is probably several thousand years too early to predict.

One must distinguish between the evidence for the membrane as an interface between two different compartments, and the evidence that the mechanisms which regulate the composition of the phases on either side of the membrane are located within it. It is both a widespread finding and a reasonable expectation that any thin membrane - biological or artificial - permits different small molecules to cross it at different rates (Mueller and Rudin, 1968; Chang, 1972), so that the simplest model of the biological membrane is one composed of curious chemistry, which could account for most of the differences in chemical composition between - on the one side - the syncytium or extracellular fluid and - on the other - the cytoplasm.

I believe that cytologists should not fall into the trap of believing that because one can detect the difference of composition of the fluids on either side of a membrane, the mechanisms causing the electrochemical gradients are themselves located within the membranes (see, for example, Harris, 1972; Andreoli, Hoffman and Fanestil, 1978). (No-one believes that the heat for Montgolfier's airship was generated within the fabric of the balloon.) However, I believe that the gradual acceptance of this non-sequitur has conditioned cytologists to seek other evidence supporting, or compatible with, the location of the regulatory mechanism within the thickness of the membrane itself. The main sources of this belief are as follows:

1. Penetration of cells with intracellular electrodes results in the abrupt recording of a voltage.
2. Injection of adenosine triphosphate into poisoned squid and cuttlefish giant axons increases efflux of sodium ions (Hodgkin and Keynes, 1955; Caldwell, Hodgkin, Keynes and Shaw, 1960).
3. The finding of sodium/potassium ion activated ATPases in fractions believed to contain membranes (Skou, 1957, 1975).
4. The recording of resting and action potentials of the expected values in squid giant axons, whose axoplasm had been extruded and replaced by known potassium chloride solutions (Baker, Hodgkin and Shaw, 1962a,b).

PROPERTIES OF THE NERVOUS SYSTEM IN THE LIGHT OF PRESENT CONCLUSIONS

5. The finding of acetylcholine, choline acetylase and cholinesterase in high concentrations 'in all types of excitable tissues investigated, in so-called "cholinergic", and "adrenergic", in motor and sensory, in peripheral and central fibres, in invertebrates and vertebrates, and in all types of muscle fibres' (Nachmansohn and Neumann, 1975, page 253). (Please see also Schlaepfer and Torack, 1966.) Subsequently, the enzyme was located mostly in invertebrate species in neuromuscular junctions, which are usually considered to be similar to synapses (Nachmansohn and Neumann, 1975, pages 253-259).
6. Examination of the chemistry, electron microscopy and low angle diffraction, of both myelin sheaths and extracted myelin, as models of the properties of concentrated samples of membranes (Schmitt, Bear and Clark, 1935; Fernandez-Moran, 1952; Finean, 1961; Stoeckenius, 1962; Autilio, Norton and Terry, 1964; Landon, 1976).
7. The chemical properties, especially the enzymology, of 'microsomal' fractions, are usually assumed to reflect the properties mainly of cell membranes and endoplasmic reticulum (Table 12). The presence in these fractions of the enzymes believed to be involved in transport is regarded as evidence that these enzymes are located in the membranes around the cell. Unfortunately, one cannot accept without reserve evidence from subcellular fractionation at present because: (a) the effects of the preparative procedures on the distribution of the enzymes have never been tested (Hillman, 1972); (b) it has yet to be shown quantitatively and rigorously that the microsomal fraction mainly or wholly consists of cell membranes and endoplasmic reticulum; (c) we have shown that the endoplasmic reticulum does not exist in living cells (Hillman and Sartory, 1980, pages 44-57).
8. Kinetic studies of transport of ions, sugars and amino acids into mammalian red cell 'ghosts', ascites cells, yeast, mitochondria, frog skin, toad bladder, slices of liver and kidney, and isolated gut have been regarded as models for the transport across membranes (Harris, 1972). Red cell 'ghosts' have been prepared by immersion in hypotonic media, which permits the egress of haemoglobin, which is due to a gross modification of the normal permeability of the cell membranes. Frog skin, toad bladder and isolated gut each contain many layers composed of different cell types sandwiched by connective tissue. Their properties are of great interest in understanding the overall movements of small molecules across the whole tissues, as well as the potential differences recordable across their thicknesses, but it seems unlikely that they can be regarded as models of single neuronal membranes.

This is an incomplete list of studies which have been regarded as evidence that the membranes themselves contain the mechanisms of transport, as well as being the sites whose permeability may change during regulation. However, if one scrutinizes attentively the results of each of these various approaches, it seems that most of them have produced data which cannot distinguish between the mechanisms causing differences in chemical composition on the two sides of a membrane being located within the wall of the membrane, or within the cytoplasm. Subcellular fractionation itself could provide the answer, if control experiments had been done satisfactorily and comprehensively, and had, indeed,

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

shown the legitimacy of the relationship between the parent tissue and the subcellular offspring.

Thus, there are two schools of thought. Both regard the semipermeability of the membranes and any channels, charged or uncharged, to be one site of regulation; they agree that the semipermeability may be influenced by the chemistry of the fluids on both sides of the membrane. However, the difference between the two schools is that most membranologists believe that the site of the biochemical machinery of regulation is within the thickness of the membrane, while I would suggest that it is a property of both the permeability of the membrane at any particular time, and of the biochemistry of the cytoplasm, the syncytium, the mitochondrioplasm and the nucleoplasm.

It seems to me quite inconceivable that the cell membranes could house the whole arsenal of: structural lipids and proteins; carriers for ions, sugars and amino acids; transport enzymes (Figure 81); microsomal enzymes; receptors for approximately 50 transmitters (Table 27); receptors for innumerable drugs; receptors for hormones (Table 29); natural antigens and agglutinogens; receptors for bacterial and viral toxins; surface antibodies to a host of bacterial and viral diseases. The molecular diameters of these substances would vary from 0.4 nm to 20 nm, and many of them are found in microsomal fractions in considerable quantity.

A few comments about each of these groups of substances are apposite.

The structural lipids and proteins are believed to be the main components that occupy the thickness of the membrane. This view was enunciated before the use of the electron microscope, as a result of studies on red cells. Calculations were made which showed that the quantity of lipids and proteins which could be extracted was enough to provide the two lipid molecules sandwiched between the two protein molecules, which is now known as the Davson-Danielli model (Gorter and Grendel, 1925; Davson and Danielli, 1942). This model was subsequently generalized to all membranes. Yet the original calculations were made for red cells, which are not claimed to have an endoplasmic reticulum. As far as I can find, the calculation has never been made for the membranes of any other mammalian cells. One probable reason for this is that it would be extremely difficult to calculate the areas of membranes of other cells. Furthermore, there seems no *prima facie* reason for believing that there is little or no lipid or protein elsewhere in cells. Yet even in the red cells the interesting calculation that there would be enough lipid to make a bilayer does not prove that the membranes in life are a bilayer, nor, of course, that they are largely composed of lipids and proteins. That being said, it seems likely membranes do contain these components, although I would suggest that more analytical and comprehensive experiments are needed to confirm the beliefs in respect of red cells, to assess them quantitatively in membranes and to permit one to generalize them to all mammalian tissues, as well as other animals and plants. If one accepts our view that the lamellae of the myelin sheath peripheral nerves are artifacts, this would cast into greater doubt the apparent congruence of the results of experiments on red cell ghosts and on extracted myelin as evidence supporting the current view of membrane structure.

The membrane is said to contain carriers. The belief in their existence arises from kinetic considerations involving separation of the region of entry and exit of particles (Osterhout, 1935; Widdas, 1952; Wilbrandt and Rosenberg, 1961; Stein, 1967,




FIGURE 81 Schematic representation of the structure of the Na^+-K^+ transporting ATPase in the membrane. This figure is reproduced by kind permission of the author and publishers, Repke, K.R.H. (1980) Structural basis of the compartmentation functions of biomembranes. In Cell Compartmentation and Metabolic Channeling, ed. Nover, L., Lynen, F., and Mothes, K. (Leipzig: Fischer Verlag and Amsterdam: Elsevier), page 37. Please note that in this model, this enzyme molecule protrudes 50 Å from the surface of the membrane

pages 148-206). In nervous tissue their existence has been inferred from kinetic studies on sodium current in giant axons (Hodgkin and Huxley, 1952a,b) and of entry of monosaccharides and amino acids into brain *in vivo* and into cerebral slices (Crone, 1961, 1965; Fishman, 1964; Le Fevre and Peters, 1966; Gilbert, 1965; Blasberg and Lajtha, 1966; Joanny, Corriol and Hillman, 1969; Joanny, Barbosa, Hillman and Corriol, 1971). Yet, the binding of a substance carried to the carrier would considerably enlarge its size, so it seems highly unlikely that it could aid transport through pores. Although carriers have always been assumed to be located in membranes, such experiments are not evidence that they are, indeed, located there.

Microsomal enzymes are legion, and have been best characterized in liver (Dixon and Webb, 1964). Since the microsomal fraction is said to consist of membranes of both the cell and the endoplasmic reticulum - presumably in different proportions in different preparations - and both membranes appear to be trilaminar and narrow on electron microscopy, they both share accommodation problems for the enzymes.

Transmitters are believed to act by combining with 'specific' receptors. Several of the criteria for receptors, such as saturability, specificity and reversibility, are also met by 'non-specific' binding of non-biological materials (Cuatrecasas, Hollenberg, Chang and Bennett, 1975; Cuatrecasas and Hollenberg, 1975).

The acetylcholine receptor has been isolated from the Torpedo electric organ, and has a molecular weight of between 250 000 and 550 000 (O'Brien, Eldefrawl and Eldefrawl, 1972; Moore, Holladay, Purett and Brady, 1974). If other neurotransmitters are of this size there would certainly be great difficulty in accommodating 'specific' receptors to each of them. The problem is gradually being diminished as the older belief in one neuron or one synapse for each transmitter is giving ground to the view that several transmitters, and therefore, presumably their 'specific' receptors, can

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

coexist in one soma or one synapse. More than one receptor has been isolated for nearly every transmitter so far examined, such as acetylcholine, adrenaline and dopamine. (Vizi and Magyar, 1984).

Three important points are worth making here. Firstly, the properties of apparent 'specificity' of binding by talc, alumina powder, microsilica, glass and cellulose acetate filters by Cuatrecasas and his colleagues (1975) bear a very important lesson - namely that 'specific' receptors are not necessary either as biological or as chemical entities. In other words, any biological mixture with its rich pot-pourri of chemicals and bonds represents a range of attractions to such a diverse population as precursors, hormones and transmitters. Secondly, we are not convinced either that 'specific' receptors exist as structures or as chemical compounds, and so we are not surprised that they are not seen. Thirdly, there is currently no consensus about whether every transmitter has one or more 'specific' receptors.

Virtually all the comments about receptors for transmitters apply equally to drug receptors. Indeed, both types of alleged chemicals are discussed together in pharmacology textbooks. The only difference is that whereas it is conceivable that cell membranes would have 'specific' receptors for neurotransmitters which occur naturally, it is extremely unlikely that they would have 'specific' receptors for any drug to which they may have been exposed for the first time, especially if it is rapidly acting. Furthermore, one must raise the question as to whether any individual really has one or more 'specific' receptors for every single effective drug in any pharmacopoeia past, present, forgotten, future and undiscovered.

The membranes are also supposed to be covered by a rich outcrop of natural antigens and agglutinogens, as well as receptors to bacterial toxins and to viruses, and antibodies to both of the latter. The antibodies for common exanthemata take a period of time of the order of days to be synthesized by the body, and may circulate in the blood as well as remain on the cell surfaces. Their effective lives in circulation vary from the very few days for the viruses for cold or influenza to the years for diphtheria and poliomyelitis.

When one assembles all these compounds which are believed to reside in or near the membrane together (Table 29), one would be led to the expectation that electron microscopy would show the surfaces of membranes to be populated by a rich tundra of mounds of a large range of sizes.

There would be little disagreement that in freeze-fracture-etch scanning electron microscopy, one usually sees an outcrop of knobs of uniform size. Nowadays these are usually attributed to the proteins of the Singer-Nicholson model (Figure 13), but where are all the other presumably protein receptors? Why can they not all be seen on scanning electron microscopy? It should be noted, en passant, that there is no direct or unequivocal evidence identifying these knobs as being due to the structural proteins such as phosphoproteins, glycoproteins or any 'extrinsic' or 'intrinsic' proteins (Guidotti, 1972; Singer, 1974) or any other specific molecules. When the molecular dimensions apparently identify a particular chemical species, the calculations have not usually taken into account the hydration of the chemical and the dehydration during preparation for electron microscopy. However, even assuming that the identification were correct, it would account for only one size of surface molecules of all those

PROPERTIES OF THE NERVOUS SYSTEM IN THE LIGHT OF PRESENT CONCLUSIONS

TABLE 29 Large molecules reported to be present within, or attached to, the membranes of mammalian neural tissue

Substance	Tissue	Molecular weight (d) or diameter (nm)	Key references
<u>Proteins</u>			
Phosphatidopeptides	Microsomal fraction of mammalian cerebral myelin		Le Baron and Folch (1956)
Encephalitogenic basic protein	Guinea-pig central nervous system		Kier, Murphy and Alford (1961)
Myelin protein	Rat sciatic nerve	8.5 nm	Adams and Bayliss (1968)
Wolfgang protein	Ox sciatic myelin		Wolfgram (1966); Eng, Chao, Gerstl, Pratt and Tavastjerna (1968)
Proteolipid apoprotein	Ox and guinea-pig cerebral myelin	34 000 d	Thorun and Mehl (1968)
Basic protein	Man, ox, rabbit, guinea-pig cerebral myelin	25 000 d	Thorun and Mehl (1968) Mehl and Halaris (1970)
Proteolipid protein	Man, ox, rabbit, guinea-pig cerebral myelin	35 000 d	Mehl and Halaris (1970)
Rhodopsin	Mammalian retina	7.5 nm x 3.5 nm, 37 000 d	Wu and Stryer (1972) Lewis, Krieg and Kirk (1974)
Periodic-acid Schiff positive protein	Microsomal fraction of rat cerebral myelin	110 000 d	Quarles, Everly and Brady (1973)
S-100	Mammalian nervous system	19 5000 d	Calissano and Bangham (1971)
<u>Enzymes</u>			
NAD nucleotidase	Microsomal fraction of vertebrate brain	36 000 d	Jacobson and Kaplan (1957)
Cholinesterase	Microsomal fraction of vertebrate brain	315 000 d	Aldridge and Johnston (1959)
Arylesterase	Microsomal fraction of vertebrate brain	223 000 d	Aldridge and Johnston (1959)
NADPH ₂ -cytochrome c reductase	Microsomal fraction of vertebrate brain	80 000 d	Guiditta and Strecker (1959)

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

Substance	Tissue		Molecular weight (d) or diameter (nm)	Key references
Alkaline phosphatase	Microsomal fraction of vertebrate brain	120 000 d		La Bella and Brown (1959)
Alkaline protease	Microsomal fraction of vertebrate brain			La Bella and Brown (1959)
Glutamine synthetase	Microsomal fraction of vertebrate brain	450 000–525,000 d		Sellinger and de Balbian Verster (1962)
Na^+ , K^+ , Mg^{2+} activated ATPase	Microsomal fraction of ox and guinea-pig brain	360 000 d 259 000 d 250 000 d 8–10 nm		Schwartz, Bachelder and McIlwain (1962) Hosie (1965) Tanaka and Strickland (1965) Maunsbach, Skriver and Jorgensen (1979)
	Synaptosomes of ox brain	280 000 d or 8.5 nm		Uesugi, Dulak, Dixon, Hexum, Dahl Perdue and Hokin (1971)
Ca^{2+} activated ATPase	Microsomal fraction of mammalian brain	102 000 d or 9.0 nm		McLennan, Seeman, Iles and Yip (1971)
Acetylcholinesterase	Microsomal fraction of guinea-pig Rat sciatic nerve	280 000 d		Schwartz, Bachelder and McIlwain (1962) Hosie (1965) Kasa (1971)
Neutral proteinase	Myelin in vertebrate central nervous system			Marks and Lajtha (1963)
Leucine aminopeptidase	Rabbit sciatic nerve	78 000–80 000 d		Adams, Davison and Gregson (1963)
Phosphatidoinositol kinase	Microsomal fraction of rat brain			Banik and Davison (1969)
ADPase	Microsomal fraction of vertebrate brain Rabbit brain			Colodzin and Kennedy (1964) Goldfischer, Essner and Novikoff (1964)
NADH-dehydrogenase Peptidase	Rat sciatic nerve			Adams (1965)
	Myelin of microsomal fraction of mammalian brain	340 000 d		Adams and Davison (1965)
	Human nervous system	340 000 d		Hallpike (1972)
$2'3'$ cyclic nucleotide 3'-phosphohydrolase	Microsomal fraction of brain	68 000 d		Kurihara and Tsukada (1967)

PROPERTIES OF THE NERVOUS SYSTEM IN THE LIGHT OF PRESENT CONCLUSIONS

Substance	Tissue	Molecular weight (d) or diameter (nm)	Key references
5' nucleotidase	Mouse brain	150 000 d	Scott (1967)
Choline acetylase	Rat sciatic nerve	68 000 d	Adams and Bayliss (1968)
Succinate dehydrogenase	Microsomal fraction of rat brain	97 000 d	Banik and Davison (1969)
Phospho-diesterase	Mammalian brain		Loten and Sneyd (1970)
Inorganic pyrophosphatase	Human nervous system	63 000-120 000 d	Hallpike (1971)
Cholesterol ester hydrolase	Myelin of microsomal fraction of rat brain	110 000 d	Eto and Suzuki (1973)
Guanylate cyclase	Mammalian brain		Quarles, Everly and Brady (1973)
Cytochrome c oxidase	Microsomal fraction of rat brain	200 000 d, 6.7 nm	Illiano, Tell, Siegel and Cuatrecasas (1973)
NADH cytochrome c reductase	Microsomal fraction of rat brain	75 000 d	Norton and Poduslo (1973)
N-acetylglucosaminyl-transferase	Ox brain myelin		Grey and Chan (1980)
Mannosylphosphoryl-transferase	Ox brain myelin		Norton and Poduslo (1973)
Adenosine 3'-phosphate, 5' sulphatophosphate-galactosyldiacylglycerol sulphotransferase	Microsomal fraction of rat brain	440 000 d	Pieringer, Subbha Rao, Mandel and Pieringer (1977)
Glyceraldehyde 3-phosphate dehydrogenase	Microsomal fraction of human brain	6.9 nm	Solomon (1977)
Monophosphoglycerate mutase	Microsomal fraction of human brain	5.0 nm	Solomon (1977)
Phosphoglycerate kinase	Microsomal fraction of human brain	4.76 nm	Solomon (1977)
UDP-galactose-ceramide galactosyltransferase	Rat brain		Constantino-Ceccarini, Castelli and De Vries (1979)

Substance	Tissue		Molecular weight (d) or diameter (nm)	Key references
Cerebroside sulpho-transferase	Microsomal fraction of rat brain			Koul and Jungalwala (1981)
UDP-galactose: ceramide galactosyltransferase	Microsomal fraction of rat brain			Koul and Jungalwala (1981)
<u>Nucleotide</u>				
RNA	Ribosomes of rodent brains	10 000 - 2 000 000 d		Jacob, Samec, Stevenin and Mandel (1967)

The Table does not include receptors for all known transmitters, antibodies, toxins and drugs, as well as those undiscovered, those to be discovered in the future, those forgotten and those buried in the literature. This list could be lengthened in proportion to the time spent on a literature search. Some of the values of molecular weights or dimensions were not necessarily derived from the substance in the same tissues as cited; many are from Dartnall and Klotz (1976). One would expect that in view of the dimensions of the molecules listed, they should be seen by freezing, scanning and transmission electron microscopy as a large outcrop of knobs of a vast range of heights, reflecting the heterogeneity of their dimensions and configurations

PROPERTIES OF THE NERVOUS SYSTEM IN THE LIGHT OF PRESENT CONCLUSIONS

listed in Table 29.

The situation is one molecule worse when one considers transmission electron microscopy. There the image of Robertson's trilaminar membrane is ubiquitous around every cell, nucleus and mitochondrion, and in the endoplasmic reticulum, cristae and Golgi membranes. It appears 6-8 nm thick. There is not even space for one protein molecule in addition to the protein layers enclosing the two molecules of lipid of Davson and Danielli's sandwich. Why does one not see Singer's mobile proteins on transmission electron microscopy?

One attempt has been made in the plant literature to explain this difficulty. Muhlethaler, Moor and Szarkowski (1965) concluded that the appearance of Robertson's unit membrane is due to uncoiling and flattening of protein molecules during fixation.

The question should be pressed as to how the vast majority of 'specific' carriers, receptors and antibodies have evaded capture and identification by the biochemists. Probably William of Occam would have explored the simpler hypothesis that one would expect the complex mixture that is biological tissue to exhibit a vast range of affinities for ions, small molecules, precursors, foreign proteins and drugs, to which they had not been previously exposed, without even extra machinery for doing so. We would thus imagine a hierarchy of affinities of the tissue from weak to strong, as follows:

- diffusion
- affinity for any cellular elements
- involvement in biosynthesis
- binding to tissues reversibly
- binding to tissues irreversibly
- having a pharmacological effect
- inducing enzymes
- producing antibodies.

It should be noted that there is no implication here that the affinity has to be for the cell membrane exclusively or any other organelle or its contents.

It would be important that all these affinities should be measured *in vivo*, or at least in isolated organs; it would be reasonable to expect that fixation, freezing and homogenization, would each affect all these kinds of affinity, and therefore affinities should only be measured in fresh tissue. In discussion, when I have questioned the belief that all these large molecules are located within the membrane (Table 29), some of my colleagues have conceded that the molecules may not actually be located within its thickness, but rather on the inside or the outside wall of the membrane. This too is unacceptable, for a similar reason as is the Singer-Nicholson model (page 132); if the molecules believed to be on either side of the membrane are each individually large enough to be detected by transmission electron microscopy, they should all be shown up by that technique.

Let us now harvest the crop of hypotheses about cell membranes, and attempt to assess them in relationship to each other.

It seems extremely unlikely that all the molecules listed in Table 29 could coexist

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

in the membrane together without being visible by electron microscopy. The following are possible explanations for failure to see them there.

1. They are washed off during subcellular fractionation and/or preparation for electron microscopy. This is unlikely because both these techniques are used to prepare, and often to identify, the proteins, enzymes, receptors etc.
2. There are many fewer (or no) such 'specific' proteins in or near the membrane. We believe that this is very likely because the total number of such 'specific' substances already shown, or believed to be capable of demonstration, or capable of being generated, is so large. It is worth recalling that the idea that each transmitter and drug has its own 'specific' receptor, and that it cannot have an action until it has combined with such a receptor, is a general hypothesis which should be tested in every particular case. However, if the binding were unstable, it might be impossible to isolate them and characterize them chemically, although many other unstable compounds, such as adenosine triphosphate or creatine phosphate have been extracted.
3. The localizations of some or all of these molecules on the membrane are wrong. It has been stressed repeatedly (page 48) that the localization of any molecule in a subcellular fraction believed to consist mainly of membranes cannot itself be regarded as certain proof that the particular molecule is in the same site in the intact tissue. The relevant and necessary experiments have not yet been done of the effects of the preparative techniques in subcellular fractionation, histo-chemistry and electron microscopy on the distribution of molecules (Hillman, 1972, 1983). There seems to be considerable reluctance to carry out these overdue but crucial experiments. Therefore, at present, we should not accept the alleged localizations of these substances listed as being present in the cell membranes.
4. It is likely that the usual thickness of the membrane as measured from electron micrographs is a gross underestimate due to shrinkage. It is possible that the real thickness is several times more in life than it appears from the measurements in red cells by the leptoscope (Gorter and Grendel, 1925; Fricke, 1925; Waugh and Schmitt, 1940), in fresh and fixed peripheral nerves by low angle diffraction and polarized light microscopy (Fernandez-Moran and Finean, 1957; Finean, 1960, 1961; Fernandez-Moran, 1962) and in dehydrated neural tissues by electron microscopy (for references, please see Sunderland, 1976, page 10; Watson, 1976, pages 102-114). If one knew the real thickness of the membrane, one might well then decide if it was, indeed, possible for it to accommodate the cohorts of molecules, which are believed to reside there.
5. It would be possible that only the enzymes and carriers believed to be involved in ion and other transport are lodged in the membrane. If this could be shown by non-disruptive procedures using reagents which one would not expect themselves to alter permeability, one would have advanced a long way on the road to showing that the membrane itself could regulate. In the classical experiments of Baker, Hodgkin and Shaw (1962a,b), they showed that placing squid axons between solutions of potassium sulphate, potassium chloride, sodium sulphate, sodium chloride or sea water, in equal concentrations on either side, produced a potential

PROPERTIES OF THE NERVOUS SYSTEM IN THE LIGHT OF PRESENT CONCLUSIONS

difference across the axon of less than 1 millivolt. If the transport system were indeed within the axon, it should have started excluding the sodium ion and concentrating the potassium ion, as does the frog skin. The only reason for it not to have done so is that the prolonged perfusion may have washed the transport enzymes out of the axon wall, or energy supplied by axoplasm is 'required' for the sodium ion pump. It would seem easier to suggest that the biochemical mechanisms which cause the asymmetrical distribution of the cations in life are located in the axoplasm. In experiments, if one washed them out, one then has to provide the chemical gradient to produce the potential difference, or the potential difference to redistribute the ions.

Hillman (1966), in reviewing these experiments, suggested that the effect of the biochemical activity in the axoplasm was to maintain the semipermeability and the excitability of the membrane.

Frog skin is frequently used as a model of active transport of sodium ions (Ussing, 1949). When immersed in water with very low concentrations of these ions, it actively moves these from a low to a high concentration. I am not aware of any such demonstration for squid and cuttlefish axons, although red cells do exclude sodium ions and exclude potassium ions against the apparent electrochemical gradients (Danowski, 1941; Harris, 1941; Flynn and Maizels, 1949; Whittam, 1958). However, these would be in a relatively natural environment in experimental conditions.

6. It is possible that the molecules are indeed present in the cell membrane, but that they do not stain by transmission electron microscopy. Nearly all protein molecules of the molecular weight of dimensions indicated in Table 29 would be within the resolution of the electron microscope, which has, of course, been used for looking at individual molecules (Cold Spring Harbor Symposium, 1972; Hayat, 1973-1977; Wisse, Daems, Molenaar and Van Duijn, 1974; Baumeister, 1978; Baumeister and Vogell, 1980).

If the regulation of the ions is not carried out by the cell membrane itself, it is possible that it is due to the cytoplasm having a high affinity for potassium ions; that is, acting as a cation exchange for sodium ions. This view was put forward as the sorption hypothesis (Shaw, Simon, Johnstone and Holman, 1956; Troshin, 1961; Kurella, 1961; Ling, 1962, 1984).

Conway (1957), citing mainly studies on frog muscle, strongly attacked the latter view on the following grounds:

- (a) the virtually complete exchangeability of potassium ions in frog muscle was indicated by influx and efflux measurements in frog muscle (Noonan, Fenn and Haege, 1941; Keynes, 1954) and in rat diaphragm (Calkins, Taylor and Hastings, 1954);
- (b) carboxylic exchange resins with fixed anionic charges should prefer potassium ions, as should powdered keratin, but they did not (Conway, 1955);
- (c) there was inadequate protein to account for ion binding;
- (d) if potassium ions, carnosine, phosphate esters and phosphocreatine are all absorbed on to protein in the cell, how could it be isosmolar with the plasma. It was better explained if they were not;

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

- (e) Ling's hypothesis (1952, 1955), did not explain the Donnan relation of potassium and chloride ions;
- (f) the latter did not explain the agreement found between the potential difference across the cell membrane and the potassium ion gradient;
- (g) it did not explain the accumulation of potassium and chloride ions when they were increased in the external fluid, nor the water entrance when these ions were accumulated from isotonic mixtures;
- (h) Conway (1957) disputed Ling's assertion that $^{24}\text{Na}^+$ entered fibres freely at 0°C when the potassium ion concentration was at or above threshold, and that metabolic inhibitors did not affect the high potassium, low sodium ion content of fibres.

Conway (1957) regarded the central fallacy of Ling as being the assumption that the smaller potassium ion could enter a zone of reduced dielectric around fixed ionic charges on protein chains farther than could the sodium ion, and therefore accumulate more in the muscle fibres. He believed that Ling underestimated the radii of the hydrated potassium and sodium ions. He also noted that Ling's calculations of radius of reduced dielectric around the ions were not as accurate as Ling believed.

Glynn (1957) added evidence from red cells against ion binding in the cytoplasm:

- (a) the distribution ratios of chloride, bicarbonate and hydroxyl ions were the same on both sides of the membrane, indicating that they would have to be bound to the same extent;
- (b) ion binding would explain the high concentration of potassium ions in cells, but not the low sodium ion concentration unless much of the cell water was bound;
- (c) freezing point determination showed osmotic equilibrium between red cells and plasma, indicating that major electrolytes must be free in solution;
- (d) Fricke and Morse (1925) had found that the specific resistance of the cell interior was 3.5 times that of the serum;
- (e) the sorption hypothesis could not explain why sodium ion efflux was increased when potassium ion concentration outside the cell was increased;
- (f) haemoglobin did not bind sodium ions;
- (g) Hodgkin and Keynes (1953) had found that $^{42}\text{K}^+$ moved inside squid axons, and Harris (1954) had found that they moved in frog muscle at the same rate as they did in extracellular fluids.

If these findings - mainly from red cells, frog muscle, and squid and cuttlefish axons - are shown to be applicable in details to mammalian neurons, then one would have to conclude that the cytoplasm could not act as a cation exchanger.

However, if they are not, this still leaves two possible mechanisms which could explain the asymmetry of the distribution of ions and small soluble molecules on either side of a membrane.

The first one has already been mentioned (page 219). It is that the biochemical reactions going on in the cytoplasm could determine the nature of the semipermeability of the cell membrane. The permeability would also reflect sooner or later changes in the

PROPERTIES OF THE NERVOUS SYSTEM IN THE LIGHT OF PRESENT CONCLUSIONS

chemical constitution, and chemical reactions going on at a particular time, in the syncytium, the mitochondrioplasm and the pars amorpha of the nucleolus.

Secondly, one should not view the idea of the cytoplasm as a cation exchange resin too narrowly. Cytoplasm is a rich mixture of chemical species and bonds - much more complex chemically even than a mixture of a few pure resins. Thus, one can think of the fluxes and concentrations of all the ions and small molecules as being totally interdependent; the concentration of each of them is determined continuously by the complete environment in which it finds itself at that instant.

The two possible mechanisms may coexist, but the first seems more likely because it would react much more quickly, and it is known that action potentials are generated and ion fluxes change rapidly. However, it should be stressed that the extremely significant experiments of Conway on muscle, Glynn on red cells, and Keynes, Hodgkin and Huxley, and all their collaborators on giant axons, show that ionic regulation occurs across cell membranes, and do not provide any evidence that the mechanism itself for the regulation is located within the cell membrane.

The pioneer work on thin films by Bangham and his colleagues, and on artificial membranes by Mueller and Rudin, have indicated the properties which may be found in artificial membranes, and may be common to biological membranes (Bangham and Horne, 1964; Mueller and Rudin, 1968; Chang, 1972; Finkelstein, 1978). The electrical properties of mixtures of natural lipids and proteins - to which antibiotics have sometimes been added - have been found to be remarkably similar to those of cell membranes examined with intracellular microelectrodes (for comparison, please see Henn and Thompson, 1969). The only constant and significant difference found was that the natural membranes had higher resistances, by two or three orders, than did the artificial membranes. Such elegant experiments clearly demonstrate what a rich variety of behaviour such 'simple' membranes have, and they should act as a constant reminder of what chemistry can do without the aid of biology.

THE EXTRACELLULAR SPACE

The extracellular fluid of the central nervous system was first measured by the use of markers, such as Na^+ , Cl^- , thiocyanate, sucrose, sulphate, inulin and ruthenium red. The use of such markers implied the assumptions that they do not enter the cells, that they do not react with the tissue, that they are not broken down, that they themselves do not affect the size of the extracellular space, that they are recovered rapidly and completely and that they themselves do not affect the biochemistry of the tissue. Na^+ and Cl^- are present inside cells; the uptake of thiocyanate depends upon the plasma level (Lajtha, 1957; Streicher, 1961). Inulin drains away into the cerebrospinal fluid (Woodward, Reed and Woodbury, 1967); it leaves tissue very slowly (Varon and McIlwain, 1961), as does radiosulphate (Woodbury, Timiras, Koch and Ballard, 1956); the efflux of inulin can be resolved into three kinetic curves, only one of which is likely to be due to passage through the extracellular space (Lund-Andersen, 1974; Lund-Andersen and Kjeldsen, 1976). Of the assumptions listed for all these markers, the beliefs that they remain extracellular and that they do not react with tissue, are both quite crucial and have rarely, if ever, been tested. The inulin concentration in the brain of a dog after it had been perfused, went up after

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

death (Rall, Oppelt and Patlak, 1962). The values of the spaces measured by these markers were from 2% to 40% in the brains of mammals under various conditions (Table 30).

Several authors have measured similar spaces in cerebral slices (Allen, 1955; Davson and Spaziani, 1959; Pappius and Elliott, 1956; Bourke and Tower, 1966a,b; Joanny and Hillman, 1963, 1964; Schousboe and Hertz, 1971; Cohen, 1974a,b; Lund-Andersen, 1974; Rodnight, 1975; Lund-Andersen and Kjeldsen, 1976). Unfortunately, the use of these preparations *in vitro* implies all the assumptions about the markers listed above. It is also complicated by the increase of weight of slices on incubation, which was usually 15-40% in the experiments of the authors cited above. This increase consisted of 'adherent' fluid and swelling fluid. The exact value of the change of volume depends upon whether initially the slice had been cut dry and weighed, or had been temporarily immersed in a Petri dish of cold incubating medium before weighing. It also depends upon the length of time for which adherent fluid is wiped off the slice (for review, please see Hillman and Wraae, 1981). For these reasons, measurements of extracellular space by markers are even less reliable *in vitro* than they are in the whole animal.

Wyckoff and Young (1954, 1956) were the first to try to assess the extracellular space by electron microscopy. They perfused spinal cords with osmium tetroxide, and concluded that there was no large extracellular space. This question was subsequently taken up by many groups including those around Luse, Gray, Horstmann, Palay, Sjostrand and van Harreveld (for reviews see van Harreveld, 1966, pages 127-158; Johnston and Roots, 1972, pages 76-92). They all saw the extracellular spaces of 10-25 nm wide, which could generally be calculated to be less than 10% of the volume. The actual values found depend upon: the species, the tissue, the age of the animal, the fixative used and its osmolarity, the pressure and temperature of the fixative, whether the tissue was frozen rapidly or slowly, and assumptions about the areas of the cell membranes and the likely numbers of cells in the tissue. However, in their calculations, authors have not taken into account the shrinkage of tissue during preparation for electron microscopy (Boyde, Bailey, Jones and Tamarin, 1977; Sleytr and Robards, 1977a,b; 1981), probably because they had no accurate way of doing so. Furthermore, in examining a particular minute sample of the central nervous system composed of cells of irregular shapes, assessments of the total areas of the membranes in that region - and, *a fortiori*, in the whole nervous system - are extremely difficult.

The volume of the extracellular space has also been calculated from impedance measurements, mainly by van Harreveld and his school (1966) (please see page 186).

Regrettably, one has to conclude that there is at the moment no satisfactory way of measuring the extracellular space. In order to do so one would have to find a marker, which would have to be shown minimally, to remain extracellular, not to react with tissues, which could be extracted completely from them, and whose measurement would be unaffected by the presence of the tissue.

However, the conclusion here that most of the central nervous system is a syncytium largely circumvents the problems of the extracellular space. The fluid bathing

PROPERTIES OF THE NERVOUS SYSTEM IN THE LIGHT OF PRESENT CONCLUSIONS

TABLE 30 Early measurements of the extracellular space in intact animals by markers

Marker	Author	Percentage
Na^+ , Cl^-	Manery and Hastings (1939)	33-40
$^{35}\text{SO}_4$	Woodbury, Timiras, Koch and Ballard (1956)	4-5
Cl^-	Woodbury, Timiras, Koch and Ballard (1956)	25
Inulin	Morrison (1959)	2
$^{35}\text{SO}_4$	Barlow, Domek, Goldberg and Roth (1961)	2-5
Thiocyanate	Streicher (1961)	5
Inulin	Rall, Oppelt and Patlak (1962)	12
$^{35}\text{SO}_4$	Kibler, O'Neill and Robin (1964)	2-6
^{14}C , sucrose, inulin and thiocyanate	Bourke, Greenberg and Tower (1965)	19-40
Inulin	Vernadakis and Woodbury (1965)	5
$^{131}\text{Iodine}$	Davson (1967)	20
Sucrose	Oldendorf and Davson (1967)	10

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

the neurons and the naked nuclei is the syncytial jelly, which acts as the extracellular environment of the neurons, the naked nuclei, the axons, the fibres, the capillary cells and the ependymal cells. However it is, in fact, an intracellular fluid. If we now go back to Virchow's concept of the neuroglia as a nerve-glue, we may regard the neuron-glia relationship in a new light, illuminated with the new attitude. The all-pervasive large volume of the syncytium may be seen as acting as a barrier between the ventricles and the capillaries on the one hand, and the neurons and naked nuclei on the other. Thus the blood-brain barrier is located partly in the syncytium and partly in the choroid plexus.

It has been repeatedly stressed that histological and electron microscopic techniques cannot be used for measuring the volumes of any tissues, which are composed in life of 60-80% water. Therefore, although our histological measurements indicate that most of the volume of the central nervous system is occupied by the syncytium, they can not give us precise assessment of the relative volumes of the syncytium, the neurons, the naked nuclei, the fibres, the ependymal cells, and the capillary cells. At present there appears to be no way of measuring any of these volumes in living people or animals, although computed axial tomography and ultrasound can measure total tissue and ventricular volumes in the living intact nervous system.

CEREBRAL OEDEMA

In the living patient or animal a measured rise of intraventricular pressure is the clearest evidence of cerebral oedema; the pressure of the cerebrospinal fluid may also rise, and papilloedema may also be seen in the fundi. Cerebral oedema may be observed at operation or at post mortem. At a cellular level it must be very hazardous to diagnose oedema in either histological or electron microscopic sections, since oedema represents the state of hydration, and embedded sections have been completely dehydrated. However, chronic oedema has been produced by tumours (Williams and Lennox, 1939; Chou, 1961; Aleu, Edelman, Katzman and Scheinberg, 1964), by cold (Hass and Taylor, 1953; Klatzo, Piraux and Laskowski, 1958; Clasen, Cooke, Tyler and Pandolfi, 1960; Bakay and Hague, 1964; Blidner and Markham, 1965), poisoning by triethyl-tin (Torack, Terry and Zimmerman, 1959; Aleu, Katzman and Terry, 1963), and several other pathological and experimental agents. Studies of swelling in the brain by electron microscopy show that the main swelling appears to occur in the astrocytes (see, for example, Gerschenfield, Wald, Zadunaisky and de Robertis, 1959; Torack, Terry and Zimmerman, 1959, 1960; Luse, 1961). Our view of the astrocytes is that their nuclei are naked nuclei, and their feet are syncytial cytoplasm (page 194). If the location of oedema in the brain is truly in the 'astrocytes', we would place it in the syncytium.

It is currently believed that membranes regulate the ions inside cells, not only by their selective permeability but by 'active transport'. Living neurons *in vivo* and *in vitro* maintain resting transmembrane potential differences, which can be altered by changing the K^+ , oxygen tension and substrate, in the medium bathing them (for reviews, see Eccles, 1957; Hillman and Hyden, 1965). Since a syncytium would contain

PROPERTIES OF THE NERVOUS SYSTEM IN THE LIGHT OF PRESENT CONCLUSIONS

no barriers in its cytoplasm, it could not easily regulate the electrolytes between one part and another. Furthermore, all agents would act on the syncytium before they reached the neurons. Therefore, oedema would be more likely to affect the syncytium before it became manifest in the neurons.

Nevertheless, it is confidently expected that in the future non-invasive radiographic techniques will indicate in what elements of the central nervous system oedema occurs in living intact animals. It may well be that oedema causes the syncytium to take up less stain than healthy tissue does. This is a plausible alternative explanation to the current view that astrocytes take up the oedema fluid - a finding which one might have expected the light microscopists to have detected before or since the electron microscopists did. Our explanation for the failure of the light microscopists to do so is firstly, that the astrocytes are difficult to identify by histological techniques, and secondly, the two kinds of microscopists are both looking at different structures when they use the term 'astrocyte'.

BLOOD-BRAIN BARRIER

There has been a long controversy as to whether the plasma or the cerebrospinal fluid is the extracellular fluid of the nervous system (van Harreveld, 1966; Davson, 1967). The identification of the syncytium as the immediate environment of the neurons diminishes the power of the controversy. Substances may travel across the blood to the neurons by two routes: either the substances in the blood in the cerebral and spinal capillaries pass through the syncytium to the neurons, or they proceed to the choroid plexus to the ventricles then cross the ependyma to the syncytium, and then reach the neurons. Obviously, the first is much quicker, and is probably the route taken by rapidly acting intravenous anaesthetics, by volatile anaesthetics, indeed by all intravenously administered drugs which act on the central nervous system soon after the few seconds that the circulation brings it to its site of action. Both of the two routes would constitute a blood-brain barrier, although the slower one is traditionally thought of as the 'blood-brain barrier' that elapses between the arrival of the substances in the blood and their penetration into the brain.

THE SYNCYTUM AND THE EXTRACELLULAR FLUID

We have concluded that the neuroglia is a syncytium in the sense of Virchow (1854, 1871). This facilitates the understanding or explanation of several other phenomena:

1. The longstanding problem of whether the blood or the cerebrospinal fluid is the effective extracellular fluid of neurons is resolved, if the neuroglial syncytium is the environment of neurons throughout the central nervous system.
2. The tissue between neurons and microglia (naked nuclei) is very resistant to stains used in neurohistology.
3. Reactive glia (naked nuclei) migrate very rapidly to sites of injury because they do not have to pass between a solid phalanx of cells.
4. The relative rapidity of spread of undifferentiated gliomas, which are also very malignant. They are naked nuclei and can migrate without difficulty through the frontier-less syncytium.

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

5. Sherrington (1925) proposed that the interaction of reflex pathways and the long duration of central excitation could best be explained by the concept of a neuronal pool; this was involved in occlusion and spatial summation. The central excitatory and inhibitory states represented a prolonged residue in the central nervous system resulting from the much briefer excitatory and inhibitory stimuli. I would suggest that the stimulus engenders 'slow' changes in the chemistry of the syncytium, which could be the site of the central excitatory and inhibitory states in the subliminal pool. This would provide a very adequate explanation for the neurophysiological phenomena.
6. The finding that the intellect and neurophysiological function of some patients with advanced hydrocephalus are not necessarily impaired (Lorber, 1980; Lorber and Priestly, 1981) would be understandable if 80-90% of the brain is a syncytial jelly, and the neurons are not connected by synapses. The hydrocephalus compresses the brain, but the neurons would be displaced (Figure 82). This suggestion assumes that thought takes place in the cerebral cortex; which seems very likely, although it has not been proved conclusively. Of course, there is abundant evidence that many neurophysiological functions reside there.

THE VESICLE HYPOTHESIS OF TRANSMISSION

From the early ideas of transmitters and some seminal experiments of Fatt and Katz (1952) on frog nerve-muscle preparations, del Castillo and Katz (1956) proposed the vesicle hypothesis for neuromuscular transmission (for summary, please see Katz, 1969). It is interesting to note that in the latter publication 13 years after the original concept, he wrote 'I should like to introduce a working hypothesis which my colleagues and I have found useful in planning our experiments. We assume that each unit packet of acetylcholine (whose release produces a miniature potential) is pre-formed within a synaptic vesicle in the nerve terminal. The vesicle may be supposed to accumulate the transmitter substance actively.'

This short section contains the following hypotheses: (a) the synaptic vesicles exist in life; (b) each contains acetylcholine; (c) each contains approximately equal quantities of acetylcholine. Item (a) cannot be proved; (b) has not been shown; (c) indirect evidence compatible with it has been brought forward, but there is no direct way of showing it.

This hypothesis has gradually been accepted as being applicable to central nervous system synapses, and also as being substantial supportive evidence for interpreting electron microscopic findings about synapses and synaptic vesicles. The criteria for transmitters have been enumerated frequently (see, for example, Hillarp, Fuxe and Dahlstrom, 1966; Werman, 1966; Barchas, Akil, Elliott, Holman and Watson, 1978).

The modern concept of transmission has been synthesized by combining evidence from the physiology of the neuromuscular junction and of the synapse, the pharmacology of the transmitters and the junctional tissues, the electron microscopy of neuromuscular junctions and synapses, and the electron microscopy and biochemistry of synaptosomes.

The large accumulation of data of varying degrees of cogency which each appear to reinforce the overall view could be dangerous if any large component contributing to




FIGURE 82 Computed axial tomogram of a young man with an intelligence quotient of 126, who had a first-class honours degree in mathematics, economics and computer studies. It shows gross hydrocephalus and an extremely narrow cerebral cortex. Published in World Medicine (1980) 15, page 21, and reproduced by kind permission of Professor John Lorber of Sheffield Royal Infirmary and the publishers

the resulting consensus should not be firmly based.

The vesicle hypothesis has been seriously questioned. Marchbanks (1975, 1978) has pointed out that synaptic clefts are rarely seen fused with the axolemma, as the hypothesis would require; depletion by prolonged stimulation is difficult to produce; the acetylcholine has not been shown to be present exclusively in the vesicles; newly synthesized acetylcholine is preferentially released before the 'stored' acetylcholine; the specific activity of the whole preparation is different from that of the isolated vesicles; it has been estimated that a 'quantum' of acetylcholine is seven times the quantity estimated to be present in the vesicle; isolated synaptic vesicles exchange acetylcholine slowly.

The incorporation of the synaptic vesicle into the pre-synaptic membrane is crucial to the vesicle hypothesis. Preparations of fractions of what are believed by the authors to be the two membranes, and examination of their chemistry, has revealed gross differences between the preparations with respect to lipids, ATPases, acetylcholinesterase, ganglioside and cholesterol, and turnover of proteins (Eichberg, Whittaker and Dawson, 1964; Hosie, 1965; Whittaker, 1970; Rodriguez de Lores Arnaiz, Alberici de Canal and De Robertis, 1971). One has to be extremely cautious in

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

comparing the chemistry of any subcellular fractions when they have been prepared using grossly different reagents and physical conditions, but the differences between these two fractions seem to be so gross that it is unlikely to be explicable solely on these grounds.

In addition, attempts to demonstrate that 'vesicle depletion' occurs by 'stimulating' preparations using potassium chloride, electrical stimulation and bungarotoxin, have been made in cat brain and superior cervical ganglion, rat superior cervical ganglion, rat and mouse neuromuscular junction, frog brain and frog neuromuscular junction. Results have varied from no change to complete depletion; sometimes the depletion has been accompanied by mitochondrial changes, sometimes the vesicles have become more or less coated, and often the changes have not been reversible (for review see Ceccarelli and Hurlbut, 1980).

Further implications of the vesicle hypothesis, as summarized by Katz (1969) do not seem to have been recognized. Firstly, no-one has shown directly that acetylcholine is present in the synaptic vesicles *in vivo* or *in vitro*, and it is difficult to conceive of experiments which could demonstrate the proposed location. Secondly, the vesicles are said to discharge the acetylcholine only when they happen to hit the pre-synaptic membrane, but not any other part of the inner membrane of the synapse. This implies that the following minimum number of events occur:

1. The vesicle hits the pre-synaptic membrane.
2. The vesicle differentiates between the inner surface of the pre-synaptic thickening and other parts of the inner surface of the membrane of the synapse; that is, the chemicals of the membrane around the synaptic vesicle must react chemically or electrostatically with constituents of the inner surface of the pre-synaptic membrane. Therefore the chemistry of the inner surface of the pre-synaptic and of the rest of the inner surface of the synaptic membrane must be different.
3. The vesicle releases the transmitter.
4. The transmitter reacts with the inner surface of the pre-synaptic membrane.
5. The membrane of the synaptic vesicle dissolves in the axoplasm or loses its ability to take up stain or is incorporated into the pre-synaptic thickening. Any of these events would occur in circumstances in which synaptic 'depletion' occurs. However, if the synaptic vesicles are believed to lose acetylcholine, which does not itself show up in electron micrographs, then the ejection of transmitter from the synaptic vesicles would not be expected to result in any change in number of vesicles, but would result either in them appearing at high magnification to have holes on their pre-synaptic aspects or to be part of 'exocytotic vesicles' in the pre-synaptic membrane. The disingenuous suggestion has been made that the synaptic vesicles pass through the pre-synaptic membrane so rapidly that they are not seen (Heuser and Reese, 1973; references given by Jones, 1975, page 192, not definitely agreed by him). Some workers have claimed to see 'pores' in the pre-synaptic membranes (Akert, Pfenninger, Sandri and Moor, 1972; Streit, Akert, Sandri, Livingstone and Moor, 1972). Others rather rarely see exocytotic vesicles (Andres, 1964; Westrum, 1965). I believe that they arise from the proximity of any

PROPERTIES OF THE NERVOUS SYSTEM IN THE LIGHT OF PRESENT CONCLUSIONS

two granules at the time of deposition of the heavy metal, which is used to stain. The deposit will not fill in the regions at which the two bodies are in contact and will appear to surround them both.

The transmitter would have to activate an extraordinarily rapid and powerful mechanism - presumably enzymatic - which could render the thick insoluble pre-synaptic membrane (consisting as they believe of protein and lipid), highly permeable to the transmitter, and very soon after that to 'switch off' that permeability. It is a central supposition of the vesicle hypothesis that the transmitter is 'packaged' in the vesicles. If the calculations and assumptions which show that there is, on average, seven times as much acetylcholine released in a quantum as could be contained in a single vesicle are true, then the majority of that transmitter must be located outside the vesicles (Marchbanks, 1978). The calculations and assumptions need to be examined critically.

6. The discharge of the transmitter would induce a change in the membrane which would cause a miniature end-plate potential. (It should be noted that, although a strong case has been made for a statistical relationship between excitation and the incidence of these miniature potentials, the proposition that the discharge of transmitter causes or is caused by the electrical transient remains a hypothesis, which is difficult to prove.) The proposed mechanism is believed to be occurring in living tissues, while the evidence is sought in metal deposits subjected to high vacuum, electron bombardment and radiation.

It should also be noted that the original relationships between the transmitter, the excitation and the miniature end-plate potentials in their physiological, pharmacological and biochemical aspects have been shown for acetylcholine and neuromuscular junctions. The evidence is rather sparse for any other transmitter or for the central nervous system. Most of the substances which have been claimed to be transmitters (Table 27) cause excitation or inhibition in neurons at low concentration, and other substances (mostly not naturally occurring) either increase or decrease the thresholds of excitation.

7. New synaptic vesicles would have to be made in the synapses, or they would become stainable again, or they would have to come from the cell body, or they would have to 'cycle' back from the synaptic cleft. When 'vesicles' are seen on either side of the synaptic membranes, and one calls them 'exocytotic' or 'endocytotic', there is no way of finding out in which direction they would have been moving if they were moving in life.
8. New transmitter has to be synthesized in the synaptic ending.
9. The transmitter has to be 'packaged', that is passed into the vesicle across the 'membrane' of the synaptic vesicle.

Considerable doubt about any calculations of the quantity of transmitter in the synaptic vesicles or any calculations based on them, or indeed the very existence in life of synaptic vesicles, must be entertained since the number of vesicles seen apparently depends upon the osmolarity of the buffer used in fixation (Bodian, 1970; Valdivia, 1971). (While these findings are of great significance empirically, it is difficult to know what role the buffering capacity can play, since the pH of the system

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

cannot be considered relevant if the tissue is to be dehydrated and impregnated with non-aqueous substances.) It has been shown that the pre- and post-synaptic thickenings are artifacts (please see page 145). A much simpler hypothesis has been proffered for the spread of excitation (please see page 197).

A GENERAL THEORY OF RECEPTORS

It is extremely unlikely that, in life, receptors for most transmitters, hormones, drugs, toxins and antibodies are found on the outside, within the 'trilaminar' membrane, or just on the inside of the cell membrane, for these reasons.

1. The cell membrane is probably solid and not trilaminar (Hillman and Sartory, 1980, page 37).
2. Many receptors for transmitters and drugs have been isolated, their molecular weights have been measured and they are physically large enough to be seen on the membranes by electron microscopy, but they are not seen by transmission electron microscopy at all; by scanning electron microscopy only, the population of protrusions from the membrane of uniform height could not possibly represent the rich repertoire of receptors of different molecular weights, dimensions and shapes claimed to be located there (page 210). If they cannot be seen because they do not stain with reagents used during the preparation for electron microscopy, then unstained pits or gaps should appear outside, within, or inside the membrane. These are not seen on scanning or transmission electron microscopy.

If one were to suppose that they were present, although they could not be seen, one would have to make the additional assumption that they had been lost during the preparation for electron microscopy. It would also be dangerous to accept the validity of the current hypothesis about receptors on the basis of such an assumption which has not been proved, and may be both unprovable and unfalsifiable.

3. It is extremely unlikely that a membrane supposed to be 7-10 nm thick could contain not merely all these receptors, but also all the carriers for all the substances which are believed to be taken up by cells by 'active' transport and by facilitated diffusion. There is plenty of evidence that sodium ions, potassium ions, calcium ions, glucose and amino acids, among others, are in continual flux between the outside and the inside of the cell. The multiple biochemical reagents believed to be involved in this flux are large molecules, and are believed to be located in or near the membrane. However, if the current belief about membrane transport were correct, the gradients of ions and other molecules would have to be maintained continuously, not merely in the region of the membrane, but also in the whole bulk of the cytoplasm. This would have to apply to most cells, including neurons of mammalian cranial nuclei, Mauthner neurons of goldfish, supramedullary neurons of puffer fish, and squid giant axons. It would be generally agreed that the cytoplasm has a huge volume relative to that of the whole cell membrane, especially if one accepts the view that the trilaminar membrane is, indeed, only 7-10 nm thick. This consideration makes the sorption hypothesis of Ling much more likely than the membrane theory (please see Ling, 1984).

PROPERTIES OF THE NERVOUS SYSTEM IN THE LIGHT OF PRESENT CONCLUSIONS

4. The fact that the addition of transmitters and many amino acids to the outside of the cell causes many neurons to be excited or inhibited, does not prove that the effect is produced in the membrane itself. It is known that these substances alter the excitability of the cells, but this could be due to any or all of the following: reaction with substances in the extracellular fluid; effects on the electrical properties of the membranes; effects on the chemical properties of the membranes; effects on the permeability of the membranes; alteration of gradients or fluxes across the membranes; effects on the cytoplasm. Although the effects of iontophoretic applications are fairly rapid, in my opinion, the temporal resolution of measuring systems currently in use is not sufficiently accurate to distinguish between which of these measurements contribute to the increase or decrease of activity, and to what extent each does. It is also highly likely that several of these effects would be interreactive, for example, that the biochemistry of the cytoplasm at any particular moment influences the properties of the cell membrane (Hillman, 1966b). Some of these possibilities could be explored in mammalian neural tissue, if one could obtain a pure neuronal membrane which one could be sure did not leak (Cummins and Hyden, 1962), or if one employed a patch clamp on such a preparation.
5. It is unlikely that rapidly acting drugs could generate receptors. If one accepts that all the receptors - if they occur in the profusion which modern pharmacologists and physiologists would have them - are unlikely to be housed comfortably in or around the membrane, one can then re-examine the concept of receptors in the nervous system. (Although this monograph is concerned only with the mammalian nervous system, the same considerations would apply to non-mammalian species, and also to tissues other than nervous systems, in which receptors are believed to play a role.)

I am not asserting that receptors may not occur on cell membranes, but that transmitters, hormones, drugs and toxins may act anywhere between the site of administration and the innermost sancta of the nervous system - the nucleolonema. We may share the common belief that drugs act near neuronal membranes. However, before they arrive there, they travel by a route with many obligatory and request stops.

1. Site of injection to plasma.
2. Plasma to plasma proteins, platelets, red or white cell membranes.
3. Blood cell membranes to their cytoplasm and organelles.
4. These membranes to capillary membranes.
- 5a. Capillary membranes to the cytoplasm and organelles of the capillary cells.
- 6a. Capillary cell membranes to the syncytium.
7. Syncytium to membranes of neurons or naked nuclei. Alternatively, they could go from 3 to 5b the membranes of the choroid plexus.
- 5b. Membranes of the choroid epithelium to their cytoplasm and organelles.
- 6b. Membranes of the cells of the choroid plexus to the cerebrospinal fluid.
- 7b. Cerebrospinal fluid to lymphocyte membranes and to proteins and other constituents of cerebrospinal fluid.

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

- 8b. Lymphocyte membranes to lymphocyte cytoplasm and organelles.
 - 9b. The latter to membranes of the ependymal cells.
 - 10b. The membranes to the cytoplasm and organelles of the ependymal cells.
 - 11b. The membranes of the ependymal cells to the syncytium.
 - 12b. Then becomes identical with 7.
 - 8a. Neuronal membranes to cytoplasm, nuclear and mitochondrial membranes.
 - 9a. Mitochondrial membranes to mitochondrioplasm and nuclear membranes to nucleoplasm.
 10. Neuronal nucleoplasm to nucleolar membrane.
 11. Nucleolar membranes to nucleolonema or pars amorpha.
- Here the journey ends.

Obviously, the concentration at which any transmitter, hormone, drug or toxin arrives at any of the sites listed depends upon the following factors: its initial chemistry; its route of administration; the vehicle; the rate of administration; its affinity and reversibility of binding with each site; the precise chemistry of each site determined by the particular physiology and health of the organism reflected at that particular site, including agents which react with or degrade it enzymatically and non-enzymatically; similar properties of any conjugated or degradation products of the original substance reacting with any of these sites.

After the transmitter, hormone, drug or toxin arrives at each of these sites, it may affect the biochemical reactions of a pathway or cycle, or the electrical properties at that site, although at these molecular dimensions a distinction between biochemical and electrical properties may be purely academic.

Incidentally, I have always regarded the debate about whether learning and memory are electrical or biochemical properties of the central nervous system in rather the same light. It seems to me to be far too premature in the current state of knowledge about these phenomena even to pose this question. Any attempt to answer it would be tantamount to trying to forecast several decades or centuries ahead whether an electrophysiologist would be a more or less likely person than a molecular biochemist to unravel this particular mystery.

The multiplicity of sites and variables involved in the effects of transmitters, hormones, drugs and toxins on the central nervous system, makes it difficult to arrive at firm conclusions about these reactions, but it is worth trying to derive a few generalizations.

Firstly, anaesthetics and transmitters, such as thiopentone, nitrous oxide, adrenalin and curare, which act after delays hardly longer than the circulation time, probably act on sites near the beginning of the journey, the capillaries or the syncytium. Their effects may be on the local haemodynamics within the capillaries or on the chemistry of the syncytium, possibly its mitochondria. Since generally drugs administered intravenously react very rapidly, it follows that the capillaries and the syncytium are the first targets of those drugs, and their reactions to them merit investigation. It also follows that most of the delay of the action of drugs administered by routes other than intravenously is due to their failure to achieve sufficient concentrations in the nervous system to produce effects; this delay would be due to their diffusion from the

PROPERTIES OF THE NERVOUS SYSTEM IN THE LIGHT OF PRESENT CONCLUSIONS

site of administration, their breakdown in the tissues, and their reaction en route with some elements of the tissue. The slower action of some drugs, when they are administered by mouth, intramuscularly or rectally, is either for the latter reasons, or because they take time to accumulate in sufficient concentrations in the syncytium, or because they reach the neurons via the choroid plexus. Since the blood-brain barrier is identified essentially by the delay of passage of substances between these two compartments (page 225), we must couple the syncytium and the choroid plexus as the two likely sites of this barrier.

Secondly, the fact that the addition of transmitters and some drugs to the outside of cells, results in changes in their excitability makes it possible, even likely, that the cell membranes are near the site of action, but does not prove conclusively that they are themselves, since the cells are so small that most of the substances added extracellularly could have penetrated to the cytoplasm within the time course that excitation or inhibition is seen to take place. On pages 231-232 we have already listed the other possible sites of action, which are not necessarily exclusive. It is an unprovable assumption that, for example, because any substance affects the permeability of a cell membrane, it might not also affect the biochemistry of the cytoplasm.

Thirdly, in contrasting the relative slowness of action, tens of minutes to hours, of many drugs in crossing the blood-brain barrier with their rapidity of action when injected intravenously, seconds to minutes (for review, see Davson, 1967, pages 82-92; Rapoport, 1976), I think one can conclude that many substances apparently act more quickly than studies of their passage across the blood-brain barrier would lead one to expect. This points to regions of high affinity, which may be too localized to detect in kinetic and dynamic curves of whole brain or spinal cord, and whose measurements may well be modified by the techniques used to make them, such as homogenization and extraction with powerful reagents.

Fourthly, substances such as thiourea, sulphaguanidine, sulphosalicylic acid, N-acetyl-4-amino-antipyrine, mecamylamine, barbital, and even Na^+ ions take tens of minutes to hours to reach a steady state in the cerebrospinal fluid (Brodie, Kurz and Schanker, 1960; Rall, Moore, Taylor and Zubrod, 1961; Davson and Pollay, 1963; Davson, 1967, page 67). On the one hand, intravenously injected thiopentone anaesthetizes patients usually within 10 seconds, and glucose takes slightly longer to revive hypoglycaemics from coma. On the other hand, passage from blood to the cerebrospinal fluid is obviously a slower process. Therefore, one may tentatively put forward the hypothesis that the rapid therapeutic effect is due to the substances taking the cerebral capillary-syncytial-neuronal route, while the slow passage across the blood-brain barrier travels via the choroid plexus-cerebrospinal circulation-ependymal-syncytial-neuronal route. The findings that many substances seem to accumulate in the brain before they do in the cerebrospinal fluid, and at higher concentrations (Davson, 1955; Rudolph and Olsen, 1956; Mayer, Maickel and Brodie, 1959; Wraae, 1980), leaves open the possibility that the substances normally travel from the blood into the brain, and from there into the cerebrospinal fluid, which is compatible with their taking the syncytial route.

It is difficult to pin-point particular sites, which play a more important role in the

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

uptake of substances into the nervous system, and maintain them on the basis of experiments which have not yet been carried out. However, on empirical grounds, we may suggest four of the list given on page 231. These would be the capillary membranes of the cerebral vessels and of the choroidal plexus, since these cells would be the first barriers from the blood. Then the next must be the syncytium, because it represents such a large volume, which molecules and ions must react with or pass through. The neuron with its membrane would be the fourth site, since iontophoretic and electrophysiological experiments show it to be so excitable. However, one must also bear in mind that the entry of particular substances into the neuronal cytoplasm would depend principally on their affinity for the syncytium, before they arrive at the membranes, and then on the permeability of membranes to them.

So far, the term 'receptor' has hardly been mentioned. I now wish to broaden the concept, while at the same time suggesting a solution to the accommodation problem for these apparently ubiquitous, populous, yet extraordinarily elusive, biochemical entities. Instead of regarding a particular neural receptor as being located exclusively on the surface of a neuronal membrane, we may regard it as the total of all the sites listed, which react with, or significantly delay the uptake, transport or action, of the corresponding transmitter, hormone, drug or toxin. The measured affinity of the particular substance for the receptor would be represented by the aggregate of the affinities for all of the main sites, depending upon the relative quantities of each. The delay before the substance acts would be due to the sum of the durations of all the reactions, both reversible and irreversible, which occur between administration and action.

If this hypothesis is correct, it would behove us to attempt to identify the reactions and delays at each of the component sites listed to see if we could thereby account for the total real delay both between the administration and arrival of any substance at its site of action, and also between the administration and the response of the tissue. For example, the rate of uptake and affinity of the components of blood for such a substance could be measured. The rate of passage through the choroid plexus of some substances is known. Neuroglial clumps could be used to measure approximately the rate of uptake and the affinity of the substance for the cerebral syncytium. Tracts from the spinal cord could be used to measure the same parameters.

In noting that the overall affinity of the substance for the nervous tissue is normally measured in homogenates, we must again draw attention to the total failure so far of those who study receptors to test the effect of homogenization on the affinity of substances for tissue or its effect on the rates of reactions within it. The truth of the supposition that the delay between administration and action of a substance is due to its reaction with different sites will only be demonstrated by showing that homogenization does not alter affinities of reactants or their rates significantly, that is, does not change the apparent number of receptors in the tissue under study. If homogenization does change them, the relevant corrections will have to be made.

If the view proposed here were accepted, one would explain the identification and extraction of receptors, for example, for acetylcholine, dopamine and noradrenalin, as extraction of the complex between these transmitters and the particular chemical components of the tissue with which they bind. However, the degree of binding could

PROPERTIES OF THE NERVOUS SYSTEM IN THE LIGHT OF PRESENT CONCLUSIONS

depend grossly on the method of separation of the receptors, including the homogenization, addition of detergents, etc. These agents could increase the degree of binding, induce an unnatural binding which was not a result of the original reaction of substance bound and the receptor, weaken the binding, reverse it or bring other molecules into contact with the reaction, which would themselves modify the binding. Once more, one must urge that the control experiments to test the effects of powerful agents on the system under study are grossly overdue.

Obviously, the more active transmitters, hormones, drugs, antigens and toxins, have the greatest affinity for a particular site. Furthermore, of course, by spreading the receptors throughout the tissue, the hypothesis proposed goes some way to solving the accommodation problems of the large number of substances which act on the nervous system. Despite this, I think that one should utter two cautions. Firstly, the idea that each of the latter agents has its own individual complement of one or more receptors is a so far unproven proportion for the vast majority of substances in the above categories. Secondly, even the suggestion made here, that proposes that receptors may be distributed throughout a much broader range of sites, does not solve the accommodation problems completely for such a large category of different reagents. I would suggest that hundreds or even thousands of these agents could share one receptor. The alternative suggestion is that the reagents themselves could rapidly induce their own receptors. This view would not be exceptional to many in the field. It seems to be totally acceptable in respect of those antigens which take days to induce antibodies, but it seems to me to be very unlikely to occur when a tissue is subjected to an unnatural agent, which it has never encountered before, and which acts within a very few seconds or minutes.

DOES NEUROGLIA HAVE A HIGH SODIUM ION CONTENT?

When experimentally a micropipette is driven into brain or spinal cord *in vivo* or *in vitro*, no significant sustained voltage is recorded as one passes from the overlying medium into the bulk of the nervous tissue, until the sudden appearance of a direct current voltage of 60-80 mV, which is believed to signal penetration of a cell. Thus the syncytium would seem to be at the same voltage as the extracellular fluid, and itself to be of high conductivity. Extracellular fluids and incubating media normally contain high concentrations of sodium salts (142-152 mmol/l), which suggests that the syncytium is at a similar electrochemical potential.

Katzman (1961) proposed that neuroglial cells had high sodium ion contents on the basis of: firstly, the excess sodium ions in the whole brain; secondly, the high concentrations of sodium ions found in tissue cultures of tumours; thirdly, the finding of Torack, Terry and Zimmerman (1960) that triethyl-tin caused an increase of cerebral sodium ions and swelling of what they called astrocytes, as seen by electron microscopy. At the time, Katzman's view was questioned (Hillman, 1966b) on the grounds that:

1. measurements of tissue sodium ions made *in vitro* were unreliable since the ability of the tissue to exclude sodium ions is grossly impaired by the preparation procedure (Hillman, Stollery and Joanny, 1974);
2. evidence from tumours which are frequently hypoxic was weak, since hypoxia raises

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

- tissue Na^+ (Hillman, 1961);
3. the fact that the triethyl-tin increased the sodium ion content in the brain at the same time as the astrocytes appeared swollen in Torack, Terry and Zimmerman's experiments (1960) was insufficient evidence either that the Na^+ went into the astrocytes or that normal astrocytes are high Na^+ cells;
 4. when Katzman was trying to account for the relatively high sodium ion content of the brain, he had not considered the Na^+ in the blood and cerebrospinal fluid, which could account for up to 10% of the total Na^+ in the brain;
 5. with the possible exception of the salt gland of the albatross, which has to exist in an inauspicious and inhospitable environment, to my knowledge no other cells with high concentrations of sodium ions have been reported in the literature (Hokin and Hokin, 1963).

Criticisms (1) to (4) still remain valid in asserting that there is insufficient cogent evidence to prove that in mammalian brains neuroglial cells are high sodium ion cells, yet one still has to explain the relatively high concentration of this cation in the whole unincubated brain.

The present hypothesis is only slightly different in asserting that the neuroglial syncytium, rather than the astrocytes, contains a high concentration of sodium ions. There are other findings which do not, however, add up to unequivocal evidence. Christensen and Hastings (1940) showed that extracted phosphatides had a high affinity for both sodium and potassium ions. Katzman and Wilson (1961) froze brain rapidly below the eutectic points of its main salts, and extracted lipids which contained a ratio of 1.6 of sodium to potassium ions compared with the overall ratio of 0.6. Ahmed and Hillman (1982) homogenized brain, dialysed and filtered out all the diffusible molecules, and found that their resultant fraction had a high affinity for sodium ions, and much less for potassium ions at the physiological range of the concentrations of these ions in the brain. This affinity was grossly affected by the presence of low concentrations of transmitters (to be published).

EPILEPSY

The generally accepted view of excitation of cells is that the arrival of an electrical stimulus or enough chemical transmitter at particular regions of the cell membrane increases permeability of the cell membrane, firstly to sodium ions, and shortly afterwards to potassium ions. Of course the cell membrane can be depolarized not only by a fall in potassium ions in the cytoplasm but also by a rise of potassium ion concentration in the extracellular fluid. Changes in the chemical composition of the syncytium, particularly a rise in potassium ions, or any other chemical change which would increase the permeability of the cell membrane, could trigger the firing. It has already been suggested that the argument between the proponents of chemical and electrical transmission could be resolved if the transmitters were regarded as conditioning agents; they may create an environment for the cell membrane in which it becomes more or less excitable (Hillman, 1961). This could occur as a consequence of changing the conductive or dielectric properties of the syncytium or the cell membrane.

In passing, a note of caution should be sounded here. The extremely important

PROPERTIES OF THE NERVOUS SYSTEM IN THE LIGHT OF PRESENT CONCLUSIONS

work on the electrical properties of giant axons by Keynes, Hodgkin, Katz, Huxley and their collaborators, has been assumed by such distinguished figures as Eccles and Davson to describe properties which are shared with mammalian neurons. The majority of neurobiologists today accept the ionic hypothesis, and they interpret their findings from electrophysiological studies, in relation to their belief in this hypothesis. Similarly, the findings of Schmitt, Fernandez-Moran, Finean and their collaborators on the biochemistry and biophysics of myelin sheaths or extracted myelin have been universally accepted as reflecting the properties of the neuronal membrane, mainly as a consequence of the belief arising from the Geren model that the myelin sheath represents a scroll of neuronal membranes.

When one discusses transport in cell membranes of any kind, the observations of Glynn and his colleagues on red cell 'ghosts' have been regarded as generally applicable to all membranes; this is despite the knowledge that these preparations have been made by subjecting the red cells to hypotonic solutions which change their permeability - albeit reversibly. However, the fact that the red cells can subsequently regain their previous properties does not necessarily mean that the properties of the red cell ghosts at the time of studying them are unaffected by the hypotonic treatment. The failure to realize this is similar to that of some electron microscopists who deeply freeze their tissues in conditions which some cryobiologists have shown to produce reversible effects. However, this does not mean that the freezing has no effect on the hydration, structure, dimensions, or chemical composition of the tissue. The point I would like to stress here most cogently is that although the properties of the squid axon, extracted myelin, or red cell ghost, may be generally similar to those of mammalian neuronal membranes, the belief that they are definitely applicable is frequently based on extrapolation, rather than on comprehensive experiment or evaluation.

Changes in the chemical composition of the syncytium or of the cytoplasm would be relatively slow compared with electrical conduction. Therefore, one could suppose that slow changes in direct current potential differences are due to changes in the chemistry of the syncytium. A large number of different chemical agents can be applied to the brain to produce convulsions *in vivo*; for example, ouabain, strychnine, penicillin, pentylenetetrazole, cardiazol, carbachol, diisopropylfluorophosphate, barbiturates, acetylcholine, methionine sulphoxime, hydrazides, allylglycine, picrotoxin, alumina, cobalt and tungstic acid (for reviews see Ward, 1969; Prince, 1972). Many of these substances induce convulsions when given to animals experimentally by mouth (Stone, 1972). Also cortisol, thyroxine, triiodothyronine, oestradiol and insulin can increase the susceptibility to experimental seizures (Woodbury, 1969). Some of these substances would be so dilute when they arrived in the nervous system that it is unlikely that their effects would be a consequence of changes in the electrical properties of the syncytium; therefore they probably either act by causing gross secondary alterations in the biochemistry of the syncytium and cytoplasm - and consequently the conductivity of the latter two compartments - or they accumulate on the cell membranes and thus modify their properties. Substances which are applied to the surface of the brain or spinal cord, or driven by current or

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

injection, are usually present at relatively high concentrations at the tips of the micropipettes.

In addition to the large number of reagents which may cause seizures, they can also result from freezing lesions, hyperbaric oxygen, carbon dioxide, wounds, scars, abscesses or tumours. These findings led Tower (1960, 1969) and Glaser (1964) to propose that seizures resulted from movements of cations in the brain. Hillman (1970) correlated findings from depolarization of brain in spreading depression, cells in cerebral slices and isolated neurons. He suggested that focal epileptic attacks originated in an efflux of potassium ions in a localized region of the brain. This would depolarize neighbouring neurons and cause them to fire. The newly depolarized area would excite the adjacent region until the wave of excitation impinged on the motor area of the parietal cortex. After the whole area had been depolarized it would take some minutes to recover completely, as happens after a patient has an epileptic attack.

If the syncytium surrounds the neurons it would follow that depolarization could result from (a) long-term changes in the chemistry of the syncytium; (b) temporary hypoxia, hypoglycaemia or ischaemia, due to arterial spasm or pressure from tumours in the syncytium adjacent to neurons; (c) lowering the threshold of excitation by decreased calcium ion or increased potassium ion concentration in the syncytium; (d) direct effects of agents applied near the outside of neurons on the permeability of their membranes. This hypothesis would predict that any agents which could enter the syncytium and also hyperpolarize neurons might prevent attacks of epilepsy. Among the agents which could do this are high concentrations of oxygen or glucose in the tissue and, possibly, ganglioside, which has been shown to increase the transmembrane potential in cells in brain slices and in isolated neurons (Hillman, 1961; Hillman and Hyden, 1965).

MYELIN AND DEMYELINATING DISEASES

The term myelin is used in five different senses: firstly, the sheath observed by light microscopy as a swelling on an axon, or in tissue-culture of peripheral nerves, (Figure 73), secondly, the thickening seen in transverse sections of the spinal white matter (Figures 39 and 40); thirdly, the characteristic lamellae seen by electron microscopy in the peripheral nerves, in the spinal cord, and also sometimes in kidney (Figure 72) and liver; fourthly, the chemical mixture extracted by organic solvents from peripheral nerves, spinal cord and brain; fifthly, the nervous tissue, mainly in the central nervous system, which does not stain in demyelinating diseases, such as multiple sclerosis and Friedreich's ataxia, and is detected by classical neural stains.

Obviously the electron microscopist believes that he is viewing essentially the same structure as the light microscopist sees, but at a higher magnification. Does he believe that 'myelin figures' in kidney and liver are identical with those in the central nervous system?

The chemical mixture extracted as myelin (Laatsch, Kier, Gordon and Alvord, 1962; Hulcher, 1963; Autilio, Norton and Terry, 1964; Uyemura, Tobari, Hirano

PROPERTIES OF THE NERVOUS SYSTEM IN THE LIGHT OF PRESENT CONCLUSIONS

and Tsukada, 1972) is an extract from a whole tissue. When dried out, it also appears lamellar with approximately the same periodicity as that seen around peripheral axons by electron microscopy. With the exception of those using phase-contrast microscopy of unfixed cells, microscopists use ethanol, xylene and propylene oxide, among other reagents, during dehydration, embedding and mounting; these are powerful chemicals which extract lipids, and yet the microscopists claim to be observing the repeating unit of lamellae high in lipids (please see page 173). The results of low angle diffraction studies of unfixed axons (Schmitt, Bear and Palmer, 1941; Fernandez-Moran and Finean, 1957; Finean, 1957, 1960) are believed to confirm and justify the studies of the histologists and electron microscopists. One might expect that the organic solvents would extract the lipids, and thus disrupt the lamellar appearance. A further problem is that the lamellae and the similarity of chemistry common to the electron microscopist and the biochemist, are the main criteria used to demonstrate the identity of the structure in the peripheral nerves and the whorls in the extracted myelin. Even if it had not been shown that the lamellae are unlikely to occur in living axons (pages 173-179), I think one should recognize the total inadequacy of the similar appearances as evidence that the biochemical properties of extracted myelin reflect those of peripheral nerves, and even less those of brain. The kind of evidence which would help one to do so would be to extract peripheral nerves with myelin extractants and subject them to powerful proteolytic and lipolytic enzymes, and then show that the lamellar appearance disappears. Another possible experiment would be to compare the chemistry of the lipids and other substances leaving the tissue during preparation for electron microscopy with those in myelin extracts, and also with those in the same tissue after preparation for electron microscopy. These experiments would be more persuasive than explanations of why the control experiments have not yet been published, or why the lamellae still appear after the tissue has been subjected to agents which extract lipids (Fleischer, Fleischer and Stoeckenius, 1967; Morowitz and Terry, 1969).

We have drawn attention to the lack of specificity of staining procedures (pages 63-91). All histopathological studies of demyelinating diseases show patches of tissue which do not stain (Blackwood, Dodds and Somerville, 1970; Greenfield, 1976; Treip, 1978). In the absence of staining, one cannot see which parts of the cells or their processes are affected. If these diseases really only affected the myelin sheaths, why are so few unmyelinated fibres, nuclei or nucleoli seen in plaques of multiple sclerosis? I would like to suggest that the term demyelinating disease is a misnomer, which has misled many neurobiologists into confining their attention to the electron microscopy of lamellae or to the chemistry of extracted myelin. The following are some more useful approaches currently in use:

1. Comparing the chemistry of plaques with that of adjacent and distant but similar unaffected tissues in the same subjects.
2. Comparing the chemistry of the blood and cerebrospinal fluid of patients with these diseases with those of matched patients without these diseases.
3. Comparing the biochemistry of metabolizing tissue slices of plaques with that of normal tissues in vitro (McIlwain, 1975).

THE CELLULAR STRUCTURE OF THE MAMMALIAN NERVOUS SYSTEM

I would also like to propose the following kinds of experiments, which to my knowledge have not yet been done:

4. Examination of unfixed plaques to see why they do not stain. Specifically, one would establish whether the structures in the plaques have degenerated and disappeared, or whether they have lost their staining properties with the procedures used. In the latter case, by comparing plaques and normal tissue at each stage of staining by bright-field and phase-contrast microscopy, as was done with neurons (Hillman, Hussain and Sartory, 1976), one could identify at which point they lost their ability to take up stain. This might well give a clue as to where to direct attention in studying the histopathology of these distressing diseases.
5. Teasing plaques under direct vision using dissecting microscopes and examining them by phase-contrast microscopy to attempt to define if any particular cells or parts of cells are affected by the conditions.
6. Comparing the findings in (1) to (4) with findings in animals from model conditions to see how accurately the models really imitate the diseases in human beings.
7. Comparing the plaques of regions of demyelination due to the different diseases by the techniques listed to see if there is, indeed, one common mechanism of demyelination, or if each disease has its unique biochemical pathology.

Of course, the use of traditional histopathological techniques for diagnosing the neurological conditions would not be affected in any way by the considerations in this section. Furthermore, I have used demyelinating diseases as an example of a histological approach to the study of neurological disease, but this approach need not necessarily be confined to this neurological condition.

BIOCHEMICAL STUDIES OF THE CENTRAL NERVOUS SYSTEM

The conclusion that most of the central nervous system consists of syncytium means that nearly all biochemical studies of whole brain or spinal cord have examined mainly the properties of the syncytium. True studies of the biochemistry of neurons have isolated these cells by hand dissection by the technique of Hyden (1959), or homogenized, filtered and subsequently picked out by the method of Roots and Johnston (1964). Since the volume of the naked nuclei is small compared with that of the total syncytium (Figures 60 and 62) one may say that study of the neuroglial clumps yields information mainly about the chemistry of the syncytium.

If the relevant control experiments had been carried out, so that one could be certain that oxidative phosphorylation occurs in mitochondria in the living intact animal, as it does in the subcellular fraction in vitro (Hillman, 1972, pages 39-40), one could attribute the high rate of cerebral oxygen uptake to the dense population of mitochondria, both in the neurons and in the syncytium. However, this would give us no clue as to what the central nervous system does with all the energy. A popular suggestion, that it is used to maintain the ionic gradients, is not adequate, since it is believed that all cells maintain the ionic gradients across the cell membranes by the

PROPERTIES OF THE NERVOUS SYSTEM IN THE LIGHT OF PRESENT CONCLUSIONS

utilization of energy, and most tissues are composed of cells whose membranes are much more closely packed together. Action potentials occupy a minute proportion of the lifetimes of excitable cells, and very small quantities of ions move during and after them, so that the total extra energy due to the restoration of the electrochemical gradients across the membranes of the cells would add an insignificant sum to the high energy expenditure of the nervous system. Perhaps it is worth wondering whether the energy is in any way used in the storage or recall of information, but at the moment I cannot think of any experiments to test such a totally adventitious hypothesis.

Many research workers have devised bulk fractions of what they claim to be neurons, astrocytes and oligodendrocytes (but not microglia) (for references please see Table 7, taken from Pevzner, 1979). Originally they called their fractions by the names of the presumed cell types, but it was often difficult to be sure of the identification of the predominant cell in fractions, or that the fractions contained no substantial proportion either of other cell types or of 'debris' - which could affect the biochemistry of the fractions significantly. Therefore, more recently, the authors have talked about 'neuron-rich', 'astrocyte-rich' and 'oligodendrocyte-rich' fractions. This change of nomenclature cannot unfortunately circumvent the real problem of the ignorance of the contribution which the particular cell type makes to the biochemical properties of the fraction as a whole. The fractions have usually been 'washed' several times. The purity of the fractions is usually judged by the subjective impression of the light or electron microscopist, and it is illustrated in publications with selected electron micrographs which are intended to show fields typical of the fraction as a whole.

Furthermore, one is not entitled to assume that the homogenization, centrifugation, and the chemistry of the suspending media used, have no influence on the biochemistry of the fraction. This difficulty is compounded by the knowledge that the different cell-rich fractions are prepared by such different physical and chemical procedures that - even if the original cell types had all shared the same properties - they would be likely to exhibit different biochemical behaviour after having been subjected to such different regimes.

One is faced with an analogous situation when trying to compare the biochemical properties of the different cell types in culture (Crain, 1976; Fedoroff and Hertz, 1977; Pevzner, 1979; Giacobini, Vernadakis and Shahar, 1980). In this situation also, the presumed different cell types are separated in different chemical environments, and are grown in different culture conditions, both of which strongly influence the chemistry of the cells. Therefore one cannot simply compare the chemical properties of such cultured cells and assume that these properties accurately reflect those of the parent tissues. However, one can probably assume that those tissues such as retina, cerebellum and muscle, in which one can recognize cell types for some time after the tissue has been isolated and cultured, represent the best model to study. We must remind ourselves that tissues do change their biochemical and staining properties in culture (Hansson, 1982). One should really favour procedures in intact animals, and those that disrupt tissues minimally. (Hillman, 1976).

12

Practical Recommendations

Throughout this monograph, and in previous publications (Hillman, 1972; Hillman and Sartory, 1980), a wide variety of experiments to control the effects of technical procedures and the agents used in them have been urged. Neither the vastness of the task, nor the failure to execute them so long after conclusions have been drawn from them, can absolve the captains of the research industry of their duty to carry them out.

In addition, it would seem appropriate to make the following recommendations:

1. More studies should be made using non-invasive techniques, wherever possible, and on tissues which metabolize at rates comparable with those *in vivo* at the time of the examination (Hillman, 1983).
2. Hand isolation, micromanipulation and teasing of preparations using no or minimal powerful chemical reagents or physical agents are to be preferred (Gray, 1931; Chambers and Chambers, 1961).
3. The specificity of neurological staining procedures should be examined comprehensively, in an attempt to prescribe what few procedures should be most encouraged by virtue of their specificity and the smallness of degree of distortion they cause. It is widely believed that, if different staining procedures show up cells of different forms, this defines genuinely different populations of cells. The assumption inherent in this belief is that the 'specific' staining procedures all distort cells to approximately the same extent. Thus, we have two interrelated tasks - viz. to define the different populations, if the neuronal and naked nuclear populations can, indeed, be subdivided, and to specify the optimal staining procedure for all such populations so defined. The latter would result in a considerable saving on expensive colour reagents, perhaps to the detriment of the aesthetics of the histologists' art and the decorations on the walls of their laboratories.
4. Neuroglia should be subject to more intensive study. The simplest preparation is the neuroglial clump, but the samples are rather small. Many microtechniques have been developed for examining the chemistry of small samples (Neuhoff, 1973; Osborne, 1974). Unfortunately, there is no information on the electrical properties of

PRACTICAL RECOMMENDATIONS

these neuroglial clumps, upon which studies would be rewarding, especially if the samples have been examined microscopically to ensure that they contain no neurons.

Further attempts should be made to study the physiology of nuclei isolated by hand dissection, as has been done with the nuclei of ganglion cells and of cells from Deiters' nucleus (Giacobini, 1957; Edstrom, 1958; Cummins and Hyden, 1962; Hillman and Sartory, 1980, page 77). The spherical nuclei are difficult to manipulate with a fine stainless steel wire, and their refractive index appears to be similar to that of the incubating medium.

5. Structures which have been shown not to exist (Table 17) should obviously not be studied by electron microscopic, biochemical or pharmacological techniques.
6. The nucleolar membrane (Hussain, Hillman and Sartory, 1974) and the nucleolonema of neurons (Estable and Sotelo, 1951; Sartory, Fasham and Hillman, 1971) should be more extensively studied by light microscopy, especially in tissue culture.
7. The structure of the membranes of the cell, nuclei and mitochondria needs to be re-examined, by non-disruptive techniques (Appendix 4, page 251).
8. The mechanism whereby particles cross cell membranes should be studied in unfixed tissues by non-disruptive techniques. It seems quite inappropriate to use histological, electron microscopic or disruptive methods, or dead tissue, to study the dynamics of a process which only occurs in living tissues. There is plenty of evidence that small molecules can diffuse into all cells, that large particles can be ingested by phagocytes, and that particles can enter the syncytium. However, knowledge is scant on the mechanisms of these movements.
9. The concepts and evidence for transport by the membrane, for transmission, and for receptors, stand in need of complete reassessment. These should start with a fundamental investigation into each element of the evidence for them. Neurobiologists should not regard the enormous volume of experiments carried out on the assumption of these concepts, as adequate evidence in support of them. I would suggest that the time has come to re-examine the validity of the concepts themselves. More vigorous attempts to falsify the hypotheses behind the interpretations should be pursued. Unfortunately, neither the number of adherents to a particular concept, nor the volume of uncritical publications to adumbrate it can be regarded as an adequate measure of its validity.
10. Intraventricular injection of antimitotic drugs must be effective against gliomas.
11. Controls for subcellular fractionation should be done urgently.
12. Receptors should be sought by light microscopy in unfixed cells.
13. Immunocytochemical experiments should be repeated in unfixed tissues.
14. Attempts should be made to find out if adult neurons become malignant.

13

Postscript

In this monograph I have tried to use only proper evidence and reasoning. The following elements are particularly important: the distinction between hypothesis and finding; the citation of the evidence of the best established authorities in a discipline (although not necessarily their interpretations of their own data); quotation of several authors, especially for dimensions; the deliberate practice of not selecting values from the literature to fit in with the hypothesis at hand; insistence on analysis of the original evidence for the hypothesis being discussed; unequivocal respect for geometry; identification of crucial control experiments; attention to the techniques and calibrations of measurements; total willingness to enter into dialogue anywhere in the world; total willingness to modify or abandon any hypotheses or conclusions as a result of such discussions.

The subject of this monograph has implications for other sciences. It represents a 'paradigm' (in the sense of Kuhn) for the way research is pursued. However, there are several other important aspects of this problem. These include what in my view are two other imperatives:

1. We have a duty to society to employ the expensive resources and skills placed at our disposal as efficiently and productively as possible. Study on apparent structures which have been shown to be artifacts, or the use of uncontrolled techniques, is an abuse of the trust society puts in research scientists, and of the esteem in which they are held.
2. Neurobiologists should be humble in relation to their relatively poor success to date. We are still far from understanding the normal physiology and biochemistry of memory and thinking, the pathology of the main psychoses, the immunology of the nervous system, the genesis of psychoses, and many other phenomena. The ill have a right to be impatient. We are hardly nearer solving these problems than Vesalius was in sixteenth-century Brussels, when he sliced off the crowns of executed criminals and looked down into their brains (please see front cover).

Appendices

APPENDIX 1. Comments by some authors on the specificity of staining procedures
(See also pages 63-91)

Authors	Page	Comments
Penfield (1924)	434	'That oligodendroglia is very difficult to stain completely has been well demonstrated by the history of these <u>adendritic</u> cells.' A much more complete description has been provided by Del Rio-Hortega, using ammoniacal silver carbonate. Even by this method the results are variable. Some undetermined factors in the preparation of the tissues for the silver bath at times cause only microglia to be stained, or microglia and classical neuroglia. When the tissue is in a more favourable condition oligodendroglia is stained, but there is also impregnation, though incomplete, of the two above elements. When the tissue is in the most favourable condition a completely selective stain of oligodendroglia is obtained and the cell prolongations are beautifully clear-cut and appear much ramified.'
Weil and Davenport (1933)	174	'Even trained technicians, however, do not always succeed in the preparation of the silver solution devised by Stern, and, besides, the use of a 10 per cent commercial formaldehyde solution in connection with 10 per cent ammoniacal silver nitrate frequently leads to the production of disturbing silver precipitates.'
Gray (1954)	401	'Nerve cells and processes are stained more easily by the dye-staining techniques, than they are by metal-staining techniques. It must be admitted that most of the techniques are not so specific, but they are certainly adequate for class demonstrations; and it must not be forgotten that much of the classical research was done by these methods rather than by metal stains.'
	403	'The literature of the hematoxylin staining techniques for nerve cells and their processes is almost as confused as the literature of the metal-staining techniques for the same purpose.'
	525	'It is difficult to decide, in the case of osmic acid, which of the staining formulas given should be regarded as fixatives.'

APPENDIX 1

Authors	Page	Comment
	537	re: demonstration of protoplasmic neuroglia in the cerebral cortex by the method of Cajal (1916), 'The successful application of this technique depends on the ability to judge the exact shade of purple which indicates a satisfactory termination of the staining process.'
	543	'Dichromate-silver method. The techniques are experimental, yielding the most brilliant results when a successful impregnation is obtained, but for which it is almost impossible to specify conditions leading to success. The length of time is entirely critical, but results cannot be predicted or even reproduced with certainty. Among the numerous pieces thus treated, one may be found which will show the required condition.'
Clark (1973)	74	'Methods that have been developed for staining a particular component are usually selective but may partially stain other components.'
	76	'Bubbles - but when small, they appear as holes 5-20 µm in diameter in cut sections and may show little compression of adjacent tissue. They may become filled (after aqueous formalin) with myelin derivatives and this material may even take a stain with basic dyes and be mistaken for degenerative changes of glia cells.'
Carleton and Drury (1962)	221	'The complexity and the intervention of factors as yet unknown make it often difficult to stain particular elements of the nervous system in the same way on different occasions.'
	232	'The Golgi methods are capricious, and it is difficult to get the same results in different species - even of the same tissue.'
	240	'When owing to post mortem change the silver methods fail, Cajal's gold technique (p. 245) is worth trying.'
Gray (1964)	411	Microglia - 'Neurons that have become dark and shrunken as a result of poor fixation or mechanical pressure may take on a similar appearance.'
McClung Jones (1967)	405	'In general the pathological forms of all three types of cells are more easily demonstrated by the metallic methods than are the normal forms. Oligodendroglia may be impregnated in the grey and white matter with equal facility. Microglia cells are studied more easily in the grey matter where they are much more numerous. When oligodendroglia is well stained microglia is usually stained also. But either cell group may be stained with exclusive selectivity. At times when using silver carbonate, astrocytes may be incompletely stained along with the two groups of smaller cells.'
Ralis, Beasley and Ralis (1973)	57	'Haematoxylin and eosin. This method has little to recommend it for demonstrating nervous tissue proper...'
Smith and Bruton (1977)	7	'The majority of "staining" techniques used today were conceived by accident: they were observed by the perceptive eye or resulted from inspired guesswork. Frequently the precise chemical constituents of the

APPENDIX 1

Authors	Page	Comment
		tissue were, and are, unknown and the "theory" of the staining mechanism empirical. Techniques based on the known chemistry of the tissue reacting to a known and understood reaction are few in number.'
	9	'In practice it must be admitted that few techniques are specific.'
	11	'Haematoxylin and eosin sequences provide a good example of the difficulties in stating "correct" staining times and the correctly stained preparation.'
Jacobson (1978)	47	'A wide spectrum of cytoplasmic densities in developing oligodendrocytes has been seen. This may make it difficult to distinguish between young astrocytes and young oligodendrocytes. To add to the difficulty, the appearance of glial cells changes with their functional states. For example, the oligodendrocyte active in myelination has more extensive cytoplasmic processes, more abundant cytoplasmic organelles, and more dispersed nuclear chromatin than the mature, resting oligodendrocyte.'
	48	'The Golgi method and the silver methods are not selective for neuroglia However the staining method used to study oligodendrocytes is not completely selective and cannot always be depended on to show those glia cells.'
Mirsky, Wendon, Black, Stolkin and Bray (1978)	121	'The neurofilaments which can be seen with the light microscope in so-called neuroblasts, neurons, and astrocytes after silver impregnation, are probably artifacts caused by clumps of neurofilaments and neurotubules.'
	258	'There are no other satisfactory histological markers for neurons; even the reduced silver methods may stain glia or fail to stain neurones under certain conditions.'

APPENDIX 2. Comments by some authors on the specificity of markers

(See also pages 88-91)

Authors	Page	Comment
Uyeda, Eng and Bignami (1972)	86	'Thus the double band obtained with extracts of tissues containing few astrocytic fibers, such as the cerebral cortex and white matter, and the basal ganglia, was lightly stained.'
	88	'The GFA protein appears to be confined to the central nervous system, as indicated by Ouchterlony double diffusion technique. It was found in normal brain tissue, brain tissue with reactive fibrillary gliosis and fibrillary astrocytoma.'
Jacobson (1978)	116	'The identification of distinct neuronal and glial phenotypes has, until recently, required overt cyto-differentiation, and thus the cells could be recognised unambiguously only when they were approaching the final stages of differentiation. This often involved the subjective impressions of the observer, and was especially subject to error in cases where cells have similar appearances during their preterminal stages of differentiation, when transitional forms might arise, or when cell differentiation might be somewhat atypical, as in tissue culture.'
Mirsky, Wenden, Black, Stolkin and Bray (1978)	258	'It is not possible to prove rigorously the specificity of tetanus toxin in all situations.'
Varon (1978)	94	'Thus, a <u>cell specific marker</u> can be best defined as a <u>property</u> that is <u>expressed</u> uniquely or prevalently in a given cell class when <u>correspondingly optimal conditions prevail</u> . This definition requires that description of a marker include the set of extrinsic circumstances under which it has been observed, and eventually the set of extrinsic circumstances which <u>condition</u> its expression.'
	96	'S-100 is an example of a molecular marker in search of a function.'
Raff, Fields, Hakamori, Mirsky, Pruss and Winter (1979)	284	'The increasing use of in vitro techniques in neurobiology has accentuated the need for cell-type specific markers, which would allow unambiguous identification of the different types of neural cells in culture.'
	290	'Although the GFAP+ cells in cultures of cerebellum, cerebral cortex, corpus callosum and meninges did not label with tetanus toxin, some process-bearing GFAP+ cells in cultures of optic nerve were weakly labelled.'
	306	'Although the use of cell-type specific markers has enabled us to study a number of properties of defined neural cell types in culture for the first time, it should be appreciated that the assignment of "positive" or "negative" for a particular property is operational and depends on the sensitivity of the assay used.'
Kennedy, Lisak and Raff (1980)	344	'Some of the GFAP+ cells were large and had large vesicular nuclei, and others were smaller and darker and had multiple branching processes. While it is likely that these were protoplasmic and fibrous astrocytes, respectively, many GFAP+ cells had an intermediate morphology and could not be so classified.'

APPENDIX 2

Authors	Page	Comment
	347	'It is still uncertain, however, whether all astrocytes are GFAP+.'
Raff, Brockes, Fields and Mirsky (1980)	18	'While it is clear that the great majority of neurones in cultures of peripheral and central nervous tissues in various species are labelled by tetanus toxin, it is not certain whether all neurones label with tetanus toxin. Therefore, one must be cautious in using tetanus toxin as a neuronal marker, particularly since some preparations of tetanus toxin label various cell types "non-specifically."'
	20	'Although the morphologies of some neural cell types are, at times, sufficiently distinctive to allow identification by this criterion alone, this is usually not the case.'
Bartlett, Noble, Pruss, Raff, Rattray and Williams (1981)	346	'All the latter cells could be identified as neurones by the fact that they bound tetanus toxin.'
	348	'Anti-Ran-2 antibodies. . . . They labelled astrocytes having diverse morphologies.'

APPENDIX 3

Appendix 3 Typical counts made with the Quantinet particle counter in examining six slides of coronal sections of the frontal lobe of rat brain. The area $\times 1000$ is the number of Pix cells in that particular field, whose mean is given in column 2 for each staining procedure. The Selector A is the setting on the particle counter chosen for each field to give the maximum contrast between the stained tissue and other elements. The overall results are given in Tables 15 and 16.

APPENDIX 4. Biological membranes and measurement of their thicknesses

An interface may be defined as a detectable region where two immiscible fluids meet.

A biological membrane is a thin solid structure separating two different fluids.

The boundaries of vacuoles and of lipid droplets in milk are interfaces. Little more need be said about them, except that most microscopic techniques can not distinguish between an interface and a membrane.

Membranes were first detected by light microscopy, usually in stained tissues, and subsequently by electron microscopy. They may be detected by (1) a difference between them and the phases on both sides of them, or (2) inference from the fact that the phases on both sides appear different, which the observer interprets as being due to a membrane being present. The differences observed under both headings are due to dissimilarity with respect to:

- (a) absorption of light of a particular wave length (colour);
- (b) emission of light after exposure to light of a particular wave length, including excitation of fluorescent emission;
- (c) diffraction at the edges (which does not permit measurement of thickness);
- (d) intensity of colour - the commonly used difference in histology and electron microscopy;
- (e) refractive index, enhanced in phase-contrast microscopy;
- (f) any of these properties consequent upon the tissue being colour stained, and the membranes and phases on either side of the membranes, having different retention of all the reagents, used throughout the whole histological procedures;
- (g) birefringence seen by polarization microscopy;
- (h) reflectivity seen by dark-ground illumination and reflected light illumination;
- (i) textures, especially with freezing techniques of electron microscopy;
- (j) different structures being present on both sides of the apparent membrane, for example, the synaptic vesicles which are believed to be on only one side of the synapse;
- (k) antibodies reacting with antigens present on, or adherent to, the cell membranes;
- (l) deposit of salts of heavy metals, such as those of silver in light microscopy, and osmium, platinum or tungsten in electron microscopy;
- (m) shrinkage artifacts around membranes.

There are several other techniques which have given more or less direct evidence for the existence of membranes:

- (n) differences of impedance, which is much higher in tissues than would be accounted for by the composition of the extracellular and cytoplasmic fluids; also, the impedance can be measured between an electrode believed to be intracellular and one in the extracellular fluid or incubating medium (Coombs, Eccles and Fatt, 1955; Frank and Fuortes, 1956);
- (o) low angle diffraction of X-rays, from which the repeating unit of the presumed membrane is calculated.

APPENDIX 4

Other information about membranes is gleaned from:

- (p) the considerable potential difference which is recorded abruptly when an electrode with a tip diameter of the order of microns is advanced into a tissue and which changes with the cation constitution of the extracellular fluid or incubating medium;
- (q) calculations based on the quantity of lipids and proteins which could be extracted from red cells and the calculated areas of those cells (Gorter and Grendel, 1925);
- (r) examination of reflections from dried erythrocytes compared with films of barium stearate using the 'leptoscope' (Waugh and Schmitt, 1940);
- (s) the ability of lipid mixtures to make monomolecular layers on salt solutions (Bangham and Horne, 1964);
- (t) the similarity of many of the physicochemical properties of well-characterized artificial membranes to those of biological membranes.

The presence of solid membranes around cells is attested by the ability with which neuron somas can be cut open (Cummins and Hyden, 1962), and the contents of squid axons can be squeezed out with a 'microsteamroller', and the axons can subsequently maintain their excitability on being refilled with potassium chloride solution (Baker, Hodgkin and Shaw, 1961). The fact that the contents of the cells of the sympathetic and dorsal root ganglia do not all flow away when the capsules are removed is further powerful, if indirect, evidence that the cells are each personally invested with a mechanical membrane, rather than a fluid interface.

Any skilled advocate knows that individual pieces of evidence vary in their cogency, and one must beware lest the quantity of evidence might be more persuasive than the quality of its individual elements.

The common belief nowadays based on electron microscopic observations and low angle diffraction measurements is that the 'unit' membrane is, say, 6-8 nm thick. It must be repeated that the electron micrographic 'trilaminar' membrane is a metal deposit on a structure which has been dehydrated, is believed to contain much lipids and has been extracted with reagents including alcohol and propylene oxide, and subjected to high vacuum, electron bombardment and X-irradiation. Many of the low angle diffraction measurements have been made on the myelin lamellae of peripheral nerves, which are shown to be artifacts (please see page 173-182).

The maximum resolution of the light microscope under optimal conditions is 20-25 nm, so that a membrane of 6-8 nm thickness could only be seen under the following circumstances:

- (a) if a metal stain was deposited along the membrane in sufficient thickness;
- (b) if an antigen-antibody reaction produced a precipitate of the required dimensions on the membrane;
- (c) if there was a large difference of refractive index on both sides of the membrane;
- (d) if the membrane was composed of multiple layers whose total thickness exceeded 20 nm.

APPENDIX 4

One must state quite categorically that if the membranes in life did have a thickness of 6-8 nm, they could not be measured by any technique of microscopy.

We have already shown elsewhere that the 'unit' membrane is an artifact, because it occurs normally to the plane of section more frequently than geometry would permit (Hillman and Sartory, 1980, page 40). Therefore, we must seek further but indirect evidence about the dimensions and the existence or non-existence of the cell membrane from non-microscopic observations (h) to (t).

The high impedance between two electrodes which can be measured when a 'resting' potential attributed to penetration of the cell membrane is recorded - like the resting potential itself - is likely to be good evidence, but cannot be conclusive until compared precisely with the impedance and the zeta potential of interfaces of droplets of the same dimensions and similar chemical compositions. Nevertheless, resistances of $0.8 \text{ M}\Omega$ found across presumed neuronal membranes (Coombs, Eccles and Fatt, 1955; Frank and Fuortes, 1956) seem high for a relatively thin interface of any kind, just as a resting potential of 50-80 mV seems high compared with zeta potentials of up to 5 mV.

Calculations of membrane thickness based on the quantities of protein and lipid which may be extracted and the surface area of the cells studied depend crucially on the completeness of the extraction of the chemicals, the accuracy of measurements of the protein and lipid, the accuracy of measurement and calculation of the total membrane area of the tissue under study (usually red cells), and the belief that the particular chemicals are almost exclusively located in the cell membranes. All of these require a pretty interplay of assumption and interpretation whose total effect is not wholly reliable. However, it would be a reasonable approach if all these uncertainties and approximations were resolved. Analogous difficulties beset the use of the leptoscope and of low angle diffraction studies, which are usually carried out on partially or completely dehydrated tissues.

The ability of lipid and proteins to make monomolecular layers and to complex in thin layers, and all the artificial membranes which have been studied, represent elegant models which can explain the biophysical properties of membranes, and they are properly used for that purpose, but in logic they cannot and do not show that the model represents the structure of the cell membranes or their dimensions. Their fundamental value in this respect is a function of the accuracy to which their physicochemical conditions are designed to reflect the total milieu of the biological environment of the real cell membrane. I have urged the value of this experimental approach elsewhere (Hillman, 1983).

It is tempting to regard the dimpling and resistance at the tip of an advancing micropipette, at what appear to be the membranes of neurons and muscle cells, as further mechanical evidence for the existence of the membranes around the cells, but unfortunately similar phenomena can result from attempts to penetrate immiscible droplets using fine 'intracellular' micropipettes.

If we now return to the light microscope and direct our attention to an attempt to see the membranes in unstained or stained tissues, we realise that we do not see the membranes, we see differences of colour, texture, refraction, and intensity of staining

APPENDIX 4

between the cytoplasm and the extracellular fluid (a) to (o); we are conditioned to believe, albeit subconsciously, that we are seeing membranes. In electron micrographs one is often inferring the existence of a membrane between two adjacent areas of different texture or intensity of staining.

The concept that the cell, nuclear or mitochondrial membranes might be liquid, fluid, quasi-liquid or quasi-fluid, is dealt with elsewhere (please see pages 25-26, 132-135).

We may therefore arrive at the following tentative conclusions. The mechanical and electrical properties present near cells make it very likely that they possess membranes, unless they are part of syncytia (please see page 108). The presence of a membrane is detected or inferred by techniques of light or electron microscopy, and confirmed by non-microscopical techniques, but so far attempts to measure membrane thickness representative of that *in vivo* have ignored known and expected effects of preparation on the measurements.

APPENDIX 5. The tissue in the electron microscope

In a preparation for electron microscopy the following types of material are present:

1. The tissue, consisting of water, proteins, lipids, carbohydrates, nucleotides, electrolytes, dissolved gases, inter alia.
2. The heavy metal stain, which consists of metals or salts of osmium, lead, uranium, tungsten or manganese.
3. The embedding media of epoxy resins, methacrylates, ice, polyesters or polyampholytes.
4. Other reagents, which have not evaporated, including fixatives, alcohols, propylene and solvents.

Each of these reagents will be grossly different in respect of the following parameters: volatility, stability, temperature coefficient of expansion or contraction, heat capacity, heat conductivity, reactivity with each of the other substances at each of the temperatures to which they are exposed, radiation emission, radiation absorption and electron density. During the preparation each of these parameters will be changing continuously due to changes in their chemical environments, temperature, electron bombardment and exposure to high pressures during freezing and low pressures in the electron microscope.

In view of the complexity of the system, it is unlikely that studies of simplified reactions like those between osmium salts and proteins or lipids can be accepted as sufficiently comprehensive models of what is occurring to tissue during preparation for electron microscopy.

The electron bombardment raises the temperature in the specimen to the equivalent of several hundred degrees (Reimer, 1965; Favard and Carasso, 1972; Grubb and Keller, 1972a,b). Such temperatures are within the range used to ash tissues for chemical measurements of metal ions, or in flame photometry to vaporize atoms. The tissue could not survive. Therefore it is a sad, if unpopular, fact of life, that the image seen through the electron microscope must be composed almost entirely of heavy metal.

It is often said that freezing techniques are independent evidence to confirm findings from transmission electron microscopy. The only step which fundamentally differs between the two techniques is that the initial fixation in the former technique is done by extreme cold, while in the latter a classical fixative such as osmic acid or glutaraldehyde is used. A large number of artifacts have been identified during the study of both biological systems and simpler systems of plastics, sucrose and even water (Sleytr and Robards, 1977, 1928a,b; Dubochet, Lepault, Freeman, Berriman and Homo, 1982; Chang, McDowell, Lepault, Freeman, Walker and Dubochet, 1983). However, until now, to our knowledge, no authors except ourselves have addressed themselves to the question of how much of what is generally regarded as the intracellular structure could be accounted for by these artifacts.

APPENDIX 6. Unsatisfactory terms

Two previous lists of terms, which in my view are misleading, loose, or meaningless, or they obscure real difficulties of interpretation, have been drawn up (Hillman, 1972, pages 115-120; Hillman and Sartory, 1980, page 99). I would respectfully refer the reader to these lists, as they are highly relevant to the considerations in this monograph. Many of the words are metaphors, and they have been taken over from other disciplines, where their meaning is often more precise and better understood. The semantic problem is a real one. How often has one heard students saying that particles enter cells by the mechanism of endocytosis, not appreciating that they are merely translating ignorance into crypto-classical Greek? Some of these terms are worthy of exposure. *An asterisk indicates a largely meaningless term.

1. *Particles are assembled or in an assembly; this means that they appear to be in a row, but the use of the term should not bear a teleological implication.
2. Concept is an idea, hypothesis, construct or fantasy, and cannot be used for a structure which is alleged to have material existence, such as a membrane. It can be used for a theoretical proposition about its properties, only until such a time as the hypothesis is tested. The longer the time that elapses before it is tested, the more urgent the testing of the hypothesis becomes.
3. *Cues are presumably movements of particles or molecules of biochemical changes which induce other changes. The term has a real meaning only in sensory physiology.
4. *Particles decorate a pore, granule or vesicle, because the author likes the look of them. They decorate, mainly because - like the mountain - they are there.
5. Endocytosis, exocytosis, pinocytosis and phagocytosis mean that particles or droplets go into and out of cells, but tell one nothing about the mechanisms causing these movements.
6. *One particle or reaction is implicated with another, if the author believes that they have a physical or biochemical relationship, but has not yet demonstrated it.
7. *Structural changes, physiological events or biochemical reactions should only be described as integrated if there is abundant evidence of their relationship in time, if there is a rationale for the alleged relationship, and if some control mechanism has been proposed or demonstrated for it.
8. A marker is a substance which should have been demonstrated to be located in a particular kind of cell, or part of a cell.
9. A model is either a physical object imitating a proposed structure and changes in it, or it is a biophysical or mathematical concept of a proposed structure or of biochemical reaction or changes in it. It is essentially a series of hypotheses whose totality is intended to mimic the structure or biochemical reaction as closely as possible. The term can not be used for the structures or the reactions themselves, since they are more closely related epistemologically to findings.
10. *Modulation means that a substance affects the excitability of a cell or the release or effect of a substance believed to be a transmitter. In using the term one should really wonder whether the modulation is special in that the modulator is being alleged to affect the reaction in a way which is more particular than simple chemistry would dictate.

APPENDIX 6

11. *An organized structure appears to have a pattern which is believed not to be an artifact of preparation.
12. *A substance packaged or parcelled in a particle is a substance which is believed to be present in that particle, and to travel in it.
13. *A putative transmitter is a substance found in the nervous system, which some authors believe to be a transmitter, and they hope to accumulate enough evidence in the future to satisfy classical criteria that it is (please see page 4).
14. A neural receptor is a particular physical or biochemical part of a cell membrane, which is believed to react with a transmitter, hormone, drug or antibody before any of them can exert their action. Few receptors have been shown, and the demonstration of the effects of one of these agents is often believed to be sufficient evidence that a receptor for it must exist on the cell membrane. Nevertheless, there are several classical criteria which define a receptor, including high affinity, competition, saturation and simple numerical relationships of their materials with the transmitter, drug, hormone or antibody.
15. *A particle or substance is restricted if it always appears in the same part of the cell. It means that it is there, and the use of the term implies that it has not moved during preparation.
16. The role or function of a particle or chemical either means the effect it has, or it means the 'purpose' for which it is there. The latter meaning is either teleological or theological.
17. *A substance which is sequestered is believed to be at a particular site; that is, it is there, temporarily or permanently.
18. *A specialization is a feature which is seen to be constant and which is believed to be the site of a particular biochemical activity.
19. *The statement that a substance is stored means that it is there, usually as a granule.
20. A transmitter is a naturally occurring substance, which is believed to be released near a neuron or a neuromuscular junction, to have an effect at relatively low concentration, to have a high affinity for the membrane, and to increase or decrease its excitability. It is also defined as having enzymes to synthesize it near its site of action. Several transmitters are believed to be broken down by enzymes in the tissues adjacent to the neurons. There are also a wide range of unnatural compounds which in low concentrations either increase or decrease the effects of transmitters.

APPENDIX 7. A few naive but important questions

Why have no systematic attempts been made to control the effects of killing animals, and of homogenising and centrifuging tissues, on the form, biochemistry and pharmacological properties, of cells?

How is the existence of a cytoskeleton seen by electron microscopy compatible with intracellular movements seen in life by light microscopy?

Why does evidence cited in discussions on the structure of central synapses, and for the vesicle hypothesis, usually originate from findings on the neuromuscular junction?

Why do immunocytochemists not use unfixed, unembedded, unstained, tissues for localising cellular constituents?

Do homogenisation and centrifugation affect the affinity of substances for receptors?

What is the difference between a particle, granule or body, being assembled, decorating, being organized, packaged, parcelled, restricted, sequestered, occupying a specialized site, stored - or just being there?

What are the nuances between enhancing, implicating, mediating, being a second messenger, modulating - and just causing?

How does one differentiate between diffusing, flowing, migrating, partitioning, being transported - and just moving?

What are the criteria used to identify bodies, deposits, free ribosomes, lysosomes, particles, precipitates, secretory granules, vesicle-like bodies - and artifacts?

What fills a bubble, droplet, lumen, pinocytotic vesicle, pore, space, shrinkage artifact, vacuole - or just a hole?

What is life?

References

- Abney E R, Bartlett P P and Raff M C (1981) Astrocytes, ependymal cells and oligodendrocytes develop on schedule in dissociated cell cultures of embryonic rat brain. *Dev Biol* 83, 301-10
- Adams C W M (1965) *Neurohistochemistry*. Elsevier, Amsterdam
- Adams C W M and Bayliss O B (1968) Histochemistry of myelin. V. Trypsin-digestible and trypsin-resistant proteins. *J Histochem Cytochem* 16, 110-14
- Adams C W M and Davison A N (1965) The myelin sheath. In *Neurohistochemistry*. ed. Adams C W M. Elsevier, Amsterdam, pp. 332-400
- Adams C W M, Davison A N and Gregson N A (1963) Enzyme activity of myelin: histochemical and biochemical evidence. *J Neurochem* 10, 383-95
- Adrian R H (1956) The effect of internal and external potassium concentrations on the membrane potential of frog muscle. *J Physiol* 133, 631-58
- Aghajanian G K and Bloom F E (1967) The formation of synaptic junctions in developing rat brain: a quantitative electron microscope study. *Brain Res* 6, 716-27
- Ahmed N and Hillman H (1982) Preparation of an insoluble fraction of ox brain and its affinity for Na^+ and K. *Neurochem Res* 7, 911-21
- Akert K, Pfenninger K, Sandri C and Moor H (1972) Freeze etching and cytochemistry of vesicles and membrane complexes in synapses of the central nervous system. In *Structure and Function of Synapses*. eds. Pappas G D and Purpura D P. Raven Press, New York, pp. 67-86
- Albot G and Jezequel A M (1962) Ultrastructure du foie et pathogenie de l'ictere au cours des hepatites virales. *Arch Mal Appar Dig Mal Nutr* 51, 505-28
- Aldridge W N and Johnston M K (1959) Cholinesterase, succinic dehydrogenase, nucleic acids, esterase and glutathione reductase in sub-cellular fractions from rat brain. *Biochem J* 73, 270-7
- Aleu F P, Edelman F L, Katzman R and Scheinberg L C (1964) Ultrastructural and biochemical analysis in cerebral edema associated with experimental mouse gliomas. *J Neuropathol* 23, 253-63
- Aleu F P, Katzman R and Terry R D (1963) Fine structure and electrolyte analyses of cerebral edema induced by alkyl tin intoxication. *J Neuropathol* 22, 403-13
- Allen J N (1955) Extracellular space in the central nervous system. *Arch Neurol Psychiatry* 73, 241-8
- Allen R D and Allen N S (1982) Video enhanced microscopy with a computer frame memory. *J Microsc* 128, 193-208
- Allen R D and Kamiya N (eds.) (1964) *Primitive Motile Systems in Cell Biology*. Academic Press, New York, pp. 1-642
- Allen R D, Metuzals J, Tasaki I, Brady S T and Gilbert S P (1982) Fast axonal transport in squid giant axon. *Science* 218, 1127-29
- Allen R D, Travis J L, Allen N S and Yilmaz H (1981) Video enhanced contrast polarization (AVEC - POL) microscopy. *Cell Motility* 1, 275-89
- Altman R (1890) *Die Elementarorganism Und Ihre Beziehungen Zu Den Zellen*. Veit, Leipzig, figure 4
- Alzheimer A (1904) Histologische Studien zur Differenzialdiagnose der progressiven Paralyse. *Histol Histopathol Arb Grosshirnrinde* 1, 18-314

REFERENCES

- Alzheimer A (1910) Beitrage zur Kenntnis der pathologischen Glia und ihre Beziehungen zu den Abbauprodukten im Nervengewebe, in Histolog u histopatholog Arbeiten. Published by Nissl F and Alzheimer A, 3, part 3
- Andreoli T E, Hoffman J F and Fanestil D D (eds.) (1978) Membrane Physiology. Plenum, New York
- Andres K H (1961) Untersuchungen über die Feinbau von Spinalganglien. Z Zellforsch Mikrosk Anat 55, 1-48
- Andres K H (1964) Mikropinozytose um Zentralnerven-system. Z Zellforsch 64, 63-73
- Aoki K and Munoz-Martinez E J (1981) Quantitative changes in myelin proteins in a peripheral neuropathy caused by Tullidora. J Neurochem 36, 1-8
- Auerbach L (1898) Nervenendigung in Centralorganen. Neurol Zentralbl 17, 445-54
- Autilio L A, Norton W T and Terry R D (1964) The preparation and some properties of purified myelin from the central nervous system. J. Neurochem 11, 17-27
- Bahr G F, and Zeitler E (1965) Determination of magnification in electron microscopy. II Means of determination of magnification. Lab Invest 14, 142/880-153/891
- Bailey P and von Bonin G (1951) The Isocortex of Man. University of Illinois Press, Urbana, Ill., p. 239
- Bailey P and Cushing H (1926) A Classification of the Tumours of the Glioma Group On a Histogenetic Basis with a Correlated Study of Prognosis. Lippincott, Philadelphia
- Bairati A (1948) Osservazioni comparate sulle glioarchitectonica. Mem Accad Soc Ist. (Bologna) 9, 613
- Bairati A (1958) Perivascular relationship of the neuroglia cells. In Biology of Neuroglia. ed. Windle W F. Thomas C C, Springfield, Ill., pp.85-98
- Bakay L and Hague I U (1964) Morphological and chemical studies in cerebral edema. I. Cold induced edema. J Neuropathol 23, 393-418
- Baker J R (1942) Plate in Bourne (1942)
- Baker J R (1948-53) The cell theory: a restatement, history and critique. Q J, Microsc Sci 89, 103-25; 90, 87-108; 93, 157-90; 94, 407-40
- Baker J R (1958) Principles of Biological Microtechnique. Methuen, London, pp. 66-88
- Baker P F, Hodgkin A L and Shaw T I (1962a) Replacement of axoplasm of giant nerve fibres with artificial solutions. J Physiol 164, 330-37
- Baker P F, Hodgkin A L and Shaw T I (1962b) The effect of changes in internal ionic concentrations on the electrical properties of perfused giant axons. J Physiol 164, 355-74
- Balazs R, Regan C, Meier E, Woodhams P L, Wilkins G P, Patel N J and Gordon R D (1980) Biochemical properties of neural cell types from rat cerebellum in tissue culture. In Neurobiology. eds. Giacobini E, Vernadakis A and Shahar A. Raven Press, New York, pp. 155-68
- Bangham A D and Horne R W (1964) Negative staining of phospholipids and their structural modification by surface active agents as observed in the electron microscope. J Mol Biol 8, 660-68
- Banik N L and Davison A N (1969) Enzyme activity and composition of myelin and subcellular fractions in the developing rat brain. Biochem J 115, 1051-62
- Barchas J D, Akil H, Elliott G R, Holman R B and Watson S J (1978) Behavioral neurochemistry: neuroregulators and behavioral states. Science 200, 964-73
- Barka T and Anderson P J (1963) Histochemistry, Theory, Practice and Bibliography. Hoeber Medical Division, Harper and Row, New York
- Barker L F (1899) The Nervous System and its Constituent Neurons. Appleton, New York, pp. 78-157
- Barlow C F, Domek N S, Goldberg M A and Roth L J (1961) Extracellular brain space measured by S³⁵ sulphate. Arch Neurol Chicago 5, 102-10
- Barnard R I (1940) Experimental changes in end-feet of Held-Auerbach in the spinal cord of the cat. J Comp Neurol 73, 235-265
- Barr M L (1939) Some observations on the morphology of the synapse in the cat's spinal cord. J Anat, Lond 74, 1-11
- Bartlett P F, Noble M D, Pruss R M, Raff M C, Rattray S and Williams C A (1981) Rat neural antigen-2 (Ran-2): A cell surface antigen on astrocytes, ependymal cells, Muller cells and lepto-meninges defined by a monoclonal antibody. Brain Res 204, 339-51
- Bastian H C (1913) The Nature and Origin of Living Matter. Fisher Unwin, London
- Bauer K F A (1953) Organisation des Nervengewebes und Neureencytumtheorie. Urban and Schwarzenberg, Munich, figure 74
- Baumeister W (1978) Biological horizons in molecular microscopy. Cytobiologie 17, 246-97

REFERENCES

- Baumeister W, Fringeli U P, Hahn M, Kopp F and Seredynski J (1976) Radiation damage in tripalmitin layers studied by means of infrared spectroscopy and electron microscopy. *Biophys J* 16, 791-810
- Baumeister W and Vogell W (eds.) (1980) *Electron Microscopy at Molecular Dimensions*. Springer-Verlag, Berlin, pp. 1-353
- Bayliss L E (1959) *Principles of General Physiology*. Vol 1. The Physico-Chemical Background. Longman, London, p. 449
- Beams H W and Kessel R G (1968) The Golgi apparatus: structure and function. *Int Rev Cytol* 23, 209-76
- Beams H W, Van Breeman Y, Newfang D M and Evans T C (1952) A correlated study of the spinal ganglion cells and associated nerve fibres with light and electron microscopy. *J Comp Neurol* 96, 249-281
- Beevor C E and Horsley V A (1890a) An experimental investigation into the arrangement of the excitable fibres of the internal capsule of the bonnet monkey (*Macacus sinicus*). *Phil Trans R Soc B* 181, 49-88
- Beevor C E and Horsley V A (1890b) A record of the results obtained by electrical excitation of the so-called motor cortex and internal capsule in an orang-outan (*Simia satyrus*). *Phil Trans R Soc B* 181, 129-58
- Benda P (1978) Rodent glial cell lines. In *Dynamic Properties of Glia Cells*. eds. Schoffeniels E, Franck G, Hertz L and Tower D P. Pergamon Press, Oxford, pp. 67-81
- Bergman R (1926) Ueber Hortegazellen und Neurophagie. *Ned Tijdschr Geneesk* 70, 2064
- Berl S, Clarke D D and Schneider D (eds.) (1975) *Metabolic Compartmentation and Neurotransmission*. Plenum Press, New York
- Berl S, Nicklas W J and Clarke D D (1978) Glial cells and metabolic compartmentation. In *Dynamic Properties of Glia Cells*. eds. Schoffeniels E, Franck G, Hertz L and Tower D B. Pergamon Press, Oxford, pp. 143-49
- Besta C (1929) Dati sul reticolo periferico della cellula nervosa, sulla rete interstiziale diffusa e sulla loro probabile derivazione da particolari elementi cellulari. *Boll Soc Ital Biol Sper* 3, 966
- Bevan-Lewis W (1889) *A Textbook of Mental Diseases With Special Reference to The Pathological Aspects of Insanity*. Griffin, London, p. 79
- Bevan-Lewis W and Clarke H (1878) The cortical lamination of the motor area of the brain. *Proc R Soc B* 27, 38-49
- Bidwell L A and Sherrington C S (1893) Focal epilepsy: trephining and removal of small haemorrhagic focus: no improvement; removal of part of leg centre after electrical stimulation; improvement. *Br Med J* 2, 988-9
- Bielschowsky M (1904) Die Silberimpregnation der Neurofibrillen. *J Psychol Neurol* 3, 169-89
- Bignami A, Eng L F, Dahl D and Uyeda C T (1972) Localisation of the glial fibrillary acidic protein in astrocytes by immunofluorescence. *Brain Res* 43, 429-35
- Bignami A, Kozak L P and Dahl D (1980) Molecular markers for the analysis of neural differentiation in culture. In *Tissue Culture in Neurobiology*. eds. Giacobini E, Vernadakis A and Shahar A. Raven Press, New York, pp. 63-73
- Birks R, Huxley H E and Katz B (1960) The fine structure of the neuromuscular junction of the frog. *J Physiol* 150, 134-44
- Blackstad T W (1963) Ultrastructural studies on the hippocampal region. In *Progress in Brain Research*. ed. Schade J P. Elsevier, Amsterdam
- Blackstad T W (1967) Cortical gray matter. In *The Neuron*. ed. Hyden H. Elsevier, Amsterdam, pp. 49-118
- Blackwood W and Corsellis J (eds.) (1976) *Greenfield's Neuropathology*. Edward Arnold, London, pp. 1-36
- Blackwood W, Dodds T C and Somerville J C (1970) *Atlas of Neuropathology*. 2nd edition. Livingstone, Edinburgh, pp. 79-89
- Blakemore W F (1969) Schmidt-Lantermann incisures in the central nervous system. *J Ultrastruct Res* 29, 496-8
- Blasberg R and Lajtha A (1966) Heterogeneity of the mediated transport systems of amino acid uptake in brain. *Brain Res* 1, 86-104
- Blinderman E E and Markham C H (1965) Fluorescein studies of cerebral oedema produced by a cryogenic probe. *J Neurosurg* 23, 38-44
- Bloom F E (1972) The formation of synaptic junctions in developing rat brain. In *Structure and Function of Synapses*. eds. Pappas G D and Purpura D P. Raven Press, New York, pp. 101-20
- Bloom W and Fawcett D W (1968) *A Textbook of Histology*. 9th edition. Saunders, Philadelphia, p. 325

REFERENCES

- Bloom W and Fawcett D W (1975) A Textbook of Histology. 10th edition. Saunders, Philadelphia, pp. 11-943
- Bodian D (1952) Introductory survey of neurons. Cold Spring Harbor Symp Quant Biol 17, 1-13
- Bodian D (1970) An electron microscopic characterization of classes of synaptic vesicles by means of controlled aldehyde fixation. J Cell Biol 44, 115-24
- Boeke J (1932) Nerve endings, motor and sensory. In Cytology and Cellular Pathology of the Nervous System. ed. Penfield W G. Vol 1, pp. 243-315
- Bohm A A and Davidoff M (1895) Lehrbuch der Histologie des Menschen einschliesslich der Mikroskopischen Technik. Oldenbourg, Wiesbaden
- Boll G (1874) Die Histologie und Histogenese der Nervosen Zentralorgan. Arch Psychiat 4, 1-138
- Bolles-Lee A (1885) The Microtomist's Vade Mecum. Churchill, London
- Bondareff W and Hyden H (1969) Submicroscopic structure of single neurons isolated from rabbit lateral vestibular nucleus. J Ultrastruct Res 26, 399-411
- Bornstein M B and Appel S H (1961) The application of tissue culture to the study of experimental "allergic" encephomyelitis. J Neuropathol Exp Neurol XX, 141-57
- Bouman L and Bok S T (1932) Histopathology of the Central Nervous System. Oosthoek Publ, Utrecht, plate 2
- Bourke R S and Tower D B (1966a) Fluid compartmentation and electrolytes of cat cerebral cortex in vitro. I, Swelling and solute distribution in mature cerebral cortex. J Neurochem 13, 1071-97
- Bourke R S and Tower D B (1966b) Fluid compartmentation and electrolytes of cat cerebral cortex in vitro. II, Sodium, potassium and chloride of mature cerebral cortex. J Neurochem 13, 1099-117
- Bourke R S, Greenberg E S and Tower D B (1965) Variations of cerebral cortex fluid spaces in vivo as a function of species brain size. Am J Physiol 208, 682-92
- Bourne G (ed.) (1942) Cytology and Cell Physiology. Clarendon Press, Oxford
- Bowsher D (1957) Pathways of absorption of protein from the cerebrospinal fluid: An autoradiograph study in the cat. Anat Rec 128, 23-39
- Boyce B B, Gray E G and Guillery R W (1960) A theory to account for the absence of boutons in silver preparations of the cerebral cortex. J Physiol 152, Proc Physiol Soc 3P
- Boyde A, Bailey E, Jones S J and Tamarin A (1977) Dimensional changes during specimen preparation for scanning electron microscopy, in Proceedings of the Workshop in Biological Specimen Preparation Techniques. March 1977, IIT Research Institute, Chicago, pp. 507-580
- Bradbury S (1967) The Evolution of the Microscope. Pergamon, Oxford, pp. 21-24
- Bradford H F (ed.) (1982) Neurotransmitter Interaction and Compartmentation. Plenum, New York
- Brady S T, Lasek R J and Allen R D (1982) Fast axonal transport in extruded axoplasm from squid giant axon. Science 218, 1129-31
- Brazier M A B (1951) The historical development of neurophysiology. In Handbook of Physiology. Section I, Neurophysiology. American Physiological Society, Washington, pp. 1-58
- Brierley J B (1957) Blood-brain barrier. Structural aspects. In Metabolism of Nervous System, ed. Richter D. Pergamon Press, London, pp. 121-35
- Brightman M W (1965a) The distribution within the brain of ferritin injected into cerebrospinal fluid compartments. Ependymal distribution. J Cell Biol 26, 99-123
- Brightman M W (1965b) The distribution within the brain of ferritin injected into cerebrospinal fluid compartments. II. Parenchymal distribution. Am J Anat 117, 193-219
- Brodal A (1981) Neurological Anatomy in Relation to Clinical Medicine. 3rd edition. Oxford University Press, New York, pp. 10-12
- Brodie B B, Kurz H and Schanker L S (1960) The importance of dissociation constant and lipid solubility in influencing the passage of drugs into the cerebrospinal fluid. J Pharmacol 130, 20-25
- Brodmann K (1903) Beitrage zur histologischen Lokalisation der Grosshirnrinde. I. Die Regio Rolandica. J Psychol Neurol, Lpz. 2, 79-107; 2. Der Calcarinatypus. J Psychol Neurol Lpz. 2, 133-59
- Brodmann K (1909) Vergleichende Lokalisationslehre der Grosshirnrinde In Ihren Prinzipien Dargestellt Auf Grund Des Zellenbaues. Barth, Leipzig
- Brownson R H (1955) Perineuronal satellite cells in the motor cortex of ageing brains. J Neuropathol 14, 424-32

REFERENCES

- Brownson R H (1956) Perineuronal satellite cells in the motor cortex of ageing brains. *J Neuropathol* 15, 190-95
- Brownson R H, Suter D B and Diller D A (1963) Acute brain damage induced by low dosage x-irradiation. *Neurology* 13, 181-91
- Bruce-Casselman W G (1959) *Histochemical Techniques*. Methuen, London, p. 164
- Bruggeller P and Mayer E (1981) EPR investigation of the nonfreezing water free water ratio in disperse systems. *J Phys Chem* 85, 4135-9
- Bullock T H, Orkand R and Grinnell A (1977) *Introduction to Nervous Systems*. W H Freeman and Company, San Francisco, p. 98
- Bunge R P (1968) Glial cells and central myelin sheath. *Physiol Rev* 48, 197-251
- Bunge M B, Bunge R P, Peterson E R and Murray M R (1967) A light and electron microscope study of long term organised cultures of rat dorsal root ganglia. *J Cell Biol* 32, 439-66
- Butschli O (1894) *Protoplasm and Microscopic Foams*. Adam and Charles Black, London, pp. 1-379
- Caldwell P C, Hodgkin A L, Keynes R D and Shaw T I (1960) Effects of injecting energy-rich phosphate compounds on the active transport of ions in the giant axons of Loligo. *J Physiol* 152, 561-90
- Calissano P and Bangham A D (1971) Effect of two brain specific proteins (S-100 and 14-3-2) on cation diffusion across artificial lipid membranes. *Biochem Biophys Res Commun* 43, 504-9
- Calkins E, Taylor I M and Hastings A B (1954) Potassium exchange in the isolated rat diaphragm; effect of anoxia and cold. *Am J Physiol* 177, 211-18
- Callan H, and Tomlin S G (1950) Experimental studies on the amphibian oocyte nuclei. *Proc R Soc B* 137, 367-78
- Callas G and Hild W (1964) Electron microscopic observations of synaptic endings in cultures of mammalian central nervous tissue. *Z Zellforsch Mikrosk Anat* 63, 686-691
- Cammermeyer J (1963) Similarities between oligodendrocyte and cerebellar granule cell nuclei in Mammalia and Aves. *Am J Anat* 112, 111-39
- Cammermeyer J (1966) Morphologic distinctions between oligodendrocytes and microglia cells in the rabbit cerebral cortex. *Am J Anat* 118, 227-48
- Cammermeyer J (1979) The life history of the microglial cell: a light microscopic study. In *Neurosciences Research*, Vol 3, eds. Ehrenpreis S and Solnitzky O C. Academic Press, New York, pp. 43-130
- Campbell A W (1905) *Histological Studies on The Localisation of Cerebral Function*. Cambridge University Press, Cambridge
- Campbell F R (1967) Nuclear elimination from the normoblast of fetal guinea pig liver as studied with electron microscopy and serial sectioning techniques. *Anat Rec* 160, 539-54
- Carleton H M (1926) *Histological Technique*. 1st edition, Oxford University Press, Oxford
- Carleton H M and Drury R A B (1962) *Histological Technique*. 3rd edition. Oxford University Press, London
- Carleton H M and Short R H D (eds.) (1960) *Schafer's Essentials of Histology*. 16th edition. Longmans, London
- Carleton H M, Drury R A B and Wallington E A (1980) *Histological Techniques*. 5th edition, Drury, London
- Causey G (1960) The Cell of Schwann. E and S Livingstone, Edinburgh, pp. 4-30
- Causey G (1962) *Electron Microscopy*. E and S Livingstone, Edinburgh, pp. 189-218
- Cavanaugh J B (1970) The proliferation of astrocytes around a needle wound in the rat brain. *J Anat* 106, 471-87
- Ceccarelli R and Hurlbut W P (1980) Vesicle hypothesis of the release of quanta of acetylcholine. *Physiol Rev* 60, 396-441
- Chambers R and Chambers E D (1961) *Explorations into the Living Cell*. Commonwealth Fund, Harvard University Press, Cambridge, Mass
- Chan-Palay V (1973) The cytology of neurons and their dendrites in the simple mammalian nucleus lateralis. *Z Anat Entwickl-Gesch* 141, 289-317
- Chang J J, McDowall A W, Lepault J, Freeman R, Walker C A and Dubochet J (1983) Freezing, sectioning and observation artefacts of frozen hydrated sections for electron microscopy. *J Microsc* 132, 109-23
- Chang T M S (1972) *Artificial Cells*. Thomas, Springfield, Ill., pp. 43-77
- Chou S N (1961) Experimental brain tumours in rabbits and associated cerebral oedema. *Lancet* 81, 78-82

REFERENCES

- Christensen H N and Hastings A B (1940) Phosphatides and inorganic salts. *J Biol Chem* 136, 387-98
- Clark G (ed.) (1973) Staining Procedures. 3rd edition. Williams and Wilkins, Baltimore, pp. 33-119
- Clarke D A (1931) Muscle counts of motor units: a study in innervation ratios. *Am J Physiol* 96, 296-304
- Clarke E and O'Malley C D (1968) The Human Brain and Spinal Cord. University of California Press, Berkeley, figure 5
- Clasen R A, Cooke P M, Tyler S and Pandolfi S (1960) Changes in intracerebral sodium chloride and water associated with experimental cerebral edema. *J Neuropathol Exp Neurol* 19, 164
- Clay R S and Court T H (1932) The History of the Microscope. Holland Press, London
- Clemente C D and Holst E A (1954) Pathological changes in neurons, neuroglia and blood-brain barrier induced by x-irradiation of heads of monkeys. *Arch Neurol Psychiatry* 17, 66-79
- Coers C (1967) Structure and organization of the myoneural junction. *Int Rev Cytol* 22, 239-67
- Cohen J and Selvendran S Y (1981) A neuronal cell-surface antigen is found in the CNS but not in peripheral neurones. *Nature* 291, 42-23
- Cohen R S, Wolosewick J J, Becker R P and Pappas G D (1984) Fine structure of synapses of the central nervous system in resinless sections. Personal Communication
- Cohen S R (1974a) The contribution of adherent films of liquid on the cut surfaces of mouse brain slices to tissue water and 'extracellular' marker spaces. *Exp Brain Res* 20, 421-34
- Cohen S R (1974b) The dependence of water content and 'extracellular' marker spaces of incubated mouse brain slices on thickness, alterations produced by slicing, and fluid spaces in intact and altered tissue. *Exp Brain Res* 20, 435-57
- Cohnheim J (1872) Untersuchungen über die Embolischen Processe. Hirschwald, Berlin, pp. 131-2
- Cold Spring Harbor Symposium on Quantitative Biology (1972) Structure and function of proteins at the three-dimensional level. Cold Spring Harbor Symp 36, pp. 1-644
- Cold Spring Harbor Symposium on Quantitative Biology (1975) The synapse. Cold Spring Harbor Symp 40, various papers
- Cold Spring Harbor Symposium on Quantitative Biology (1982) Organisation of the cytoplasm. Cold Spring Harbor Symp 46, most papers, 2 volumes, pp. 1-1047
- Cole K S (1937) Electric impedance of marine egg membranes. *Trans Faraday Soc* 33, 966-72
- Cole W V (1955) Motor endings in the striate muscle of vertebrates. *J Comp Neurol* 102, 671-715
- Cole W V (1957) Structural variations of nerve endings in the striated muscles of the rat. *J Comp Neurol* 108, 445-63
- Colodzin M and Kennedy E P (1964) Biosynthesis of diphosphoinositides in the brain. *J Biol Chem* 240, 3771-80
- Conel J Le Roy (1939-63) The Post Natal Development of the Human Cerebral Cortex. Vols 1-6, Harvard University Press, Cambridge, Mass
- Conradi S (1976) Functional anatomy of the anterior horn motor neuron. In The Peripheral Nerve. ed. Landon D N. Chapman and Hall, London, pp. 279-354
- Constantinides P (1974) Functional Electronic Histology. Elsevier Scientific Publishing Co., Amsterdam
- Constantino-Ceccarini E, Castelli A and De Vries G H (1979) Characterization and developmental changes of UDP-galactose-ceramide galactosyl transferase in a rat CNS axolemma-enriched fraction. Differences and similarities of the enzyme associated with the microsomal and myelin fractions. *J Neurochem* 32, 1175-82
- Conway E J (1955) Evidence for a redox pump in active transport of cations. *Int Rev Cytol* 4, 377-96
- Conway E J (1957) Nature and significance of concentration relations of potassium and sodium ions in skeletal muscle. *Physiol Rev* 37, 84-132
- Cook D J (1983) Optical illusions in microscopy. *Microscopy* 34, 533-52
- Coombs J S, Eccles J C and Fatt P (1955) The specific ionic conductances and ionic movements across the motoneuronal membrane that produce the inhibitory post synaptic potential. *J Physiol* 130, 326-73
- Cosslett V E (1982) Resolution and contrast in the electron microscope: an historical review. *J. Microsc* 128, 23-31
- Costero I and Pomerat C M (1951) Cultivation of neurons from the adult human cerebral and cerebellar cortex. *Am J Anat* 89, 405-68

REFERENCES

- Cottrell G A and Usherwood N R (eds.) (1977) *Synapses*. Blackie, Glasgow, pp. 1-384
- Couteaux R (1958) Morphological and cytochemical observations on the post-synaptic membrane at motor end-plates and ganglionic synapses. *Exp Cell Res*, Academic Press, New York, Suppl 5, 292-322
- Couteaux R (1960) Motor end-plate structure. In *Structure and Function of Muscle*, ed. Bourne G H. Vol 1, Academic Press, New York, pp. 337-80
- Cox W (1891) Impregnation des centralen Nervensystems mit Quecksilbersalzen. *Arch Mikrosk Anat* 37, 16-21
- Cragg B G (1967) The density of synapses and neurones in the motor and visual areas of the cerebral cortex. *J Anat* 101, 639-54
- Cragg B G (1972) The development of synapses in cat visual cortex. *Invest Ophthalmol* 11, 377-85
- Crain S M (1956) Resting and action potentials of cultured chick embryo spinal ganglion cells. *J Comp Neurol* 104, 285-330
- Crain S M (1976) *Neurophysiological Studies in Tissue Cultures*. Raven Press, New York, pp. 1-280
- Crain S M, Peterson E R, and Bornstein M B (1968) Formation of functional inter-neuronal connection between explants of various mammalian central nervous tissues during development in vitro. In *Ciba Foundation Symposium, Growth of the Nervous System*. eds. Wolstenholme G E W and O'Connor, M. Churchill, London, pp. 13-31
- Creutzfeldt H G and Metz A (1926) Ueber Gestalt und Tätigkeit der Hortegazellen bei pathologischen Vorgangen. *Z Gesamte Neurol Psychiatr* 106, 18
- Crile G W, Hosmer H R and Rowland A F (1922) The electrical conductivity of animal tissue under normal and pathological conditions. *Am J Physiol* 60, 59-106
- Crone C (1961) Om Diffusionen av Nogle Organiske Non Electrolyter Fra Blod Til Hjernevaev. Munksgaard, Copenhagen, p. 180
- Crone C (1965) Facilitated transfer of glucose from blood into brain tissue. *J Physiol* 181, 103-13
- Cruickshank B, Dodds T C and Gardner D L (1968) *Human Histology*. E and S Livingstone Ltd, Edinburgh, pp. 240-61
- Cuatrecasas P and Hollenberg M D (1975) Binding of insulin and other hormones to nonreceptor materials. Saturability, specificity and apparent "negative" cooperativity. *Biochem Biophys Res* 62, 31-41
- Cuatrecasas P, Hollenberg M D, Chang K-J and Bennett V (1975) Hormone receptor complexes and their modulation of membrane function. *Rec Prog Horm Res* 31, 37-94
- Culling C F A (1975) *Handbook of Histopathological and Histochemical Techniques*. Butterworths, London, pp. 1-669
- Cummins J E and Hyden H (1962) Adenosine triphosphatase levels and adenosine triphosphatas in neurons, glia and neuronal membranes of the vestibular nucleus. *Biochim Biophys Acta* 60, 271-83
- Currie D N (1980) Identification of cell type by immunofluorescence in defined cell cultures of cerebellum. In *Tissue Culture in Neurobiology*. eds. Giacobini E, Vernadakis A and Shahar A. Raven Press, New York, pp. 75-87
- Currie D N and Kelly J S (1981) Glial versus neuronal uptake of glutamate. *J Exp Biol* 95, 181-93
- Curtis B A, Jacobson S and Marcus E M (1972) *An Introduction to the Neurosciences*. Saunders, Philadelphia, pp. 53-66
- Dahlstrom A and Fuxe K (1965a) Evidence for the existence of monoamine containing neurons in the central nervous system. I. Demonstration of monoamines in the cell bodies of brain stem neurons. *Acta Physiol Scand (Suppl)* 232, 1-55
- Dahlstrom A and Fuxe K (1965b) Evidence for the existence of monoamine neurons in the central nervous system. II. Experimentally induced changes in the intraneuronal amine levels of bulbospinal neuron system. *Acta Physiol Scand (Suppl)* 247, 1-36
- Dancer J B (1840), quoted by Henderson H S (1978) Presidential Address. *Microscopy* 33, 257-63
- Danielli J F (1942) The cell surface and cell physiology. In *Cytology and Cell Physiology*. ed. Bourne G, Clarendon Press, Oxford, pp. 68-98
- Danowski T S (1941) The transfer of potassium ions across the human blood cell membrane. *J Biol Chem* 139, 693-705
- Dartnall D W and Klotz I M (1976) Subunit constitution of proteins, in *Handbook of Biochemistry and Molecular Biology*. 3rd edition. Proteins II, ed. Fasman G D. CRC Press, Cleveland, pp. 325-37
- Darwin C (1875) *Insectivorous Plants*. John Murray, London, pp. 413-15

REFERENCES

- Davson H (1955) A comparative study of aqueous fluid and cerebrospinal fluid in the rabbit. *J Physiol* 129, 111-33
- Davson H (1967) Physiology Of The Cerebrospinal Fluid. Churchill, London, pp. 262-70
- Davson H and Danielli J F (1936) The Permeability of Natural Membranes. 1st edition. Cambridge University Press, Cambridge
- Davson H and Danielli J F (1943) The Permeability of Natural Membranes. 2nd edition. Cambridge University Press, Cambridge
- Davson H and Eggleton M G (1968) Starling Lovatt Evans Principles of Human Physiology. 14th edition, Churchill, London, p. 1170
- Davson H and Pollay M (1963) The turnover of ^{24}Na in the cerebrospinal fluid and its bearing on the blood-brain barrier. *J Physiol* 167, 247-55
- Davson H and Spaziani E (1959) The blood-brain barrier. *J Physiol* 149, 135-43
- De Brabander M and de Mey J. eds. (1980) Microtubules and Microtubule Inhibitors. Elsevier/North Holland, Amsterdam
- De Duve C (1963) The lysosome concept. In Lysosomes. eds. De Reuck A S V and Cameron M P. Churchill, London, pp. 1-35
- Deiters O F C (1865) Untersuchungen über Gehirn und Ruckenmark des Menschen und der Saugethiere. Vieweg F u Sohn, Braunschweig, pp. 1-318
- del Castillo J and Katz B (1956) Biophysical aspects of neuromuscular transmission. In Progress in Biophys. Biophys Chem 6, ed. Butler J A V. Pergamon, London, pp. 122-170
- Del Rio-Hortega P (1917) Noticia de un nuevo y facil metodo para la coloracion de la neuroglia y del tejido conjuntivo. *Trab Lab Invest Biol Univ Madrid* 15, 367-78
- Del Rio-Hortega P (1919a) El 'tercer elemento' de los centros nerviosos. *Bol Soc Esp Biol* 9, 69-120
- Del Rio-Hortega P (1919b) Poder fagocitario y movilidad de la microglia. *Bol Soc Esp Biol* 9, 154-67
- Del Rio-Hortega P (1920) La microglia y sus transformacion en celulas en bastoncito y cuerpos granuloadiosos. *Trab Lab Invest Biol Univ Madrid* 18, 37-82
- Del Rio-Hortega P (1928) Tercera aportacion al conocimiento morfologico del interpretacion functional de la oligodendroglia. *Mem Real Soc Esp Hist Nat* 14, 1-122
- Del Rio-Hortega P (1932) Microglia. In Cytology and Cellular Pathology of the Nervous System. Vol 2, ed. Penfield W. Hoeber, New York, pp. 482-534
- Dempsey E W and Luse S (1958) Fine structure of the neuropil in relation to neuroglia cells. In Biology of Neuroglia. ed. Windle W F. C Thomas, Springfield, Ill., pp. 99-129
- De Robertis E (1959) Submicroscopic morphology of the synapse. *Int Rev Cytol* 8, 61-96
- De Robertis E (1963) Ultrastructure and cytochemistry of the synaptic region. *Science* 156, 907-14
- De Robertis E and Bennet H S (1955) Some features of the sub-microscopic morphology of synapses in frog and earthworm. *J Biophys Biochem Cytol* 1, 47-58
- De Robertis E, Gerschenfeld H M and Wald F (1960) Ultrastructure and function of glial cells. In Structure and Function of the Cerebral Cortex. eds. Tower D B and Van Schade J. Elsevier, Amsterdam, pp. 69-79
- Deutsch K (1973) An electron microscopical study of the cerebral cortex of the calf. *Zbl Vet Med A*, 20, 692-8
- Deutsch K and Hillman H (1977) The effect of six fixatives on the areas of rabbit neurons and rabbit and rat cerebral slices. *J Microsc* 110, 303-9
- De Vieussens R (1684) Neurographia Universalis. Certe, Lyons.
- Dimpfel W, Neale J H and Haberman E (1975) ^{125}I -labelled tetanus toxin as a neuronal marker on tissue cultures derived from embryonic CNS. *Naunyn Schmiedebergs Arch Exp Pathol Pharmakol* 290, 329-33
- Dittman L, Sensenbrenner M, Hertz L and Mandel P (1973) Respiration by cultivated astrocytes and neurons from the cerebral hemispheres. *J. Neurochem* 2, 191-8
- Dixon M and Webb E C (1964) Enzymes. 2nd edition. Longmans, London, pp. 623-36
- Dodgson M (1948) Physical nature of mouse brain. *Nature* 162, 253
- Driest E and Muller H O (1935) Elektronmikroskopische Aufnahmen (Elektronenmikrogramme) von Chitinobjekten. *Z Wiss Mikrosk* 52, 53-7
- Drury R A B and Wallington E A (1967) (eds.) Carleton's Histological Technique. 4th edition. Oxford University Press, New York, pp. 1-401
- Dubochet J, Lepault J, Freeman R, Berriman J A and Homo J C (1982) Electron microscopy of frozen water and aqueous solution. *J Microsc* 128, 219-37
- Du Bois-Reymond (1899) quoted in Eccles J C (1964) The Physiology of Synapses. Springer, Berlin, p. 7

REFERENCES

- Dupouy G (1972) Performance and applications of the Toulouse 3 million volt electron microscope. *J Microsc* 97, 3-28
- Dusser de Barenne J G (1910) Die segmentare Strychninvergiftung der dorsalen Ruckenmarksmechanismen; ein Beitrag zur Dermatometrie der hinteren Extremitaten des Hundes. *Folia Neuro-biol, Lpz*, 5, 342-59
- Dusser de Barenne J G (1916) Experimental researches on sensory localizations in the cerebral cortex. *Q J Exp Physiol* 9, 355-90
- Dustin P (1978) *Microtubules*. Springer, Berlin, pp. 1-452
- Dutta C R, Siegesmund K A and Fox C A (1963) Light and electron microscopic observations of an intranucleolar body in nerve cells. *J Ultrastruct Res* 8, 542-51
- Ebner F F and Colonnier M (1975) Synaptic patterns in the visual cortex of turtle. An electron microscopic study. *J Comp Neurol* 160, 51-80
- Eccles J C (1957) *The Physiology of Nerve Cells*. Johns Hopkins Press, Baltimore, pp. 218-19
- Eccles J C (1961) The mechanism of synaptic transmission. *Ergebn Physiol* 51, 299-430
- Eccles J C (1964) *The Physiology of Synapses*. Springer, Berlin, pp. 1-26
- Edstrom J E (1958) Quantitative determination of ribonucleic acid in the micromicrogram range. *J Neurochem* 3, 100-6
- Ehrenberg C G (1833) Nottwendigkeit einer feineren mechanischen Zerlegung des Gehirns und der Nerven vor der chemischen, dargestellt aus Beobachtungen von C G Ehrenberg. *Poggendorffs Ann Phys* 28, 449-73
- Eichberg J, Whittaker V P and Dawson R M C (1964) The distribution of lipids in subcellular particles of guinea-pig brain. *Biochem J* 92, 91-100
- Elvin L-G (1961a) The ultrastructure of the nodes of Ranvier in cat sympathetic nerve fibres. *J Ultrastruct Res* 5, 374-87
- Elvin L-G (1961b) Electron microscopic investigation of the plasma membrane and myelin sheath of autonomic nerve fibres in the cat. *J Ultrastruct Res* 5, 388-407
- Elvin L-G (1968) The structure and composition of motor, sensory and autonomic nerves and nerve fibres. In *The Structure and Function of Nervous Tissue*. ed. Bourne G H, Vol 1, Academic Press, New York, pp. 325-77
- Elliott H C (1969) *Textbook of Neuroanatomy*. J B Lippincott, Philadelphia
- Elliott T R (1904) The action of adrenaline. *J Physiol* 31, 20-21P
- Emmelot P and Benedetti E L (1960) Changes in the fine structure of rat liver cells brought about by dimethylnitrosamine. *J Biophys Biochem Cytol* 7, 393-6
- Emmelot P and Benedetti E L (1961) Some observations on the effect of liver carcinogens on the fine structure and function of the endoplasmic reticulum of rat liver cells, in *Protein Biosynthesis*. ed. Harris R J C. Academic Press, New York, pp. 99-123
- Eng L F, Chase F C, Gerstl B, Pratt D and Tavastjerna M G (1968) The maturation of human white matter myelin. Fractionation of the myelin membrane proteins. *Biochemistry* 7, 4455-6
- Eng L F, Vanderhaeghen J T, Bignami A and Gerstl B (1971) An acidic protein isolated from fibrous astrocytes. *Brain Res* 28, 351
- Entin T (1954) A study of synapses in the visual area of the cerebral cortex. *Arch Anat Histol Embryol, Leningrad*, 31, 25
- Epstein M H and O'Connor J S (1965) Respiration of single cortical neurons and surrounding neuropile. *J Neurochem* 12, 389-95
- Erlanger J and Gasser H S (1937) *Electrical Signs of Nervous Activity*. University of Pennsylvania Press, Philadelphia, pp. 1-221
- Estable C and Sotelo J R (1951) Una nueva estructura celular: el nucleolonema. *Publ Inst Invest Sci Biol* 1, 105-23
- Estable-Puig J F, De Estable R F, Tobias C and Haymaker W (1964) Degeneration and regeneration of myelinated fibers in the cerebral and cerebellar cortex following damage from ionizing particle radiation. *Acta Neuropathol* 4, 175-90
- Etcheverry G J and De Iraldi A P (1968) Ultrastructure of neurones in the arcuate nucleus of the rat. *Anat Rec* 160, 239-53
- Eto Y and Suzuki K (1973) Cholesterol ester metabolism in rat brain, a cholesterol ester hydrolase specifically localized in the myelin sheath. *J Biol Chem* 248, 1986-91
- Evans W H (1974) Nucleotide pyrophosphatase, a sialoglycoprotein located on the hepatocyte surface, *Nature* 250, 391-4
- Farquhar M C and Hartmann J F (1957) Neuroglial structure and relationships as revealed by electron microscopy. *J Neuropathol Exp Neurol* 16, 18-39

REFERENCES

- Fatt P and Katz B (1952) Spontaneous subthreshold activity at motor nerve endings. *J Physiol* 117, 109-28
- Favard P and Carasso N (1972) The preparation and observation of thick biological sections in the high voltage electron microscope. *J Microsc* 97, 59-82
- Fawcett D W (1981) *The Cell*. Saunders, Philadelphia, pp. 30-456
- Fedoroff S (1978) The development of glial cells in culture. In *Dynamic Properties of Glial Cells*. eds. Schoffeniels E, Franck G, Hertz L and Tower D B. Pergamon Press, Oxford, pp. 83-92
- Fedoroff S and Hertz L (1977) *Cell, Tissue and Organ Cultures in Neurobiology*. Academic Press, New York, pp. 1-689
- Fernandez-Moran H (1950a) Electron microscope observations on the structure of the myelinated nerve fiber sheath. *Exp Cell Res* 1, 143-9
- Fernandez-Moran H (1950b) Sheath and axon structures in the internode portion of vertebrate myelinated nerve fibres. *Exp Cell Res* 1, 309-40
- Fernandez-Moran H (1952) The submicroscopic organisation of vertebrate nerve fibres. *Exp Cell Res* 3, 282-59
- Fernandez-Moran H (1957) Electron microscopy of the nervous tissue. In *Metabolism of the Nervous System*. ed. Richter D. Pergamon Press, London, pp. 1-34
- Fernandez-Moran H (1962) Correlation of electron microscopy and X-ray diffraction data in ultrastructural studies of lipoprotein membrane systems. In *Symposia of The International Society for Cell Biology*. ed. Harris R J C. Academic Press, New York, pp. 411-28
- Fernandez-Moran H (1967) Membrane ultrastructure in nerve cells. In *Neurosciences*. eds. Quarton G C, Melnechuk T and Schmitt F O. Rockefeller University Press, New York, pp. 281-304
- Fernandez-Moran H and Finean J B (1957) Electron microscope and low angle x-ray diffraction studies of the nerve myelin sheath. *J Biophys Biochem Cytol* 3, 725-48
- Ferrier D (1876) *The Functions of the Brain*. Smith Elder, London
- Field E J (1957) Histogenesis of compound granular corpuscles in the mouse brain after trauma and a note on the influence of cortisone. *J Neuropathol Exp Neurol* 16, 48-56
- Fine R E and Bray D (1971) Actin in growing nerve cells. *Nature, London, New Biol*, 234, 115-18
- Finean J B (1954) The effects of osmium tetroxide fixation on the structure of myelin in sciatic nerve. *Exp Cell Res* 6, 283-92
- Finean J B (1957) The role of water in the structure of peripheral nerve myelin. *J Biophys Biochem Cytol* 3, 95-102
- Finean J B (1960) Electron microscope and X-ray diffraction studies of the effect of dehydration on the structure of nerve myelin. 11. Optic nerve. *J Biophys Biochem Cytol* 8, 13-37
- Finean J B (1961) *Chemical Ultrastructure In Living Tissues*. C C Thomas, Springfield, Ill., p. 111
- Finkelstein A (1978) Lipid bilayer membranes. Their permeability properties as related to those of cell membranes. In *Membrane Physiology*. eds. Andreoli T E, Hoffman J E and Fanestil D D. Raven, New York, pp 205-16
- Fischbach G D and Nelson P G (1977) Cell culture in neurobiology. In *Handbook of Physiology*. ed. Kandel E R. Sec 1, Vol 1, Am Physiol Soc, Bethesda, pp. 719-774
- Fisher R A (1935) *The Design of Experiments*. Oliver and Boyd, Edinburgh, pp. 1-252
- Fishman R A (1964) Carrier transport of glucose between blood and cerebrospinal fluid. *Am J Physiol* 206, 836-44
- Flechsig P (1876) *Die Leitungsbahnen in Gehirn und Rückenmark des Menschen auf Grund Entwicklungsgeschichtlicher Untersuchungen*. Englemann, Leipzig, pp. 1-382
- Flechsig P (1881) Zur Anatomie und Entwicklungsgeschichte der Leitungsbahnen im Grosshirn des Menschen. *Arch Anat Physiol, Anat Abt*, pp. 12-75
- Fleischer S, Fleischer B and Stoeckenius B (1967) Fine structure of lipid depleted mitochondria. *J Cell Biol* 32, 193-208
- Flemming W (1882) *Zellsubstanz, Kern und Zellteilung*. Vogel, Leipzig.
- Flynn F and Maizels M (1949) Cation control in human erythrocytes. *J Physiol* 110, 301-18
- Fonnum F (ed.) (1978) *Amino Acids as Chemical Transmitters*. Raven Press, New York
- Fontana F (1781) *Traité sur le Venin de la Vipere*. Florence, Vol 2, pp. 207-8
- Fontana F (1784) Lettre de M. L'Abbe Fontana à M. Gibelin, à Aix en Provence, datée de Florence du 10 juillet, 1782. *J Physique* 24, 417-21
- Forel A H (1887) Eine hirnanatomische Betrachtungen und Ergebnisse. *Arch Psychiat Nervkrankh* 18, 162-98

REFERENCES

- Foster M and Sherrington C S (1897) A Textbook of Physiology. 7th edition. Macmillan, London, Vol III, p. 929
- Frank K and Fuortes M G F (1955) Potentials recorded from the spinal cord with micro-electrodes. *J Physiol* 130, 625-54
- Frank K and Fuortes M G F (1956) Stimulation of spinal motoneurons with intracellular electrodes. *J Physiol* 134, 451-70
- Franke W W (1974) Nuclear envelopes. *Phil Trans R Soc Lond B* 268, 67-93
- Franke W W, Schmid E, Osborn M and Weber K (1978) Different intermediate sized filaments distinguished by immunofluorescence microscopy. *Proc Natl Acad Sci, USA* 75, 5034-8
- Frey T G and Chan S H P (1980) The structure of cytochrome c-oxidase. Immunolectron microscopy and image analysis of two dimensional crystals. In *Electron Microscopy at Molecular Dimensions*. eds. Baumeister W and Vogell W. Springer, Berlin, pp. 81-90
- Freygang W H and Landau W M (1955) Some relations between resistivity and electrical activity in the cerebral cortex of the cat. *J Comp Physiol* 45, 377-92
- Fricke H (1925) The electric capacity of suspensions with special reference to blood. *J Gen Physiol* 9, 137-52
- Fricke H and Curtis H J (1935) The dielectric constant and resistance of colloidal solutions. *Physiol Rev* 47, 974-5
- Fricke H and Morse S (1925) The electrical resistance and capacity of blood for frequencies between 800 and 47½ million cycles. *J Gen Physiol* 9, 153-167
- Friede R L (1965) Enzyme histochemistry of neuroglia. In *Progress in Brain Research*, 15, eds. de Robertis E D P and Carrea R, Elsevier, Amsterdam, pp. 35-47
- Friede R L and Samorajski T (1970) Axon caliber related to neurofilaments and microtubules in sciatic nerve fibers of rats and mice. *Anat Rec* 167, 379-87
- Fritsch G and Hitzig E (1870) Ueber die elektrische Erregbarkeit des Grosshirns. *Arch Anat Physiol Wiss Med* 37, 300-32
- Fulton J F (1949) *Physiology of the Nervous System*. 3rd edition. Oxford University Press, New York, pp. 36-53
- Gallyas F (1963) Silver impregnation method for microglia. *Acta Neuropathol* 3, 206-9
- Garrison F H (1924) *An Introduction to the History of Medicine*. 3rd edition. Saunders, Philadelphia
- Gasser H S (1958) Comparison of the structure, as revealed with the electron microscope, and the physiology of the unmedullated fibers in the skin nerves and in the olfactory nerves. *Exp Cell Res Suppl* 5, Academic Press, New York, pp. 3-17
- Gatenby J and Painter T S (1937) *The Microtomist's Vade-Mecum*. 10th edition, Churchill, London, pp. 150-562
- Gauthier G F (1976) The motor end-plate: structure. In *The Peripheral Nerve*. ed. Landon, D N. Chapman and Hall, London, pp. 464-94
- Geiger R S (1957) In vitro studies on the growth properties of brain cortex cells of adult individuals. In *Ultrastructure and Cellular Chemistry of Neural Tissue*. ed. Waelsch H. Vol II, Cassell, London, pp. 83-99
- Gelfan S (1964) Neuronal interdependence. In *Organization of The Spinal Cord*, eds. Eccles J C and Schade J P. *Progress in Brain Research*. Elsevier, Amsterdam, 2, pp. 238-60
- Gent W L G, Gregson N A, Gammack D B and Raper J H (1964) The lipid-protein unit in myelin. *Nature*, London, 204, 553-5
- Geren B B (1954) The formation from the Schwann cell surface of myelin in the peripheral nerves of chick embryos. *Exp Cell Res* 7, 558-62
- Gerlach J von (1858) *Mikroskopische Studien aus dem Gebiete der Menschlischen Morphologie*. Enke, Erlangen
- Gerlach J von (1872a) Über die grauen Substanz des menschlichen Grosshirns vorläufige Mittheilung. *Zbl Med Wiss* 10, 273-5
- Gerlach J von (1872b) The spinal cord. In Stricker S (1870-73) *Manual of Human and Comparative Histology*. New Sydenham Society, London, 3 vols
- Gerschenfield H M, Wald F, Zadunaisky J A and De Robertis E D P (1959) Function of astroglia in the water-ion metabolism of the central nervous system. An electron microscope study. *Neurology* 9, 412-25
- Gersh I (1959) Fixation and staining. In *The Cell*. eds. Brachet J and Mirsky A E. Academic Press, New York, 1, 21-60
- Ghandour M S, Derer P, Labourdette G, Delaunoy J P and Langley O K (1981) Glial cell markers in the reeler mutant mouse: a biochemical and immunohistological study. *J Neurochem* 36, 195-200

REFERENCES

- Gheurens J, Lowenthal A, Karcher D and Noppe M (1978) Similarities between the nervous tissue proteins α -albumin and glial fibrillary acidic protein. In Dynamic Properties of Glia Cells. eds. Schoffeniels E, Franck G, Hertz L and Tower D B. Pergamon Press, Oxford, pp. 257-66
- Giacobini E (1957) The intracellular distribution of cholinesterase in the nerve cell studied by means of a quantitative micromethod. *Acta Physiol Scand* 42, Suppl 145, 1-49
- Giacobini E (1961) Localization of carbonic anhydrase in nervous system. *Science* 134, 1524-5
- Giacobini E, Vernadakis A and Shahar A eds. (1980) Tissue Culture in Neurobiology. Raven Press, New York, pp. 349-426
- Gibson W C (1940) Degeneration and regeneration of sympathetic synapses. *J Neurophysiol* 3, 237-47
- Gilbert J C (1965) Mechanism of sugar transport in brain slices. *Nature*, London, 205, 87-8
- Giuditta A and Strecker H J (1959) Alternate pathways of pyridine nucleotide oxidation in cerebral tissue. *J Neurochem* 5, 50-61
- Glaser G H (1964) Sodium and seizures. *Epilepsia* 5, 97-111
- Glauert A M (1979) Recent advances of high voltage electron microscopy in biology. *J Microsc* 117, 93-101
- Glees P (1944) The anatomical basis of cortico-striate connexions. *J Anat Lond* 78, 47-51
- Glees P (1946) Terminal degeneration within the central nervous system as studied by a new silver method. *J Neuropathol Exp Neurol* 5, 54-9
- Glees P (1955) Neuroglia, Morphology and Function. Blackwell, Oxford
- Glees P (1973) The neuroglial compartments at light microscopic and electron microscopic levels. In Metabolic Compartmentation in the Brain. eds. Balazs R and Cremer J E. Macmillan, London, pp. 209-31
- Glees P and Hasan M (1973) Neuronal soma-somatic contacts in the lateral geniculate body of the monkey in relation to its synaptic organization. *Acta Anat* 84, 541-51
- Glees P and Le Gros Clark W E (1941) The termination of optic fibres in the lateral geniculate body of the monkey. *J Anat*, London, 75, 295-309
- Glees P and Meller K (1968) Morphology of neuroglia. In The Structure and Function of Nervous Tissue. ed. Bourne G H. Vol 1, Structure 1, Academic Press, New York, pp. 301-23
- Glynn I M (1957) The ionic permeability of the red cell membrane. *Prog Biophys Biophys Chem*, eds. Butler J A V and Katz B, 8, 241-307
- Goldfischer S, Essner E and Novikoff A B (1964) The localization of phosphatase activities at the level of ultrastructure. *J Histochem Cytochem* 12, 72-95
- Goldman R, Pollard T and Rosenbaum J eds. (1976a) Cell Motility. Book A. Motility of Muscle and Non-muscle cells. Cold Spring Harbor Conf on Cell Proliferation. Vol 3
- Goldman R, Pollard T and Rosenbaum J eds. (1976b) Cell Motility. Book B, Actin, Myosin and Associated Proteins. Cold Spring Harbor Conf on Cell Proliferation. Vol 3
- Goldman R, Pollard T and Rosenbaum J eds. (1976c) Cell Motility. Book C. Microtubules and Related Proteins. Cold Spring Harbor Conf on Cell Proliferation. Vol 3
- Goldstein A, Aronow L and Kalman S M (1968) Principles of Drug Action. The Basis of Pharmacology. Hoeber Medical Division, Harper and Row, New York, pp. 1-98
- Golgi C (1873) Sulla struttura della sostanza grigia dell cervello. *Gass Med Ital Lombarda*, 33, 244-6
- Golgi C (1875) Sulla fina struttura dei bulbi offectoria. *Riv sper Freniatria Med Legal* 1, 66-78
- Golgi C (1879) Di una nuova reazione apparentemente nera delle cellule nervose cerebrali ottenuta con bichloruro di mercurio. *Arch Sci Med* 3, 1-7
- Golgi C (1879) Un nuovo processo di tecnica microscopica. *Rendiconti dell Instituto Lombardo*, 12, 206-10
- Golgi C (1883) Recherches sur l'histologie des centres nerveux. *Arch Ital Biol* 3, 285-317
- Golgi C (1885) Sulla fina antomia della sistema nervoso. *Opera Omnia*, Milan (1903)
- Golgi C (1886) Professor Golgi's method of black colouring of the central nervous organs. *Alien Neurol* 7, 127-31
- Golgi C (1894) Untersuchungen Über Den Feineren Bau Des Centralen Und Peripherischen Nervensystems. Translated by Teuscher R, Fisher, Jena

REFERENCES

- Golgi C (1898) Sur la structure des cellules nerveuses. *Arch Ital Biol* 30, 60-71.
 Sur la structure des cellules nerveuses des ganglions spinaux. *Arch Ital Biol* 30, 278-86
- Golgi C (1899) De nouveau sur la structure des cellules nerveuses des ganglions spinaux. *Arch Ital Biol* 31, 273-80
- Golgi C (1959) Revue historique. Recherches sur l'histologie des centres nerveux. *Arch Ital Biol* 97, 276-99
- Goltz F L (1881) Trans International Medical Congress, London, 2-9 August 1881. Kolckmann, London, Vol 1, pp. 218-43
- Goltz F L (1892) Die Hund ohne Grosshirn. Siebente Abhandlung über die Verrichtung des Grosshirns. *Pflug Arch Ges Physiol* 51, 571-614
- Gorter M D and Grendel F (1925) On bimolecular layers of lipoids on the chromocytes of the blood. *J Exp Med* 41, 439-43
- Grafstein B and Forman D S (1980) Intracellular transport in neurons. *Physiol Rev* 60, 1167-1283
- Gray E G (1959) Axosomatic and axodendritic synapses of the cerebral cortex: an electron microscope study. *J Anat Lond* 93, 420-33
- Gray E G (1961a) The granule cells, mossy synapses and Purkinje spine synapses of the cerebellum: light and electron microscopic observations. *J Anat Lond* 95, 345-56
- Gray E G (1961b) Ultrastructure of synapses of the cerebral cortex and of certain specializations of neuroglial membranes. *J Anat, Suppl. Arnold, London*, pp. 54-73
- Gray E G (1964) Tissue of the central nervous system. In *Electron Microscopic Anatomy*. ed. Kurtz, S M. Academic Press, New York, pp. 369-417
- Gray E G (1966) Problems of interpreting the fine structure of vertebrate and invertebrate synapses. *Int Rev Gen Exp Zool* 2, 139-70
- Gray E G and Guillory R W (1966) Synaptic morphology in the normal and degenerating nervous system. *Int Rev Cytol* 19, 111-82
- Gray E G and Whittaker V P (1960) The isolation of synaptic vesicles from the central nervous system. *J Physiol, Lond*, 153, 35P-37P
- Gray E G and Whittaker V P (1962) The isolation of nerve endings from brain: an electron microscopic study of cell fragments derived by homogenization and centrifugation. *J Anat Lond* 96, 79-87
- Gray E G, Gordon-Weeks P R and Burgoine R D (1982) Nerve terminal ultrastructure: a role for neurotubules?, in *Neurotransmitter Interaction and Compartmentation*. ed. Bradford H F. Plenum Press, New York, pp. 1-13
- Gray J (1931) A Textbook of Experimental Cytology. Cambridge University Press, Cambridge, pp. 1-516
- Gray P (1954) The Microtomist's Formulary and Guide. The Blakiston Company, Inc., New York, pp. 269-621
- Gray P (1958) Handbook of Basic Microtechnique. McGraw-Hill, New York, p. 63
- Greenfield J G (1976) Neuropathology. eds. Blackwood W and Corsellis J A N. 3rd edition. Arnold, London, pp. 21-36
- Gregson N (1981) Review of the Living Cell. *Biochem Soc Trans* 9, 598
- Grubb D T and Keller A (1972a) Beam induced radiation damage in polymers and its effects on the image formed in electron microscope. *Proc Fifth Eur Congr Electron Microscopy*. pp. 554-60
- Grubb D T and Keller A (1972b) Origin of contrast effects in the electron microscopy of polymers. Part 2, Polyethylene spherulites. *J Mat Sci* 7, 822-35
- Gruner J E and Zahnd J P (1967) Sur la maturation synaptique dans le cortex visuel du lapin. in *Regional Development of the Brain In Early Life*. ed. Minkowski A. Blackwell, Oxford, pp. 125-33
- Guidotti G (1972) Membrane proteins. *Ann Rev Biochem* 41, 731-52
- Guth L (1956) Regeneration in the mammalian peripheral nervous system. *Physiol Rev* 36, 441-78
- Haapanen L, Kolmodin G M and Skoglund C R (1958) Membrane and action potentials of spinal interneurones in the cat. *Acta Physiol Scand* 43, 315-48
- Hallpike J F (1971) Proteolytic activity and basic protein loss in experimental myelin breakdown and multiple sclerosis. M.D. Thesis, London University
- Hallpike J F (1972) Enzyme and protein changes in myelin breakdown and multiple sclerosis. *Progress in Histochemistry and Cytochemistry* 3, pp. 179-215
- Ham A W (1974) Histology. 7th edition, Lippincott, Philadelphia, pp. 103-540
- Ham A W (1976) Histology. 8th edition, Lippincott, Philadelphia, pp. 467-517

REFERENCES

- Hamberger A (1963) Difference between isolated neuronal and vascular glia with respect to respiratory activity. *Acta Physiol Scand* 58, Suppl 203, 1-58
- Hamberger A, Hansson H-A and Sellstrom A (1975) Scanning and transmission electron microscopy on bulk prepared neuronal and glial cells. *Exp Cell Res* 92, 1-10
- Hamberger A and Sellstrom A (1975) Techniques for separation of neurons and glia and their application to metabolic studies. In *Metabolic Compartmentation and Neurotransmission*. eds. Berl S, Clarke D D and Schneider D. Plenum, New York, pp. 145-66
- Hammarberg C (1895) *Studien Uber Klinik Und Pathologie der Idiotie Nebst Untersuchungen Uber die Normale Anatomie der Hirnrinde*. Akad Buchdruckerei, Uppsala
- Han S S and Holmstedt J O V (1981) *Human Microscopic Anatomy*. McGraw-Hill, New York, pp. 138-66
- Hankin E H (1890) A cure for tetanus and diphtheria. *Nature*, 43, 121-3
- Hannover A (1840) Die Chromosaure, ein vorzugliches Mittel beim mikroskopischen Untersuchungen. *Arch Anat Physiol*, 549-58
- Hansson E (1982) Primary Astroglial Cultures. Docent thesis, Institute of Neurobiology, University of Goteborg, Sweden, pp. 1-71
- Hansson E, Sellstrom A, Persson L I and Ronnback L (1980) Brain primary culture - a characterization. *Brain Res* 188, 233-46
- Hansson H A and Sourander P (1964) Studies on cultures of mammalian retina. *Z Zellforsch* 62, 26-47
- Hardesty I (1904) On the development and nature of neuroglia. *Am J Anat* 3, 229-68
- Hardy W B (1899) On the structure of cell protoplasm. *J Physiol* 24, 158-210
- Harford J B, Waechter C J, Saul R and De Vries G H (1979) Evidence for the bio-synthesis of mannosylphosphoryldolichol by an axolemma-enriched membrane preparation from bovine white matter. *J Neurochem* 32, 91-8
- Harris E J (1954) Ion transport in frog muscle. *J Physiol* 124, 248-53
- Harris E J (1972) *Transport and Accumulation In Biological Systems*. 3rd edition. Butterworth, London
- Harris J E (1941) The influence of the metabolism of human erythrocytes on their potassium content. *J Biol Chem* 141, 578-95
- Harrison R G (1907) Observations on the living developing nerve fibre. *Proc Soc Exp Biol Med* 4, 140-3; *Anat Rec* 1, 116-18
- Harrison R G (1910) The outgrowth of the nerve fibre as a mode of protoplasmic movement. *J Exp Zool* 9, 787-846
- Hartmann J (1953) An electron optical study of sections of the central nervous system. *J Comp Neurol* 99, 201-49
- Hartmann J (1962) Identification of neuroglia in electron micrographs of normal nerve tissue. In *IV. Internationaler Kongress fur Neuropathologie*. Proceedings. ed. Jacob H. Georg Thieme, Stuttgart, 2, pp. 32-35
- Hartridge H and Haynes F (1930) *Histology for Medical Students*. Oxford University Press, London, pp. 122-369
- Hass G M and Taylor C B (1953) Quantitative studies of experimental production and treatment of acute closed cerebral injury. *Arch Neurol Psychiatry* 69, 145-70
- Hassler R and Stephan H (eds.) (1966) *Evolution of The Forebrain*. George Thieme, Stuttgart
- Hayat M A (ed.) (1973-77) *Electron Microscopy of Enzymes*. Vols 1-5, Van Nostrand Reinhold, New York
- Hayat M A (1970-78) *Techniques Of Electron Microscopy, Biological Applications*. Vols 1-8, Van Nostrand Reinhold, New York
- Heimer L and Ekholm R (1967) Neuronal argyrophilia in early degenerative states: A light and electron microscopic study of Glees and Nauta techniques. *Experientia* 23, 237-9
- Held H (1897) Beitrage zur Structur der Nervenzellen und ihrer Fortsatze. *Arch Anat Physiol Anat Abt* 21, 204-94
- Held H (1909) Ueber die Neuroglia marginalis des menschlichen Grosshirnrinde. *Monatsschr Psychol Neurol* 26, Erg 360-416
- Henle J (1871) *Handbuch der Nervenlehre des Menschen*. Vieweg, Braunschweig.
- Henn F A and Thompson T E (1969) Synthetic lipid bilayer membranes. *Ann Rev Biochem* 38, 241-62
- Henrikson C K and Vaughn J E (1974) Fine structural relationships between neurites and radial glial processes in developing mouse spinal cord. *J Neurocytol* 3, 659-75
- Herdson P B and Kaltenbach J P (1965) Electron microscope studies on enzyme activity and the isolation of thiohydantoin-induced myelin figures in rat liver. *J Cell Biol* 25, 485-93

REFERENCES

- Herman L, Eber L and Fitzgerald P J (1962) Liver cell degeneration with ethionine administration, Proceedings of the 5th International Congress for Electron Microscopy. ed. Breese S S Jr. Academic Press, New York, 2, pp. VV-6
- Herrera A (1869-1942) quoted in Bergier J (1957) The Secrets of Living Matter. Barrie and Rockliff, London, pp. 6-8
- Hertl M (1957) Zum Nucleolus-problem. Z Zellforsch Mikrosk Anat 46, 18-51
- Hertwig G (1931) Einfluss der Fixierung auf das Kernung Zellvolumen. Z Mikrosk Anat Forsch 23, 484-504
- Hertz L (1965) Possible role of glia: a potassium mediated neuronal-neuroglial-neuronal impulse transmission system. Nature, Lond, 206, 1091-92
- Hertz L (1966) Neuroglial localization of potassium and sodium effects on respiration in brain. J Neurochem 13, 1373-87
- Hertz L (1977) Drug-induced alterations of ion distribution at the cellular level of the central nervous system. Pharmacol Rev 29, 35-65
- Hertz L (1979) Functional interactions between neurons and astrocytes. I. Turnover and metabolism of putative aminoacid transmitters. Prog Neurobiol 13, 277-323
- Hertz L, Schousboe A, Boehler N, Mukerji S and Fedoroff S (1978) Kinetic characteristics of the glutamate uptake into normal astrocytes in culture. Neurochem Res 3, 1-14
- Hess A (1953) The ground substance of the central nervous system as revealed by histochemical staining. J Comp Neurol 98, 69-91
- Heuser J E and Reese T S (1973) Evidence for recycling of synaptic vesicle membrane during transmitter release at the frog neuromuscular junction. J Cell Biol 57, 315-44
- Hild W (1957) Myelinogenesis in cultures of mammalian central nervous tissue. Z Zellforsch 46, 71-95
- Hild W (1959) Myelin formation in cultures of mammalian central nervous tissue. In Korey S R, ed. The Biology of Myelin. Cassell, London, pp. 188-99
- Hild W, Chang J J and Tasaki I (1958) Electrical responses of astrocytic glia from the mammalian central nervous system cultivated in vitro. Experientia 14, 220-1
- Hild W and Tasaki I (1962) Morphological and physiological properties of neurons and glial cells in tissue vulture. J Neurophysiol 25, 277-304
- Hildebrand C (1971) Ultrastructural and light microscopic studies of the nodal region on large myelinated fibres of the adult female spinal cord white matter. I. The nodes of Ranvier. Acta Physiol Scand Suppl 364, 81-109
- Hill J (1770) The Construction of Timber From Its Early Growth. Hill, London, pp. 4-10
- Hillarp N A, Fuxe K and Dahlstrom A (1966) Central monoamine neurons, in International Symposium on Mechanisms of Release of Biogenic Amines. Stockholm, 1965, Pergamon Press, Oxford, pp. 31-57
- Hillman H (1961) Membrane potentials in guinea-pig cerebral cortex slices in vitro. Their dependence on substrate and oxygen. The effect of clupein and of ganglioside preparations. J Neurochem 8, 257-61
- Hillman H (1966a) Growth of processes from single isolated dorsal root ganglion cells of young rats. Nature 209, 102-3
- Hillman H (1966b) The role of potassium and sodium ions as studied in mammalian brain. Internat Rev Cytol 20, 125-137
- Hillman H (1970) Chemical basis of epilepsy. Lancet, 2, 23-4
- Hillman H (1972) Certainty and Uncertainty in Biochemical Techniques. Surrey University Press, Blackie, Glasgow
- Hillman H (1976) Towards a classification of the validity of experimental evidence in biology. Acta Biotheoretica 25, 153-62
- Hillman H (1982a) Cell structure: some questions not answered by the electron microscopists. School Science Review 222, 173-77
- Hillman H (1982b) Problems of light microscopy in biological systems, with approaches to solving them. Microscopy 34, 448-64
- Hillman H (1983) Some fundamental theoretical and practical problems associated with neurochemical techniques in mammalian studies. Neurochem Int 5, 1-13
- Hillman H (1985) The anatomical synapse in mammals by light and electron microscopy. Medical Hypotheses 17, 1-32
- Hillman H and Deutsch K (1978a) Area changes in slices of rat brain during preparation for histology or electron microscopy. J Microsc 114, 77-84
- Hillman H and Deutsch K (1978b) Mitochondria in isolated neuroglial clumps and neuronal cell bodies of rabbit. Microscopy 33, 308-9
- Hillman H and Hyden H (1965) Membrane potentials in isolated neurons in vitro from Deiters' nucleus of rabbit. J Physiol 177, 398-410

REFERENCES

- Hillman H and Khallawan S A (1970) The growth of new processes from isolated dorsal and sympathetic cell bodies of rat and rabbit. *Tissue and Cell* 2, 249-59
- Hillman H and Richards P (1983) Types of light microscopy. *Microscopy* 34, 630-6
- Hillman H and Sartory P (1980a) The Living Cell - a Re-examination of its Fine Structure. Packard, Chichester, pp. 1-112
- Hillman H and Sartory P (1980b) A re-examination of the fine structure of the living cell and its implications for biological research. *School Science Review* 62, 241-52
- Hillman H and Sheikh K (1968) The growth in vitro of new processes from vestibular neurons isolated from adult and young rabbits. *Exp Cell Res* 50, 315-22
- Hillman H and Sjuve S (1980) The appearances of rat tissues under phase contrast microscopy and stained with haematoxylin and eosin. *Microscopy* 34, 136-46
- Hillman H and Wraae O (1981) Practical and theoretical problems associated with the use of cerebral slices, in *Chemisms of The Brain*. eds. Rodnight R, Bachelard H S; and Stahl W L. Churchill-Livingstone, Edinburgh, pp. 37-46
- Hillman H, Campbell W J and McIlwain H (1963) Membrane potentials in isolated and electrically stimulated mammalian cerebral cortex. *J Neurochem* 10, 325-39
- Hillman H, Deutsch K, Allen J and Sartory P (1977) The histology of neuroglial clumps in rabbit brain. *Microscopy* 33, 175-80
- Hillman H, Hussain T and Sartory P (1973) Autofluorescence of isolated unfixed rabbit Deiters' neurons and surrounding neuroglial clumps. *Experientia* 29, 1113-15
- Hillman H, Hussain T and Sartory P (1976) The effect of haematoxylin and eosin staining and embedding on the appearance of isolated neurons. *Microscopy* 33, 77-84
- Hillman H, Stollery S and Joanny P (1974) The mechanism of the exclusion of sodium ions and the concentration of potassium ions by guinea-pig cerebral cortex slices. *J Neurochem* 22, 573-9
- Hirano A (1969) The fine structure of brain in edema. In *The Structure and Function of Nervous Tissue*, ed. Bourne G H, Vol 2, Structure 2, Academic Press, New York, pp. 69-135
- Hirsch H E (1972) Differential determination of hexosaminidases A and B of two forms of β -galactosidase, in the layers of the human cerebellum. *J Neurochem* 19, 1513-17
- His W (1887) Zur Geschichte des menschlichen Rückenmarkes und des Nervenzellen. Abh K Sachsischen Ges Wiss, Math-Phys Classe 13, 477-514
- His W (1888) Über die embryonale Entwicklung der Nervenbahnen. *Anat Anz* 3, 499-506
- Hitzig G (1874) Untersuchungen Über Das Gehirn, Hirschwald, Berlin, pp. 1-276
- Hodgkin A L (1951) The ionic basis of electrical activity in nerve and muscle. *Biol Rev* 26, 339-409
- Hodgkin A L and Huxley A F (1952a) Currents carried by sodium and potassium ions through the membrane of the giant axon of *Loligo*. *J Physiol* 116, 449-72
- Hodgkin A L and Huxley A F (1952b) The dual effect of membrane potential on sodium conductance in the giant axon of *Loligo*. *J Physiol* 116, 497-506
- Hodgkin A L and Keynes R D (1953) The mobility and diffusion coefficient of potassium in giant axons from *Sepia*. *J Physiol* 119, 513-28
- Hodgkin A L and Keynes R D (1955) Active transport of cations in giant axons from *Sepia* and *Loligo*. *J Physiol* 128, 28-60
- Hogue M (1964) Human fetal brain cells in tissue cultures: their identification and motility. *J Exp Zool* 106, 85-108
- Hogue M (1950) Brain cells from human fetuses and infants, cultured in vitro after death of the individuals. *Anat Rec* 108, 457-75
- Hokin L E and Hokin M R (1963) Biological transport. *Ann Rev Biochem* 32, 553-78
- Holmes S (1905) Silver Blaze, quoted in *Memoirs of Sherlock Holmes* by Conan-Doyle A. George Newnes, London
- Holmgren E (1900) Weitere Mitteilungen über die 'Saftkanalchen' der Nervenzellen. *Anat Anz* 18, 290-6
- Holzer W (1923) Ueber die Bestandteile des Heldschen Gliasyncitiums. *Zeitschr Ges Neurol Psychiatr* 87, 167-75
- Hooke R E (1665) *Micrographia Or Some Physiological Descriptions of Minute Bodies Made By Magnifying Glasses, With Observations and Inquiries Thereupon*. Martyn and Allestry, London, pp. 1-246
- Hopwood D (1969) Fixatives and fixation, a review. *Histochem J* 1, 323-60
- Hopwood D (1970) The reactions between formaldehyde, glutaraldehyde, and osmium tetroxide and their fixation effects on bovine serum albumin and on tissue blocks. *Histochemie* 24, 50-64
- Horne R W and Harris J R (1981) The electron microscope in biology. *School Science Review*, 63, 53-69

REFERENCES

- Hosie R J A (1965) The localization of adenotriphosphatases in morphologically characterized subcellular fractions of guinea-pig brain. *Biochem J* 96, 404-12
- Hosli L, Andres P F and Hosli E (1979) Effects of 4-aminopyridine and tetraethyl-ammonium on the depolarization of GABA on cultured satellite glial cells. *Neurosci Letters* 11, 193-96
- Hosli L, Andres P F and Hosli E (1980) Neurone-glia interaction: indirect effects of amino acid transmitters on cultured glial cells. In *Tissue Culture in Neurobiology*. eds. Giacobini E, Vernadakis A and Shahar A. Raven Press, New York, pp. 373-83
- Hosokawa H (1963) General characteristics of the neuroglia. *Morphology of Neuroglia*. Glia Research Group. ed. Nakai J. Igaku Shoin Ltd, Tokyo, pp. 1-12
- Hulcher F H (1963) Physical and chemical properties of myelin. *Arch Biochem Biophys* 100, 237-44
- Humason G L (1967) *Animal Tissue Techniques*. 2nd edition. W H Freeman and Company, San Francisco, pp. 1-516
- Hussain T S, Hillman H and Sartory P (1974) A nucleolar membrane in neurons. *Microscopy* 32, 348-53
- Hyden H (1959) Quantitative assay of compounds in isolated fresh nerve cells and glial cells from control and stimulated animals, *Nature*, 184, 433-5
- Hyden H (1961) The neuron. In *The Cell*. eds. Brachet J and Mirsky A E. Vol IV, Academic Press, New York, pp. 215-323
- Hyden H (1961) Satellite cells in the nervous system, *Sci Am* 205, 62-70
- Hyden H (1962) A molecular basis for neuron-glia interaction. In *Macromolecular Specificity and Biological Memory*. ed. Schmitt F O. MIT Press, Cambridge, Mass, pp. 55-69
- Hyden H (1964) RNA - a functional characteristic of the neuron and its glia. In *Brain Function*. ed. Brazier M A. Vol 2, University of California Press, pp. 29-68
- Hyden H (ed.) (1967) *The Neuron*. Elsevier, Amsterdam
- Hyden H (1967) Dynamic aspects of the neuron-glia relationship. In *The Neuron*. ed. Hyden H. Elsevier, Amsterdam, pp. 179-219
- Hyden H and Hamberger C A (1945) Cytochemical changes in the cochlear ganglion caused by acoustic stimulation and trauma. *Acta Oto-Laryngol* (Stockholm), 61, Suppl 1
- Hyden H and McEwen B (1966) A glial protein specific for the nervous system. *Proc Natl Acad Sci* 55, 354-58
- Hyden H and Pigozzi A (1960) A cytophysiological study of the functional relationship between oligodendroglial cells and cells of Deiters' nucleus. *J Neurochem* 6, 57-72
- Illiano G, Tell P E, Siegel M I and Cuatrecasas P (1973) Guanosine 3'-5'-cyclic monophosphate and the action of insulin and acetylcholine. *Proc Natl Acad Sci, USA*, 70, 2443-7
- International Congress on Cell Biology (1981) ed. Schweiger H. Springer, Berlin, papers on pp. 275-358
- Iversen L L (1975) *Handbook of Psychopharmacology*. Vol 4, Amino acid neurotransmitters. Plenum Press, New York
- Jacob M, Samec J, Stevenin J and Mandel P (1967) Polysomes and polysomal RNA of rat brain. *J Neurochem* 14, 169-78
- Jacobson K B and Kaplan N O (1957) Distribution of enzymes cleaving pyridine nucleotides in animal tissues. *J Biophys Biochem Cytol* 3, 31-43
- Jacobson M (1978) *Developmental Neurobiology*. Plenum Press, New York, pp. 27-146
- Jacque C M and Baumann N A (1978) Glial cell markers in dysmyelinating mutants. In *Dynamic Properties of Glial Cells*. eds. Schoffeniels E, Franck G, Hertz L and Tower D B. Pergamon Press, Oxford, pp. 275-8
- Jacque C, Jorgensen O S, Baumann N A and Bock E (1981) Brain specific antigens in the quaking mouse during ontogeny, personal communication
- Jakob A (1927) *Anatomie und Histologie des Grosshirns*. Leipzig, 1 and 2
- Jastrowitz M (1870-71) *Encephalitis und Myelitis des ersten Kindesalters*. Arch Psychiatr 2, 389-414; 3, 162-213
- Joanny P, Barbosa E, Hillman H and Corriol J (1971) The uptake and efflux of glycine from rat cerebral-cortex slices. *Biochem J* 125, 255-60
- Joanny P and Hillman H (1963) Substrates and the potassium and sodium levels of guinea-pig: cerebral cortex slices in vitro: effects of application of electrical pulses, or inhibitors and of anoxia. *J Neurochem* 10, 655-64

REFERENCES

- Joanny P and Hillman H (1964) Further studies on the potassium and sodium concentrations of mammalian cerebral slices in vitro. *J Neurochem* 11, 413-22
- Joanny P, Corriol J and Hillman H (1969) Uptake of monosaccharides by guinea-pig cerebral cortex slices. *Biochem J*, 112, 367-71
- Johnston P V and Roots B I (1972) *Nerve Membranes*. Pergamon Press, Oxford, p. 36
- Jones D G (1975) *Synapses and Synaptosomes, Morphological Aspects*. Chapman and Hall, London, pp. 1-258
- Jones D G and Brearley R F (1972a) Further studies on synaptic junctions. I. Ultrastructural features in intact rat cerebral cortex. *Z Zellforsch Mikrosk Anat* 125, 415-31
- Jones D G and Brearley R F (1972b) Further studies on synaptic junctions. II. A comparison of synaptic ultrastructure in fractionated and intact cerebral cortex. *Z Zellforsch Mikrosk Anat* 125, 432-47
- Jones E G and Powell T P S (1970) Electron microscopy of the somatic sensory cortex of the cat. *Phil Trans R Soc Lond B* 257, 1-21
- Jorgensen O S (1981) Neuronal membrane D2 protein during rat brain ontogeny. *J Neurochem* 37, 939-46
- Joseph J (1954) Nuclear population changes in degenerating posterior columns of rabbits' spinal cord. *Acta Anat* 21, 356-65
- Kai M and Hawthorne J N (1967) quoted in *Enzyme Cytology*. ed. Roodyn D B. Academic Press, London, pp. 378-406
- Kaiserman-Abramof I R and Peters A (1972) Some aspects of the morphology of Betz cells in the cerebral cortex of the cat. *Brain Res* 43, 527-46
- Karlsson U (1966) Three-dimensional study of neurons in the lateral geniculate nucleus of the rat. II. Environment of perikarya and proximal parts of their branches. *J Ultrastruct Res* 16, 482-504
- Karnovsky M J and Roots L (1964) A direct-colouring thiocholine method for cholinesterases. *J Histochem Cytochem* 12, 219-21
- Kasa P (1971) Ultrastructural localization of choline acetyltransferase and acetylcholinesterase in central peripheral nervous system. *Histochemistry of Nervous Transmission*. *Prog Brain Res* 34, 337-44
- Kates M and Kuksis A (1980) *Membrane Fluidity: Biophysical Techniques and Cellular Regulation*. Humana Press, Clifton, NJ, pp. 1-445
- Katz B (1969) *The Release of Neural Transmitter Substances*. Sherrington Lecture, Liverpool University Press, Liverpool
- Katzman R (1961) Electrolyte distribution in mammalian central nervous system. Are glia high sodium cells? *Neurology* 11, 27-36
- Katzman R and Wilson C E (1961) Extraction of lipid and lipid-cation from frozen brain tissue. *J Neurochem* 7, 113-27
- Kelly J S and Dick F (1978) GABA in glial cells of the peripheral and central nervous system. In *Dynamic Properties of Glial Cells*. eds. Schoffeniels E, Franck G, Tower D B and Hertz L. Pergamon Press, Oxford, pp. 183-92
- Kennedy P G E, Lisak R P and Raff M C (1980) Cell type-specific markers for human glial and neuronal cells in culture. *Lab Invest* 43, 342-51
- Kernohan J W and Sayre G P (1952) *Tumours Of The Central Nervous System*. Armed Forces Institute Of Pathology, Washington
- Kershman J (1938) The medulloblast and the medulloblastoma. *Arch Neurol Psychiatry* 40, 937-67
- Kershman J (1939) Genesis of microglia in the human brain. *Arch Neurol Psychiatry* 41, 24-50
- Key E A H and Retzius G M (1876) *Studien in Anatomie des Nervensystems und des Bindegewebes*. Samson and Wallin, Stockholm
- Keynes R D (1954) The ionic fluxes in frog muscle, *Proc R Soc London, B* 142, 359-82
- Kibler R F, O'Neill R P and Robin E D (1964) Intracellular acid base relations of dog brain with reference to the brain extracellular volume. *J Clin Invest* 43, 431-43
- Kies M W, Murphy J B and Alford E C (1961) Studies on the encephalitogenic factor in guinea pig central nervous system. In *Chemical Pathology of the Nervous System*. ed. Folch-Pi J. Pergamon Press, Oxford, pp. 197-204
- King L S (1937) Method for rapid impregnation of microglia and oligodendroglia in material fixed in formaldehyde. *Arch Neurol Psychiatry* 38, 362-4
- King J S (1968) A light and electron microscopical study of perineuronal glial cells and processes in the rabbit neocortex. *Anat Rec* 161, 111-24
- Kirkman H and Severinghaus A E (1938) A review of the Golgi apparatus. *Anat Rec* 70, 413-31; 70, 557-73; 71, 79-103

REFERENCES

- Kitai S T and Bishop G A (1981) Horse radish peroxidase. Intracellular staining of neurons. In *Neuroanatomical Tract Tracing Methods*. eds. Heimer L and Robards M J. Plenum, New York, pp. 263-77
- Klatzo I and Miquel J (1960) Observations on pinocytosis in nervous tissue. *J Neuropathol Exp Neurol* 19, 475-87
- Klatzo I, Miquel J, Ferris P J, Prockop L D and Smith D E (1964) Observations on the passage of the fluorescein labelled serum proteins (FLSP) from the cerebrospinal fluid. *J Neuropathol Exp Neurol* 23, 18-35
- Klatzo I, Piraux A and Laskowski E J (1958) The relationship between edema, blood-brain barrier and tissue elements in a local brain injury. *J Neuropathol* 17, 548-64
- Klestadt B (1915) Experimentelle Untersuchungen über die resorptive Funktion des Epithels des Plexus choroideus und des Ependyms der Seitenventrikel. *Zentralbl Allg Pathol Pathol Anat* 26, 161-6
- Knoll M and Ruska E (1932) Das Elektronenmikroskop. *Z Physik* 78, 318-39
- Koelle G B and Friendenwald J A (1949) A histochemical method for localising cholinesterase activity. *Proc Soc Exp Biol Med*, New York, 70, 617-22
- Koenig H (1964) RNA metabolism in the nervous system. Some RNA dependent functions of neurons and glia. In *Morphological and Biochemical Correlates of Neural Activity*. eds. Cohen M M and Snider R S. Hoeber, New York, pp. 39-56
- Koenig H, Bunge M B and Bunge R P (1962) Nucleic acid and protein metabolism in white matter. *Arch Neurol* 6, 177-93
- Konigsmark B W and Sidman R L (1963) Origin of brain macrophages in the mouse. *J Neuropathol Exp Neurol* 22, 643-76
- Korr H (1980) Proliferation of different cell types in the brain. *Adv Anat Embryol Cell Biol* 61, 1-72
- Koul O and Jungalwala F B (1981) UDP-Galactose:ceramide galactosyltransferase of rat central-nervous-system myelin. *Biochem J* 194, 633-7
- Krnjevic K and Schwartz S (1967) Some properties of unresponsive cells in the cerebral cortex. *Exp Brain Res* 3, 306-19
- Kruger L and Maxwell D S (1966) Electron microscopy of oligodendrocytes in normal rat cerebrum. *Am J Anat* 118, 411-36
- Krypsin-Exner W (1952) Ueber die Architektonik der Glia im Zentral nervensystem des Menschen und der Säugetiere. *Proc First Int Cong Neuropathol*, Rome, 3, 504-10
- Kuffler S W and Nicholls J G (1976) *From Neuron to Brain*. Sinauer Associates, Mass, pp. 253-88
- Kuhne W (1862) Über die peripherischen Endorgane der motorischen Nerven. Engelmann, Leipzig, pp. 17-18
- Kuntz A and Hamilton J W (1938) Afferent innervation of the skin. *Anat Rec* 71, 387-400
- Kurella G A (1961) Polyelectrolytic properties of protoplasm and the character of resting potentials, in *Membrane Transport and Metabolism*. eds. Kleinzeller A and Kotyk A. Czechoslovak Academy of Sciences, Prague, pp. 54-68
- Kurihara T and Tsukada Y (1967) The regional and subcellular distribution of 2'-3'-cyclic nucleotide 3'phosphohydrolase in the central nervous system. *J Neurochem* 14, 1167-74
- Laatsch B A, Kier M W, Gordon S and Alvord E C (1962) The encephalomyelic activity of myelin isolated by ultracentrifugation. *J Exp Med* 115, 777-88
- LaBella F S and Brown J H U (1959) Cell fractionation of anterior pituitary glands from beef and pig. *J Biophys Biochem Cytol* 5, 17-24
- Lajtha A (1957) The development of the blood-brain barrier. *J Neurochem* 1, 216-27
- Landon D N (1976) *The Peripheral Nerve*. Chapman and Hall, London
- Landon D N and Hall S (1976) The myelinated nerve fibre. In *The Peripheral Nerve*. ed. Landon D N. Chapman and Hall, London, pp. 1-105
- Lanterman A J (1877) Ueber den feineren Bau der Markhaltigen Nervenfasern. *Arch Mikrosk Anat Entwicklungsmech* 13, 1-8
- Lassek A M (1948) The pyramidal tract. Basic considerations of corticospinal neurons. *Res Publ Ass Nerv Ment Dis*, pp. 106-128
- La Valette St George V (1867) Über die Genese der Samenkörper. *Arch Mikroskop Anat Entwicklungsmech* 3, 263-73
- Lazarides E (1980) Intermediate filaments as mechanical integrators of cellular space. *Nature* 283, 249-56
- Le Baron F N and Folch J (1956) The isolation from brain tissue of a trypsin-resistant fraction containing combined inositol, and its relation to neurokeratin. *J Neurochem* 1, 101-8
- Lee J C and Olszewski J (1960) Penetration of radioactive albumin from cerebrospinal fluid into brain tissue. *Neurology* 10, 814-22

REFERENCES

- Leeson T S, and Leeson C R (1979) A Brief Atlas of Histology. Saunders, Philadelphia, pp. 89-104
- Leeson T S and Leeson C R (1981) Histology. Saunders, Philadelphia, pp. 216-53
- Leeuwenhoek A von (1719) Epistolae physiologicae super compluribus naturae arcanis Epistola XXXII, Beman, Delft, pp. 310-17
- Le Fevre P E and Peters A A (1966) Evidence of mediated transfer of monosaccharides from blood to brain in rodents. *J Neurochem* 13, 35-46
- Leonhardt H (1951) Plasmatische Abteilung. Ein modus der Farbstoffausscheidung aus dem Gehirn. (Zugleich ein Beitrag Zur Physiologie des Neurencytiums). *Z Anat Entwicklungsgesch* 115, 555-69
- Levi-Montalcini R, Meyer A and Hamberger V (1954) In vitro experiments on the effects of mouse sarcomas 180 and 37 on the spinal and sympathetic ganglia of the chick embryo. *Cancer Res* 14, 49-57
- Lewis A J (1976) Mechanisms of Neurological Disease. Little, Brown and Company, Boston, pp. 41 and 45
- Lewis M R and Lewis W H (1915) Mitochondria and other cytoplasmic structures in tissue cultures. *Am J Anat* 17, 339-402
- Lewis M S, Krieg L C and Kirk W (1974) The molecular weight and detergent binding of bovine rhodopsin. *Exp Eye Res* 18, 29-40
- Lewis W H (1923) Observation on cells in tissue cultures with dark field illumination. *Anat Rec* 26, 15-29
- Lewis W H (1931) Pinocytosis. *Bull Johns Hopkins Hosp* 49, 17-27
- Lewis W H and McCoy C C (1922) The survival of cells after the death of the organism. *Bull Johns Hopkins Hosp* 33, 284-93
- Leyton A S F and Sherrington C S (1917) Observations on the excitable cortex of the chimpanzee, orang-utan and gorilla. *Q J Exp Physiol* 11, 135-222
- Li C L, Bak A F and Parker L O (1968) Specific resistivity of the cerebral cortex and white matter. *Exp Neurol* 20, 533-43
- Li C L, Okujava V M and Bak A F (1971) Some electrical measurements of cortical elements and neuronal responses to direct stimulation with particular reference to input resistance. *Exp Neurol* 31, 263-76
- Lindsay R M (1979) Adult rat astrocytes support survival of both NGF-dependent and NGF-insensitive neurones. *Nature* 282, 80-2
- Lindsay R M, Barber P C, Sherwood M R C, Zimmer J and Raisman G (1982) Astrocyte cultures from adult rat brain. Derivation, characterization and neurotrophic properties of pure astroglial cells from corpus callosum. *Brain Res* 243, 329-43
- Ling G (1952) The role of phosphate in the maintenance of the resting potential and selective ionic accumulation in frog muscle cells. In *Phosphorus Metabolism*. eds. McElroy W D and Glass B. Vol 2, Johns Hopkins Press, Baltimore, pp. 748-97
- Ling G (1955) Muscle electrolytes. *Am J Physiol Med* 34, 89-101
- Ling G (1962) A Physical Theory of The Living State. The Association Induction Hypothesis. Ginn (Blaisdell), Boston, Mass, pp. 1-680
- Lisak R P, Abramsky O, Dorfman S H, George J, Manning M C, Pleasure D E, Saida T and Silberberg D H (1979) Antibodies to galactocerebroside bind to oligodendroglia in suspension culture. *J Neurol Sci* 40, 65
- Lister J (1830) On some properties in achromatic object-glasses application to the improvement of the microscope. *Phil Trans R Soc* 1-2, 187-200
- Lodin Z, Mares V, Karasek J and Skrivanova P (1967) Studies of the effect of fixation on nervous tissue. II. Changes of size of nuclei of nervous cells after fixation and further histological treatment of nervous tissue. *Acta Histochem* 28, 297-312
- Lorber J (1980) 'Is your brain really necessary?' *World Med* 15, 21-4
- Lorber J and Priestley B L (1981) Children with large heads: a practical approach to diagnosis in 557 children, with special reference to 109 children with megalencephaly. *Dev Med Child Neurol* 23, 494-504
- Lorente de Nò R (1934) Studies on the structure of the cerebral cortex. I. The area entorhinalis. *J Psychol Neurol* 45, 381-438
- Lorente de Nò R (1938) Limits of variation on the synaptic delay of motoneurons. *J Neurophysiol* 1, 187-94
- Lorente de Nò R (1938) Synaptic stimulation of motoneurons as a local process. *J Neurophysiol* 1, 195-206
- Loten E G and Sneyd J G J (1970) An effect of insulin on adipose-tissue adenosine 3':5'-cyclic monophosphate phosphodiesterase. *Biochem J* 120, 187-933
- Lubinska L (1964) Axoplasmic streaming in regenerating and normal nerves. In *Mechanisms in Neural Regeneration. Progress in Brain Research*. eds. Singer S J and Van Schade J P. Elsevier, Amsterdam, pp. 1-66

REFERENCES

- Lucy J A (1973) Review of 'Certainty and Uncertainty in Biochemical Techniques.' *Biochem Educ* 1, 31-2
- Lumsden C E (1958) Histological and histochemical aspects of normal neuroglia cells. In *Biology of Neuroglia*. ed. Windle W F. C C Thomas, Springfield, Ill., pp. 141-161
- Lumsden C E (1968) Nervous tissue in culture. In *The Structure and Function of Nervous Tissue*. ed. Bourne G H. Vol 1, p. 104
- Lund-Andersen H (1974) Extracellular and intracellular distribution of inulin in rat brain cortex slices. *Brain Res* 65, 239-54
- Lund-Andersen H and Kjeldsen C S (1976) Uptake of glucose analogues by rat brain cortex slices: a kinetic analysis based upon a model. *J Neurochem* 27, 361-8
- Luse S A (1956) Electron microscopic observations of the central nervous system. *J Biophys Biochem Cytol* 2, 531-42
- Luse S A (1961) Brain ultrastructure in hydration and dehydration. *Arch Neurol* 4, 139-52
- Luse S A (1962) Ultrastructure of the brain and its relation to transport of metabolites. In *Ultrastructure and Metabolism of the Nervous System*, Association for Research In Nervous and Mental Disease, eds. Korey S R, Pope A and Robins E. Williams and Wilkins, Baltimore, Vol XL, pp. 1-26
- Lux H D and Pollen D A (1966) Electrical constants of neurons in the motor cortex of the cat. *J Neurophysiol* 29, 207-20
- Madge E and Scheibel A B (1958) Neurons and neuroglia cells as seen with the light microscope, In *Biology of Neuroglia*, ed. Windle W F. Charles C Thomas, Springfield, Ill., pp. 5-23
- Mallory F B (1900) A contribution to staining methods. *J Exp Med* 5, 19-20
- Manery J F and Hastings A B (1933) The distribution of electrolytes in mammalian tissues. *J Biol Chem*, 127-676
- Mann G (1902) *Physiological History, Methods and Theory*. Clarendon Press, Oxford, pp. 180-306
- Mannen H (1963) Astroglia and oligodendroglia. In *Morphology of Neuroglia*. Glia Research Group. ed. Nakai J. Igaku Shoin, Tokyo, pp. 5-10
- Marchbanks R M (1975) The subcellular origin of the acetylcholine released at synapses. *Int J Biochem* 6, 303-13
- Marchbanks R M (1978) The vesicular hypothesis questioned. *Trends Neurosci* 1, 83-4
- Marchi V and Algeri G (1885-86) Sulla degenerazioni discendenti consecutive a lesioni sperimentale in diverse zone delle corteccia cerebrale. *Riv sper Freniatra Med Legal* 11, 492-4; 12, 208-52
- Marks N and Lajtha A (1963) Protein breakdown in the brain: subcellular distribution and properties of neural and acid proteinases. *Biochem J* 89, 438-47
- Marsland T A, Glees P and Erikson L B (1954) Modification of the Glees silver impregnation for paraffin sections. *J Neuropathol Exp Neurol* 13, 587-91
- Matthews M A (1974) Microglia and reactive 'M' cells of degenerating central nervous system: Does similar morphology and function imply a common origin? *Cell Tiss Res* 148, 477-91
- Matthews M A and Kruger L (1973) Electron microscopy of non-neuronal cellular changes accompanying neural degeneration in thalamic nuclei of the rabbit. II. Reactive elements within the neuropil. *J Comp Neurol* 148, 313-46
- Maunsbach A B, Skriver E and Jorgensen P L (1979) Ultrastructure of purified Na⁺-K⁺ATPase membranes. In *Sodium-Potassium ATPases*. eds. Skou J C and Norby J G. Academic Press, London, pp. 3-13
- Mauthner L (1860) Beitrage zur naheren Kenntniss der morphologischen Elemente der Nervensystems. Sachser Akad Wiss Wien, Math- Naturw Kl 39, 583-90
- Maxwell D S and Kruger L (1965) The fine structure of astrocytes in the cerebral cortex and their response to focal injury produced by heavy ionizing particles. *J Cell Biol* 25, 141-57
- Mayer S, Maickel R P and Brodie B B (1959) Kinetics of penetration of drugs and other foreign compounds into cerebrospinal fluid and brain. *J Pharmacol* 127, 205-11
- Maynard E A, Schultz R L and Pease D C (1957) Electron microscopy of the vascular bed of rat cerebral cortex. *Am J Anat* 100, 409-33
- McClung C E M (1928) *Handbook of Microscopical Techniques*. Hoeber, New York
- McClung-Jones R (1967) *McClung's Handbook of Microscopical Techniques*. 3rd edition. Hafner Publ Co, New York, pp. 492-505
- McIlwain H (1975) Preparing neural tissues for metabolic studies in isolation. In *Practical Neurochemistry*. ed. McIlwain H. Churchill-Livingstone, London, pp. 106-26

REFERENCES

- McLennan D H, Seeman P, Iles G H and Yip C C (1971) Membrane formation by the adenosine triphosphatase of sarcoplasmic reticulum. *J Biol Chem* 246, 2702-10
- McMahan U J, Spitzer N C and Peper K (1972) Visual identification of nerve terminals in living isolated skeletal muscle. *Proc Soc B*, 181, 421-30
- McManus J F A and Mowry R W (1960) *Staining Methods*. Harper and Row, New York, pp. 67-353
- Meek G A (1976) *Practical Electron Microscopy for Biologists*. 2nd edition, Wiley, New York
- Mehl E and Halaris A (1970) Stoichiometric relation of protein components in cerebral myelin from different species. *J Neurochem* 17, 659-68
- Meier E, Regan C, Balazs R and Wilkin G P (1982) Specific recognition of the neuronal cell surface by an antiserum raised against plasma membrane preparations of immature rat cerebellum. *Neurochem Res* 7, 1031-43
- Meller K, Breipohl W and Glees P (1968) The cytology of the developing molecular layer of mouse motor cortex: An electron microscopical and Golgi impregnation study. *Z Zellforsch Mikrosk Anat Abt Histochem* 86, 171-83
- Meryman H T and Kafig E (1955) The study of frozen specimens, ice crystals and ice crystal growth by electron microscopy. *Naval Med Res Inst Rept* 13, 529-44
- Metchnikoff E (1893) *Lectures on Comparative Pathology of Inflammation*. Translated by Starling F A. Kegan Paul, Trench and Trubner, London
- Metz A (1926) Die drei Gliazellarten und der Eisenstoffwechsel. *Z Ges Neurol Psychiat* 100, 428-49
- Metz A and Spatz H (1924) Die Hortegaschen Zellen (= das sogenannte "dritte Element") und über ihre funktionelle Bedeutung. *Z Ges Neurol Psychiat* 89, 138-70
- Meyer A and Meyer M (1945) Boutons terminaux in the cerebral cortex. *J Anat* 79, 180-5
- Meynert T (1867) Der Bau des Gross-Hirnrinde und seine ortliche Verscheidenheiten, nebst einen pathologisch-anatomischen Corollarium. *Vjschr Psychiat* 1, 77-99; 125-217; (1868) 2, 88-113; 381-403
- Michell R H, Finean J B and Coleman R (1982) Widely accepted modern views of cell structure are fundamentally correct. A reply to Hillman and Sartory, *School Science Review*, 63, 434-41
- Mikata A and Luse S A (1964) Ultrastructural changes in the rat liver produced by N-2-fluorenyldiacetamide. *Am J Pathol* 44, 455-79
- Millhouse E O (1981) The Golgi methods. In *Neuroanatomical Tract Tracing Methods*. eds. Reimer L and Robards M J. Plenum, New York, pp. 311-44
- Minckler J (1942) Pathologic alterations in surface relationships and morphology at the human synapse. *Am J Pathol* 18, 1061-1104
- Minkowski M (1923) Etude sur les connexions anatomiques des circonvolutions rolandiques, pariétales et frontales. *Schweiz Arch Neurol Psychiat* 12, 71-104
- Mirsky R, Wendon L M B, Black P, Stolkin C and Bray D (1978) Tetanus toxin: a cell surface marker for neurons in culture. *Brain Res* 148, 251-9
- Moll R, Franke W W, Schiller D L, Geiger B and Krepler R (1982) The catalog of human cytokeratins: patterns of expression in normal epithelia tumours and cultured cells. *Cell* 31, 11-24
- Moonen G, Cam Y, Sensenbrenner M and Mandel P (1975) Variability of the effects of serum-free medium, dibutyryl-cyclic AMP or theophylline on the morphology of cultured new-born rat astroblasts. *Cell Tissue Res* 163, 365-72
- Moor H (1969) Freeze etching. *Int Rev Cytol* 26, 391-412
- Moore B W (1972) Chemistry and biology of two proteins, S-100 and 14-3-2, specific to the nervous system. *Int Rev Neurobiol* 15, 215-25
- Moore R Y and Bloom F E (1978) Central catecholamine neuron systems. Anatomy and physiology of the dopamine systems. *Ann Rev Neurosci* 1, 129-69
- Moore R Y and Bloom F E (1979) Central catecholamine neuron systems. Anatomy and physiology of the norepinephrine systems. *Ann Rev Neurosci* 2, 113-63
- Moore W M, Holladay L A, Purett D and Brady R N (1974) On the conformation of the acetylcholine receptor proteins from *Torpedo Nobiliana*. *FEBS Lett* 45, 145-9
- Morgan J F, Morton H J and Parker R C (1950) Nutrition of animal cells in tissue culture. *Proc Soc Exp Biol* 73, 1-8
- Mori S and Leblond C P (1969) Identification of microglia in light and electron microscopy. *J Comp Neurol* 135, 57-80
- Morowitz H J and Terry T (1969) Characterization of the plasma membrane of *Mycoplasma Laidlawii*. V. Effects of selective removal of protein and lipid. *Biochim Biophys Acta* 183, 276-94

REFERENCES

- Morrison A B (1959) The distribution of intravenously injected inulin in the fluids of the nervous system of the dog and rat. *J Clin Invest* 38, 1769-77
- Mueller P and Rudin D P (1968) Resting and action potentials in bimolecular lipid membranes. *J Theoret Biol* 18, 222-8
- Mugnaini E, Osen K K, Dahl A L, Friedrich V L and Korte K (1980) Fine structure of granule cells and related interneurons (termed Golgi cells) in the cochlear nuclear complex of cat, rat and mouse. *J Neurocytol* 9, 537-70
- Mugnaini E and Walberg F (1964) Ultrastructure of neuroglia. *Ergebn Anat Entwickl Gesch* 37, 194-236
- Muhlethaler K, Moor H and Szarkowski J W (1965) The ultrastructure of the chloroplast lamellae. *Planta* 67, 305-23
- Muller G (1930) Zur Frage der Alterbestimmung histologischer Veränderungen im menschlischen Gehirn unter Berücksichtigung der örtlichen Verteilung. *Z Ges Neurol Psychiat* 124, 1-112
- Muller K J and McMahan U J (1976) The shapes of sensory and motor neurons and the distribution of their synapses in ganglia of the leech. A study using intracellular injection of horse radish peroxidase. *Proc R Soc Lond B* 194, 481-99
- Murai S (1973) Experimental studies on the cerebral metabolism in head injury. With special reference to respiratory function of mitochondria and carbonic anhydrase activity of rabbits cerebral cortex fractions containing isolated, metabolically active neuronal and glial cells. *Nippon Univ Med J* 23, 583-605
- Murray M R (1958) Response of oligodendrocytes to serotonin. In *Biology of Neuroglia*. ed. Windle W F, C C Thomas, Springfield, Ill., pp. 176-90
- Murray M R (1965) Nervous tissue in vitro. In *Cells and Tissues in Culture: Methods, Biology and Physiology*. ed. Willmer E N. Vol 2, pp. 373-455
- Nachmansohn D and Neumann E (1975) *Chemical and Molecular Basis of Nerve Activity*. Academic Press, New York, pp. 1-403
- Nagata Y, Mikoshiba K and Tsukada Y (1974) Neuronal cell body enriched and glial cell enriched fractions from young and adult rat brains: preparation and morphological and biochemical properties. *J Neurochem* 22, 493-503
- Nageotte J (1932) Sheaths of the peripheral nerves. Nerve degeneration and regeneration. In *Cytology and Cellular Pathology of the Nervous System*. ed. Penfield W. Paul B Hoeber, Vol 1, pp. 191-239
- Nairn R C (ed.) (1969) *Fluorescent Protein Tracing*. 3rd edition, Livingstone, Edinburgh
- Nakai J (1963) *Morphology of Neuroglia*. Igaku Shoin, Tokyo
- Nathaniel E J H and Nathaniel D R (1966) Fine structure of the neurons of the posterior horn in the rat spinal cord. *Anat Rec* 155, 629-41
- Nauta W J H and Ebbeson S O E (eds.) (1970) *Contemporary Research Methods in Neuroanatomy*. Springer-Verlag, New York, (several papers)
- Nauta W J H and Feirtag M (1979) The organization of the brain. *Sci Am*, September 1979, pp. 88-111
- Nelson P G (1973) Electrophysiological studies of normal and neoplastic cells in tissue culture. In *Tissue Culture of the Nervous System*. ed. Sato G. Plenum, New York, pp. 135-60
- Nelson P G and Lux H D (1970) Some electrical measurements of motoneuron parameters. *Biophys J* 10, 55-73
- Neuhoff V (1973) *Micromethods in Molecular Biology*. Springer, Berlin
- Niessing K, Scharrer E, Scharrer B and Oksche A (1980) *Handbuch der Microskopischen Anatomie des Menschen*. 4. Band Nervensystem. 10 Teil. *Neuroglia* 1, pp. 1-156
- Nishii B (1929) Zur Kenntnis der diffusen Sarkomatose des Nervensystems. *Arb Neurol Inst, Wien*, 31, 116-29
- Nissl F (1894) Über die sogenannten Granula der Nervenzellen. *Neurol Zentralbl* 13, 676-85, 781-9
- Nissl F (1899) Ueber einige Beziehungen zur Nervenzellenerkrankungen und gliosen, Erscheinungen bei verschieden Psychosen. *Arch Psychiat* 32, 656-76
- Noonan T R, Fenn W O and Haege L (1941) The distribution of injected radioactive potassium in rats. *Am J Physiol* 132, 474-88
- Norenberg M D and Martinez-Hernandez A (1979) Fine structural localization of glutamine synthetase in astrocytes of rat brain. *Brain Res* 161, 303-10
- Norton W T and Poduslo S E (1973) Characteristics of concomitantly isolated myelin, in *Research Methods in Neurochemistry*. eds. Marks N and Rodnight R. Plenum, New York, Vol 5, p. 19

REFERENCES

- Novikoff A B (1967) Enzyme localization and ultrastructure of neurons. In *The Neuron*, ed. Hyden H. Elsevier Publishing Company, Amsterdam, pp. 255-318
- Novikoff A B and Goldfischer S (1961) Nucleoside diphosphatase activity in the Golgi apparatus and its usefulness in cytological studies. *Proc Natl Acad Sci USA* 47, 802-10
- Nurnberger J I (1958) Direct enumeration of cells of the brain. In *Biology of Neuroglia*. ed. Windle W F. pp. 193-210
- Nurnberger J I and Gordon W H (1957) The cell density of neural tissues. In *Ultrastructure and Cellular Chemistry of Neural Tissue*. ed. Waelsch H. Cassell, London, pp. 100-38
- Obata K (1972) The inhibitory action of γ -amino butyric acid, a probable synaptic transmitter. *Int Rev Neurobiol* 15, 167-87
- Obersteiner H (1890) *The Anatomy of the Central Nervous Organs in Health and Disease*. Translated by Hill A. Griffin, London
- O'Brien R D, Eldefrawi M E and Eldefrawi A T (1972) Isolation of acetylcholine receptors. *Pharmacol Rev* 12, 19-34
- Ochoa J (1976) The unmyelinated nerve fibre. In *The Peripheral Nerve*. ed. Landon D N. Chapman and Hall, London, pp. 106-58
- Oehmichen M (1978) Mononuclear Phagocytes in the Central Nervous System. Translated by Clarkson M M. Springer, Berlin, pp. 1-173
- Oehmichen M, Gruninger H, Saebich R and Narita Y (1973) Microglia und Pericyten als Transformationsformen der Blut-Monocyten mit erhaltener Proliferations Fahigkeit. *Acta Neuropathol* 23, 200-18
- Oldendorf W H and Davson H (1967) Brain extracellular space and sink action of cerebrospinal fluid. *Arch Neurol Chicago* 17, 196-205
- Orkand R K (1977) Glial cells. In *Handbook of Physiology*. ed. Kandel E R. American Physiological Society, Bethesda, Vol 1, pages 855-75
- Osborne N N (1974) *Micro-chemical Analysis Of Nervous Tissue*. Pergamon Press, Oxford
- Osborne N N (1981) Communication between neurones. Current concepts. *Neurochem Int* 3, 3-16
- Osterhout W J V (1935) How do electrolytes enter the cell? *Proc Natl Acad Sci, USA* 21, 125-32
- Overton E (1895) Über die osmotischen Eigenschaften der lebenden Pflanzen und Tierzelle. *Vjschr Naturforsch Ges, Zurich* 40, 159-201
- Overton E (1902) Beiträge zur allgemeinen Muskel-und Nerven physiologie. *Pfl Archiv Ges Physiol* 92, 115-280
- Packman P M, Blomstrand C and Hamberger A (1971) Disc electrophoretic separation of proteins in neuronal, glial and subcellular fractions from cerebellar cortex. *J Neurochem* 18, 479-87
- Palade G E (1953) An electron microscopic study of the mitochondrial structure. *J Histochem Cytochem* 1, 188-211
- Palade G E (1954) Electron microscopic observations of interneuronal and neuromuscular synapses. *Anat Rec* 118, 335-6
- Palay S L (1954) Electron microscope study of the cytoplasm of neurons. *Anat Rec* 118, 336
- Palay S L (1956a) Structure and function in the neuron. In *Progress in Neurobiology*, Vol 2. *Ultrastructure and Cellular Chemistry of Neural Tissue*. ed. Waelsch H. Hoeber, New York, pp. 31-44
- Palay S L (1956b) Synapses in the central nervous system. *J Biophys Biochem Cytol* 2 (Suppl) 193-202
- Palay S L (1958a) An electron microscopical study of neuroglia. In *Biology of Neuroglia*, ed. Windle W F. C C Thomas, Springfield, Ill. pp. 24-49
- Palay S L (1958b) The morphology of synapses in the central nervous system. *Exp Cell Res, Suppl* 5, 275-93
- Palay S L (1958c) The morphology of secretion. In *Frontiers of Cytology*. ed. Palay S L. Yale University Press, New Haven, pp. 305-42
- Palay S L (1964) The structural basis for neural action. In *Brain Function*, Vol 2, RNA and Brain Function, Memory and Learning. ed. Brazier M A B, University of California Press, Berkeley, pp. 69-108
- Palay S L and Chan-Palay V (1972) The structural heterogeneity of central nervous tissue, in *Metabolic Compartmentation in the Brain*. eds. Balazs R and Cremer J E. Macmillan, London, pp. 187-207

REFERENCES

- Palay S L and Chan-Palay V (1977) General morphology of neurons and neuroglia. In *Handbook of Physiology, Section 1, Vol 1, Pt. I, Nervous System. Cellular Biology of Neurons.* Williams and Wilkins, Baltimore, pp. 5-37
- Palay S L and Palade G E (1955) The fine structure of neurons. *J Biophys Biochem Cytol* 1, 69-88
- Palay S L, Billings-Gagliardi S and Chan-Palay V (1974) Neuronal perikarya with dispersed, single ribosomes in the visual cortex of Macaca mulatta. *J Cell Biol* 63, 1074-89
- Palmgren A (1948) A rapid method for selective silver staining of nerve fibres and nerve endings in mounted paraffin sections. *Acta Zool* 29, 377-93
- Pappas G D and Purpura D P (eds.) (1972) *Structure and Function of Synapses.* Raven Press, New York, pp. 1-308
- Pappas G D and Waxman S G (1972) Synaptic fine structure-morphological correlates of chemical and electrotonic transmission. In *Structure and Function of Synapses.* eds. Pappas G D and Purpura D P. Raven Press, New York, pp. 1-43
- Pappenheimer J R, Hersey S R, Jordan E F and Downer J de C (1962) Perfusion of the cerebral ventricular system in anaesthetized goats. *Am J Physiol* 203, 763-74
- Pappius H and Elliott K A C (1956) Water distribution in incubated slices of brain and other tissues. *Canad J Biochem Physiol* 34, 1007-22
- Parsons D F (1974) Structure of wet specimens in electron microscopy, *Science* 186, 407-14
- Parsons D F (1978) High voltage microscopy of cells and membranes in the hydrated states. In *Frontiers of Biological Engineering: from Electrons to Tissues.* eds. Dutton P L, Leigh J S and Scarpa A. Vol 1, Academic Press, New York, p. 734
- Patay R (1934) Une nouvelle technique de coloration trichromique pour l'histologie. *Bull d'Histologie Appliquee a la Physiologie et a la Pathologie et de Technique Microscopique*, 11, 408-9
- Pearse A G E (1972) *Histochemistry.* Vol 2, 3rd edition, Longman, London
- Pearse A G E (1980) *Histochemistry.* Vol 1, 2nd edition, Longman, London
- Penfield W (1924) Oligodendroglia and its relation to classical neuroglia. *Brain* 47, 430-52
- Penfield W (1928) Neuroglia and microglia. The interstitial tissue of the central nervous system. In *Special Cytology.* ed. Cowdry E V. Hoeber, New York, Vol 2, pp. 1032-69
- Penfield W (1930) A further modification of Del Rio-Hortega's method of staining oligodendroglia. *Am J Pathol* 6, 445-7
- Penfield W (1932) Neuroglia: normal and pathological. In *Cytology and Cellular Pathology of the Nervous System.* ed. Penfield W. Hoeber, New York, Vol 2, pp. 421-79
- Penfield W and Erickson T C (1941) Epilepsy and Cerebral Localisation. A Study of the Mechanism, Treatment and Prevention of Epileptic Seizures. C C Thomas, Springfield, Ill., pp. 1-623
- Pepeu G and Kuhar M J (eds.) (1980) *Receptors for Neurotransmitters and Peptide Hormones.* Raven Press, New York
- Peters A (1968) The morphology of axons of the central nervous system. In *The Structure and Function of Nervous Tissue.* ed. Bourne G H, Vol 1, Structure 1, Academic Press, New York, pp. 141-86
- Peters A and Vaughn J E (1967) Microtubules and filaments in the axons and astrocytes of early postnatal rat optic nerves. *J Cell Biol* 32, 113-19
- Peters A, Palay S L and Webster H de F (1976) *The Fine Structure of the Nervous System.* Saunders Co, Philadelphia, pp. 1-395
- Peters A, Proskauer C C and Kaiserman-Abramof I R (1968) The small pyramidal neuron of the rat cerebral cortex. *J Cell Biol* 39, 604-29
- Peterson E R and Crain S M (1970) Innervation in cultures of foetal rodent skeletal muscle by organotypic explants of spinal cord from different animals. *Z Zellforsch* 106, 1-21
- Peterson E R and Murray M R (1955) Myelin sheath formation in cultures of avian spinal ganglia. *Am J Anat* 96, 319-56
- Pevzner L Z (1978) Phasic interrelations between neuronal and neuroglial macromolecular metabolism. In *Dynamic Properties of Glia Cells.* eds. Schoffeniels E, Franck G, Hertz L and Tower D B. Pergamon Press, Oxford, pp. 223-30
- Pevzner L Z (1979) *Functional Biochemistry of the Neuroglia.* Consultants Bureau, New York, pp. 1-306
- Phalen G S and Davenport H A (1937) Pericellular end-bulbs in the central nervous system of vertebrates. *J Comp Neurol* 68, 67-82
- Pickel V M (1981) Immunocytochemical methods. In *Neuroanatomical Tract Tracing Methods.* eds. Heimer L and Robards M J. Plenum, New York, pp. 483-509

REFERENCES

- Pieringer J, Subba Rao G, Mandel P and Pieringer R A (1977) The association of the sulphogalactosylglycerolipid of rat brain with myelination. *Biochem J* 166, 421-8
- Polak M (1956) Sobre una variante a la tecnica de Rio-Hortega para la impregnacion de celulas reticuloendotheliales normales y patologicas micro y oligodendroglia "barrera epitelial argentofila" (Polak), elementos neuroglicos blastomatoses etc, *Arch Histol Norm Pat* 6, 220-2
- Pomerat C M (1958) Functional concepts based on tissue culture studies of neuroglia cells. In *Biology of Neuroglia*. ed. Windle W F. C C Thomas. Springfield, Ill., p. 162
- Pomerat C M and Costero I (1956) Tissue cultures of cat cerebellum. *Am J Anat* 99, 211-47
- Pomerat C M, Hendleman W J, Raiborn C W and Massey J F (1967) Dynamic activities of nervous tissue in vitro. In *The Neuron*. ed. Hyden H. Elsevier, Amsterdam, pp. 119-78
- Porter K R (1966) Cytoplasmic microtubules and their function. In *Principles of Biomolecular Organization*. eds. Wolstenholme G E W and O'Connor, M. Churchill, London, pp. 308-45
- Porter K R, Claude A and Fullam E F (1945) A study of tissue cells by electron microscopy. *J Exp Med* 81, 233-46
- Prince D A (1972) Topical convulsant drugs and metabolic antagonists. In *Experimental Models of Epilepsy*. eds. Purpura D P, Penry J K, Tower D B, Woodbury D M and Walter R D. Raven Press, New York, pp. 51-84
- Privat A (1978) Morphological approaches to the problems of neuroglia. Some comments in *Dynamic Properties of Glia Cells*, ed. Schoffeniels E, Franck G, Hertz L and Tower D B. Pergamon Press, Oxford, pp. 55-64
- Pruijns W M (1927) Über Mikroglia, ihre Herkunft, Funktion und ihr Verhältnis zu anderen Gliaelementen. *Z Ges Neurol Psychiat* 108, 298-315
- Purkinje J E (1838) Neueste Untersuchungen aus der Nerven und Hirn-Anatomie, in Bericht Über Die Versammlung Deutscher Naturforscher Und Aerzte In Prag im September, 1837. Hasse, Prague, pp. 177-180
- Puszkin S and Berl S (1972) Actinomysin-like protein from brain. *Biochim Biophys Acta* 256, 695-709
- Quarles R H, Everly J L and Brady R O (1973) Evidence for the close association of a glycoprotein with myelin in rat brain. *J Neurochem* 21, 1177-91
- Radin N S, Brenkert A, Arora R C, Sellinger O Z and Flangas A L (1972) Glial and neuronal localization of cerebroside-metabolizing enzymes. *Brain Res* 39, 163-9
- Raff M C, Brockes J P, Fields K L and Mirsky R (1980) Neural cell markers: the end of the beginning. Development and chemical specificity of neurons. In *Progress in Brain Research*. eds. Cuenod M, Kreutzberg G W and Bloom F E, Elsevier/North Holland Biomedical Press, Amsterdam, Vol 51, pp. 17-22
- Raff M C, Fields K L, Hakamori S I, Mirsky R, Pruss R and Winter J (1979) Cell-type specific markers for distinguishing and studying neurons and major classes of glial cells in culture. *Brain Res* 174, 283-308
- Raff M C, Mirsky R, Fields R P, Lisak R P, Dorfman S H, Silberberg D H, Gregson N A, Leibowitz S and Kennedy M C (1978) Galactocerebroside is a specific cell-surface marker for oligodendrocytes in culture. *Nature* 274, 813-16
- Raghavan S S, Rhoads D B and Kanfer J N (1972) Acid hydrolases in neuronal and glial enriched fractions of rat brain. *Biochim Biophys Acta* 268, 755-60
- Raine C S and Wisniewski H (1970) On the occurrence of microtubules within mature astrocytes. *Anat Rec* 167, 303-8
- Raisman G and Matthews M R (1972) Degeneration and regeneration of synapses. In *The Structure and Function of Nervous Tissue*. ed. Bourne G H. Academic Press, London, Vol 4, pp. 61-104
- Ralis H M, Beasley R A and Ralis Z A (1973) *Techniques in Neurohistology*. Butterworths, London, pp. 1-162

REFERENCES

- Rall D P (1964) The structure and function of the cerebrospinal fluid. In The Cellular Functions of Membrane Transport. ed. Hoffman J F. Prentice-Hall, Englewood Cliffs, NJ, p. 269
- Rall D P, Moore E, Taylor N and Zubrod C G (1961) The blood-cerebrospinal fluid barrier in man. *Arch Neurol* 4, 318-22
- Rall D P, Moore E, Taylor N and Zubrod C G (1961) The blood-cerebrospinal fluid barrier in man. *Arch Neurol* 4, 318-22
- Rall D P, Oppelt W W and Patlak C S (1962) Extracellular space of brain as determined by diffusion of inulin from the ventricular system. *Life Sciences* 2, 43-48
- Ramon Y Cajal S (1892) El nuevo concepto de la histología de los centros nerviosos. *Revista de Ciencias Medicas de Barcelona*, 18, 457-476, in Neurological Classics in Modern Translation. eds Rottenburg D A and Hochberg F (1977) Hafner Press, New York, pages 7-29
- Ramon Y Cajal S (1909-1911) *Histologie du Systeme Nerveux de L'Homme et des Vertébres.* 2 Volumes, Maloine, Paris
- Ramon Y Cajal S (1913a) Sobre un nuevo proceder de impregnacion de la neuroglia y sus resultados en los centros nerviosos del hombre y animales. *Trab Lab Invest Biol Univ Madr* 11, 219-237
- Ramon Y Cajal S (1913b) Contribucion al conocimiento de la neuroglia del cerebro humano. *Trab Lab Invest Biol Univ Madr* 11, 255-315
- Ramon Y Cajal S (1932-1933) La retine des Vertébres, Suppl 1-44. *Trav Lab Rech Biol Univ Madr* 28, Plate VII
- Ramon Y Cajal S (1954) Neuron Theory or Reticular Theory. Translated by Ubeda Purkiss M and Fox C A. Consejo Superior de Investigaciones Cientificas. Instituto Ramon Y Cajal, Madrid, pages i-xv, 1-144
- Ramon-Moliner E (1958) A study of neuroglia: the problem of transitional forms. *J Comp Neurol* 110, 157-171
- Ramon-Moliner E (1968) The morphology of dendrites, in The Structure and Function of Nervous Tissue. ed Bourne G H. Academic Press, New York, Vol 1, pages 205-267
- Ranck J B (1963a) Specific impedance of rabbit cerebral cortex. *Exp Neurol* 7, 144-53
- Ranck J B (1963b) Analysis of specific impedance of rabbit cerebral cortex. *Exp Neurol* 7, 153-74
- Ranson S W and Clark S L (1948) The Anatomy of The Nervous System. 1st edition. Saunders, Philadelphia, pp. 101-3
- Ranson S W and Clark S L (1959) The Anatomy of The Nervous System. 2nd edition. Saunders, Philadelphia, pp. 125-6
- Ranvier L-A (1871) Contributions a l'histologie et a la physiologie des nerfs périphériques. *Compt Rend Hebd Acad Sci, Paris*, 73, 1168-71
- Rapaport S I (1976) Blood Brain Barrier in Physiology and Medicine. Raven Press, New York
- Rapoport A and Stempak J (1968) Ultrastructure of the ventral horn cell in the albino rat. *Anat Rec* 161, 361-76
- Rash J E and Hudson C S (eds.) (1979) Freeze Fracture: Methods, Artifacts and Interpretations. Raven Press, New York
- Reade J B (1837) On a new method of illuminating microscopic objects. In *Micrographia Containing Practical Essays on Reflecting, Solar, Oxy, Hydrogen Gas Microscopes, Micrometers, Eyepieces, etc.* Whittaker, London, pp. 227-31
- Reese T S and Shepherd G M (1972) Dendro-dendritic synapses in the central nervous system. In *Structure and Function of the Synapses*, eds. Pappas G D and Purpura D P. Raven Press, New York, pp. 121-36
- Reil J C (1809) Untersuchung über den Bau des grossen Gehirns im Menschen. Vierte Fortsetzung XIII. *Arch Physiol*, Halle, 9, 136-46
- Reimer L (1965) Irradiation changes in organic and inorganic objects. *Lab Invest* 14, 344/1082-358/1096
- Reisner J H (1965) Determination of magnification in electron microscopy. I Instrumental factors influencing estimate of magnification. *Lab Invest* 14, 137/875-141/880

REFERENCES

- Remak R (1836) Verlaufige Mittheilung mikroskopischer Beobachtungen über den innern Bau der Cerebrospinalnerven und über die Entwicklung ihrer Formelemente. *Arch Anat Physiol*, 145-161
- Remak R (1838) *Observations Anatomicae et Microscopicae de Systematis Nervosi Structura*. Reimerianis, Berlin (thesis)
- Repke K R H (1980) Structural basis of the compartmentalization functions of bio-membranes. In *Cell Compartmentation and Metabolic Channeling*. eds. Nover L, Lynen F and Mothes K. Fisher, Jena and Elsevier, Amsterdam, pp. 33-46
- Rhodin J A G (1963) *An Atlas of Ultrastructure*. Saunders, Philadelphia
- Rhodin J A G (1975) *An Atlas of Histology*. Oxford University Press, New York
- Robards A W (1981) Review of The Living Cell. *Proc Microsc Soc* 16, 317-20
- Roberts E (1962) γ -Aminobutyric acid and neuronal function. In *Ultrastructure and Metabolism of the Nervous System*. Association for Research in Nervous and Mental Diseases, Williams and Wilkins, Baltimore, Vol XL, pp. 288-99
- Roberts K and Hyams J S (eds.) (1979) *Microtubules*. Academic Press, London, pp. 1-595
- Robertson J D (1959) The ultrastructure of cell membranes and their derivatives. In *Biochemical Symposia*, Vol 16, The Structure and Function of Subcellular Components. Cambridge University Press, Cambridge, pp. 3-43
- Robertson J D (1969) Molecular structure of biological membranes. In *Handbook of Molecular Cytology*. ed. Lima de Faria A. North Holland, Amsterdam, pp. 1403-4
- Robertson J D (1972) The unit membrane of cells and mechanisms of myelin formation. In *Ultrastructure and Metabolism of The Nervous System*, Vol XL, pages 94-158, Research Publications, ARNMD, USA
- Robertson J D (1980) *The Anatomy of Biological Interfaces in Membrane Physiology*. eds. Andreoli T E, Hoffman J F and Fanestil D D. Plenum Medical, New York, pp. 1-26
- Robertson W F (1899) Normal and pathological history of the nerve cell. *Brain* 22, 203-327
- Rodnight R (1975) Constituents of neural tissues: fluids, some electrolytes, amino acids and ammonia. In *Practical Neurochemistry*. ed. McIlwain H. Churchill-Livingstone, Edinburgh, pp. 17-32
- Rodriguez de Lores Arnaiz G, Alberici de Canal M and De Robertis E (1971) Turnover of proteins in subcellular fractions of rat cerebral cortex. *Brain Res* 31, 179-84
- Romeis B (1968) *Mikroskopische Technik*. Oldenbourg, Munich
- Roots B I (1978) A phylogenetic approach to the anatomy of glia. In *Dynamic Properties of Glia Cells*. eds. Schoffeniels E, Franck G, Hertz L and Tower D B, Pergamon Press, Oxford, pp. 45-54
- Roots B I and Johnston P V (1964) Neurons of ox brain nuclei: their isolation and appearance by light and electron microscopy. *J Ultrastruct Res* 10, 350-61
- Roots B I and Johnston P V (1965) Isolated rabbit neurons: electron microscopical observations. *Nature*, 207, 315-16
- Ross K F A (1953) Cell shrinkage caused by fixatives and paraffin-wax embedding in ordinary cytological preparations. *Q J Microsc Sci* 94, 125-39
- Ross L L (1964) Peripheral nervous tissue. In *Electron Microscopic Anatomy*. ed. Kurtz S M, Academic Press, New York, pp. 341-67
- Rottenberg D A and Hochberg F H (eds.) (1977) *Neurological Classics in Modern Translation*. Hafner Press, New York
- Rouiller C and Simon G (1962) Contribution de la microscopie electronique au progrès de nos connaissances en cytologie et en histo-pathologie hépatique. *Rev Intern Hepatol* 12, 167-206
- Rozsa G, Morgan C, Szent-Gyorgyi A and Wyckoff R W G (1959) The electron microscopy of myelinated nerve. *Biochem Biophys Acta* 6, 13-27
- Rudolph G G and Olsen N S (1956) Transfer of potassium between blood, cerebrospinal fluid and brain tissue. *Am J Physiol* 186, 157-60
- Ruska E (1980) The Early Development of Electron Lenses and Electron Microscopy, translated by Mulvey T. Hirzel, Stuttgart, pp. 2-8
- Russell D S and Rubinstein L J (1977) *Pathology Of Tumours Of The Nervous System*. 4th edition. Edward Arnold, London
- Rydberg E (1932) Cerebral injury in new-born children consequent on birth trauma; with an inquiry into the normal and pathological anatomy of the neuroglia. *Acta Pathol Microbiol Scand* 10 (Suppl), 1-247

REFERENCES

- Salomon J C (1962) Modifications des cellules du parenchyme hépatique du rat sous l'effet de la thioacetamide. *J Ultrastruct Res* 7, 293-307
- Sartory P, Fasham J and Hillman H (1971) Microscopical observations of the nucleoli in unfixed rabbit Deiters' neurons. *Microscopy* 32, 93-100
- Satake M (1972) Some aspects of the protein metabolism of the neuron. *Int Rev Neurobiol* 15, 189-213
- Sauer F C (1935a) Mitosis in the neural tube. *J Comp Neurol* 62, 377-405
- Sauer F C (1935b) The cellular structure of the neural tube. *J Comp Neurol* 63, 13-23
- Schade J P and Ford D H (1965) Basic Neurology. Elsevier, Amsterdam, p. 101
- Schlaepfer W W and Torack R M (1966) The ultrastructural localization of cholinesterase activity in the sciatic nerve of the rat. *J Histochem Cytochem* 14, 27-50
- Schleiden M J and Schwann T (1838) Contributions to Phytogenesis. English translation, Sydenham Society, London, published in 1847
- Schmidt W J (1936) Doppelbrechung und Feinbau der Markscheide der Nervenfasern. *Z Zellforsch* 23, 657-76
- Schmidt H D (1874) On the construction of the dark or double bordered nerve fibre. *Mon Microsc J Lond* 11, 200-21
- Schmitt F O (1936) Nerve ultrastructure as revealed by x-ray diffraction and polarized light studies. *Cold Spring Harbor Symposia* 4, 7-12
- Schmitt F O (1939) The ultrastructure of protoplasmic constituents. *Physiol Rev* 19, 270-302
- Schmitt F O (1958) Axon-satellite cell relations in peripheral nerve fibres. *Exp Cell Res, Suppl* 5, 33-57
- Schmitt F O and Bear R S (1939) The ultrastructure of the nerve axon sheath. *Biol Rev* 14, 27-50
- Schmitt F O, Bear R S and Clark G L (1935) X-ray diffraction studies on nerve. *Radiology* 25, 131051
- Schmitt F O, Bear R S and Palmer K J (1941) X-ray diffraction studies on the structure of the nerve myelin sheath. *J Cell Comp Physiol* 18, 31-42
- Schnapp B and Mugnaini E (1975) The myelin sheath: Electron microscopic studies with thin sections and freeze-fracture. *Golgi Centennial Symposium Proceedings*, ed. Santini M, Raven Press, New York
- Schoffeniels E, Franck G, Tower D B and Hertz L (eds.) (1978) Dynamic Properties of Glial Cells. Pergamon Press, Oxford, pp. 1-467
- Schousboe A (1978) Metabolism and function of GABA in the mammalian brain. Possible role of glia cells. University of Copenhagen, thesis, pp. 1-83
- Schousboe A and Hertz L (1971) Effects of potassium on indicator spaces and fluxes in slices of brain cortex from adult and new born rats. *J Neurochem* 18, 67-77
- Schousboe A, Nissen C, Bock E, Sapirstein V S, Juurlink B H K and Hertz L (1980) Biochemical development of rodent astrocytes in primary cultures. In *Tissue Culture in Neurobiology*, eds. Giacobini E, Vernadakis A and Shahar A, Raven Press, New York, pp. 397-409
- Schousboe A, Svenneby G and Hertz L (1977) Uptake and metabolism of glutamate in astrocytes cultured from dissociated mouse brain hemispheres. *J Neurochem* 29, 999-1005
- Schroder P (1914) Die vordere Zentralwindung bei Lasionen der Pyramidenbahn und bei amyotropischer Lateralsklerose. *Monatschr Psychiatr Neurol* 35, 1-25
- Schultz R L (1964) Macrogliad identification in electron micrographs. *J Comp Neurol* 122, 281-95
- Schultz R, Maynard E A and Pease D C (1957) Electron microscopy of neurons and neuroglia of the cerebral cortex and corpus callosum. *Am J Anat* 100, 369-407
- Schurch-Rathgeb Y and Monard D (1980) Dissociated brain cell cultures and established cell lines as tools to study the cellular regulation of neurite extension, in *Tissue Culture in Neurobiology*, eds. Giacobini E, Vernadakis A and Shahar A, Raven Press, New York, pp. 321-32
- Schwann T (1838) Ueber die Analogie in der Structur und dem Wachsthum der Thiere und Pflanzen. *Froriep's Neu Notizen* 5, p. 228
- Schwann T (1839) Mikroskopische Untersuchungen über die Uebereinstimmung in der Struktur und dem Wachsthum der Thiere und Pflanzen. Reimer, Berlin
- Schwartz A, Bachelard H S and McIlwain H (1962) The sodium-stimulated adenosine-triphosphatase activity and other properties of cerebral microsomal fractions and subfractions. *Biochem J* 84, 626-37
- Scott T G (1967) The distribution of 5'-nucleotidase in the brain of the mouse. *J Comp Neurol* 129, 97-114

REFERENCES

- Seil F J and Leiman A L (1977) Spontaneous versus driven activity in intracerebellar nuclei: a tissue culture study. *Exp Neurol* 54, 110-27
- Sellinger O Z and de Balbian Verster F (1962) Glutamine synthetase of rat cerebral cortex: intracellular distribution and structural latency. *J Biol Chem* 237, 2836-44
- Sellinger O Z and Santiago J C (1972) Unequal development of two mitochondrial enzymes in neuronal cell bodies and glial cells of rat cerebral cortex. *Neurobiol* 2, 133-46
- Semenza G and Carafoli E (eds.) (1977) Biochemistry of Membrane Transport. FEBS Symposium No. 42, Springer, Berlin
- Sensenbrenner M, Labourdette G, Delaunoy J P, Pettman B, Devilliers G, Moonen G and Bock E (1980) Morphological and biochemical differentiation of glial cells in primary culture. In *Tissue Culture in Neurobiology*, (eds.) Giacobini E, Vernadakis A, and Shahar A. Raven Press, New York, pp. 385-95
- Shahar A, Grunfeld Y, Spiegelstein M Y and Monzain R (1975) Myelinization in long-term cultures of dissociated mammalian neurons. *Brain Res* 88, 41-51
- Shantha T R, Manocha S L, Bourne G H and Ariens-Kappers J (1969) The morphology and cytology of neurons, in *The Structure and Function of Nervous Tissue*. Vol II, ed. Bourne G H. Academic Press, New York, pp. 1-67
- Sharpey-Schafer E (1885) *Essentials of Histology*. Longmans, London
- Shaw F H, Simon S E, Johnstone B M and Holman M E (1956) The effect of changes of the environment on the electrical and ionic pattern of muscle. *J Gen Physiol* 40, 263-88
- Shelanski M L (1973) Chemistry of the filaments and tubules of brain. *J Histochem Cytochem* 21, 529-39
- Shelanski M L, Liem R K H, Letterier J F and Keith K H (1981) The cytoskeleton and neuronal disease. Papers presented at the International Congress on Cell Biology. Berlin, August 31-September 5 1980, ed. Schweiger H G. Springer, Berlin, pp. 428-39
- Sherrington C S (1897) See Foster and Sherrington (1897)
- Sherrington C S (1906) *The Integrative Action of the Nervous System*. Constable, London, p. 16
- Sherrington C S (1925) Remarks on some aspects of reflex inhibition. *Proc R Soc B*, 97, 519-45
- Sholl D A (1953) Dendritic organization in the neurons of the visual and motor cortices of the cat. *J Anat* 87, 387-406
- Sholl D A (1956) *The Organization of the Cerebral Cortex*. Methuen, London, pp. 4-5
- Singer C (1921) Steps leading to the invention of the first optical apparatus. In *Studies in the History and Method of Science*. ed. Singer C. Vol II, Clarendon Press, Oxford, pp. 385-413
- Singer S J (1974) The molecular organization of membranes. *Ann Rev Biochem* 43, 805-33
- Singer S J and Nicholson G (1972) The fluid mosaic model of the structure of cell membranes. *Science* 175, 720-31
- Sinha A K and Rose S P R (1972) Compartmentation of lysosomes in neurones and neuropil and a new neuronal marker. *Brain Res* 39, 181-96
- Sjostrand F S (1950) Electron microscopic demonstration of a membrane structure isolated from nerve tissue. *Nature, Lond* 165, 482-3
- Sjostrand F S (1953a) Electron microscopy of mitochondria and cytoplasmic double membranes. *Nature* 171, 31-3
- Sjostrand F S (1953b) The lamellated structure of the nerve myelin sheath as revealed by high resolution electron microscopy. *Experientia* 9, 68-9
- Skou J C (1957) The influence of some cations on an adenosine triphosphate from peripheral nerves. *Biochim Biophys Acta* 23, 394-401
- Skou J C (1975) Identification and isolation of the components of the sodium pump. *Q Rev Biophys* 7, 401-34
- Sleytr U B and Robards A W (1977a) Plastic deformation during freeze cleaving: a review. *J Microsc* 110, 1-23
- Sleytr U B and Robards A W (1977b) Freeze fracturing: a review of methods and results. *J Microsc* 111, 77-100
- Sleytr U B and Robards A W (1982) Understanding the artefact problem in freeze-fracture replications: a review. *J Microsc* 126, 101-22
- Smart I and Leblond C P (1961) Evidence for division and transformation of neuroglia cells in the mouse brain as derived from radioautography after injection of thymidine-H³. *J Comp Neurol* 116, 349-67

REFERENCES

- Smith A and Bruton J (1977) A Colour Atlas of Histological Staining Techniques. Wolfe, London, pp. 19-173
- Smythies J R, Gibson W C and Purkis V A (1957) The distribution and morphology of boutons terminaux in the human cerebrum. *J Comp Neurol* 108, 173-223
- Solomon A K (1977) Reflections on the membrane mediated linkage between cation transport and glycolysis in human red cells, in Membrane Transport Processes. Vol 1, Raven Press, New York, pp. 31-59
- Somogyi P and Cowey A (1981) Combined Golgi and electron microscopic study on the synapses formed by double bouquet cells in the visual cortex of the cat and monkey. *J Comp Neurol* 195, 547-66
- Sotelo C and Palay S L (1968) The fine structure of the lateral vestibular nucleus in the rat. *J Cell Biol* 36, 151-79
- Spatz H (1918) Beitrag zur normalen Histologie des Rückenmarks des neugeborenen Kaninchens. *Histol Histopath Arb* 6, 478-604
- Sprague J M and Hongchien H A (1964) The terminal fields of dorsal root fibers in the lumbosacral spinal cord of the cat, and the dendritic organization of the motor nuclei. In Organization of the Spinal Cord. Progress in Brain Research, eds. Eccles J C and Schade J P, Elsevier, Amsterdam, Vol 2, pp. 120-54
- Stahl W L, Spence A M, Coates P W and Broderson S H (1978) Studies on cellular localization of Na^+ , K^+ -ATPase activity in nervous tissue. In Dynamic Properties of Glia Cells. eds. Schoffeniels E, Franck G, Hertz L and Tower D P. Pergamon Press, Oxford, pp. 371-81
- Stampfli R (1954) Saltatory conduction in nerves. *Physiol Rev* 34, 101-12
- Steere R L (1957) Electron microscopy of structural details in frozen biological specimens. *J Biophys Biochem Cytol* 3, 45-60
- Stein W D (1967) The Movement Of Molecules Across Cell Membranes. Academic Press, New York, pp. 1-369
- Steiner J W and Baglio C L (1963) Electron-microscopy of the cytoplasm of parenchymal liver cells in α -naphthylisocyanate-induced cirrhosis. *Lab Invest* 12, 765-90
- Steiner J W, Mujai K and Phillips M J (1964) Electron microscopy of membrane particle arrays in liver cells of ethionine-intoxicated rats. *Am J Pathol* 44, 169-214
- Stenger R J (1964) Regenerative nodules in carbon tetrachloride-induced cirrhosis. A light and electron microscopic study of lamellar structures encountered therein. *Am J Pathol* 44, 31A
- Stenwig A E (1972) The origin of brain macrophages in traumatic lesions, Wallerian degeneration and retrograde degeneration. *J Neuropathol Exp Neurol* 31, 696-704
- Sternberger L A (1974) Immunocytochemistry. Prentice Hall, Englewood Cliffs, NJ
- Stieda L (1899) Geschichte der Entwicklung der Lehre von den Nervenzellen und Nervenfasern während des XIX Jahrhunderts. 1. Teil: Von Sommering bis Deiters, Gustav Fischer, Jena pp. 1-118
- Stilling B and Wallach J (1842) Untersuchungen über den Bau des Nervensystems, Vol 1. Untersuchungen über den Bau des Nervensystems. Vol 2. Untersuchungen über die Textur des Rückenmarks. Wigand, Leipsiz.
- Stirling W (1881) Histological references to the structure of nerve fibres. *J Anat Physiol* 15, 446-7
- Stoeckenius W (1962) The molecular structure of lipid-water systems and cell membrane models studies with the electron microscope. In The Interpretation Of Ultrastructure. ed. Harris R J C. Academic Press, New York, pp. 349-67
- Stolinski C and Breathnach A S (1975) Freeze Fraction Replication of Biological Tissues: Techniques, Interpretation and Applications. Academic Press, London, pp. 1-173
- Stone W E (1972) Systemic chemical convulsants and metabolic derangements. In Experimental Models of Epilepsy. eds. Purpura D P, Penty J K, Tower D B, Woodbury D M and Walter R D. Raven Press, New York, pp. 407-32
- Stowell R E (1941) Effect on tissue volume of various methods of fixation, dehydration and embedding. *Stain Technol* 16, 67-73
- Streicher E (1961) Thiocyanate space of rat brain. *Am J Physiol* 201, 334-6
- Streit P, Akert K, Sandri C, Livingstone R B and Moor H (1972) Dynamic ultra-structure of presynaptic membranes at nerve terminals in spinal cord of rats. Anaesthetized and unanaesthetized preparations compared. *Brain Res* 48, 11-26
- Stricker S and Unger L (1879) Untersuchungen über den Bau der Grosshirnrinde. *Weiner Sitzungsber* 80, 450-9
- Struck D K and Lennarz W (1976) Utilization of exogenous GDP-mannose for the synthesis of mannose-containing lipids and glycoproteins by oviduct cells. *J Biol Chem* 251, 2511-19

REFERENCES

- Sultzmann R (1962) Die mikroskopische Morphologie der zentralen markhaltigen Nervenfaser. *Wis Z Friedrich-Schiller Univ Jena* 11, 197-245
- Sunderland S Sir (1978) Nerves and Nerve Injuries. Churchill Livingstone, Edinburgh, pp. 3-15
- Symposium on Nerve Growth Factor (1964) *Ann NY Acad Sci* 118, 1-147
- Symposium on Quantitative Electron Microscopy (1965) (Several papers), *Lab Invest* 14, 1-602
- Szentagothai J (1964) Propriospinal pathways and their synapses. In *Organization of the Spinal Cord*. eds. Eccles J C and Schade J P. *Progress in Brain Research*, Elsevier, Amsterdam, Vol 2, pp. 155-77
- Szentagothai-Schimert J and Albert A (1955) The synaptology of Clarke's column. *Acta Morphol Hung* 5, 43-51
- Taft A E (1938) Intercellular substance of cerebral cortex (Nissl's cerebral grey matter). Physiological significance. *Arch Neurol Psychiatry Chicago* 40, 313-21
- Taft A and Ludlum S (1929) On the nature of two forms of neuroglia. *J Nerv Ment Dis* 70, 365-8
- Tanaka R and Strickland K P (1965) Role of phospholipid in the activation of Na^+ , K^+ -activated adenosine triphosphatase of beef brain. *Arch Biochem Biophys* 111, 583-92
- Tani E and Ametani T (1970) Substructure of microtubules in brain nerve cells as revealed by ruthenium red. *J Cell Biol* 46, 159-65
- Tellyesniczky K (1898) Ueber die Fixierungs- (Haertungs) -Fluessigkeiten. *Arch Micr Anat* 52, 202-20
- Thorun W and Mehl E (1968) Determination of molecular weights of microgram quantities of protein components from biological membranes and other complex mixtures: gel electrophoresis across linear gradients of acrylamide. *Biochem Biophys Acta* 160, 132-4
- Tiegs O W (1927) A critical review of evidence on which is based the theory of discontinuous synapses of the spinal cord. *Austral J Exp Biol Med Sci IV*, 193-212
- Tiegs O W (1929) On the neurofibrillary structure of nerve cells. *Austral J Exp Biol Med Sci*, 6, 111-18
- Torack R M, Terry R D and Zimmerman H M (1969) Swelling produced by triethyltin poisoning and its comparison with that in the human brain. *Am J Pathol* 36, 273-87
- Tower D B (1960) Neurochemistry of Epilepsy. C C Thomas, Springfield, Ill.
- Tower D B (1969) Neurochemical mechanisms. In *Basic Mechanisms of the Epilepsies*. eds. Jasper H H, Ward A A Jr, and Pope A. Little, Brown and Company, Boston, pp. 611-38
- Tower D B (1978) General perspectives and conclusions of the symposium on dynamic properties of glial cells. In *Dynamic Properties of Glial Cells*. eds. Schoffeniels E, Franck G, Hertz L and Tower D B. Pergamon Press, Oxford, pp. 443-60
- Tower S S (1940) Units for sensory reception in cornea, with notes on nerve impulses from sclera, iris and lenses. *J. Neurophysiol* 3, 486-500
- Treip C S (1978) A Colour Atlas of Neuropathology. Wolfe, London, p. 153
- Triggle D J (1967) Chemical Aspects of the Autonomic Nervous System. Academic Press, London
- Troshin A S (1966) Problems of Cell Permeability. Pergamon Press, Oxford, pp. 1-549
- Truex R C and Carpenter M B (1969) Human Neuroanatomy. 6th edition. Williams and Wilkins, Baltimore
- Uesugi S, Dulak N C, Dixon N C, Hexum J F, Dahl T D, Perdue J F and Hokin L E (1971) Studies on the characterization of Na-K transport adenosine triphosphatase. *J Biol Chem* 246, 531-43
- Ungerstedt U (1971) Stereotaxic mapping of the monoamine pathways in the rat brain. *Acta Physiol Scand (Suppl)* 367, 1-48
- Urechia G I and Elekes N (1925) Contribution a l'etude des plaques seniles. Role de la microglia. *Bull Acad Med* 94, 43
- Ussing H H (1949) Transport of ions across cellular membranes. *Physiol Rev* 29, 127-55
- Uyeda C T, Eng L F and Bignami A (1972) Immunological study of the glial fibrillary acidic protein. *Brain Res* 37, 81-9
- Uyemura K, Tobari C, Hirano S and Tsukada T (1972) Comparative studies on the myelin proteins of bovine peripheral nerve and spinal cord. *J Neurochem* 19, 2607-14

REFERENCES

- Valdivia O (1971) Methods of fixation and the morphology of synaptic vesicles. *J Comp Neurol* 142, 257-74
- Valentin G G (1836) Über den Verlauf und die letzten Ende der Nerven. *Nova Acta Phys med Acad Caes Leopold Carol Nat Curiosorum*, Breslau 18, 51-240
- Van der Loos H (1960) On dendro-dendritic junctions in the cerebral cortex, in *Structure and Function of the Cerebral Cortex*. eds. Tower D B, and Schade J P. *Proceedings of the Second International Meeting of Neurobiologists*. Amsterdam, pp. 36-42
- Van der Loos H (1965) Fine structure of synapses in the cerebral cortex. *Neurosci Res Program Bull* 3, 22-4
- Van der Loos H (1967) The history of the neuron. In *The Neuron*. ed. Hyden H. Elsevier, Amsterdam, pp. 1-47
- Van der Loos H (1974) Dendrodendritic junctions. In *Dynamic Patterns of Brain Cell Assemblies. Report of NRP Work Session*, 14-16 May 1972, eds. Katchalsky A K, Rowland V and Blumenthal R, *Neurosci Res Prog Bull* 12, 86-90
- Van der Loos H and Glaser E M (1972) Autapses in neocortex cerebri: synapses between a pyramidal cell's axon and its own dendrites. *Brain Res* 48, 355-60
- Vanhalla-Pertulla T and Grimley P M (1970) Loss of proteins and other macromolecules during preparation of cell cultures for high resolution autoradiography. *J Histochem Cytochem* 18, 565-73
- Van Harreveld A (1947) On the force and size of motor units in the rabbit's sartorius muscle. *Am J Physiol* 151, 96-106
- Van Harreveld A (1966) *Brain Tissue Electrolytes*. Butterworth, London, p. 69
- Van Harreveld A, Murphy T and Novel K W (1963) Specific impedance of rabbit's cortical tissue. *Am J Physiol* 205, 203-207
- Varon S (1978) Macromolecular glial cell markers. In *Dynamic Properties of Glia Cells*. eds. Schoffeniels E, Franck G, Hertz L and Tower D P. Pergamon Press, Oxford, pp. 93-103
- Varon S and McIlwain H (1961) Fluid content and compartments in isolated cerebral tissue. *J Neurochem* 8, 262-75
- Varon S, Raiborn W R Jr, Seto T and Pomerat C M (1963) A cell line trypsinized adult rabbit brain tissue. *Z Zellforsch* 59, 35-46
- Vaughn J E and Peters A (1968) A third neuroglial cell type - an electron microscopic study. *J Comp Neurol* 133, 269-75
- Vaughn J E, Hinds P L and Skoff R P (1970) Electron microscopic studies of Wallerian degeneration in rat optic nerves. I. The multipotential glia. *J Comp Neurol* 140, 175-206
- Vernadakis A and Culver B (1979) Neural tissue culture: a biochemical tool, in *Biochemistry of Brain*. ed. Kumar S. Pergamon Press, Oxford, pp. 407-52
- Vernadakis A and Woodbury D M (1965) Cellular and extracellular spaces in developing rat brain. *Arch Neurol*, Chicago 12, 284-93
- Vernadakis A, Parker K and Norenberg M (1980) Changes in biochemical properties of glial cells in culture: in vitro differentiation? In *Tissue Culture in Neurobiology*. eds. Giacobini E, Vernadakis A and Shahar A. Raven Press, New York, pp. 441-26
- Vesalius A (1553, 1555) *De Humani Corporis Fabrica*, Johannes Oporinus, Basel
- Virchow R (1846) Ueber das granulierte Aussehen der Wandungen des Gehirnventrikels. *Allg Z Psychiatrie Berlin*, 3, 242-50
- Virchow R (1854) Über eine im Gehirn und Rückenmark des Menschen aufgefundene Substanz mit der Chemische Reaction der Cellulose. *Virchows Arch Pathol Anat Physiol* 6, 135-8
- Virchow R (1856) Gesammelte Abhandlungen zur Wissenschaftlichen Medizin. Meidinger, Frankfurt am Main, p. 890
- Virchow R (1871) *Die Cellularpathologie in Ihrer Begründung auf Physiologische und Pathologische Gewebelehre*. 4th edition. Hirschwald, Berlin
- Virchow R (1863, published 1971) *Cellular Pathology*. Dover Publications, New York, p. 313
- Vizi E S and Magyar K (eds.) (1984) *Regulation of Transmitter Function: Basic and Clinical Aspects*. Akademai Kiado, Budapest, pp. 129-73
- Voeller K, Pappas G D and Purpura D P (1963) Electron microscope study of development of cat superficial neocortex. *Exp Neurol* 7, 108-130
- Vogt O (1903) Zur anatomischen Gliederung des Cortex cerebri. *J Psychol Neurol Lpz* 2, 160-80
- Vogt O and Vogt C (1919) Ergebnisse unserer Hirnforschung. *J Psychol Neurol Lpz* 25, 277-462

REFERENCES

- Von Economo C (1926) Ein Koeffizient fur die Organisationshohei. *Klin. Wochenschr* 5, 593-5
- Von Economo C (1929) *The Cytoarchitectonics of the Human Cerebral Cortex*. Oxford University Press, London, p. 23
- Von Economo C and Koskinas G N (1925) Die Cytoarchitectonik der Hirnrinde des Erwachsenen Menschen. Springer, Vienna, pp. 1-810
- Von Gudden B (1874) Uber die Kreuzung der Fasern im Chiasma nervorum opticorum. *Albrecht von Graefe's Arch Ophthal* 20, 249-68
- Von Gudden B A (1875) Uber ein neues Microtom. *Arch Psychiatrie* 5, 229-34
- Von Kirsche W (1977) Historischer Uberblick zur Entdeckung der Ultrastruktur der interneuronalen Synapsen und Bemerkungen zum gegenwärtigen Stand. Ein Beitrag aus Anlass der Wiederkehr des 125. Geburtstages von Ramon Y Cajal. *Z Mikrosk Anat Forsch Lpz* 91, 595-686
- Vrbova G, Gordon T and Jones R (1978) *Nerve Muscle Interaction*. Chapman and Hall, London, pp. 48-51
- Waldeyer-Hartz H (1888) Das Gorilla- Ruckenmark. *K. Preufs Akad d Wissenschafts, Phys Abh III* Taf 13
- Waldeyer-Hartz H (1891) Über einige neuere Forschungen im Gebiete der Anatomie des Centralnervensystems. *Deutsch med Wochenschr* 17, 1213-18; 1244-46; 1267-9; 1287-9; 1331-2; 1352-6
- Walker C E (1928) Artefacts as a guide to the chemistry of the cell. *Proc R Soc B* 103, 397-403
- Walker C E and Allen M (1927) On the nature of the Golgi bodies in fixed material. *Proc R Soc B* 101, 468-83
- Ward A A Jr (1969) The epileptic neuron: chronic foci in animals and man. In *Basic Mechanisms of the Epilepsies*, eds. Jasper H H, Ward A A Jr and Pope A. Little, Brown and Company, Boston, pp. 263-88
- Wassle H (1982) Morphological types and central projections of ganglion cells in the cat retina. In *Progress in Retinal Research*, Vol 1, eds. Osborne N N and Chader G J, Pergamon Press, Oxford, pp. 125-52
- Watson W E (1976) *Cell Biology of The Brain*. Chapman and Hall, London
- Waugh D F and Schmitt F O (1940) Investigations of the thickness and ultrastructure of cellular components by the analytical leptoscope. *Cold Spring Harbor Symp Quant Biol* 8, 233-41
- Weakley B E (1981) *Beginners Handbook in Biological Transmission Electron Microscopy*. 2nd edition. Churchill-Livingstone, London, pp. 1-264
- Weber A (1955) Gonflements ou stases de l'axoplasme, en des zones localisees du tissu nerveux central ou peripherique. *Acta Anat* 23, 33-48
- Wechsler W (1970) The development and structure of peripheral nerves in vertebrates, in *Handbook of Clinical Neurology*. eds. Vinken P J and Bruyn G W. Vol 7, Diseases of Nerves, pp. 1-39
- Weddell G and Glees P (1941) The early stages in the degeneration of cutaneous nerve fibres. *J Anat Lond* 76, 65-93
- Weigert C (1882) Über eine neue Untersuchungsmethode des Centralnervensystems. *Z Med Wiss* 20, 753-7, 772-4
- Weigert C (1885) Ueber Schnittserien von Celloidinpräparaten des Zentralnervensystems zum Zwecke der Markscheidenfarbung. *Z Wiss Mikroskop Mikroskop. Tech.* 2, 490-5
- Weigert C (1885) Eine Verbesserung der Haematoxylin-Blutlaugensalz-methode für das Centralnervensystem. *Fortschritte der Medizine* 3, pp. 236-9
- Weigert C (1895) Beiträge zur Kenntnis der normalen menschlichen Neuroglia. *Abh Senckenberg Naturforsch Ges*, 19, 65-209
- Weil A and Davenport H A (1933) Staining of oligodendroglia and microglia in celloidin sections. *Arch Neurol Psychiatry* 30, 175-8
- Weir D M (ed.) (1973) *Handbook of Experimental Immunology*. 2nd edition, Blackwell, Oxford
- Weiss P (1961) The concept of perpetual neuron growth and proximo-distance substance convection. In *Regional Neurochemistry*. eds. Kety S S and Elkes J. Pergamon Press, Oxford, pp.
- Weiss P and Hiscoe H B (1948) Experiments on the mechanism of nerve growth. *J Exp Zool* 107, 315-95
- Weiss P and Wang H (1936) Neurofibrils in living ganglion cells of the chick cultivated in vitro. *Anat Res* 67, 105-17

REFERENCES

- Wendell-Smith C P and Blunt M J (1965) Possible role of neuroglia. *Nature* 208, 600-2
- Werman R (1966) Criteria for the identification of central nervous system transmitter. *Comp Biochem Physiol* 18, 745-66
- Westrum L E (1965) On the origin of synaptic vesicles in the cerebral cortex. *J Physiol Lond* 179, 4P-6P
- Wherrett J R and McIlwain H (1962) Gangliosides, phospholipids, protein and ribonucleic acid in subfractions of cerebral microsomal material. *Biochem J* 84, 232-7
- White G W (1966) Introduction To Microscopy. Butterworth, London, pp. 29-36
- White F P, Dutton G R and Norenberg M D (1981) Microvessels isolated from rat brain: Localization of astrocyte processes by immunohistochemical techniques. *J Neurochem* 36, 328-32
- Whittaker V P (1970) The vesicle hypothesis. In *Excitatory Synaptic Mechanisms*. eds. Andersen P and Jansen J K S. Universitetsforlaget, Oslo, pp. 67-76
- Whittam R (1958) Potassium movements and ATP in human red cells. *J Physiol* 140, 479-7
- Widdas W F (1952) Inability of diffusion to account for placental glucose transfer in the sheep and consideration of the kinetics of a possible carrier transfer. *J Physiol Lond* 118, 23-39
- Wilbrandt W and Rosenberg T (1961) The concept of carrier transport and its corollaries in pharmacology. *Pharmacol Rev* 13, 109-83
- Wilkin G P, Csillag A, Balazs R, Kingsbury A E, Wilson J E and Johnson A L (1981) Localization of high affinity (³H) glycine transport sites in the cerebellar cortex. *Brain Res* 216, 11-33
- Williams C A and Chase M W (eds.) (1967-77) Methods In Immunology and Immunochemistry. 4 Vols, Academic Press, New York
- Williams D and Lennox W G (1939) The cerebral blood-flow in arterial hypertension, arteriosclerosis and high intracranial pressure. *Q J Med* 8, 185-94
- Windle W F (1958) Nature of the ground substance in relation to extracellular space. In *Biology of Neuroglia*. ed. Windle W F. Thomas, Springfield, Ill., pp. 108-29
- Windle W F (1969) Textbook of Histology. McGraw-Hill, New York, p. 221
- Wisse E, Daems W Th, Molenaar I and van Duijn P (eds.) Electron Microscopy and Cytochemistry. North Holland-American Elsevier, Amsterdam
- Wolfe L S (1962) Lipid-protein complexes in membranes of nervous tissue. *Canad J Biochem Biophys* 40, 1261-71
- Wolfgram F (1966) A new proteolipid fraction of the nervous system. I. Isolation and aminoacid analyses. *J Neurochem* 13, 461-70
- Wolman M and Hestrin-Lerner S (1960) A histological contribution to the study of the molecular morphology of myelin sheath. *J Neurochem* 5, 114-20
- Wolosewick J J and Porter K R (1976) Stereo high voltage electron microscopy of WI-38 cells. *Anat Rec* 181, 511-12
- Wolosewick J J and Porter K R (1979) Microtrabecular lattice of the cytoplasmic ground substance. Artifact or reality. *J Cell Biol* 82, 114-39
- Woodbury D M (1969) Mechanisms of action of anticonvulsants. In *Basic Mechanisms of the Epilepsies*. eds. Jasper H H, Ward A A Jr and Pope A. Little, Brown, Boston, pp. 647-81
- Woodbury D M, Timiras P S, Koch A and Ballard A (1956) Distribution of radiochloride, radiosulphate and inulin in brain of rats. *Fed Proc Fed Am Soc Exp Biol* 15, 501
- Woodbury L A, Hecht H H and Christopherson R A (1951) Membrane resting and action potentials of single cardiac muscle fibres of frog ventricle. *Am J Physiol* 164, 307-18
- Woodward D L, Reed D J and Woodbury D M (1967) The extracellular space of rat cerebral cortex. *Am J Physiol* 212, 367-70
- Woollard H H, Weddell G and Harpman J A (1940) Observations on the neurohistological basis of cutaneous pain. *J Anat Lond* 74, 413-40
- Wraae O (1980) Kinetic Studies of the Lithium Ion in Rat Brain in vitro and in vivo. Ph.D. thesis, University of Odense, pp. 1-240
- Wu C W and Stryer L (1972) Proximity relationships in rhodopsin. *Proc Natl Acad Sci USA* 69, 1104-8
- Wuerker R B (1970) Neural filaments and glial filaments. *Tissue and Cell*, 2, 1-10
- Wuerker R B and Kirkpatrick J B (1972) Neuronal microtubules, neurofilaments and microfilaments. *Int Rev Cytol* 33, 45-75
- Wuerker R B and Palay S L (1969) Neurofilaments and microtubules in anterior horn cells of the rat. *Tissue and Cell* 1, 387-402

REFERENCES

- Wyckoff R W and Young J Z (1954a) The organisation within neurons. *J Anat* 88, 569P
- Wyckoff R W and Young J Z (1954b) The nerve cell surface. *J Anat* 88, 569P
- Wyckoff R W and Young J Z (1956) The motoneurone surface. *Proc R Soc B*, 144, 440-50
- Yamada K M, Spooner B S and Wessels N K (1971) Ultrastructure and function of growth cones and axons of cultured nerve cells. *J Cell Biol* 49, 614-35
- Young J Z (1945) History of the shape of a nerve fibre. In *Essays on Growth and Form*, presented to D'Arcy Wentworth Thompson. eds. LeGros-Clark W E and Medawar P B. Clarendon Press, Oxford, pp. 41-94
- Yusuf H K M and Dickerson J W T (1978) Disialoganglioside G_{D1A} of rat brain subcellular particles during development. *Biochem J* 174, 655-7
- Zacks S I (1964) *The Motor End-Plate*. Saunders, Philadelphia
- Zaunick R (ed.) (1961) Purkyne-Symposion. Nova Leopoldina Abhandl der Deutschen Akad der Naturforsch Leopoldina. 151, Barth, Leipzig, pp. 1-230
- Zernike F (1934) Begungstheorie des Schneidenverfahrens und seiner verbesserter form der phasenkontrastmethode. *Physica* 1, 689-704
- Zulch K J (1965) *Brain Tumours. Their Biology and Pathology*. 2nd American edition. Heinemann Medical, London
- Zulch K H (1979) International Histological Classification of Tumours. No. 21. *Histological Typing of Tumours of the Central Nervous System*. World Health Organisation. Geneva, pp. 1-188

Author Index

- Abney E R 43,85,89,90,91
Abramsky O 91
Adams C W M 11,34,205,213,214,215
Adrian R H 187
Aghajanian G K 184
Ahmed N 236
Akert K 150,154,155,228
Akil H 17,155,226
Albert A 153
Alberici de Canal M 227
Albot C 177
Aldridge W N 213
Aleu F P 224
Alford E C 213
Algeri G 22
Allen J 34,92,121
Allen J N 222
Allen M 14
Allen N S 127
Allen R D 127,198
Altmann R 13
Alvord E C 238
Alzheimer A 13,18,183
Ametani T 131
Anderson P J 11
Andreoli T E 208
Andres K H 23,228
Andres P F 79
Aoki K 176
Appel S H 42,43
Ariens-Kappers J 21,39,52-53
Aronow L 207
Arora R C 88
Auerbach L 15
Autilio L A 209,238
Bachelard H S 214
Baglio C M 177
Bahr G F 48
Bailey E 47,222
Bailey P 168-169,206
Bairati A 39
Bak A F 186
Bakay L 224
Baker J R 6,8,14,33,63
Baker P F 208,218,252
Balazs R 89,90,91
Ballard A 221,223
Bangham A D 213,221,252
Banik N L 214,215
Barber P C 42,43
Barbossa E 211
Barchas J D 17,155,226
Barka T 11
Barker L F 6,52-53,152
Barlow C F 223
Barnard R I 138,152
Barr M L 138,152,153
Bartlett P P 43,85,89,90,91,249
Bastian H C 200
Bauer K F A 6,11,39
Baumann N A 89,90,91
Baumeister W 48,219
Bayliss L E 207
Bayliss O B 213,215
Beams H W 14,25
Bear R S 19,24,209,239
Beasley R A 54-61,74-77,93,157,246
Becker H N 131
Beevor C E 22
Benda P 88,89,90
Benedetti E L 177
Bennett V 15,211
Bergman R 39
Berl S 25,88,196
Berriman J A 255
Besta C 39
Bevan-Lewis W 21,183
Bidwell L A 22
Bielschowsky M 10
Bignami A 74-77,88,89,90,248
Billings-Gagliardi S 131
Birks R 154
Bishop G A 50,185,186
Black P 90,247,248
Blackstad T W 153,154
Blackwood W 66-69,74-77,239
Blakemore W F 19
Blasberg R 211
Blindermann E E 224
Blomstrand C 89
Bloom F E 154,184,205
Bloom W 54-61,66-67,74-77,150,
155,168,176,181

AUTHOR INDEX

- Blunt M J 38
 Bock E 88,89,90
 Bodian D 153,229
 Boechler N 89
 Boeke J 156
 Bohm AA 10
 Bok S T 164
 Boll G 17,37,39
 Bolles-Lee A 10
 Bondareff W 64-65
 Bornstein M B 23,42,43
 Bouman L 164
 Bourke R S 222,223
 Bourne G H 6,14,20,39,52-53
 Bowsher D 185
 Boycott B B 135,141,153
 Boyde A 47,222
 Bradbury S 6,10
 Bradford H F 196
 Brady R N 211
 Brady R O 213,215
 Brady S T 127
 Bray D 25,90,247,248
 Brazier M A B 6
 Brearley R F 150
 Breathnach A S 26,202
 Brenkert A 88
 Brieppohl W 184
 Brierley J B 183
 Brightman M W 185
 Brockes J P 249
 Brodal A 138,150,153,154,155
 Broderson S H 88
 Brodie B B 233
 Brodmann K 21
 Brown J H U 214
 Brownson R H 37,78
 Bruce-Casselman W G 63
 Bruggeller P 202
 Bruton J 74-77,246
 Bullock T H 153
 Bunge M B 23,39
 Bunge R P 6,23,39
 Burgoyne R D 154
 Butschli O 200
 Caldwell P C 208
 Calissano P 213
 Calkins E 219
 Callan H 26
 Callas G 23
 Cam Y 43
 Cammermeyer J 38,39,56-57,60-61,
 184
 Campbell A W 21
 Campbell F R 177
 Campbell W J 187,188
 Carafoli E 25,132
 Carasso N 255
 Carleton H M 10,72,74-77,246
 Carpenter M B 39,58-59,68-69
 Castelli A 215
 Causey G 176,180
 Cavanaugh J B 207
 Ceccarelli R 228
 Chambers E D 27,242
 Chambers R 27,242
 Chan S H P 215
 Chan-Palay V 66-67,131
 Chang J J 23,255
 Chang K J 211
 Chang T M S 208,221
 Chao F C 213
 Chase M W 26
 Chou S N 224
 Christensen H N 236
 Christopherson R A 187
 Clark G 74-77,245
 Clark G L 19,209
 Clark S L 52-53,54-59
 Clarke D A 22
 Clarke D D 88,196
 Clarke E 6,21
 Clarke H 21
 Clasen R A 224
 Claude A 25,26
 Clay T H 10
 Clemente C D 39
 Coates P W 88
 Coers C 156
 Cohen J 43,90,91
 Cohen R S 131
 Cohen S R 222
 Cohnheim J 18
 Cold Spring Harbor Symposium on
 Quantitative Biology 1972 219
 Cold Spring Harbor Symposium on
 Quantitative Biology 1975 156
 Cold Spring Harbor Symposium on
 Quantitative Biology 1982 26
 Cole K S 24
 Cole W V 156
 Coleman R 194,201
 Colodzin M 214
 Colonnier M 17
 Conel J Le Roy 143,156,185,186
 Conradi S 153
 Constantanides P 66-71
 Constantino-Ceccarini E 215
 Conway E J 219,220,221
 Cook D J 201
 Cooke P M 224
 Combs J S 186,251,253
 Corsellis J 66-69,74-77
 Corriol J 211
 Cosslett A V 46
 Costero I 23,41,85,148,149,176,181
 Cottrell G A 134
 Court T H 10
 Couteaux R 138,152,156
 Cowdry E W 79
 Cowey A 154
 Cox W 10
 Cragg B G 47,153
 Crain S M 23,43,46,241
 Creutzfeldt H G 39
 Crile G W 186
 Crone C 211
 Cruickshank B 74-77
 Csillag A 90
 Cuatrecasas P 211,212,215
 Culling C F A 56-61,74-77

AUTHOR INDEX

- Culver B 42
 Cummings, see Hill, J
 Cummins J E 78,133,187,231,243,
 252
 Currie D N 39,43,89,90,91
 Curtis B A 56-59,66-71,138,140,150,
 153,176
 Curtis H J 24
 Cushing H 206
 Daems W Th 219
 Dahl A L 66-71
 Dahl D 89,90
 Dahl T D 214
 Dahlstrom A 17,90,226
 Dancer J B 180
 Danielli J F 24,25,132,133,210
 Danowski T S 219
 Dartnall D W 216
 Darwin C 10
 Davenport H A 74-77,92,138,152,
 153,245
 Davidoff M 10
 Davison A N 34,214,215
 Davson H 17,24,25,38,132,133,210,
 222,223,225,233,237
 Dawson R M C 227
 De Balbian Verster F 214
 De Brabander M 198
 De Duve C 25
 De Estable R F 39
 De Iraldi A P 64-65
 De May J 198
 Deiters O F C 8,9,13,17,18,20,37,
 39,52,53,165,187
 Delaunoy J P 88,89,90
 Del Castillo J 17,226
 Del Rio Hortega P 10,18,24,36,39,
 58-59,74-77,159,183,184,245
 De Mey J 198
 Dempsey E W 38
 Derer P 88,90
 De Robertis E D P 15,38,64,65,
 68-69,136,139,153,155,176,224,227
 Deutsch K 33,34,47,92,94,121,122,
 176
 De Vieussens R 21
 Devilliers G 88,89
 De Vries G H 215
 Dick F 90
 Diller D A 37
 Dimpfel W 90
 Dittman L 43
 Dixon M 211
 Dixon N C 214
 Dodds T C 74-77,239
 Dodgson M 18,183,184
 Domek N S 223
 Dorfman S H 43,89,91
 Downer J de C 185
 Driest E 25
 Drury R A B 72,74-77,246
 Dubochet J 255
 De Bois-Reymond E 17
 Dulak N C 214
 Dupouy G 26,47
 Dusser de Barenne J G 22
 Dustin P 198
 Dutta C R 21
 Dutton G R 88,90
 Ebbeson S O E 156
 Eber L 177
 Ebner F F 17
 Eccles J C 150,153,155,186,196,224,
 237,251,253
 Ecclesiastes iii
 Edelman F L 224
 Edstrom J E 243
 Eggleton M G 17
 Ehrenberg C G 7
 Eichberg J Jnr 227
 Ekholm R 154
 Eldefrawl A T 211
 Eldefrawl M E 211
 Elekes N 39
 Elfvin L G 26,176
 Elliott G R 17,155,226
 Elliott H C 52-53,56-59,60-61,66-67
 Elliott K A C 222
 Elliott T R 17
 Emmelot P 177
 Eng L F 74-77,89,119,213,248
 Entin T 153
 Epstein M H 27
 Erickson T C 22
 Erikson L B 74-77,92
 Erlanger J 32
 Essner E 214
 Estable C 18,20,243
 Estable-Puig J F 39
 Etcheverry G J 64-65
 Eto Y 215
 Evans T C 25
 Evans W H 215
 Everly J L 213,215
 Fanestil D D 208
 Farquhar M C 38,64-65,68-69
 Fasham J 20,128,159,160,243
 Fatt P 17,185,186,226,251,253
 Favard P 255
 Fawcett D W 13,20,54-61,66-67,74-77,
 150,155,168,176,181
 Fedoroff S 41,42,43,89,241
 Feirtag M 183,193
 Fenn W O 219
 Fernandez-Moran H 20,26,34,209,218,
 237,239
 Ferrier D 21,22
 Ferris P J 185
 Field E J 39
 Fields K L 39,41,43,85,89,90,91,248,
 249
 Fine R E 25
 Finean J B 34,194,201,202,209,218,
 237,239
 Finkelstein A 221
 Fischbach G D 39,148
 Fisher R A 11
 Fishman R A 211

AUTHOR INDEX

- Fitzgerald P J 177
 Flangas A L 88
 Flechsig P 22
 Fleischer B 239
 Fleischer S 239
 Flemming W 18
 Flynn F 219
 Folch J 213
 Fonnum F 196
 Fontana F 6,19
 Ford D H 176
 Forel A H 12
 Forman D S 24
 Fox C A 21
 Franck G 41,78
 Frank K 186,187,251,253
 Franke W W 26,40
 Freeman R 255
 Frey T G 215
 Freygang W H 185,186
 Fricke H 24,218,220
 Friede R L 78
 Friedenwald J S 156
 Friedrich V L 66-71
 Fringeli U D 48
 Fritsch G 21,22
 Fullam E F 25,26
 Fulton J F 6
 Fuortes M G F 186,187,251,253
 Fuxe K 17,90,226
 Galilei G iii
 Gallyas F 92,93
 Gardner D L 74-77
 Garrison F H 6
 Gasser H S 32,176
 Gatenby J B 74-77
 Gauthier G F 156
 Geiger B 40
 Geiger R S 42,43,80,85,148
 Gelfan S 138,152
 George J 91
 Geren B B 20,173,237
 Gerlach W 12,37,39,201
 Gerschenfeld H M 38,64-65,68-69,
 174,176,224
 Gersh I 33
 Gerstl B 89,213
 Ghandour M S 88,90
 Gheuens J 89
 Giacobini E 78,88,187,241,243
 Gibson W C 138,152,153
 Gilbert J C 211
 Gilbert S P 127
 Glaser E M 151
 Glaser G H 238
 Glauert A M 47
 Glees P 6,10,11,36,37,38,39,54-55,
 66-69,74-77,92,120,135,152,154,162,
 176,184,203,204
 Glynn I M 220,221,237
 Goldberg M A 223
 Goldfischer S 14,214
 Goldman R 198
 Goldstein A 207
 Golgi C 10,11,12,14,17,18,36,37,39,
 156,159,201
 Goltz F L 22
 Gordon N W 170
 Gordon R D 89,91
 Gordon S 238
 Gordon T 156
 Gordon-Weeks P R 154
 Gorter M D 24,210,218,252
 Grafstein B 24
 Gray E G 38,56-57,60-61,64-71,
 135,141,146,150,153,154,155,222,246
 Gray J 27,72,74-77,93,242
 Gray P 27,74-77,93,245,246
 Greenberg E S 223
 Greenfield J G 54-55,70-71,74-77,
 203,239
 Gregson N A 5,43,89,91,214
 Grendel F 24,210,218,252
 Grey S 215
 Grimley P M 179
 Grinnell A 153
 Grubb D T 255
 Gruner J E 184
 Grunfeld Y 43,148
 Gruninger H 184
 Guiditta A 213
 Guidotti G 212
 Guillery R W G 135,141,150,153,
 154,155
 Guth L 204
 Haapanen L 187
 Haberman E 90
 Haege L 219
 Hague I U 224
 Hahn M 48
 Hakamori S I 39,41,43,85,89,91,248
 Halaris A 213
 Hall S 54-55,176
 Hallpike J F 214,215
 Ham A W 56-61,66-67,70-71,120,194
 Hamberger A 6,32-33,39,56-57,
 66-67,78,88,89,119,187
 Hamberger C A 20,56-57
 Hamburger V 203
 Hamilton J W 156
 Hammarberg C 21
 Han S S 66-67,131
 Hankin E H 10
 Hannover A 7
 Hansson E 43,89,90,241
 Hansson H A 39,40,56-57,66-67
 Hardesty I 11,181
 Hardy W E 200
 Harford J B 215
 Harpman J A 156
 Harris E J 207,208,209,220
 Harris J E 219
 Harris J R 120,194,201
 Harrison R G 23
 Hartmann J F 26,38,64-65,68-69
 Hartridge H 74-77
 Hasan M 154
 Hass G M 224
 Hassler R 184
 Hastings A B 219,223,236
 Hayat M A 26,219

AUTHOR INDEX

- Haymaker W 39
 Haynes F 74-77
 Hecht H H 187
 Heimer L 154
 Held H 11,13,15,16,39,136,183
 Henderson H S 179
 Hendleman W J 148,181
 Henle J 17
 Henn F A 221
 Henrikson C K 17
 Herdson P B 177
 Herman L 177
 Herrera A 200
 Hersey S R 185
 Hertl M 20
 Hertwig G 34
 Hertz L 27,41,43,78,88,89,90,119,
 121,187,222,241
 Hess A 11,39,183
 Hestrin-Lerner S 34
 Heuser J E 228
 Hexum J F 214
 Hild W 23,42,43,46,73,79,148,149,
 180,181,186
 Hildebrand C 19
 Hill J 7
 Hillarp N-Å 17,90,226
 Hillman H 2,4,5,11,14,20,23,26,27,
 30,31,32,33,34,35,36,47,48,92,95,
 120,121,122,126,128,132,134,135,
 147,151,158,159,160,168,171,172,173,
 187,188,194,198,200,202,204,205,209,
 211,218,219,222,224,230,231,235,236,
 238,240,241,242,243,253,256
 Hinds P L 184
 Hirano S 176,238
 Hirsch H E 88
 His W 12
 Hiscoe H B 24
 Hitzig G 21,22
 Hochberg F H 6
 Hodgkin A L 197,208,211,218,220,221,
 237,252
 Hoffman J F 208
 Hogue M 23,183
 Hokin L E 214,236
 Hokin M R 236
 Holladay L A 211
 Hollenberg M D 211
 Holman M E 17,219
 Holman R B 17,155,226
 Holmes S 81
 Holmgren E 14
 Holmstedt J O V 66-67,131
 Holst E A 39
 Holzer W 11,93,172
 Homo J C 255
 Hongchien H A 138,152
 Hooke R E 6
 Hopwood D 179
 Horne R W 120,194,201,221,252
 Horsley V A 22
 Horstmann E 222
 Hosie R J A 214,227
 Hosli E 79
 Hosli L 79
 Hosmer H R 186
 Hosokawa H 54-57,60-61,64-71,
 74-77
 Hudson C S 202
 Hulcher F H 238
 Humason G L 74-77
 Hurlbut W P 228
 Hussain T S 20,29,30,32,34,35,121
 159,168,240,243
 Huxley A F 211,221,237
 Huxley H E 154
 Hyams J S 198
 Hyden H 5,6,18,20,27-30,32,39,
 52-53,64-65,78,89,92,121,128,
 133,146,147,150,153,155,165,187,
 224,231,238,242,243,252,253
 Iles G H 214
 Illiano G 215
 Ilmaz Y 127
 International Congress of Cell Biology
 1981 26
 Iversen L L 196
 Jacob M 216
 Jacobson K B 213
 Jacobson M 89,130,184-185,203-204,
 246,248
 Jacobson S 56-59,66-71,138,140,
 150,153,176
 Jacque C M 89,90,91
 Jakob A 39
 Jastrowitz M 39
 Jezequel A M 177
 Joanny P 211,222,235
 Johnson A L 90
 Johnston M K 213
 Johnston P V 52-53,60-61,64-71,
 78,131,146,150,152,153,154,155,
 176,222,240
 Johnstone B M 219
 Jones D G 155,228
 Jones E G 131,150,195
 Jones R 156
 Jones S J 47,222
 Jordan E F 185
 Jørgensen O S 89,90
 Jørgensen P L 214
 Joseph J 39
 Jungalwala F B 216
 Juurlink B H J 88,89
 Kafig E 26
 Kaiserman-Abramof I R 131,153
 Kalman S M 207
 Kaltenbach J P 177
 Kamiya N 198
 Kanfer J N 88
 Kaplan N O 213
 Karasek J 34
 Karcher D 89
 Karlsson U 150,153
 Karnovsky M J 156
 Kasa P 214
 Kates M 25,132
 Katz B 17,154,196,197,226,228,237

AUTHOR INDEX

- Katzman R 224,235-236
 Keith K H 126
 Keller A 255
 Kelly J S 39,43,90,91
 Kennedy E P 214
 Kennedy M C 43,89,90,91
 Kennedy P G E 37,85,248
 Kernohan J W 206
 Kershman J 24
 Kessel R G 14
 Key E A H 18
 Keynes R D 208,219,220,221,237
 Khallawan S A 172,204
 Kibler R F 223
 Kier M W 213,238
 King J S 131
 King L S 39
 Kingsbury A E 90
 Kirby J 3,99
 Kirk W 213
 Kirkman H 14
 Kirkpatrick J B 25,26,131
 Kirsche W von 6
 Kitai S T 50,185,186
 Kjeldsen C S 221,222
 Klatzo I 24,185,224
 Klestadt B 185
 Klotz I M 216
 Knoll M 25
 Koch A 221,223
 Koelle G B 156
 Koelliker A 19
 Koenig H 39,78
 Kolmuddin G M 187
 Konigsmark B W 184
 Kopp F 48
 Korr H 184,203
 Korte K 66-71
 Koskinas G N 21,22,183
 Koul O 216
 Kozak L P 89,90
 Krepler R 40
 Krieg L C 213
 Krnjевич K 79
 Kruger L 38,184
 Krypsin-Exner W 78
 Kuffler S W 38,39
 Kuhar M J 196
 Kuhne W 14
 Kuksis A 25,132
 Kuntz A 156
 Kurella G A 219
 Kurihara T 214
 Kurz H 233
 Laatsch B A 238
 La Bella F S 214
 Labourdette G 88,89,90
 Lajtha A 211,214,221
 Landau W M 186
 Landon D N 20,54-55,176,180,209,
 210
 Langley O K 88,90
 Lantermann A J 19
 Lasek R J 22,127
 Laskowski E J 24,224
 Lassek A M 22
 La Vallette St. George V 13
 Lazarides E 26,126
 Le Baron F N 213
 Leblond C P 38,39,60-61,70-71,184
 Lee J C 185
 Leeson C R 74-77,138,176
 Leeson T S 74-77,138,176
 Leeuwenhoek A van 6,7,19
 Le Fevre P G 211
 Legros-Clark W E 152
 Leibowitz G 43,89,91
 Leiman A L 43
 Lennartz W 215
 Lennox W G 224
 Leonhardt H 24
 Lepault J 255
 Letterier J F 126
 Levi-Montalcini R 203
 Lewis A J 37,38,39,58-59,66-69,162
 Lewis M R 23
 Lewis M S 213
 Lewis W H 23
 Leyton A S F 22
 Li C L 186
 Liem R K H 126
 Lindsay R M 42,43
 Ling G N 219,220,230
 Lisak R P 37,43,85,89,91,248
 Lister J 7
 Livingstone R B 228
 Lodin Z 34
 Lorber J 226,227
 Lorente de No R 22,138,152
 Loten E G 215
 Lowenthal A 89
 Lubinska L 24
 Lucy J A 5
 Ludlum S 11,183
 Lumsden C E 39,42,85,154
 Lund-Andersen H 221,222
 Luse S A 38,64-65,68-69,177,222,
 224
 Lux H D 186
 Madge E 138
 Magyar K 212
 Maickel R P 233
 Maizels M 219
 Mallory F B 36,74-77,92
 Mandel P 43,215,216
 Manery J F 223
 Mann G 10,47,63,74-77
 Mannen H 54-57,74-77
 Manning M C 91
 Manocha S L 20,23,39,42,52-53
 Marchbanks R M 227,228,229
 Marchi V 22
 Marcus E M 56-59,66-71,138,140,
 150,153,176
 Mares V 34
 Markham C H 224
 Marks N 214
 Marsland T A 74-77,92
 Martinez-Hernandez A 88
 Massey J F 148,181

AUTHOR INDEX

- Matthews M A 38,184
 Matthews M R 154
 Maunsbach A B 214
 Mauthner L 20
 Mayer E 202
 Mayer S 233
 Maynard E A 38,64-65,68-69
 Maxwell D S 38,184
 McClung C E M 10,27
 McClung-Jones R 27,63,74-77,246
 McCoy C C 23
 McDowell A N 255
 McEwen B 89
 McIlwain H 187,188,214,221,239
 McLennan D H 214
 McMahan U J 50,156
 McManus J F A 74-77
 Meek G A 26
 Mehl E 213
 Meier E 89,90,91
 Meller K 38,66-69,176,184
 Meryman H T 26
 Metchnikoff E 24
 Metuzals J 127
 Metz A 39
 Meyer A 138,152,203
 Meyer M 138,152
 Meynert T 21
 Michell R H 194,201
 Mikata A 177
 Mikoshiba K 88
 Millhouse E O 74-77
 Minckler J 152
 Minkowski M 22
 Miquel J 24,185
 Mirsky R M 39,41,43,85,89,90,91,
 247,248,249
 Molenaar I 219
 Moll R 40
 Monard D 89
 Monzain R 43,148
 Moonen G 43,88,89
 Moor H 26,150,154,155,217,228
 Moore B W 89
 Moore E 233
 Moore R Y 205
 Moore W M 211
 Morgan C 20
 Morgan J F 128
 Mori S 38,39,60-61,70-71
 Morowitz H J 239
 Morrison A B 223
 Morse S 220
 Morton H J 128
 Mowry R W 74-77
 Mueller P 208,221
 Mugnaini E 38,66-71,175
 Muhlethaler K 217
 Mujai K 177
 Mukerji S 89
 Muller G 78
 Muller H O 25
 Muller K J 50
 Munoz-Martinez E J 176
 Murai S 88
 Murphy J B 213
 Murphy T 186
 Murray M R 23,42,43
 Nachmansohn D 197,209
 Nagata Y 88
 Nageotte J 181
 Nairn R C 26
 Nakai J 203
 Narita Y 184
 Nathaniel D R 64-65
 Nathaniel E J H 64-65
 Nauta W J H 156,183,193
 Neale J H 90
 Nelson P G 39,78,148,186
 Neuhoff V Von 242
 Neumann E 197,209
 Newfang D M 25
 Nicholls J G 38,39
 Nicholson G 25-26,132
 Nicklas W J 88
 Niessing K 6
 Nishii B 206
 Nissen C 88,89
 Nissl F 10,18,24
 Nobel K W 186
 Noble M D 43,90,91,249
 Noonan T R 219
 Noppe M 89
 Norenberg M D 88,90
 Norton W T 209,215,238
 Novikoff H B 14,176,214
 Nurnberger J I 39,170
 Obata K 90
 Obersteiner H 183
 O'Brien R D 211
 Ochoa J 176
 O'Connor J S 27
 Oehmichen M 24,184,203
 Oksche A 6
 Okujava V M 186
 Oldendorf W H 223
 Olsen N S 233
 Olszewski J 185
 O'Malley C D 6,21
 O'Neill R P 223
 Oppelt W W 222,223
 Orkand R K 38,39,153
 Osborn M 26
 Osborne N N 155,242
 Osen K K 66-71
 Osterhout W J V 210
 Overton E 24
 Packman P M 89
 Painter T S 74-77
 Palade G E 13,15,25,26,64-65
 Palay S L 6,13,15,17,20,25,26,38,
 54-55,62,63,64-71,78,131,150,154,
 155,158,166,180,222
 Palmer K J 239
 Palmgren A 74-77
 Pandolfi S 224
 Pappas G D 131,134,150,154,155,184

AUTHOR INDEX

- Pappenheimer J R 185
 Pappius H M 222
 Parker K 88
 Parker L O 186
 Parker R C 128
 Parsons D F 26,47
 Patay R 92
 Patel N J 89,91
 Patlak C S 222,223
 Pavlin R 187
 Pearse A G E 11,63,205
 Pease D C 38,64-65,68-69
 Penfield W 11,22,36,39,54-59,74-77,
 159,245
 Pepeu G 156,196
 Perdue J F 214
 Persson L I 43
 Peters A A 6,13,17,20,38,54-55,62,
 63,64-71,78,131,150,153,154,155,158,
 166,174,176,180,211
 Peterson E R 23
 Pettman B 88,89
 Pevzner L Z 41,44-45,78,88,119,241
 Pfenninger K 150,154,155,228
 Phalen G S 138,152,153
 Phillips M J 177
 Pickel V M 46,205
 Pieringer J 215
 Pieringer R A 215
 Pigon A 32,92,187
 Piraux A 24,224
 Pleasure D E 91
 Poduslo S E 215
 Polak M 39
 Pollard T 198
 Pollay M 233
 Pollen D A 186
 Pomerat C M 23,31,40,41,42,43,85,
 148,149,176,181
 Popper K 4,14
 Porter K R 25,26,131,168
 Powell T P S 131,195
 Pratt D 213
 Priestley B L 226
 Prince D A 237
 Privat A 38
 Prockop L D 185
 Proskauer C C 131
 Pruijs W M 39
 Pruss R M 39,41,43,85,89,90,91,
 248,249
 Purett D 211
 Purkinje J E 8,9,20,27,187
 Purkis V A 153
 Purpura D P 134,184
 Puszkin S 25
 Quarles R H 213,215
 Radin N S 88
 Raff M C 37,39,41,43,85,89,90,91,
 119,248,249
 Raghaven S S 88
 Raiborn C W 181
 Raiborn W R 31,42,43,148
 Raine C S 131
 Raisman G 42,43,154
 Ralis H M 54-61,74-77,93,157,246
 Ralis Z A 54-61,74-77,93,157,246
 Rall D P 185,222,223,233
 Ramon-Moliner E 37,39,161
 Ramon y Cajal S iii,6,10,12,13,17,
 18,21,32,36,52-55,149,153,156,180,
 245
 Ranck J B 186
 Ranson S W 52-53,54-59
 Ranvier L A 19
 Rapoport A 64-65,233
 Rash J E 202
 Rattray S 43,90,91,249
 Reade J B 179
 Reed D J 221
 Reese T S 154,228
 Regan C 89,90,91
 Reil J C 7
 Reimer L 255
 Reisner J H 48
 Remak R 8,19
 Repke K R H 25,132,211
 Retzius G M 18
 Rhoads D B 88
 Rhodin J A G 13,20,131,176
 Richards P R 23
 Robards A 5,202-203,222,255
 Roberts E 88,90
 Roberts K 198
 Robertson J D 20,24-25,34,46-47,
 132,135,176
 Robertson W F 6
 Robin E D 223
 Rodnight R 222
 Rodriguez de Lores Arnaiz G 227
 Romeis B 6,72,93
 Ronnback L 43
 Roots B I 38,52-53,60-61,64-71,
 78,131,146,150,152,153,154,155,176,
 222,240
 Roots L 156
 Rose S P R 88
 Rosenbaum J 198
 Rosenberg T 207,210
 Ross K F A 33
 Ross L L 64-65
 Roth L J 223
 Rottenberg D A 6
 Rouiller C 177
 Rowland A F 186
 Royal Microscopical Society 14
 Rozsa G 20
 Rubinstein L J 206-207
 Rudin D O 208,221
 Rudolph G G 233
 Ruska E 6,25
 Russell D S 206-207
 Rydberg E 39
 Saebich R 184
 Saida T 91
 Salomon J C 177
 Samec J 216
 Sandri C 150,154,155,228
 Santiago J C 88
 Sapirstein V S 88,89

AUTHOR INDEX

- Sartory P 2,5,14,20,26,30,32,34,35,
36,92,120,121,126,128,132,135,159,
160,173,194,198,209,230,240,242,243,
253,256
 Satake M 89
 Sauer F C 184
 Saul R 215
 Sayre G P 206
 Schade J P 176
 Schanker L S 233
 Scharrer B 6
 Scharrer E 6
 Scheibel A B 138
 Scheinberg L C 224
 Schiller D L 40
 Schlaepfer W W 209
 Schleiden M J 6
 Schmid E 26
 Schmidt H D 19
 Schmidt W J 19
 Schmitt F O 19,24,180,209,218,239,
252
 Schnapp B 38
 Schneider D D 196
 Schoffeniels E 41,78
 Schousboe A 88,89,90,119,222
 Schroder P 22
 Schultz R L 38,39,64-65,68-69
 Schurch-Rathgeb Y 89
 Schwann T 6,20,179
 Schwartz A 214
 Schwartz S 79
 Scott T G 215
 Seeman P 214
 Seil F J 43
 Sellinger O Z 88,214
 Sellstrom A 39,43,56-57,66-67,88,89
 Selvendran S Y 43,90,91
 Semenza G 25,132
 Sensenbrenner M 43,88,89
 Seredynski J 48
 Seto T 31,42,43,148
 Severinghaus A E 14
 Shahar A 43,78,148,241
 Shantha T R 20,39,52-53
 Sharpey-Schafer E 10
 Shaw F H 219
 Shaw T I 208,218,252
 Sheikh K 204
 Shelanski M L 26,126
 Shepherd G M 154
 Sherrington C S 15,22,226
 Sherwood M R 42,43
 Sholl D A 39,52-53,138,142,156,164,
194
 Short R H D 74-77
 Sidman R L 184
 Siegel M I 215
 Seigesmund K A 21
 Silberberg D H 43,89,91
 Simon G 177
 Simon S E 219
 Singer C 6
 Singer S J 25-26,132,212
 Sinha A K 88
 Sjostrand F S 13,15,20,25,222
 Sjuve S 35,94,95
 Skoff R P 184
 Skoglund C R 187
 Skou J C 208
 Skrivanova P 34
 Skriver E 214
 Sleytr U B 202-203,222,255
 Smart I 184
 Smith A 74-77,246
 Smith D E 185
 Smythies J R 153
 Sneyd J G J 215
 Solomon A K 215
 Somerville J C 239
 Somogyi P 154
 Sotelo C 64-65
 Sotelo J R 18,20,243
 Sourander P 40
 Spatz H 37,39
 Spaziani E 222
 Spence A M 88
 Spiegestein M Y 43,148
 Spinoza B iii
 Spitzer N C 156
 Spooner B S 131
 Sprague J M 138,152
 Stahl W L 88
 Stampfli R 179
 Steere R L 26
 Stein W D 207,210
 Steiner J W 177
 Stempak J 64-65
 Stenger R J 177
 Stenwig A E 184
 Stephan H 184
 Sternberger L A 26
 Stevenin J 216
 Stieda L 6
 Stilling B 8
 Stirling W 19
 Stoeckenius W B 209,239
 Stolinski C 26,202
 Stolkin C 90,247,248
 Stollery S 235
 Stone W E 237
 Stowell R E 33
 Strecker H J 213
 Streicher E 221,223
 Streit P 228
 Stricker S 18,39
 Strickland K P 214
 Struck D K 215
 Stryer L 213
 Subba Rao G 215
 Sulzmann R 39
 Sunderland S 180,181,218
 Suter D B 37
 Suzuki K 215
 Svenneby G 90
 Symposium on Nerve Growth Factor
1964 203
 Symposium of Quantitative Electron
Microscopy 1965 63
 Szarkowski J W 217
 Szentagothai or Schimert J 138,153
 Szent-Gyorgyi A 20

AUTHOR INDEX

- Taft A E 11,183
 Tamarin A 47,222
 Tanaka R 214
 Tani E 131
 Tasaki I 23,42,43,46,73,79,127,148,
 149,186
 Tavastjerna M G 213
 Taylor C B 224
 Taylor I M 219
 Taylor N 233
 Tell P E 215
 Tellyesniczky K 33
 Terry R D 209,224,235-236,238
 Terry T 239
 Thompson T E 221
 Thorun W 213
 Tiegs O W 6
 Timiras P S 221,223
 Tobari C 238
 Tobias C 39
 Todd A 17
 Tomlin S G 26
 Torack R M 209,224,235-236
 Tower D B 41,78,88,222,223,238
 Tower S S 156
 Travis L 127
 Treip C S 162,203,206-207,239
 Triggle D J 196
 Troshin A S 219
 Truex R C 39,58-59,68-69
 Tsukada T 214,239
 Tsukada Y 88
 Tyler S 224
 Uesugi S 214
 Unger L 18,39
 Ungerstedt U 205
 Urechia G I 39
 Usherwood N R 134
 Ussing H H 219
 Uyeda C T 74-77,89,248
 Uyemura K 238
 Valdivia O 153,155,229
 Valentin G G 7,8,20
 Van Breeman Y 25
 Vanderhaeghen J T 89
 Van der Loos H 6,27,150,151
 Van Duijn P 219
 Vanha-Pertulla T 179
 Van Harreveld A 22,186,222,225
 Varon S 31,42,43,88,89,90,119,221,
 248
 Vaughn J E 17,66-69,131,176,184
 Vernadakis A 42,78,88,223,241
 Vesalius A front cover, 244
 Virchow R 5,8,10,11,18,183,225
 Vizi E S 212
 Voeller K 184
 Vogell W 219
 Vogt C 21
 Vogt O 21
 Von Bonin G 168-169
 Von Economo C 21,22,183
 Von Gerlach J 10,11,12
 Von Gudden B A 7,21,22
 Vrbova G 156
 Waechter C J 215
 Walberg E 66-71
 Wald F 38,64-65,68-69,176,224
 Waldeyer-Hartz H W G 12,171
 Walker C A 255
 Walker C E 14
 Wallach J 8
 Wallington E A 74-77
 Wang H 23
 Ward A A Jnr 237
 Wassle H 141,144
 Watson S J 17,155,226
 Watson W E 39,218
 Waugh D F 218,252
 Waxman S G 150,154,155
 Weakley B E 26
 Webb E C 211
 Weber A 138
 Weber K 26
 Webster H de F 6,13,17,20,38,
 54-55,62,63,66-71,78,150,154,158,
 166,180
 Wechsler W 180
 Weddell G 156,204
 Welgert C von 11,17,18,22,74-77
 Weil A 74-77,92,245
 Weir D M 26
 Weiss P 23,24
 Wendell-Smith C P 38
 Wendorf M B 90,247,248
 Werman R 17,226
 Wessells N K 131
 Westrum L E 228
 White F P 88,90
 White G W 36
 Whittaker V P 154,155,227
 Whittam R 219
 Widdas W F 210
 Wilbrandt W 207,210
 Wilkin G P 89,90,91
 William of Occam (Ockham) 4
 Williams C A 26,43,89,90,91,249
 Williams D 224
 Wilson J E 90
 Windle W F 11,138,183,203
 Winter J 39,41,43,85,89,91,248
 Wisniewski H 131
 Wisse E 219
 Wolfgram F 213
 Wolman M 34
 Wolosewick J J 25,131,168
 Woodbury D M 221,223,237
 Woodbury L A 187
 Woodhams P L 89,91
 Woodward D L 221
 Woppard H H 156
 Wraae O 222,233
 Wu C W 213
 Wuerker R B 25,26,131
 Wyckoff R W G 20,135-138,141,152,
 153,154,183,222
 Yamada K M 131
 Yip C C 214
 Young J Z 19,78,135-138,141,152,
 153,154,183,222

AUTHOR INDEX

- Zacks S I 156
Zadunaisky J A 38,224
Zahnd J P 184
Zaunick R 6
Zeitler E 48
Zernicke F 23
Zimmer J 42,43
Zimmerman H M 224,235-236
Zubrod C G 233
Zulch K J 203,206-207

Subject Index

- Abducens nucleus neurons 150,155
Aberrations of lenses 7,36
Ablation experiments 22
Abscesses, cerebral 203-204,238
N-acetyl- β -galactosaminidase 45
Acetylcholine
distribution in nervous tissue 209
in neuroglial cells 206
receptors 211,212,234
in synaptic vesicles 226,227,228
as transmitter 155,156,196,209,229
Acetylcholinesterase 44,209
Acetylneuraminate cytidyltransferase 44
Achromatic objectives 7
Acid DNase 44
Acid hydrolases in lysosomes 158
Acid phosphatase 44
Acoustic microscopy 23
Acoustic nerve ganglion 138
Action potentials 21,23,27,32,46,208
Active transport, criteria for 208,
224,230
Actomyosin filaments 25,126,198-200
Adenosine
in neuroglial cells 206
as transmitter 206
Adenosine triphosphate 208,218
as transmitter 155,218
Adhesive fluid in vitro 222
Adrenalin (epinephrine) as transmitter
17,155,212
Adrenergic fibres 209
Adrenocorticotrophic hormone as
transmitter 155
Affinity for cellular elements 207,
217,233,234
effects of homogenisation on 234
Agglutinin 210
Agnostic views in science 4
Albumin 19
Alkaline DNases 44
Alkaline phosphatases 44
Alphaglycerophosphate dehydrogenase 44
Aminoacids 24
transport 211
Amoeboid movements 18
Anaesthetics 24,225,233
Anaplasia 206
Angiotensin as transmitter 155
Aniline dyes for staining 10
Anopteral microscopy 23,32
Anterior horn cells, see ventral horn
cells
Anterior white commissures 116,154
Antibiotics, effect on transport 221
Antibodies 26,40,46,80,199,210,212
as markers 90
production 217
Antigen - antibody reactions 46,251,252
Antigens 25,40,46,132,199,210,212
Antimitotic drugs into ventricles 243
Arabinosidase 45
Arey's discs 36
Artifacts 2,23,127
aberrations 36
astrocytic foot 165-168
cracks 36
dehydration 34,36
in egg albumin 19,197
of electron microscopy 46-48,127,
189,255
endoplasmic reticulum 189,209
Golgi body 14,189
light microscopy 189
list of, and likely origin 127
myelin lamellae 210
nuclear pores 158-159
pre- and post synaptic thickenings
203
shrinkage 36,251
staining 19,123
study of 244
synapses 196-198
synaptic vesicles 154
trilaminar appearance, explanation
126,133
unit membrane 198
Artificial membranes 208-209,221,252,
253
Arylsulphatase 45
Ascites cells 209
Aspartate as transmitter 79,155
Aspartate aminotransferase 44
Assumptions inherent in techniques
4,27,30,199,230,243

SUBJECT INDEX

- Astroblasts 206
 Astrocytes, see also reactive astrocytes
 appearances by electron microscopy 64-67, 166
 appearances by light microscopy 54-56, 162-194
 attachment to blood vessels 18, 159, 164
 electrical properties 23, 46, 79
 enzymes in 44-45
 inability to distinguish from other cells 37-39, 42, 43, 62, 63, 187-188
 markers of 86, 88-91
 in oedema 224, 225
 original descriptions 17-18, 32
 proliferation 203-204
 staining of 36, 74-77, 92
 swelling 224
 tissue culture 36, 74-77, 85, 92
 tumours of 206
 Astrocytic feet 95, 164-167, 191, 194
 Astrocytoma 207
 ATP as transmitter 155
 ATPase 45, 78, 208, 211, 227
 Auditory cortex 154
 Autofluorescence of neurons 168
 Autolysis in culture 41
 Automatic photography 23
 Autoradiography 23
 Axodendritic synapses 150
 Axonal flow, axoplasmic transport, axonal transport 24, 34, 191
 Axons
 horse radish peroxidase for seeing 50
 in invertebrate preparations 40
 in mechanical integrity of nervous system 201
 myelin sheath 115, 173-177
 relation to dendrites 156
 original descriptions 6, 7, 13
 relation to neuromuscular junction 156
 relation to oligodendrocyte 165
 relation to Schwann cell 20, 181-182, 189
 staining of 74-77
 structure of 115-118
 teasing out 31, 32
 in thick sections 50
 in tracts 21
 white matter 22, 115, 169
 Axoplasm 6, 7, 115, 181, 189, 208, 219
 Axosomatic synapses 150
 Bacterial toxin receptors 210
 Basket cells of cerebellum 113, 136, 189
 Beta endorphins as transmitters 155
 Beta-galactosidase 45
 Beta-glucosaminidase 45
 Beta-glucosidase 45
 Beta-mannosidase 45
 Bicarbonate ions 220
 Bichromate staining 10
 Bielschowsky's stain 74-77, 157
 Bimolecular layer 24, 210
 Binding of non-biological substances 211
 Biochemical changes in culture 46
 Biochemical markers 41, 44-46, 119
 Biochemical studies 27
 Biochemistry of brain, dominated by syncytium 240-241
 Biological membranes and their thickness 251-254
 Biopsy 23
 Biosynthesis 217
 Birefringence and its measurement 24, 251
 Blood brain barrier 224, 225, 233
 Blood vessels 156, 201
 Bombesin as transmitter 155
 Brachial nerve 115
 Bright field illumination 32, 35
 Brownian movements
 of intracellular particles 34, 157, 176, 199
 in syncytium 183
 Bulk or enriched fractions of cell types 241
 Bungarotoxin 228
 Butyrylcholinesterase 44
 Cajal's stain 74-77, 93, 162
 Calcium ions 196
 Calibrations in electron microscopy 48
 Capacity of cell membranes 24
 Capillaries
 in nervous system 33, 72, 94, 114, 164, 172, 189, 205, 224
 for mechanical integrity 201
 Carbon fourteen 223
 Carbonic anhydrase 45, 88
 Carbon dioxide causing fits 238
 Carbowax 94
 Carmine staining 10
 Carnosine 219
 as transmitter 155
 Carriers for aminoacids, ions, sugars 210, 211, 218
 Catecholamines 17, 46, 156, 205
 Cathepsins 45
 Cation exchange as mechanism for ion concentration 201, 219
 Cell bodies or somas 17, 27, 30
 Cell compartments 158
 Cell, history of 6-7
 Cell membranes
 absence around microglia 95
 absence in syncytium 92
 carriers in 211
 chemical and physical properties 24-25, 135
 Davson-Danielli hypothesis 25, 132
 definition of 251
 dimensions of 132, 253
 effects of staining 35
 electronmicroscopical appearances 46-48
 fluid or solid 134
 around neurons 18, 120, 231
 pores in 211
 proteins in 26, 132

SUBJECT INDEX

- relation to microsomes 132
- resolution of 46
- Singer-Nicholson hypothesis 26,134
- structure of 211,243
- transport across 207-221
- Cell motility 24
- Cell pulsation in culture 41
- Cellularity 6,206
- Central canal 201
- Central excitatory, inhibitory states 226
- Centres 22-23
- Centromeres 19
- Cerebellar cortex 43,148,150,155,185
- Cerebellar white matter 12,136
- Cerebellum
 - basket cells in 136
 - cytoskeleton in 131
 - development of 185
 - electrical activity in culture 149
 - granule cells 153
 - intracellular filaments, fibrils, tubules 131
 - migration of cells during development 184
 - mitochondria in 94
 - myelin in 176
 - myelin sheaths in 180
 - Purkinje cells 113
 - stellate cells 153
 - synapses in 131,148,155
 - tissue culture 148-149
 - white matter 115,117,169-171
- Cerebral commissures 136
- Cerebral cortex
 - arborisation 143,185
 - cytoarchitectonics 21,22,183
 - cytoskeleton in 131
 - embedded, unstained 95
 - intracellular filaments, fibrils, tubules 131
 - localisation 22
 - motor tracts from 22,169
 - stained appearance 114
 - synapses in culture 148
 - synapses, density and dimensions 138,152
 - synaptic clefts 150
 - tissue culture 43,85,148
 - white matter 169
- Cerebral myelin 213
- Cerebral oedema 224-225
- Cerebral slices 187,188,211,222,239
- Cerebral thrombosis 2
- Cerebroside sulphotransferase 44
- Cerebrospinal fluid 165,224,233,236, 239
- Cerebrum 8,12,21,29,78
- Changes of cells during culture 41
- Chelating agents 196
- Chemical hypothesis of nerve conduction 197
- Chemical hypothesis of transmission 198
- Chloride ions 220,221,223
- Cholecystokinin as transmitter 155
- Cholesterol 227
- Choline acetylase 209
- Choline acetyltransferase 44
- Cholinergic fibres 209
- Cholinesterases
 - in nervous tissue 209,213,227
 - at neuromuscular junctions 156
- Choroid plexus 185,224,233,234
- Chromatic aberration of lenses 36
- Cilia 172
- Cinemicrography 23
- Cisternae 158
- Classification of cell types 205
- Clones 80
- Cochlear nerve 175
- Colour stains 46,47
- Colpoids 23,200
- Commissures 115-116,201
- Computed axial tomography 224, 225,227
- Concept 256
- Conduction in nerves 21,197
- Conductivity of tissues 41,236
- Control experiments 4
 - difficulty of statistics in electron microscopy 47
 - early experiments 10,23,193
 - effect of homogenisation on affinities 80,234
 - insufficient in histochemistry, histology, immunocytochemistry, electron microscopy 4,11,36,73,80,199,205,244
 - absence in subcellular fraction studies 240
 - test of specificity of stains 194,242
 - for techniques 242
- Convulsions 236-238
- Cornea 156
- Corpus callosum 43,85
- Corrections of lenses 36
- Cortical mapping 21,22
- Corticosteroids as transmitters 155
- Coverslip corrections 36
- Cranial nerves 22
- Cranial nerve nuclei 5,20,29,30, 124,230
- Cresyl violet staining 74-77,93,94, 97,112,113,183
- Criteria for extracellular markers 221
- Critical point drying 202
- Critique of our own experiments 119-125
- Cryofixation 202-203
- Cryomicroscopy 23
- Cues 256
- 2',3' cyclic nucleotide-3-phosphohydrolase 45
- Cytoarchitectonics 21
- Cytochrome oxidase 44,78
- Cytoplasm
 - appearance by electron microscopy 64-71
 - appearance by light microscopy 52-61
 - as a cation exchange resin 221
 - compartments in 158
 - early description of 6,8
 - effects of staining on 34-35
 - endoskeleton or cytoskeleton 25,157

SUBJECT INDEX

- fibres in 23
 Golgi apparatus 14
 intracellular movement 157
 low refractive index 157
 mitochondria in 148
 of neurons 157,189-190
 staining of 92-95
 low viscosity in life 126,157
Cytoskeleton or endoskeleton (see also endoplasmic reticulum)
 absence in living cells 190
 appearances by electron microscopy 25-26,64-71
 as artifact 19,36,126,127,157
 compartments in 158
 dimensions of elements 131
 fluorescence and 40
 Golgi apparatus and 14
 incompatibility with intracellular movement 23,126,198-200
 not seen by light microscopy 126-127
 not seen in living tissues 40
 relation to microsomal fraction 132
 as precipitate 36,200
 as single strands 131
 trabeculae, filaments, tubules and fibrils 131
- Dark ground or dark field microscopy** 23,32,118
Davson-Danielli model of membranes 24-25,132-135,158,210,217
Decortication 22
Dedifferentiation in culture 41,46
Degeneration studies of tracts 22,23
Dehydration during preparation for microscopy 24,26,34-36,40,123,169, 212
Deiters' cells, see microglia 10,17,18
Deiters' nucleus, cells of 18,92,128-130, 153,154,160,243
Del-Rio Hortega's stain 75-77,161,164
Demyelinating diseases 238-240
Demyelination by toxic agents 177
Denaturation during preparation 34
Dendrites
 absence of connections preterminally 136,141
 arborisation 143,185
 broken during isolation of neurons 27
 of cerebellar basket cells 113
 different connections 17
 early descriptions 13
 number observed in section 156
 relation to processes 41,50,204
 structure and staining 52-61,64-71, 74-77
 tissue culture 80
Dense core vesicles 19
Depolarisation 238
Depolarising agents 196,238
Deposits 19,48
Development of brain, migration of cells 184-185
Diapedesis from blood vessels 203
Diaphragm 219
- Dichromate stains** 46
Dielectric properties of syncytium 236
Diffusion
 during biochemical preparation 34
 within cells 157,207,217,236
 facilitated 230
Dimethoxyphenylethylamine as transmitter 155
Disruptive techniques, undesirability of 243
DNA polymerase 44
Donnan equilibrium 220
Dopadecarboxylase 45
Dopamine
 in neuroglial cells 206
 as transmitter 155,206,234
Dorsal cochlear nucleus neurons 150
Dorsal root ganglia
 cell structure 31,45,85,116,171-172, 252
 filaments and tubules 131
 isolation 31,108,115
 synapses on 148,152
 tissue culture 43,148
DPX medium 35
Drugs causing fits 237
Drug receptors 1,210,218,230-235
- Ectoderm** 205
Eggs 24
Electrical conduction between neurons, hypothesis 196,197
Electrical properties of neuroglia 23,78
Electrical stimulation of synaptosomes 228
Electron microscopy 255
 appearance of cells 64-71,83-84
 artifacts in 46,255
 bombardment by electrons 35,197, 255
 calibration in 48
 cell membranes 46
 cytoskeleton by 40
 difficulties of 47-48,201-203
 early findings 25-26
 high voltage 26
 myelin sheaths 19-20
 neurons and naked nuclei 108-111
 preparation for 30
 rapid freezing 202,255
 ribosomes and lysosomes 19
 synaptic connections 17
 temperature rise in 255
 thick and thin sections 2
 wet chambers 26
Embedding 30,34,35
Embryonic tissue 23,24,41,81,205
End-feet, see synapses
Endocytosis 19,127,159,227,256
 of vesicles 127,229
Endoplasmic reticulum
 appearance by electron microscopy 64-71
 as artifact 127,134,189-190
 compartmentation 158-159
 dimensions 26

SUBJECT INDEX

- early description 25
- relation to Golgi body 14
- relation to microsomal fraction 48, 132, 134, 209
- Endorphins 46
- Endoskeleton, see cytoskeleton
- Endothelial cells 81, 172
- End-plates, see neuromuscular junctions
- Enzymes
 - and activities 44-46, 78, 132, 158
 - induction 217
 - as markers 44, 46, 79-80, 88-91
 - on membranes 213-216
- Ependymal cells
 - migration 184, 185
 - appearance 10, 114, 172, 189, 191, 205, 224
 - tumours of 206
- Epidermal cells 205-206
- Epilepsy 236-238
 - agents causing it experimentally 237
 - foci 22, 62, 236-238
 - movements 62
- Epoxy resins in electron microscopy 30, 48, 95, 100
- Eukaryotes, compartments in 158
- Exanthemata, antibodies to 212
- Excitability of cells in culture 21, 22, 23, 41, 78, 148
 - of neurons 27, 73, 78, 79, 234
- Excitation
 - passing between neurons 79, 141, 238
 - across sarcomeres 156
- Exocytosis 19, 127, 159, 256
 - of vesicles 228, 229
- Experimental lesions 22
- Explants 23
- Extirpation of nuclei or centres 22
- Extracellular compartment 158, 221-224
- Extracellular fluid 35, 92, 158, 184, 221-224
- Extracellular space markers
 - in cerebral slices 222
 - by electron microscopy 222
 - estimated by impedance 222
 - markers 221, 223
- Extracted myelin 209, 239
- Falsifiability of hypotheses 133
- Femoral nerve 115
- Ferritin moving into parenchyma 185
- Fibres, fibrils
 - in culture 23, 204
 - myelinated 19
 - orientation of 127
 - preterminal 141
 - in reticular theory 12-13
 - staining of 10, 52-61
 - structure 189
 - in syncytium 11
 - tracts 191
 - white matter 21, 115
- Fibrin 74-77, 204
- Fibroblasts 81, 131, 168
- Filaments 12
- Fixation 7, 23, 26, 40, 121, 217, 229, 255
- Flagellar movements 172
- Fluidity and fluid definition 134
- Fluid mosaic model 25-26, 132-135, 254
- Fluorescence of cells 26, 40, 199
- Fourth ventricle, floor of 107, 172
- Free nerve endings 156
- Free nuclei 170
- Freezing lesions causing fits 238
- Freezing techniques in electron microscopy 25, 26, 40, 132, 201-203, 212, 251, 255
- Friedreichs ataxia 238
- Frog skin as model for active transport 209, 219
- Frontal cortex and lobe 94, 95, 96, 100, 102, 114
- Galactocerebrosidase 45
- Galactosyltransferase 44
- Gallyas's staining procedure 92, 93, 99, 100-108, 124
- Gamma aminobutyric acid
 - in neuroglial cells 206
 - as transmitter 155
- Gammahydroxybutyrate as transmitter 155
- Gangliogliomas 207
- Ganglion cells, dorsal and sympathetic 7, 17, 23, 152, 171-172, 189, 243
- Ganglion cells of retina 141
- Ganglioneuromas 207
- Gangliosides 227, 238
- Gastrin as transmitter 155
- Gemistocytic astrocytes 207
- Genotype for enzyme formation 79, 80
- Geometrical considerations
 - of cells 49-51
 - of cell membrane 47, 194
 - cytoskeleton 134, 189-190
 - dendrites 141
 - freezing techniques 203
 - myelin lamellae 173-176
 - orientation of section 2, 49-50, 51
 - of synapses 144-148, 151
- Geren model of myelination 20, 174, 177-179, 191
- Germinal layers in developing brain 185
- Giant axons of squid and cuttlefish 40, 208, 221, 230, 252
- Glare in microscopy 30
- Glia, see neuroglia
- Glial fibrillary acid protein 86, 89, 90
- Glial filaments 25, 74-77, 131, 204
- Glial processes 41, 74-77, 80, 131, 204
- Glioblastoma 18
- Gliomas
 - classification of 206-207
 - spread of 225, 243
 - structure of 18, 225
- Gliosis 18, 24, 203-204
- Glucocerebrosidase 45
- Glucose 238
- Glucose phosphate dehydrogenase 44
- Glucose phosphate isomerase 45
- Glutamate as transmitter 79, 155

SUBJECT INDEX

- Glutamate decarboxylase 45,88
 Glutamate dehydrogenase 44
 Glutamate synthetase 86
 Glutamine synthetase 45
 Glutaraldehyde fixation 100
 Glycine as transmitter 155
 Glycoproteins 212
 Gold stain 10
 Goldfish Mauthner neurons 40
 Golgi body or apparatus 13-14,19,25,
 26,36,127,157,189-190,200,201
 as compartment 158
 Golgi and Golgi-Cox stains 36,62,63,
 74-77,93,141,142,164
 Granule cells in cerebellum 153
 Granules, apparently on nuclear
 membranes 159
 Granules, fluorescent 141
 Granules in tissues 10,15,18,19,20,21,
 23,24,34,136,158,258
 Ground substance of brain 11,168,
 183-188
 Guanylate cyclase 45

 Haematoxylin and eosin staining 19,35,
 46,74-77,92,93,94,95,96,99,100-108,
 112,113,114,116,124
 Haemoglobin 220
 Hand dissection of neurons 5,9,27,50
 Heavy metal stains in electron
 microscopy 10,46-48,133,156,159,
 251,255
 Heirarchy of affinities 217
 Hemispherectomy 22
 Hexokinase 44
 High voltage electron microscopy 26,47
 Hippocampus 154
 Histamine as transmitter 155
 Histiocytes 203
 Histochemistry 11,23
 Histology of stained tissues 22,33-40,
 82-83
 History of cellular structure 6-26
 Holzer's stain 74-77,93,172
 Homogenisation and homogenates 20,80
 Hormones causing fits 237
 Hormone receptors 210
 Horse radish peroxidase 50,141,185
 Hydrocephalus 226-227
 Hydrolase ratios in neurons and
 neuroglia 44-45
 Hydroxyl ions 220
 5-Hydroxymethyltryptamine as
 transmitter 155
 Hyperbaric oxygen causing fits 238
 Hypoglycaemia 233,238
 Hypotheses 17,119,196,218,243,244
 Hypoxia causing depolarisation 196,
 208,224,235,238

 Identification of cell types 21,37-39,
 49-66,74-77
 Illumination 28,80
 Immunization 10
 Immunocytochemistry 23,26,40,46,80,
 199,205-206

 Immunofluorescence 40,80,199
 Impedance measurements of extracellular
 space 222,251
 Inducing enzymes 217
 Infections of brain 5,18,24,203-204
 Inflammation of brain 203-204
 Inhibition 141
 Injury 224-225
 Insoluble fraction of brain 236
 Integrity of central nervous system
 200-201
 Intellect, seat of 226
 Interface between two phases 208,251
 Interference microscopy 23,32
 Internuncial neurons 152
 Intracellular electrodes and recording
 73,186-188,208
 Intracellular fibrils 157
 Intracellular fluids 34,94
 Intracellular ground substance 168
 Intracellular injections 23
 Intracellular movements
 actomyosin filaments 199
 axonal transport 24
 imitated by colpoids 200
 in compartments 157
 effects of histological preparation 34
 incompatibility with cytoskeleton 157,
 198
 nucleolonema 20
 in nucleolus 128-130
 tissue culture 41,81
 Intracellular organelles 50,94,157
 Intracellular recording in culture 73,
 149
 Intracellular viscosity 158,189-190
 Intracortical connections 168-169
 Intramembrane nuclear membrane
 space 158
 Intramembrane space 158
 Intramitochondrial space, see also
 mitochondrioplasm 158
 Intranucleolar movement 4,159
 Intraventricular injection of drugs 243
 Intraventricular pressure 224
 Inulin as extracellular marker 221-222,
 223
 moving into parenchyma 185
 Invertebrate neurons, study of 40
 Inverted microscopy 23
 Iodine one three one 223
 Ion binding 219
 Ion transport 209,221
 Iontophoretic application of transmitters
 194-197,231,237
 Irradiation during electron microscopy
 35,46,197,252
 Irreversible binding to tissues 217
 Ischaemia causing fits 238
 Isocitrate dehydrogenase 44
 Isocortex, migration of cells 184
 Isolated gut, transport in 209
 Isomerasers, ratios in neurons and
 neuroglia 45

 Janus green B-neutral red 94,122

SUBJECT INDEX

- Junctional folds 14
- Kidney as control 94, 95, 96-98, 120, 176, 177, 178
myelin, figures in 178
slices 209
- Killing animals 4, 196
- Labilisers 196
- Lactate dehydrogenase 44
- Lamellae of myelin sheath 210, 239
- Laminae of cerebral cortex 21
- Laminar flow in axons 200
- Lateral geniculate nucleus neurons 150, 152, 153, 154, 155, 176
- Lead or lead salts in staining 46
- Leech neurons and neuroglia 40
- Leptoscope 218, 252
- Liases, ratios in neurons and neuroglia 45
- Ligases, ratios in neurons and neuroglia 45
- Light microscopy 1, 23, 25, 26, 32, 33-40, 52-61
studies dependent upon 1
- Lipids
as markers 91
cell membranes 24, 25, 132, 210, 252
chemical 24, 26, 72
extraction during preparation 24, 26, 132, 179, 236, 239, 253
stains 47, 72, 197
structural 210
in synaptosomes 227
- Lipolytic enzymes 239
- Liver staining compared with brain 92, 94, 95, 96-98
myelin figures in 176
microsomal enzymes in 211
slices 209
- Lobster stretch receptors 40
- Low angle diffraction of nerves 19, 24, 34, 202, 209, 239, 251
- Lungs 22
- Luteinising hormone releasing factor as transmitter 155
- Lymph 23
- Lysosomes
acid hydrolases in their absence 200
as artifact 157, 189
as compartment 158
early description 25
membrane around 47
compared with particles 14, 19
- Macrophages 85
- Magnesium ions 154
- Malate dehydrogenase 44
- Mallory's phosphotungstic acid
haematoxylin stain 36, 74-77, 92, 93, 94, 95, 96, 99, 102-108, 112, 113, 114, 120
- Mammillary body 154
- Manic depression 2
- Markers for different cells 41, 44, 80, 86, 88-91, 256
their specificity 119, 248-249
- Marsland, Glees and Erikson's stain 36, 74-77, 92, 93-108, 114, 120, 124, 136
- Mauthner neurons 40, 230
- May-Grunwald Giemsa stain 168
- Measurements (see also shrinkage)
of extracellular space 223
of membrane thickness 251-254
- Mechanists 208
- Medial gastrocnemius nerve 176
- Medial mammillary nucleus 154
- Median septal nucleus 154
- Medulla oblongata 29, 30, 94, 95, 138, 203, 205
- Melatonin as transmitter 155
- Membranes, see also cell membranes
carriers 210
channels or pores 210, 211
potentials 224, 252
semipermeability 219
transport 207-221, 237, 243
- Memory, neurobiology of 78, 244
- Meninges 201, 205
- Mesaxons, internal and external 20, 177
- Mesentery 168
- Mesoderm 205
- Metal deposits in electron microscopy 48
- Metastases 22
- Methionine sulphoxime 238
- 5-Methoxydimethyltryptamine as transmitter 155
- 5-Methoxytryptamine as transmitter 155
- Methylene blue staining 29, 33, 183
- Microchemistry of neurons 23
- Microdissection of neurons 5, 8, 9, 27-32, 242
- Microdissection of peripheral nerves 32, 242
- Microfibrils, microfilaments, microtubules and microtrabeculae 25, 26, 127, 131, 157, 189
- Microglia
amoeboid movements 24, 183
appearance by electron microscopy 68-71, 108, 109
appearance by light microscopy 58-61, 161, 164, 185
appearance in tissue culture 85
relation to, astrocytes 95
as Deiters' cells 10
not distinguishable from other cells 37-39
in gliosis 18, 203
in histology 96-98, 100, 114
no markers for 88
as naked nuclei 92, 94, 169-170
nature of 92, 124, 165
as phagocytes 79, 169, 184
quantimetry 104, 150
identity with reactive astrocytes 79
rim of cytoplasm as artifact 95, 99
specificity of stains 36, 72, 74-77
staining properties 92

SUBJECT INDEX

- in tissue cultures 85
- in tumours 206-207
- Micromanipulation** 23,242
- Microscopy and microscopic techniques** 23,40
- Microsomal enzymes 210,211
- Microsomal fraction 48,132,134,209, 213,214
- Microspectrography** 23
- Microsurgery** 23
- Microtechniques** 23,242
- Microtomy** 7,23
- Microtrabeculae 25,127,131,157,189
- Microtubules 25,26,121,189
- Middle popliteal nerve 176
- Migration of cells during development 184,185
- Miniature end plate potentials 226,229
- Mitochondria**
 - cristae of 13,25,48,158,189-190
 - early descriptions 13
 - fraction 48,240
 - loss during preparation 34,35,121
 - membrane 120
 - movements 34
 - in neuroglial clumps 94,122
 - intramembrane space 158,189
 - intramitochondrial space 158
 - in neurons 172,189,201
 - only structures in cytoplasm 157
 - distinguished from synapses 147-148
 - in syncytium 94,121,189,201
 - in tissue culture 23
 - transport, in and out 209
- Mitochondrial membrane 189,194,235
- Mitochondrial movement 34,157,199,200
- Mitochondrioplasm 35,157,158,190, 210,221
- Mitoses in tumours 206,207
- Models 47,132-133,256
- Modulator** 256
- Molecular sizes in or on membrane 210-216
- Monoamine oxidase 44
- Monosaccharide transport 211
- Motor area of cortex 153,154
- Motor nerves 22
- Mounting of sections 23
- Mucopolysaccharides 34
- Multiple sclerosis plaques 2,238-240
- Muscle 22,156,219,220,221
- Muscle fibres 22,187,209
- Muscle innervation 22
- Myelin and demyelinating diseases** 238-240
- Myelin extracted 209,237
- Myelin figures in liver and kidney 176-178,238
- Myelin lamellae 20,170,171,173-179, 191,210,238,239
- Myelin sheath 173-182,237,238
 - demyelinating diseases 238
 - early descriptions of 19
 - electron microscopy 20
 - geometrical problems 173-179
 - lamellae as artifacts 173-182,179,252
- low angle diffraction 34,209
- model for membranes 24-25
- stained appearance 118
- stains for 74-77
- in unfixed axons 32,115
- Myelinated nerve 7,13,19,20,31, 115,118
- Myelination 22,23,41,180,238
- Myeloarchitectonics** 21
- Myosin and myosin filaments 198-200
- Naive questions** 258
- Naked nuclei, see microglia
- Nerve fibres 12,204-205
- Nerve Growth Factor** 5,8,183
- Neural explants in eye 40
- Neural stains 23,74,93,104
- Neural tube 184
- Neurites 203-204
- Neuroepithelial germinal cells 205
- Neurofibrils 74-77,131,157,189
- Neurofilaments 25,74-77,131,157,189
- Neuroglia**
 - appearances by electron microscopy 64-71
 - appearances by light microscopy 49,54-61
 - cell types 17-18,159-165,203-204
 - clumps 30,32-33,35,78,92,94,121, 168,187,234,240-243
 - difficulty in interpreting cell types 37-39,42-46,72-73,161
 - difficulty in differentiating from neurons 42,62-63,205
 - early descriptions of 5,8,10
 - electrical properties 73,242
 - enzymes in 44-45
 - identification 1
 - intracellular nature 94
 - movements 24
 - phagocytes 79
 - high sodium ion content 235-238
 - specificity of stains 36
 - supporting or satellite cells 78
 - synapses with axons 17
 - syncytium 11,183-188
 - in tissue culture 4,149
- Neuroglial fibres 156
- Neuroglial fractions 241
- Neuroglial nuclei 34,74-77
- Neuroglial processes 156,161,164, 203-204
- Neurological disease 22,36,81,205
- Neuromuscular activity 23,41
- Neuromuscular junction or end-plates 14,17,22,74-77,136,146,156,157, 197,209,226,228,229
- Neuron diagram 190,191
- Neuron-glial relationship 78-79,224
- Neuronal membrane 133
- Neuronal theory 12
- Neurons**
 - appearance by electron microscopy 64-65,109,110
 - appearance by light microscopy 52-54,126-135,205

SUBJECT INDEX

- changes during histology 35-36
 cytoarchitectonics 21
 difficulties in distinguishing from neuroglia 17,37-39,42-46,62,63,205
 difficulties of identification 36
 early observations 7,10,13,17,18
 invertebrate 40
 isolated, respiration of 27
 markers for 88-91
 membrane potentials 27
 memory and 78
 microdissection or isolated 5,27-32, 92
 in serial sections 102-104
 somas 27-32,136-139,141,152,153, 156,159
 stained 96-98
 staining of 10,74-77
 staining with neuroglial stains 124
 tissue culture 23,40-42,85,148,149, 156
 transition forms with neuroglia 18
 action of transmitters 156
 tumours, rare in 206
 unstained but embedded 95,100
 Neurotensin as transmitter 155
 Neurotubules 131,189
 Nissl body or granules in neurons 18,52-53,74-77
 Nodes of Ranvier 19,32,115,117,179,191
 Nodose ganglion 131
 Non-disruptive techniques 218,241
 Non-invasive techniques 20,40,242
 Non-specific binding 211
 Non-specificity of receptors 40
 Noradrenalin as transmitter 155,234
 Nuclear fraction 170
 Nuclear membranes 25,120,158,194,243
 Nuclear pores or pore apparatuses 26, 135,158-159
 Nuclear rotation 34
 Nuclei, see also microglia
 appearances by electron microscopy 64-71
 appearances by light microscopy 52-61
 early descriptions 7,18,20
 free nuclei 170
 isolated 169
 effects of preparation 34,35
 staining 74-77
 Nucleolar associated chromatin 20
 Nucleolar membrane 8,20,31,34,159,160, 189,243
 Nucleoli
 absence in naked nuclei 124
 appearances by electron microscopy 64-71,110
 appearances by light microscopy 52-61
 early descriptions of 7,8,20
 compared with granules 18,19
 movement within 41,128-130,160
 staining of 74-77,134
 structure of 159,160,189
 Nucleololi 18,20,21
 Nucleolonema of neurons 18,20,34,52-61, 64-71,158,159,160,243
 Nucleoplasm 8,18,34,35,64-71,158, 159,190,210
 of naked nuclei 169
 Nucleotides 24,216
 Occam's Razor 4,108,207,217
 Occipital cortex and lobe 21,94, 95,100,131,138,142,153,154
 Octopamine as transmitter 155
 Oculomotor nucleus neurons 150, 154,155
 Oestrogens as transmitters 155
 Oil-water partition coefficients 24
 Olfactory glomerulus 154
 Oligodendrocytes or oligodendroglia
 appearance by electron microscopy 68-69,163
 appearance by light microscopy 56-59,79,161-162
 difficulty of demonstration in normal tissue 165
 difficulty to distinguish from other cells 37-39,42-46,62-63,72, 187-188
 early descriptions of 18
 enzymes in 44
 possible role in gliosis 204
 markers for 88-91
 quantimetry 104
 in neuroglial clumps 32,33
 on peripheral nerves 181
 specificity of staining 36,37
 tissue culture 85
 tumours of 206
 Optic nerve 22,24,43,85,131,176
 Organ culture 80
 Organised structure 257
 Orientation of sections 51
 Osmium salts as used in electron microscopy 10,30,46-47,72,93,95, 100,178,182
 Osmosis 27,34
 Oxidative phosphorylation 13,158, 240
 Oxidoreductases, ratios in neurons and neuroglia 44
 Oxygen tension on potential difference 224,238
 Oxygen uptake of isolated neurons and neuroglia 78
 Oxytocin as transmitter 155
 Packaging 229,258
 Palmgren's stain 31,74-77,93,112, 113,116,117,136,168
 Papilloedema 224
 Paradigm in the sense of Kuhn 244
 Paramecium 172
 Parietal cortex and lobe 21,94,95, 100,109-111,131
 Pars amorpha of nucleolus 34,158, 221
 Particles 14,35,258
 in an assembly 256
 as decorations 256
 Patay's triple stain 74-77,92,93,102-108, 114

SUBJECT INDEX

- Pathology 33
 Penfield's stain 74-77
 Pericyte 168
 Perineurium 168
 Peripheral nerves 7, 19, 20, 32, 40, 171
 Permeability of membranes 24
 Permeabilisation in cytochemistry 26, 40, 80
 Phagocytes 24, 184
 in gliosis 203
 Phagocytosis 24, 199, 200, 256
 in syncytium 169
 Pharmacological effects of drugs 217
 Phase contrast microscopy 251
 early description 23
 ganglion cells 171
 neurons seen by 30, 31
 nucleolar membrane by 159
 nucleoloneema 20, 128-130
 peripheral nerves 32, 115
 plaques 240
 stained tissue 96-98
 stained with osmium salts 100
 of tissue culture 23, 85, 180
 unstained embedded tissue 35, 98
 Phenylethylamine as transmitter 155
 Phenylethanolamine as transmitter 155
 Philosophy behind this monograph 4-5, 86
 Phosphate esters 219
 Phosphatides 236
 Phosphocreatine or creatine phosphate 218, 219
 Phosphogluconate dehydrogenase 44
 Phospholipase A 44
 Phosphoproteins 212
 Photomicrography 23, 179-180
 Physiognomists 22
 Physiological synaptic activity 41
 Picrate stain 13, 46
 Pineoblasts and pineocytes 206
 Pinocytosis 256
 in neurons 23, 157, 199, 200, 243
 in parenchyma 185
 Plant cells 6, 24
 Plasma proteins into parenchyma 185
 Plasticity 22
 Polarising microscopy 23, 32, 251
 Pools, metabolic 158
 Popper's philosophy 4
 Postscript 244
 Potassium ions
 stimulation depletes vesicles 228
 in epilepsy 236-237
 effect on resting potentials 186, 208, 218, 219, 220
 in hypoxia 224
 on oxygen uptake 78
 Practical recommendations 240-243
 Pre- and post synaptic fibres in ganglia 198
 Pre- and post synaptic thickenings 15, 17, 198, 228
 Precipitates during preparation for microscopy 36, 46, 156, 159, 197
 Prepyriform cortex 150, 154, 155
 Preterminal fibres 136, 139, 197
 Processes of cells 18, 52-61, 64-71, 100
 Proctolin as transmitter 155
 Prokaryotes, compartments in 158
 Prolactin as transmitter 155
 Proportion of tissue occupied by cells 104-108, 183
 Prostaglandins as transmitters 155
 Proteins 24, 25, 26, 34, 80, 88-91
 as markers 88-91
 believed to be in membranes 132, 210, 213-216, 252, 253
 in ribosomes 158
 staining of 47, 197
 structural 210, 212, 217
 synthesis 158, 227
 Proteolytic enzymes 31, 158, 177, 239
 Puffer supramedullary neurons 40
 Purines as transmitters 155
 Psychological disease 22, 46, 244
 Purkinje cells of cerebellum 113, 120, 123, 136, 153, 185, 189
 in culture 149
 Pus 204
 Putative transmitters 156, 196
 defined 257
 Pyramidal neurons 142, 153, 195
 Pyramidal tract 22, 176
 Pyriform cortex neurons 150
 Quantimetry 104-108, 121, 124, 250
 Radiation in electron microscopy 229
 Reactive astrocytes 79, 203, 207, 225
 Receptors
 for drugs, transmitters, hormones and toxins 1, 132, 210, 211, 257
 a general theory of 230-235, 243
 Red cells 24, 25, 132, 220, 221, 252
 Red cell ghosts 209, 237
 Reflection microscopy 23, 251
 Reflexes
 at synapses 141
 without synapses 226
 Refractive index
 of cytoplasm 157, 251, 252
 of incubating media 36, 252
 Resistance and impedance of
 brain 186
 cell membranes 24, 186, 221
 cerebral cortex 186
 cerebrospinal fluid 186
 cytoplasm 220
 serum 220
 spinal cord 186
 syncytium 186
 white matter 186
 Respiration of neurons 27
 Resting potentials
 in brain 186-188
 in membranes 253
 in neurons 27, 73, 208, 236
 Reticular formation 138, 139, 140, 153, 155
 Reticular theory 12

SUBJECT INDEX

- Reticulum 19
 Retina 15, 40, 113, 141, 144, 206, 213
 Reversibility of binding 217
 Rhodopsin 213
 Ribosomes 19, 25, 157, 158, 200
 RNA 216
 Ruthenium red as extracellular marker 221
 S 100 86, 89, 213
 Salt gland of albatross 237
 Saltatory conduction 179
 Sarcomas 206
 Sarcomeres 22, 156
 Satellites or supporting cells 78, 111, 187-188
 Schizophrenia 2
 Schmidt-Lantermann clefts 19, 32, 115, 171, 179, 191
 Schwann cell 20, 168, 174, 180
 cytoplasm 115
 nuclei 168, 177, 178, 182, 191
 sheath 115, 177-179, 182, 191
 Sciatic nerve 31, 115, 118, 176, 213, 214
 Semipermeability of membranes 219
 Septal region 103-104
 Sequestration of particles 257
 Sensory cortex 154
 Sensory fibres 22
 Serial sections 8, 22, 34, 47, 49-50, 99-104
 Serotonin as transmitter 155
 Serum 220
 Shrinkage during histology and electron microscopy 8, 24, 26, 33, 34, 35, 36, 50, 80, 100, 121, 123, 135, 156, 251
 Sialyl transferase 44
 Silver stains 10, 14, 16, 23, 46, 136, 140, 169, 197
 Singer-Nicholson model of membranes 25, 132-135, 158, 212, 217, 254
 Skin, free nerve endings in 156, 176
 Sleep factor delta as transmitter 155
 Sodium ions
 epilepsy 236
 as markers 221
 in neuroglia 235-236
 transport 208, 211, 218, 219, 220
 Somatostatin as transmitter 155
 Sorbtion hypothesis 219, 230
 Specialisation 257, 258
 Specific cell markers 79-81, 86, 88-91, 248-249
 Specific of staining procedures 36, 63-91, 120, 151, 242, 245-247
 Spherical aberration 36
 Spinal cord
 anterior or ventral horn cells 29, 112, 131, 136, 138
 anterior white commissure 116
 early observations 7, 15
 explants of 23
 fibres in 171
 Golgi apparatus in 14
 light microscopy 171
 microdissection of cells in 29
 microglia in 95
 myelination in 22
 neuroglia in 94
 nucleolar membrane in 20
 quantimetry 108
 synapses in 15, 137
 tissue culture 85, 148
 white matter 22, 115, 117, 169
 Spread of gliomas 206-207
 Spreading depression of Leao 238
 Stabilisers of membranes 196
 Staining and staining procedures 2, 10, 21, 23, 26, 35, 36, 46, 94, 123, 205
 their unreliability 245-257
 used here 93
 Statistics 11, 81
 Stellate cells in cerebellum 153
 Stimulation of brain cells 21
 Stomachs 22
 Storage of transmitters 258
 Streaming 199
 Structure of nervous system (summary) 189-192
 Strychnine neuronography 22
 Subarachnoid space 185
 Subcellular fractionation 1, 48, 132, 158, 169, 170, 209, 210, 218, 228
 Subcellular organelles 27, 35, 48
 Subfornical organ neurons 150, 154, 155
 Subjectivity in experiments 36, 244
 Substance P as transmitter 155
 Succinate, cytochrome C reductase 44
 Succinate dehydrogenase 44, 78
 Sucrose as extracellular marker 221, 223
 Sudan III as stain 182
 Sulphate as marker 221, 223
 Superior cervical ganglion 138, 228
 Supramedullary neurons of puffer 40, 230
 Sympathetic ganglia 31, 138, 152, 171-172, 252
 Synapses
 as artifacts 135-156, 197-198
 relation to dendrites 156
 density of 152, 153, 154, 155
 dimensions of 151, 153, 154, 155
 early observations 8, 9, 15-17
 electron microscopy of 145-155
 fraction of (synaptosomes) 80, 134, 226
 relation to neuromuscular junction 14, 209
 particles within 19
 preterminal fibres, absence of 136-146
 pre- and post synaptic thickenings 15
 in tissue culture 23
 in transmission 194-197, 226-230
 Synaptic activity, defined physiologically 172
 Synaptic cleft 149, 150, 197, 227
 Synaptic knobs 189
 Synaptic vesicles 15, 17, 19, 141, 154, 157, 226-230

SUBJECT INDEX

- Syncytium of neuroglia
 relation to astrocytes 224
 as blood-brain barrier 224,233
 different concepts of 6,11,39,201
 during development 184
 early views on 11-13,18
 environment of neurons 224,238
 relation to extracellular fluid 225-226
 fibres in 116
 few membranes in 210
 mitochondria in 94
 relation to neuroglial clumps 92,
 121-122
 determines permeability 221
 poor staining of 92,101
 structure of 165-169,183-188
 most of tissue 108
 tumours in 207
 viscous jelly in life 189
- Taurine as transmitter 155
 Teasing tissues 240,242
 Television microscopy 23
 Teloglia or telodendroglia 141,197
 Temporal cortex or lobe 21,154
 Testosterone as transmitter 155
 Tetanus and tetanus toxin 10,86
 as marker 90-91
 Tetrahydroisoquinolines as transmitters 155
 Thalamus 43,152
 Thiocyanate as extracellular marker 221,223
 Thought, neurobiology of 226,244
 Three-dimensional model 126,194
 Thyrotropic releasing hormone as transmitter 155
 Thyroxine as transmitter 155
 Time lapse photography 23,40,181
 Tissue culture 42-46
 appearance of cells in 80-81,84,85
 biochemical studies of 41,241
 cell identification in 37,40,42,73,78,
 85
 changes during culture 41
 conduction in 41
 disappearance of cells in 41
 dissociated cells 81
 early cultures 23-24,42-46
 de-differentiation in 41
 difficulty of distinguishing parent cells 42,43
 intracellular movements 23,40,81
 as living tissues 40-42
 myelination in 177,180,238
 physiological properties in 41,73-74
 retina in 40
 Schwann cells in culture 180
 Tissue extracts 63
 Tissue in electron microscopy 255
 Tissue transplantation 23
 Toad bladder 209
 Toluidine blue staining 183
 Toxic agents inducing myelin figures 177
 Toxins 78,91,196,212
- Tracheal epithelium 172
 Tracts 21-23,191,201
 Transferases 44
 Transmission 14,17,191,194-198,
 226,243
 Transmitters
 in absence of synapses 194,196,197
 in cells 80
 defined 257
 listed 155
 as markers 90
 movements during preparation 205
 in neuroglial cells 205-206
 and pre-synaptic membrane 228
 receptors for 132,210,230,231,235
 Transport across membranes 209,230
 Transport enzymes 210,211
 Trapezoid body 16,152
 Triethyl tin poisoning 224,235,236
 Trilaminar appearance of membranes 24,126-127,132,152
 Trypsin for separating cells 31-32
 Tryptamine as transmitter 155
 Tryptolines as transmitter 155
 Tube length corrections 36
 Tumours of central nervous system
 causing fits 2,38
 classification 1,5,206
 clinical history 81,206-207
 in culture 81
 neurological and physiological function with 22,36
 pattern of cells in diagnosis 62
 relation to parent cells 41,80
 sodium ion concentration 235
 spread of 225
- Tungsten salts in staining 46
 Tyramine as transmitter 155
- Ultrasound for ventricular volume 224
 Ultraviolet microscopy 23,40
 Unfixed tissue 34
 Unit membrane 24,25,47,132,135,
 217,252
 as artifact 33,198,252,253
- Unmyelinated fibres
 in brain 168
 in periphery 13,131
- Unsatisfactory terms 256-257
 Unstained tissue 31,32,40,47,95
 Utricularia 10
- Vacuoles 20,21,35
 Vacuum during electron microscopy 35,46
- Vascular glia 32-33
 Vasoactive intestinal peptide as transmitter 155
 Vasopressin as transmitter 155
 Ventral cochlear nucleus 154
 Ventral or anterior horn cells
 cytoskeleton in 131
 density and dimensions of synapses on 152,153,154,155
 dissection of 29,30
 myelination in 22

SUBJECT INDEX

- nucleolar membrane in 20
- stained by most procedures 112,120,123
- synapses on 137,138,140,148,152-154
- synaptic clefts 150
- Ventricles 10,185,191,224
- Vertical illumination microscopy 23,41
- Vesicles 19
- Vesicle hypothesis of transmission 17,
194,226-230
- Video-enhanced contrast microscopy 127
- Viral toxin receptors 1,210,212
- Viscera, free nerve endings in 156
- Viscosity of brain 187
- Viscosity of cytoplasm 157
- 'Vital' staining 23
- Vitalists 208
- Vitrification during freezing for electron
microscopy 202-203
- Warm stages 23
- Water, see also dehydration and shrinkage
 - in cells 34,202
 - in cell membranes 113-134
- Weigert Pal's stain 74-77,93,117
- Weil and Davenport's stain 74-77,92,
93,94,98,99,102-108,120,124,197
- Wet chambers in electron microscopy 26
- White matter 21-23,115,117,121,136,
169-171
- Wolfgram protein 213
- X-ray microscopy 23,251
- Yeast 209