

## Study of Possible Correlations Between Prenatal Brain Development and Placental Weight<sup>1</sup>

S. ZAMENHOF, L. GRAUEL and E. VAN MARTHENS

Department of Medical Microbiology and Immunology,  
Department of Biological Chemistry,  
Brain Research Institute and Mental Retardation Center, School of Medicine,  
University of California, Los Angeles, Calif.

One of several factors, studied in our laboratory, that influence DNA synthesis and cell number in prenatal brain [4, 6, 12], is placental development [review in 13]. WINICK [10] and STEPHAN and CHOW [5] have demonstrated that maternal dietary restriction results in placentas of lower cell number, and lower weight, respectively. The newborns of such mothers have lower neonatal brain cell number [10, 12] but the relationship between the placental and the cerebral development has not been established.

Several reports indicate a positive correlation between placental weight and birth weight [review in 3]. The development of the placenta, especially from the point of view of placental cell number (DNA) has been the subject of a recent study by WINICK and NOBLE [8] and by WINICK *et al.* [9].

The object of this work is the study of correlations between various neonatal parameters, especially placental weights and brain development in *normal* animals (rabbits). In particular, the aim of the study was to establish whether, on a statistical basis, an individual with a heavier term placenta is also likely to have a higher neonatal cerebral weight or cerebral cell number.

### *Materials and Methods*

11 6-month-old does of the New Zealand White strain, weighing 3-4 kg, were mated and 80 fetuses and their (fetal) placentas were removed by cesarean section on day 30 of gestation; an additional 10 does were used for cesarean sections between days 20 and 28

<sup>1</sup> Presented in part at the Annual Meeting of the Perinatal Research Society, September 1970.

providing 7–20 fetuses for each of the days. The fetuses were weighed, then cerebral hemispheres (without cerebellum and olfactory lobes) were dissected out, and their wet weight as well as the wet weight of the corresponding placentas, were determined. Total cerebral DNA was determined as described previously [4, 11]. From this DNA content one can calculate the total number of cerebral cells [13]: neonatal cerebral DNA 1,804  $\mu\text{g}$  (mean value, figure 3); DNA content of diploid rabbit cell [7]  $5.3 \times 10^{-6} \mu\text{g}$ ; total neonatal rabbit cerebral cell number  $3.4 \times 10^8$ .

In the rat, neurons (neuroblasts) cease to proliferate around birth and the neonatal cell number (as measured by neonatal DNA) is indicative of the final neuron number [for a discussion see 4, 6, 12, 13]. To our knowledge, the time of cessation of neuron proliferation in the rabbit has not been reported; thus, for the time being, the neonatal cerebral DNA determined in this work, indicates only the total cerebral cell number. As in the case of term in the rat and hatching in the chick [13] we now find that this number (DNA content) also reaches a plateau around term in rabbit cerebrum: 45% at 20 days, 67% at 22, 79% at 24, 88.5% at 25.5, 97.5% at 27.5, assuming 100% at 30 days. This time, then, is convenient for DNA determination, because a small error in the estimated fetal age does not result in




Fig. 1–3. Correlations between parameters at 30 days of pregnancy (rabbit). Numbers 1 to 9 indicate offspring of the same litter, with A and B representing litter number 10 and 11, respectively. Since the graphs illustrate the spread around the slope, identical values for littermates were eliminated due to technical difficulties in plotting.


Fig. 1. Placental weight minus its mean (g) vs body weight minus its mean (g). Equality of slopes,  $F=1.89$  on 9 and 54 degrees of freedom (DF) with probability,  $p=0.07$ . Common slope regression coefficient,  $b=0.109$ ,  $t=14.66$  on 63 DF, probability (zero slope error),  $p<0.0001$ .

an appreciable difference in cerebral DNA content. Day 30 was chosen as close enough to term (31.5), without the risk of natural birth and the loss of the placenta; the corresponding values are referred to as 'neonatal'.

The following statistical method was utilized [14] to determine if significant linear correlations exist between any of the following measured parameters: placental weight, body weight, cerebral weight, and cerebral DNA. A slope was estimated for each individual litter of pups and an *F* test for equality of slopes applied to demonstrate the constancy of the linear relationships amongst all of the litters. If the slopes were indeed similar, yielding  $p > 0.05$ , a regression coefficient, *b*, was estimated for the entire group. The degree of correlation was determined by testing the equality of the regression coefficient to zero, with  $p < 0.01$  constituting significance.

### *Results and Discussion*

The problem of quantification in the placenta is a difficult one because of the number of factors involved. The permeability of the placenta, the exchange surface, and the maternal blood flow to the placenta are all involved, to a not well defined degree. It has been reported that maternal blood flow is greater to the placentas of larger fetuses [2]



*Fig. 2.* Placental weight minus its mean (g) vs cerebral weight minus its mean (g).  $F = 1.56$  9 and 54 DF,  $p = 0.15$ ;  $b = 7.785$ ,  $t = 5.73$ , 63 D F,  $p < 0.0001$ .




Fig. 3. Cerebral DNA minus its mean ( $\mu\text{g}$ ) vs cerebral weight minus its mean (g).  $F=1.06$ , 10 and 58 DF,  $p=0.406$ ;  $b=682.41$ ,  $t=5.56$ , 68 DF,  $p<0.0001$ .

and that blood flow to the uterus increases in proportion to the placental weight [1].

In our work a sample of 20 placentas were dried to constant weight; their water content so determined was practically identical ( $83.35\% \pm 0.63\%$  SD). Thus, for all practical purposes, the weights of placentas are also indicative of the dry weights of their tissues and of the blood therein. This parameter then, the placental weight, was chosen in the present work, bearing in mind the limitations involved.

Our results (fig. 1-3) indicate that highly significant correlations can be demonstrated between 'neonatal' body weight and placental weight, cerebral weight and placental weight, and cerebral DNA (cell number) and cerebral weight. The correlation between cerebral DNA and body weight may be considered as being on the borderline of significance ( $p=0.0186$ ); the correlation between cerebral DNA and placental weight for this size sample was not significant ( $p=0.0589$ ).

Younger fetuses show the same trend of correlations down to at least the 20th day.

In conclusion, on a statistical basis, an individual animal with a heavier term placenta is also likely to have a higher neonatal cerebral weight; the latter is also likely to have a higher number of neonatal cerebral cells (DNA), although for the size of the sample studied, a direct significance of correlation between placental weight and cerebral DNA could not be demonstrated.

### *Acknowledgements*

This study was supported by grants HD-04612 and NS-08723 from the US Public Health Service, OEG 9-70-0043 (057) 00705 from the US Office of Education, and 409 from the Nutrition Foundation. Computing assistance was obtained from the Health Sciences Computing Facility UCLA sponsored by NIH Special Resources grant RR-3. We thank Dr. ALAN FORSYTHE, Department of Biomathematics, for the statistical interpretation.

### *Summary*

Correlations between various neonatal parameters, especially placental weights and brain development in normal animals (rabbits) have been studied. It was found that, on a statistical basis, an individual animal with a heavier term placenta is also likely to have a higher neonatal cerebral weight; the latter is also likely to have a higher number of neonatal cerebral cells (DNA).

### *References*

1. BARRON, D. H.: Homeostasis of the foetus; in FISHBEIN's Congenital malformations, p. 247 (Lippincott, Philadelphia 1960).
2. DUNCAN, S. L. B. and LEWIS, B. V.: Maternal placental and myometrial blood flow in the pregnant rabbit. *J. Physiol., Lond.* 202: 471 (1969).
3. McLAREN, A.: Genetic and environmental effects on foetal and placental growth in mice. *J. Reprod. Fertil.* 9: 79 (1965).
4. MARGOLIS, F. L.: DNA and DNA-polymerase activity in chicken brain regions during ontogeny. *J. Neurochem.* 16: 447 (1969).
5. STEPHAN, J. K. and CHOW, B. F.: The fetus and placenta in maternal dietary restriction. *Fed. Proc.* 28: 915 (1969).
6. VAN MARTHENS, E. and ZAMENHOF, S.: Deoxyribonucleic acid of neonatal rat cerebrum increased by operative restriction of litter size. *Exp. Neurol.* 23: 214 (1969).
7. VENDRELY, R.: The deoxyribonucleic acid content of the nucleus; in CHARGAFF's and DAVIDSON's The nucleic acids, vol. 2, pp. 155-180 (Academic Press, New York 1955).
8. WINICK, M. and NOBLE, A.: Quantitative changes in ribonucleic acids and protein during normal growth of rat placenta. *Nature, Lond.* 212: 34 (1966).
9. WINICK, M.; NOBLE, A., and COSCIA, A.: Cellular growth in human placenta. I. Normal placental growth. *Pediatrics* 39: 248 (1967).
10. WINICK, M.: Cellular growth in intrauterine malnutrition. *Pediatr. Clin. N. Am.* 17: 69 (1970).

11. ZAMENHOF, S.; BURSZTYN, H.; RICH, K., and ZAMENHOF, P.J.: The determination of deoxyribonucleic acid and of cell number in brain. *J. Neurochem.* 11: 505 (1964).
12. ZAMENHOF, S.; VAN MARTIENS, E., and MARGOLIS, F.L.: DNA (cell number) and protein in neonatal brain; alteration by maternal dietary protein restriction. *Science* 160: 322 (1968).
13. ZAMENHOF, S. and VAN MARTIENS, E.: Hormonal and nutritional aspects of prenatal brain development; in PEASE's Cellular aspects of growth and differentiation in nervous tissue. UCLA Forum in Medical Sciences No.14 (University of California Press, Berkeley 1971).
14. Computer program BMDPIV: Analysis of covariance, H. S. Computing Facility (UCLA).

Author's address: Dr. S. ZAMENHOF, Mrs. I. GRAUEL and Dr. E. VAN MARTIENS, Department of Medical Microbiology and Immunology, University of California School of Medicine, Los Angeles, CA 90024 (USA)