

This is an extremely exciting time to be a cosmologist as Roman, Rubin, and Euclid are all starting, or already have started, to take data and will soon contribute to the growing number of intriguing results in cosmology like the evolving dark energy results from DESI or the H_0 and S_8 tensions. Understanding the redshift evolution of cosmological parameters and the growth of structure in the Universe will be key to disentangling some of these mysteries which is why I am particularly excited to study galaxy clusters. Galaxy clusters can give us strong constraints on $\sigma_8(z)$, $\Omega_M(z)$, and $f_{\text{gas}}(z)$ all of which are fundamental to probe the growth of structure in the Universe.

While my ultimate goal is to use these instruments to study galaxy clusters via weak lensing and constrain the redshift evolution of cosmological parameters, there are a number of important intermediate steps that must be performed. Over the next few years, I plan to validate the science pipelines used in cluster cosmology by measuring any inherent biases, work on novel methods of mitigating weak lensing systematics, and perform early science measurements on clusters using these instruments. My previous experiences with cluster and shear calibration equip me with the necessary tools to successfully carry out this proposal.

Previous and Current Research *Blending* During my PhD, I focused on the weak lensing systematic of **unrecognized blends** which are objects (typically galaxies) that are mistakenly identified as a single entity due to severe amounts of overlap between sources (see Fig. 1 for an example). The repercussions of unrecognized blends via detection bias alone have shown shifts in cosmology parameters at the level of a few percent [**ben'clustering, blends'2011, kidsblends'2025**], and an early study including the impact on shape and flux measurements on cosmic shear found up to a 2σ shift in S_8 [**erfan**].

For 3x2pt cosmological analysis, expensive image simulation is used to measure the bias induced by unrecognized blends on redshift and shear calibration. Directly identifying unrecognized blends at the catalog level is an appealing solution that I pursued during my PhD.

In [**unrec'liang**], we showed that unrecognized blends can be readily detected at the catalog level using machine learning techniques that assign an unrecognized blend probability based on measured fluxes. While we showed that **it is possible to remove unrecognized blends from the catalog using observational data alone**, it was unclear if such a removal would ultimately improve our cosmology. I investigated this by studying the impact of our blend removal method on shear calibration using the AnaCal shear framework. We showed that not only does the bias reduce, **the bias can be almost totally removed** with a 30-40% sample cut [**unrec'shear**]. Given that LSST will be systematics limited, a 30% cut to remove a source of bias may be worthwhile.

Clusters Outside of blending I have lead efforts to produce some of the first weak lensing results with the Vera Rubin Observatory by studying the galaxy cluster Abell 360 [**a360**]. For that work I specifically worked on the **source sample selection, photo-z validation with DESI, shear**




Figure 1: Example unrecognized blend in the COSMOS field with observed magnitudes and estimated photometric redshifts attached. Dark blue is the object as detected and measured by Subaru, dashed-dotted red and purple are detections and measurements from HST.

measurement with shapeHSM and AnaCal (Fig. 2), using SSI to evaluate cuts, and cluster mass estimation. As the scope of this project grew, my responsibilities also extended beyond purely scientific as I took on the role of data manager, daily meeting coordinator, and writing coordinator. Along with being my first foray into clusters, this was my first time leading a large collaborative effort.




Figure 2: Tangential shear signal measured around Abell 360 using a variety of shape measurements and source selection. [paper in prep.]

looking into cluster weak lensing. I will validate and measure cluster weak lensing profiles along with working on reducing weak lensing systematics.

Goal 1: Cluster measurements for Roman and Rubin

Precise ensemble measurements of galaxy clusters require an estimate of both statistical and systematic bias. For clusters, this is traditionally done with N-body or hydrodynamic simulations that can capture physics-induced bias (like miscentering of optical clusters due to triaxiality), but is computationally expensive and can ignore biases coming from image processing. I propose to work on both direct image generation and synthetic source injection (SSI) to validate shear and redshift measurements coming from science pipelines.

Direct image generation has been the standard technique to calibrate any biases in cosmic shear [macrann, hsc] as it is able to directly test for systematic biases inherent in the tools and algorithms being used. There are a number of improvements we need to make to these image simulations to properly test clusters, including varying background redshifts, introducing cluster members, intra-cluster light, and cluster triaxiality. Introducing these systematics into our simulations sequentially will allow us to understand the bias introduced by each feature to inform future work on mitigation strategies.

While image generation is a powerful tool, it relies on correctly understanding and implementing all the systematics that exist in data. Alternatively, SSI allows us to characterize the existing data and test the measurement pipeline directly on synthetic sources. In the Abell 360 work, this technique was used to understand the effect of various cuts on the source sample and validate star-galaxy separation along with flux measurement. Repeating this work for Roman and Rubin clusters on a larger scale will allow us to validate measurements made on the real data.

Performing both image simulation and SSI gives us an understanding of what causes biases

Following my efforts on Abell 360, I have been working on testing the combination of AnaCal and RAIL which produce shears and redshift estimates respectively by measuring shear profiles around clusters in the Euclid Deep Field South [publication in prep.] In this work we have **measured tangential shear signals around clusters that were undetectable in the base Rubin catalog** due to cleaner sample selection and higher quality shear estimates, highlighting the power of the combination of AnaCal and RAIL. The combination of AnaCal and RAIL is currently being used for HSC Year 6 cosmic shear and this is one of the first cluster analysis with these tools.

Research Proposal During my postdoc, I plan to focus on the early data from Roman and Rubin primarily

and the total level of bias present in the data. Improving on the existing image simulation tools and developing the pipelines to measure biases via SSI are necessary steps for future data releases with more stringent systematics requirements. My experiences with image simulation to validate blend removal and Abell 360 cluster measurements make cluster validation a natural next step that I am qualified to work on and lead.

With a validated pipeline in hand, we will have confident (or at least properly weighted) cluster weak lensing masses. For Rubin Data Preview 2 there are ~ 100 X-Ray detected clusters by eRosita and ~ 400 SZ detected clusters by ACT, which are more than enough to produce a LSST mass calibration for those external samples. The previous ACT calibration uses a similar number of clusters [citation], so it is possible to produce competitive results using this early data alone. Producing cosmology contours from calibration constitutes another project that would require estimating a covariance, modeling shear profiles (either hierarchically or stacked), redshift distribution estimation, and running cosmology chains, all of which are quite new to me. I am excited to learn and build towards a tomographic understanding of the cosmos.

Goal 2: Mitigate weak lensing systematics

I will continue to work on mitigating weak lensing systematics, especially blending. Extending my previous work on blend removal in [unrec'shear] to include redshift is the next natural step towards understanding the effects of this methodology on cosmology. Completing a redshift study would enable a re-analysis of previous data, like DES Y3 or HSC Y3, providing a comprehensive look at the effect of unrecognized blends in weak lensing.

While my previous work on blending has focused on removing these undetected objects, it may be possible to model the effects of blending for better mitigation without incurring a statistical cost. Using the machine learning probability from [unrec'liang] as a prior, we can treat certain objects as a superposition of two galaxies which can lead to more accurate posteriors for redshift and shear.

For redshift distributions, the self-organizing map (SOM) which maps measured fluxes to a 2D surface, is an effective tool for estimating ensemble distributions $n(z)$ but can also be extended to account for blends as outlined in Fig 3. In that example, we use the blending probability λ to combine two redshift probability distributions: the distribution assuming a single galaxy and the distribution assuming a superposition of two galaxies. In preliminary work, I have shown that this can produce more accurate posterior estimates than a single-galaxy SOM approach. For shears, accounting for cross-redshift blends can be done at the image processing level by improving deblending which can drive down shape noise. Both of these are relatively straight-forward to show as proof of concept, but would require much larger tests to validate.

The best solution to unrecognized blends would be to directly detect them which can be done through joint processing of space and ground based data. While there are strong solutions on how to best combine photometry, there is work to be done on combining shape measurements in order to maximize weak lensing signal. During my postdoc, I plan to contribute to the on-going efforts to enable joint-processing for both Metadetect and AnaCal. This will mostly rely on simulated datasets, like the existing Roman-Rubin simulations, but can be tested on data using deep field overlap of clusters fields which I am well equipped to study.

This is an extremely exciting time to be doing cosmology and I have been at the forefront working with some of the first images taken by Rubin. I want to build on my experiences from my PhD by working on weak lensing systematics and clusters so that we can better probe the universe




Figure 3: Example of modeling blends when producing redshift estimates via a SOM. Instead of only using the cell of the best fit, a weighted SOM can be made from synthetic blends that can be used to produce a more accurate PDF.

tomographically.