




The sspm R package for spatially-explicit surplus production population models

Valentin Lucet ^{1*} and Eric J. Pedersen ^{1,2*}

¹ Department of Biology, Concordia University, Montreal, CA ² Department of Biology, Memorial University of Newfoundland, St. John's, CA * These authors contributed equally.

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Open Journals](#) 

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Statement of need

Population models are important tools for making management decisions, especially in fisheries, where predictive methods like Surplus Production Models (SPMs) are widely used. Fisheries analysts and managers often lack user-friendly, flexible tools to implement and apply SPMs. In addition, SPMs are rarely spatially explicit and usually cannot account for relevant ecosystem drivers. Therefore, there is a need for tools that implement spatially explicit surplus production models (SSPMs). The Northern Shrimp stock in the Newfoundland and Labrador Shelves is an example of a stock in need of an SSPM that can integrate important spatially-structured ecosystem drivers.

Summary

Population modelling is an exercise of interest within environmental sciences and adjacent fields. Early population models such as the logistic model assumed that while the abundance of a population might change over time, the conditions governing parameters affecting that rate of change, such as the maximum rate of growth or the carrying capacity of the population, stay constant over time ([Gotelli, 2008](#)). Modern population models increasingly acknowledge the non-stationary nature of wild populations and work to incorporate environmental fluctuations into dynamic models ([Thorson et al., 2015, 2017](#)). Population models designed to answer applied resource management questions, such as fisheries assessment models, increasingly address how the dynamics of stocks vary across space and time.

Resource managers are becoming increasingly interested in how variation in ecosystem factors such as predator abundance and abiotic variables impact the spatiotemporal variability of population parameters, such as productivity ([Szuwalski & Hollowed, 2016](#); [Zhang et al., 2021](#)). Further, treating spatially structured stocks as single unstructured stocks can lead to substantially biased estimates of population change ([Thorson et al., 2015](#)). However, stock models that explicitly incorporate spatial dynamics and time-varying ecosystem variables are still rare in fisheries science, despite the push for more ecosystem-based management methods in fisheries management ([Berkes, 2012](#); [Crowder et al., 2008](#); [Tam et al., 2017](#)).

Surplus production models (SPMs) are one of the classic models used in fisheries and are based on modelling changes in the total biomass of a stock in a given location over time as a function of current stock abundance and fishing pressure ([Walters et al., 2008](#)). Classically, SPMs assume single unstructured stocks with purely logistic dynamics ([Walters et al., 2008](#)) and, as such, have been of limited use for modelling more complex stocks. They are useful in data-poor contexts where the age structure of the population is not accessible or when age or length structure do not change substantially over time ([Prager, 1994](#); [Punt, 2003](#)). SPMs typically model spatially constant productivity. They also assume that populations are only affected by past abundance and fishing, which ignores stressors like climate change which

42 affect growth rates independently of fishing pressure.

43 In the context of climate change and shifting ranges, fisheries productivity is likely to be a
 44 moving target (Karp et al., 2019), and managers need better methods that account for varying
 45 productivity (Szuwalski & Hollowed, 2016). The Northern Shrimp (*Pandalus borealis*) in
 46 the Newfoundland and Labrador Shelves, which has undergone several periods of large-scale
 47 biomass change in the last two decades, despite a relatively constant harvest regime, is a prime
 48 example of a population thought to be affected by environmental conditions (DFO, 2019).
 49 These populations currently lack a population model to understand the drivers of this change
 50 and to predict how fishing pressure and changing environmental conditions may affect future
 51 abundance, which managers are advised to account for.

52 Population models like SPMs usually fall under the two following categories: process-based
 53 and statistical models. Process-based models often rely on differential or difference equations
 54 and are based on replicating the underlying processes (e.g., predation, recruitment, dispersal)
 55 driving population dynamics. Statistical models instead fit a regression model to time series
 56 of population abundances, abundance indices, or productivities, with some assumed error
 57 distribution for variation around predictions.

58 We have chosen a statistical approach to fitting SPMs. Statistical models allow for estimation
 59 of parameter uncertainty and ranges of model predictions and for flexibly incorporating potential
 60 ecosystem drivers into models (Plagányi et al., 2014). Statistical models also allow for straight-
 61 forward estimation of spatial variation in population parameters such as maximum productivity
 62 or density dependence from data, in the absence of theory predicting how these parameters
 63 should vary.

64 In this paper, we use a statistical approach to fitting SPMs using Generalized Additive Models
 65 (GAMS), estimated using the mgcv R package (Wood, 2017) as the backend. We apply this
 66 approach to the population of Northern Shrimp of the Newfoundland and Labrador Shelves,
 67 leveraging the smoothing properties of GAMs to account for varying productivity across time
 68 and space. The resulting model is a spatial SPM (SSPM), implemented via an R package:
 69 `sspm`.

70 The R package `sspm` is designed to make SSPMs simpler to estimate and apply to any spatially
 71 structured stock. The basic model this package implements was first used to model time-
 72 varying production in Newfoundland and Labrador Northern Shrimp stocks (E. J. Pedersen
 73 et al., 2022). However, the general modelling approach used here will work for any spatially
 74 structured fishery with sufficient data. It includes a range of features to manipulate harvest
 75 and biomass data. Those features are organized in a stepwise workflow, whose implementation
 76 is described in more detail in Figure 1 and in the next section.

77 Although it was developed in a fisheries context, the package is suited to model spatially-
 78 structured population dynamics in general.

79 Package design

80 The package follows an object oriented design, making use of the S4 class systems to model a
 81 stepwise workflow: (Figure 1).

82 The key workflow steps are:

- 83 ■ Discretization and aggregation of spatially structured observations into discrete patches,
 84 with a range of methods of discretization (random or custom sampling, Voronoi tessella-
 85 tion, or Delaunay triangulation).
 - 86 1. Provided boundary data in the form of a shapefile is converted into a `sspm_boundary`
 87 object using `sspm_as_boundary()` to define the boundary/region of interest.
 - 88 2. The region within the boundary is discretized into patches with the
 89 `sspm_discretize()` function, creating a `sspm_discrete_boundary` object.

- 90 ■ Spatiotemporal smoothing of biomass and environmental predictors using GAMs.
- 91 3. The `spm_as_dataset()` function turns user-provided data frames of raw observa-
- 92 tions into `sspm_dataset` objects that explicitly track locations, data types, and
- 93 aggregation scales for each input. `sspm` recognizes three types of data: **trawl** (i.e.,
- 94 biomass estimates from scientific surveys), **predictors**, and **catch** (i.e., harvest).
- 95 4. The `spm_smooth()` function uses GAMs to calculate spatially smoothed yearly
- 96 estimates of biomass and environmental predictors for each patch from trawl-level
- 97 data, based on the spatial structure from the `sspm_discrete_boundary` object. The
- 98 user specifies a GAM formula with custom smooth terms. The output is another
- 99 `sspm_dataset` object with a `smoothed_data` slot which contains the smoothed
- 100 predictions for all patches.
- 101 ■ Computation of surplus production based on biomass density and fishing effort.
- 102 5. The `spm_aggregate_catch()` function aggregates catch into patches and years and
- 103 calculates patch-specific productivity for each year as the ratio of estimated biomass
- 104 density plus catch from the next year divided by estimated biomass density of the
- 105 current year. The result is returned as a `sspm_dataset`.
- 106 6. The `sspm()` function combines productivity and predictor datasets into a single
- 107 dataset. Additionally, the user may create lagged versions of predictors with
- 108 `spm_lag()` and split data into testing and training sets for model validation with
- 109 `spm_split()` at this stage.
- 110 ■ Fitting of SSPMs to productivity estimates with GAMs.
- 111 7. The `spm()` function is used to fit a SSPM model to the output of step 6, using a
- 112 GAM model with custom syntax able to model a range of SSPMs. The output is
- 113 an `sspm` object.
- 114 ■ Visualization of results, and one-step-ahead projections of biomass for model validation
- 115 and scenario-based predictions.
- 116 8. Plots can be generated with the `plot()` method. Predictions from the fitted model
- 117 can be obtained using the built-in `predict()` method, including confidence and
- 118 prediction intervals




Figure 1: The sspm workflow. Gray cylinders represent raw, unprocessed sources of data. Each blue diamond shape represents a function processing a raw input and validating it, or producing an intermediate package object, represented as a green object. Secondary objects like formulas, which must be created by the user, are represented by a purple document shape. Finally, outputs are represented by a red document shape. The steps of the workflow as described above are denoted by dotted lines and corresponding step number.

Connections to other surplus-production based stock assessment approaches

The sspm package uses a model-based, random-effects based approach to estimate the effects of ecosystem drivers on surplus production across space and time. Our approach is conceptually related to the stochastic stock assessment approaches used by the spict (M. W. Pedersen & Berg, 2017) and jabba (Winker et al., 2018) R packages for surplus production modelling, in that we assume that biomass dynamics can be modelled as effectively a logistic growth model with both process and measurement error. While sspm does not currently have the capacity to model biomass dynamics as a continuous-time process, as with spict, or incorporate prior parameter information on catchability or biomass dynamics, as in jabba, sspm can model spatially and temporally varying productivity, which is currently not possible in these models.

The sspm package can be viewed as a spatiotemporal Model of Intermediate Complexity [a 'MICE-in-space' model; Thorson et al. (2019)] that can incorporate effects of other species

and ecosystem drivers as well as changes in fishing pressure on stock status. Our approach is closely connected to approaches used by other modern model-based spatial abundance estimation software, such as the VAST R package (Thorson et al., 2019) and the sdmTMB R package (Anderson et al., 2022). Our method shares the same approach as both VAST and sdmTMB of using spatially explicit models to estimate local biomass density (Figure 1 steps 1, 2, and 4), then aggregating up from those models to predict aggregate stock-level metrics such as total biomass and productivity (Figure 1 steps 8). The multiplicative surplus production model used by sspm is also conceptually similar to the vector-autoregressive model for biomass changes used by these two packages, as both VAST and sdmTMB can model local temporal changes as autoregressive processes on the link-scale of a generalized linear model. The sspm package cannot, however, model the dynamics of multiple species simultaneously; multi-species modelling would require generating a separate surplus production model (Figure 1 steps 5 and 6) for each species of interest.

One major difference between the sspm package and other model-based spatiotemporal modelling packages is its special-purpose nature. The default `spatial_smooth` function uses a computationally simpler (although somewhat less flexible) Intrinsic Conditional Autoregressive (ICAR) model (Rue & Held, 2005) for modelling spatial variation in covariates and biomass, as compared to the more complex spatial random effects possible with VAST and sdmTMB. This has the advantage of computational speed and less user knowledge of how to set up complex spatial grids, although it is less flexible. This means that sspm should be easier to adapt to novel fisheries than more complex packages that require more user modelling knowledge. Further, it is possible to specify alternative spatial smoothers than the ICAR model in the `spatial_smooth` function via the `bs=` argument, although this functionality has not been well-tested and should be considered experimental.

The other benefit of sspm, relative to other modelling packages, is the ability to model productivity rates directly (Figure 1 steps 5 and 6), rather than implicitly via an auto-regressive processes as used in VAST or sdmTMB. This means that it is possible in sspm to model nonlinear relationships between environmental covariates and productivity, or to easily include factors such as time-lagged effects of predictors on productivity in a given year. This approach does, however, sacrifice the ability to propagate measurement error into uncertainty about rates of change. One of the future directions for development of this package is to include variance propagation methods into the surplus production modelling step.

Acknowledgements

This research was supported by the Canadian Fisheries and Oceans Canada's (DFO) Sustainable fisheries Science Fund and by a Discovery Grant from the Canadian Natural Sciences and Engineering Research Council (NSERC) to E. J. Pedersen. We thank Fonya Irvine and John-Philip Williams for their help in testing the package and providing feedback on model implementation.

References

- Anderson, S. C., Ward, E. J., English, P. A., & Barnett, L. A. K. (2022). *sdmTMB: An R package for fast, flexible, and user-friendly generalized linear mixed effects models with spatial and spatiotemporal random fields* (p. 2022.03.24.485545). bioRxiv. <https://doi.org/10.1101/2022.03.24.485545>
- Berkes, F. (2012). Implementing ecosystem-based management: Evolution or revolution? *Fish and Fisheries*, 13(4), 465–476. <https://doi.org/10.1111/j.1467-2979.2011.00452.x>
- Crowder, L. B., Hazen, E. L., Avissar, N., Bjorkland, R., Latanich, C., & Ogburn, M. B. (2008). The impacts of fisheries on marine ecosystems and the transition to ecosystem-based

- management. *Annual Review of Ecology, Evolution, and Systematics*, 39(1), 259–278.
<https://doi.org/10.1146/annurev.ecolsys.39.110707.173406>
- DFO. (2019). *An assessment of Northern Shrimp (Pandalus borealis) in Shrimp Fishing Areas 4–6 and of Striped Shrimp (Pandalus montagui) in shrimp fishing area 4 in 2018* (Technical Report No. 2019/027; p. 23). Canadian Science Advisory Secretariat (CSAS).
- Gotelli, N. J. (2008). *A Primer of Ecology* (Fourth Edition). Oxford University Press. ISBN: 978-0-87893-318-1
- Karp, M. A., Peterson, J. O., Lynch, P. D., Griffis, R. B., Adams, C. F., Arnold, W. S., Barnett, L. A. K., deReynier, Y., DiCosimo, J., Fenske, K. H., Gaichas, S. K., Hollowed, A., Holsman, K., Karnauskas, M., Kobayashi, D., Leising, A., Manderson, J. P., McClure, M., Morrison, W. E., ... Link, J. S. (2019). Accounting for shifting distributions and changing productivity in the development of scientific advice for fishery management. *ICES Journal of Marine Science*, 76, 1305–1315. <https://doi.org/10.1093/icesjms/fsz048>
- Pedersen, E. J., Skanes, K., Le Corre, N., Koen-Alonso, M., & Baker, K. (2022). *A new spatial ecosystem-based surplus production model for SFA 4-6 Northern Shrimp* [Canadian Science Advisory Secretariat (CSAS) Research Document]. https://www.dfo-mpo.gc.ca/csas-sccs/Publications/ResDocs-DocRech/2022/2022_062-eng.html
- Pedersen, M. W., & Berg, C. W. (2017). A stochastic surplus production model in continuous time. *Fish and Fisheries*, 18(2), 226–243. <https://doi.org/10.1111/faf.12174>
- Plagányi, É. E., Punt, A. E., Hillary, R., Morello, E. B., Thébaud, O., Hutton, T., Pillans, R. D., Thorson, J. T., Fulton, E. A., Smith, A. D. M., Smith, F., Bayliss, P., Haywood, M., Lyne, V., & Rothlisberg, P. C. (2014). Multispecies fisheries management and conservation: Tactical applications using models of intermediate complexity. *Fish and Fisheries*, 15(1), 1–22. <https://doi.org/10.1111/j.1467-2979.2012.00488.x>
- Prager, M. H. (1994). A suite of extensions to a nonequilibrium surplus-production. *Fishery Bulletin*, 92, 374–389.
- Punt, A. E. (2003). Extending production models to include process error in the population dynamics. *Canadian Journal of Fisheries and Aquatic Sciences*, 60(10), 1217–1228. <https://doi.org/10.1139/f03-105>
- Rue, H., & Held, L. (2005). *Gaussian Markov random fields: Theory and applications*. CRC press.
- Szuwalski, C. S., & Hollowed, A. B. (2016). Climate change and non-stationary population processes in fisheries management. *ICES Journal of Marine Science*, 73(5), 1297–1305. <https://doi.org/10.1093/icesjms/fsv229>
- Tam, J. C., Link, J. S., Rossberg, A. G., Rogers, S. I., Levin, P. S., Rochet, M.-J., Bundy, A., Belgrano, A., Libralato, S., Tomczak, M., Wolfshaar, K. van de, Pranovi, F., Gorokhova, E., Large, S. I., Niquil, N., Greenstreet, S. P. R., Druon, J.-N., Lesutiene, J., Johansen, M., ... Rindorf, A. (2017). Towards ecosystem-based management: Identifying operational food-web indicators for marine ecosystems. *ICES Journal of Marine Science*, 74(7), 2040–2052. <https://doi.org/10.1093/icesjms/fsw230>
- Thorson, J. T., Adams, G., & Holsman, K. (2019). Spatio-temporal models of intermediate complexity for ecosystem assessments: A new tool for spatial fisheries management. *Fish and Fisheries*, 20(6), 1083–1099. <https://doi.org/10.1111/faf.12398>
- Thorson, J. T., Jannot, J., & Somers, K. (2017). Using spatio-temporal models of population growth and movement to monitor overlap between human impacts and fish populations. *Journal of Applied Ecology*, 54(2), 577–587. <https://doi.org/10.1111/1365-2664.12664>
- Thorson, J. T., Skaug, H. J., Kristensen, K., Shelton, A. O., Ward, E. J., Harms, J. H., & Benante, J. A. (2015). The importance of spatial models for estimating the strength of

- 227 density dependence. *Ecology*, 96(5), 1202–1212. <https://doi.org/10.1890/14-0739.1>
- 228 Walters, C. J., Hilborn, R., & Christensen, V. (2008). Surplus production dynamics in declining
229 and recovering fish populations. *Canadian Journal of Fisheries and Aquatic Sciences*,
230 65(11), 2536–2551. <https://doi.org/10.1139/F08-170>
- 231 Winker, H., Carvalho, F., & Kapur, M. (2018). JABBA: Just Another Bayesian Biomass
232 Assessment. *Fisheries Research*, 204, 275–288. [https://doi.org/10.1016/j.fishres.2018.03.](https://doi.org/10.1016/j.fishres.2018.03.010)
233 010
- 234 Wood, S. N. (2017). *Generalized Additive Models: An Introduction with R, 2nd Edition* (2nd
235 ed.). CRC Press. ISBN: 978-1-4987-2837-9
- 236 Zhang, F., Reid, K. B., & Nudds, T. D. (2021). The longer the better? Trade-offs in
237 fisheries stock assessment in dynamic ecosystems. *Fish and Fisheries*, 22(4), 789–797.
238 <https://doi.org/10.1111/faf.12550>

DRAFT