

Basic Notions — Spectral Sequences

Pedro Núñez

University of Freiburg

10th December 2020

Idea is easy to motivate



Notation gets ugly real quick



“You could have invented spectral sequences”

- ▶ This is the title of the expository article [[Cho06](#)].

“You could have invented spectral sequences”

- ▶ This is the title of the expository article [Cho06].
- ▶ In the spirit of [Cho06], we will go through an easy example in detail and hopefully convince ourselves by the end of the talk that the definition of a spectral sequence makes sense.

“You could have invented spectral sequences”

- ▶ This is the title of the expository article [Cho06].
- ▶ In the spirit of [Cho06], we will go through an easy example in detail and hopefully convince ourselves by the end of the talk that the definition of a spectral sequence makes sense.
- ▶ **Example:** let k be a field and consider a cochain complex C^\bullet of finite dimensional k -vector spaces

$$\cdots \rightarrow 0 \rightarrow C^0 \xrightarrow{d^0} C^1 \xrightarrow{d^1} C^2 \rightarrow 0 \rightarrow \cdots$$

“You could have invented spectral sequences”

- ▶ This is the title of the expository article [Cho06].
- ▶ In the spirit of [Cho06], we will go through an easy example in detail and hopefully convince ourselves by the end of the talk that the definition of a spectral sequence makes sense.
- ▶ **Example:** let k be a field and consider a cochain complex C^\bullet of finite dimensional k -vector spaces

$$\cdots \rightarrow 0 \rightarrow C^0 \xrightarrow{d^0} C^1 \xrightarrow{d^1} C^2 \rightarrow 0 \rightarrow \cdots$$

- ▶ **Goal:** compute the cohomology H^\bullet of the complex C^\bullet

$$H^\bullet := H^0(C^\bullet)[0] \oplus H^1(C^\bullet)[-1] \oplus H^2(C^\bullet)[-2].$$

Starting point: a filtration on our complex

- ▶ If we do not have any extra information about C^\bullet , there is not much more we can do besides applying the definition

$$H^\bullet = \ker(d^0)[0] \oplus \frac{\ker(d^1)}{\operatorname{im}(d^0)}[-1] \oplus \frac{C^2}{\operatorname{im}(d^1)}[-2].$$

Starting point: a filtration on our complex

- ▶ If we do not have any extra information about C^\bullet , there is not much more we can do besides applying the definition

$$H^\bullet = \ker(d^0)[0] \oplus \frac{\ker(d^1)}{\operatorname{im}(d^0)}[-1] \oplus \frac{C^2}{\operatorname{im}(d^1)}[-2].$$

- ▶ But C^\bullet has often some useful filtration $F^\bullet C^\bullet$, say

$$\begin{array}{ccccccc}
 C^\bullet = F^0 C^\bullet & \cdots 0 & \longrightarrow & C^0 & \xrightarrow{d^0} & C^1 & \xrightarrow{d^1} & C^2 & \longrightarrow & 0 \cdots \\
 \cup & & & \cup & & \cup & & \cup & & \\
 F^1 C^\bullet & \cdots 0 & \longrightarrow & F^1 C^0 & \longrightarrow & F^1 C^1 & \longrightarrow & F^1 C^2 & \longrightarrow & 0 \cdots \\
 \cup & & & \cup & & \cup & & \cup & & \\
 0 = F^2 C^\bullet & \cdots 0 & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & 0 \cdots
 \end{array}$$

How could a filtration on C^\bullet potentially help?

How could a filtration on C^\bullet potentially help?

1. $0 \subseteq F^1 C^\bullet \subseteq C^\bullet$ induces a filtration $0 \subseteq F^1 H^\bullet \subseteq H^\bullet$ given by

$$F^1 H^i := \operatorname{im} (H^i(F^1 C^\bullet) \rightarrow H^i(C^\bullet)) \text{ for all } i \in \{0, 1, 2\}.$$

How could a filtration on C^\bullet potentially help?

1. $0 \subseteq F^1 C^\bullet \subseteq C^\bullet$ induces a filtration $0 \subseteq F^1 H^\bullet \subseteq H^\bullet$ given by

$$F^1 H^i := \operatorname{im} (H^i(F^1 C^\bullet) \rightarrow H^i(C^\bullet)) \text{ for all } i \in \{0, 1, 2\}.$$

2. We can then recover H^i as the associated graded vector space

$$\operatorname{gr}(F^\bullet H^i) = F^1 H^i \oplus (H^i / F^1 H^i) \cong H^i \text{ for all } i \in \{0, 1, 2\}.$$

How could a filtration on C^\bullet potentially help?

1. $0 \subseteq F^1 C^\bullet \subseteq C^\bullet$ induces a filtration $0 \subseteq F^1 H^\bullet \subseteq H^\bullet$ given by

$$F^1 H^i := \operatorname{im} (H^i(F^1 C^\bullet) \rightarrow H^i(C^\bullet)) \text{ for all } i \in \{0, 1, 2\}.$$

2. We can then recover H^i as the associated graded vector space

$$\operatorname{gr}(F^\bullet H^i) = F^1 H^i \oplus (H^i / F^1 H^i) \cong H^i \text{ for all } i \in \{0, 1, 2\}.$$

3. $F^\bullet C^\bullet$ being a filtered complex means that $F^1 C^\bullet \subseteq C^\bullet$ is a subcomplex, so that $F^1 C^\bullet$ and $C^\bullet / F^1 C^\bullet$ are complexes as well.

How could a filtration on C^\bullet potentially help?

1. $0 \subseteq F^1 C^\bullet \subseteq C^\bullet$ induces a filtration $0 \subseteq F^1 H^\bullet \subseteq H^\bullet$ given by

$$F^1 H^i := \operatorname{im} (H^i(F^1 C^\bullet) \rightarrow H^i(C^\bullet)) \text{ for all } i \in \{0, 1, 2\}.$$

2. We can then recover H^i as the associated graded vector space

$$\operatorname{gr}(F^\bullet H^i) = F^1 H^i \oplus (H^i / F^1 H^i) \cong H^i \text{ for all } i \in \{0, 1, 2\}.$$

3. $F^\bullet C^\bullet$ being a filtered complex means that $F^1 C^\bullet \subseteq C^\bullet$ is a subcomplex, so that $F^1 C^\bullet$ and $C^\bullet / F^1 C^\bullet$ are complexes as well.

- **Q1:** Can we compute the cohomologies of $F^1 C^\bullet$ and $C^\bullet / F^1 C^\bullet$?

How could a filtration on C^\bullet potentially help?

1. $0 \subseteq F^1 C^\bullet \subseteq C^\bullet$ induces a filtration $0 \subseteq F^1 H^\bullet \subseteq H^\bullet$ given by

$$F^1 H^i := \operatorname{im} (H^i(F^1 C^\bullet) \rightarrow H^i(C^\bullet)) \text{ for all } i \in \{0, 1, 2\}.$$

2. We can then recover H^i as the associated graded vector space

$$\operatorname{gr}(F^\bullet H^i) = F^1 H^i \oplus (H^i / F^1 H^i) \cong H^i \text{ for all } i \in \{0, 1, 2\}.$$

3. $F^\bullet C^\bullet$ being a filtered complex means that $F^1 C^\bullet \subseteq C^\bullet$ is a subcomplex, so that $F^1 C^\bullet$ and $C^\bullet / F^1 C^\bullet$ are complexes as well.

- **Q1:** Can we compute the cohomologies of $F^1 C^\bullet$ and $C^\bullet / F^1 C^\bullet$?
- **Q2:** Are these cohomologies related to $F^1 H^\bullet$ and $H^\bullet / F^1 H^\bullet$?

Possible scenario I: the good

- ▶ Suppose $C^\bullet = C_1^\bullet \oplus C_2^\bullet$ with $H^\bullet(C_1^\bullet)$ and $H^\bullet(C_2^\bullet)$ computable.

Possible scenario I: the good

- ▶ Suppose $C^\bullet = C_1^\bullet \oplus C_2^\bullet$ with $H^\bullet(C_1^\bullet)$ and $H^\bullet(C_2^\bullet)$ computable.
- ▶ Then we are done, because $H^\bullet(C^\bullet) = H^\bullet(C_1^\bullet) \oplus H^\bullet(C_2^\bullet)$.

Possible scenario I: the good

- ▶ Suppose $C^\bullet = C_1^\bullet \oplus C_2^\bullet$ with $H^\bullet(C_1^\bullet)$ and $H^\bullet(C_2^\bullet)$ computable.
- ▶ Then we are done, because $H^\bullet(C^\bullet) = H^\bullet(C_1^\bullet) \oplus H^\bullet(C_2^\bullet)$.
- ▶ To rephrase this with filtrations we take

$$F^1 C^\bullet := C_1^\bullet \quad (\Rightarrow \quad C^\bullet / F^1 C^\bullet = C_2^\bullet.)$$

Possible scenario I: the good

- ▶ Suppose $C^\bullet = C_1^\bullet \oplus C_2^\bullet$ with $H^\bullet(C_1^\bullet)$ and $H^\bullet(C_2^\bullet)$ computable.
- ▶ Then we are done, because $H^\bullet(C^\bullet) = H^\bullet(C_1^\bullet) \oplus H^\bullet(C_2^\bullet)$.
- ▶ To rephrase this with filtrations we take

$$F^1 C^\bullet := C_1^\bullet \quad (\Rightarrow \quad C^\bullet / F^1 C^\bullet = C_2^\bullet.)$$

- ▶ Then we have

$$F^1 H^\bullet := \text{im}(H^\bullet(F^1 C^\bullet) \rightarrow H^\bullet(C^\bullet)) = H^\bullet(C_1^\bullet) = H^\bullet(F^1 C^\bullet)$$

and similarly $H^\bullet / F^1 H^\bullet = H^\bullet(C_2^\bullet) = H^\bullet(C^\bullet / F^1 C^\bullet)$.

Possible scenario I: the good

- ▶ Suppose $C^\bullet = C_1^\bullet \oplus C_2^\bullet$ with $H^\bullet(C_1^\bullet)$ and $H^\bullet(C_2^\bullet)$ computable.
- ▶ Then we are done, because $H^\bullet(C^\bullet) = H^\bullet(C_1^\bullet) \oplus H^\bullet(C_2^\bullet)$.
- ▶ To rephrase this with filtrations we take


$$F^1 C^\bullet := C_1^\bullet \quad (\Rightarrow \quad C^\bullet / F^1 C^\bullet = C_2^\bullet.)$$

- ▶ Then we have

$$F^1 H^\bullet := \text{im}(H^\bullet(F^1 C^\bullet) \rightarrow H^\bullet(C^\bullet)) = H^\bullet(C_1^\bullet) = H^\bullet(F^1 C^\bullet)$$

and similarly $H^\bullet / F^1 H^\bullet = H^\bullet(C_2^\bullet) = H^\bullet(C^\bullet / F^1 C^\bullet)$.

So in this case we can carry out our naive strategy of computing each graded piece of the induced filtration in cohomology separately:



Possible scenario II: the bad

Suppose we know nothing about C^\bullet . The *stupid filtration*

$$\begin{cases} F^1 C^\bullet := \cdots \rightarrow 0 \rightarrow 0 \rightarrow C^1 \xrightarrow{d^1} C^2 \rightarrow 0 \rightarrow \cdots \\ F^2 C^\bullet := \cdots \rightarrow 0 \rightarrow 0 \rightarrow 0 \rightarrow C^2 \rightarrow 0 \rightarrow \cdots \end{cases}$$

induces the cohomology filtration

$$\begin{cases} F^1 H^\bullet := \operatorname{im}(H^\bullet(F^1 C^\bullet) \rightarrow H^\bullet(C^\bullet)) = H^1(C^\bullet)[-1] \oplus H^2(C^\bullet)[-2] \\ F^2 H^\bullet := \operatorname{im}(H^\bullet(F^2 C^\bullet) \rightarrow H^\bullet(C^\bullet)) = H^2(C^\bullet)[-2] \end{cases}$$

whose graded pieces are $H^0(C^\bullet)[0]$, $H^1(C^\bullet)[-1]$ and $H^2(C^\bullet)[-2]$.
So in this case computing $\operatorname{gr}^i H^\bullet(C^\bullet)$ is as hard as computing $H^i(C^\bullet)$ and we just wasted our time with this detour.

Possible scenario III: the ugly

Suppose now that we are somewhere in between the good and the bad scenarios. Namely, assume that we have some filtration $0 \subsetneq F^1 C^\bullet \subsetneq C^\bullet$ which does not correspond to a direct sum decomposition but such that we can compute the graded pieces $\operatorname{gr}^i(H^\bullet(C^\bullet))$ of the filtration induced in cohomology.

References



Timothy Y. Chow.

You could have invented spectral sequences.

Notices Amer. Math. Soc., 53(1):15–19, 2006.

4