Basic Notions — Spectral Sequences

Pedro Núñez

University of Freiburg

10th December 2020

"You could have invented spectral sequences"¹



Notation gets ugly very soon



▶ Only finite dimensional vector spaces over ℚ for simplicity.

- ▶ Only finite dimensional vector spaces over ℚ for simplicity.
- ▶ (Cochain) complexes denoted with capital letters C and not C•.

- ▶ Only finite dimensional vector spaces over ℚ for simplicity.
- ightharpoonup (Cochain) complexes denoted with capital letters C and not C^{\bullet} .
- ▶ Cohomology again denoted just H(C) and not $H^{\bullet}(C)$.

- ▶ Only finite dimensional vector spaces over ℚ for simplicity.
- ightharpoonup (Cochain) complexes denoted with capital letters C and not C^{\bullet} .
- ▶ Cohomology again denoted just H(C) and not $H^{\bullet}(C)$.
- ► Cohomology is regarded as a copmlex with 0 differentials

$$H(C) = \bigoplus_{i \in \mathbb{Z}} H^i(C)[-i] := \left(\cdots H^{-1}(C) \xrightarrow{0} H^0(C) \xrightarrow{0} H^1(C) \cdots\right).$$

- ▶ Only finite dimensional vector spaces over ℚ for simplicity.
- ightharpoonup (Cochain) complexes denoted with capital letters C and not C^{\bullet} .
- ▶ Cohomology again denoted just H(C) and not $H^{\bullet}(C)$.
- ► Cohomology is regarded as a copmlex with 0 differentials

$$H(C) = \bigoplus_{i \in \mathbb{Z}} H^i(C)[-i] := \left(\cdots H^{-1}(C) \xrightarrow{0} H^0(C) \xrightarrow{0} H^1(C) \cdots \right).$$

▶ If it is clear to which C we refer, denote H(C) simply by H.

- Only finite dimensional vector spaces over Q for simplicity.
- ightharpoonup (Cochain) complexes denoted with capital letters C and not C^{\bullet} .
- ▶ Cohomology again denoted just H(C) and not $H^{\bullet}(C)$.
- ► Cohomology is regarded as a copmlex with 0 differentials

$$H(C) = \bigoplus_{i \in \mathbb{Z}} H^{i}(C)[-i] := \left(\cdots H^{-1}(C) \xrightarrow{0} H^{0}(C) \xrightarrow{0} H^{1}(C) \cdots\right).$$

- If it is clear to which C we refer, denote H(C) simply by H.
- (Decreasing) filtrations are always finite for simplicity:

$$F^{\bullet}A: \quad 0 = F^nA \subseteq \ldots \subseteq F^0A = A.$$



- ▶ Only finite dimensional vector spaces over ℚ for simplicity.
- ightharpoonup (Cochain) complexes denoted with capital letters C and not C^{\bullet} .
- ▶ Cohomology again denoted just H(C) and not $H^{\bullet}(C)$.
- ► Cohomology is regarded as a copmlex with 0 differentials

$$H(C) = \bigoplus_{i \in \mathbb{Z}} H^{i}(C)[-i] := \left(\cdots H^{-1}(C) \xrightarrow{0} H^{0}(C) \xrightarrow{0} H^{1}(C) \cdots\right).$$

- If it is clear to which C we refer, denote H(C) simply by H.
- (Decreasing) filtrations are always finite for simplicity:


$$F^{\bullet}A: \quad 0 = F^nA \subseteq \ldots \subseteq F^0A = A.$$

▶ The graded object associated to a filtration $F^{\bullet}A$ is denoted

$$\operatorname{gr} A = \bigoplus_{p \in \mathbb{N}} \operatorname{gr}^p A := \bigoplus_{p \in \mathbb{N}} (F^p A / F^{p+1} A).$$



GOAL: compute cohomology with the help of a filtration



Filtration induced in cohomology

A filtration on the complx $F^{\bullet}C$ induces a filtration in cohomology

$$F^pH := \operatorname{im}(H(F^pC) \to H(C)) \subseteq H(C) = H,$$

where $H(F^pC) \to H(C)$ is induced by the inclusion map $F^pC \hookrightarrow C$.

BABY EXAMPLE: 2 step filtration on a length 1 complex

BABY EXAMPLE: filtration induced in cohomology

We first look at the map induced by $F^1C \hookrightarrow C$ in cohomology:

$$H(F^{1}C) = \ker \left(d^{0}|_{F^{1}C}\right) \left[0\right] \oplus \left(\frac{F^{1}C^{1}}{\operatorname{im}\left(d^{0}|_{F^{1}C}\right)}\right) \left[-1\right]$$

$$\downarrow \qquad \qquad \downarrow$$

$$H(C) = \ker \left(d^{0}\right) \left[0\right] \oplus \left(\frac{C^{1}}{\operatorname{im}\left(d^{0}\right)}\right) \left[-1\right]$$

BABY EXAMPLE: filtration induced in cohomology

We first look at the map induced by $F^1C \hookrightarrow C$ in cohomology:

$$\begin{array}{ll} H(F^1C) & = & \ker\left(d^0|_{F^1C}\right)\left[0\right] \oplus \left(\frac{F^1C^1}{\operatorname{im}\left(d^0|_{F^1C}\right)}\right)\left[-1\right] \\ & \downarrow & & \downarrow \\ H(C) & = & \ker(d^0)\left[0\right] \oplus \left(\frac{C^1}{\operatorname{im}\left(d^0\right)}\right)\left[-1\right] \end{array}$$

The filtration on cohomology was given by its image, hence

$$F^{1}H = \ker \left(d^{0}|_{F^{1}C}\right)[0] \oplus \left(\frac{F^{1}C^{1} + \operatorname{im}(d^{0})}{\operatorname{im}(d^{0})}\right)[-1]$$

BABY EXAMPLE: graded cohomology pieces

Recall that $\operatorname{gr}^0 H := F^0 H / F^1 H = H / F^1 H$, hence (after \cong -theorem)

$$\operatorname{gr}^{0} H = \left(\frac{\ker(d^{0})}{\ker(d^{0}|_{F^{1}C})}\right) [0] \oplus \left(\frac{C^{1}}{F^{1}C^{1} + \operatorname{im}(d^{0})}\right) [-1]$$

BABY EXAMPLE: graded cohomology pieces


Recall that $\operatorname{gr}^0 H := F^0 H / F^1 H = H / F^1 H$, hence (after \cong -theorem)

$$\operatorname{gr}^0 H = \left(\frac{\ker(d^0)}{\ker(d^0|_{F^1C})}\right) [0] \oplus \left(\frac{C^1}{F^1C^1 + \operatorname{im}(d^0)}\right) [-1]$$

Similarly, $\operatorname{gr}^1 H = F^1 H / 0 = F^1 H$, thus

$$\operatorname{gr}^{1} H = \ker \left(d^{0}|_{F^{1}C} \right) [0] \oplus \left(\frac{F^{1}C^{1} + \operatorname{im}(d^{0})}{\operatorname{im}(d^{0})} \right) [-1]$$

BABY EXAMPLE: define $E_0^{p,q} := (\operatorname{gr}^p C)^{p+q}$ and visualize in \mathbb{Z}^2



References



Timothy Y. Chow.

You could have invented spectral sequences.

Notices Amer. Math. Soc., 53(1):15–19, 2006.

-



The Stacks project authors.

The stacks project.

https://stacks.math.columbia.edu, 2020.

(document)