

Basic Notions — Spectral Sequences

Pedro Núñez

University of Freiburg

10th December 2020

“You could have invented spectral sequences”¹



¹Title of the expository article [Cho06]

Notation gets ugly very soon



Notation and conventions [Sta20]

- ▶ Only finite dimensional vector spaces over \mathbb{Q} for simplicity.

Notation and conventions [Sta20]

- ▶ Only finite dimensional vector spaces over \mathbb{Q} for simplicity.
- ▶ (Cochain) complexes denoted with capital letters C and not C^\bullet .

Notation and conventions [Sta20]

- ▶ Only finite dimensional vector spaces over \mathbb{Q} for simplicity.
- ▶ (Cochain) complexes denoted with capital letters C and not C^\bullet .
- ▶ Cohomology again denoted just $H(C)$ and not $H^\bullet(C)$.

Notation and conventions [Sta20]

- ▶ Only finite dimensional vector spaces over \mathbb{Q} for simplicity.
- ▶ (Cochain) complexes denoted with capital letters C and not C^\bullet .
- ▶ Cohomology again denoted just $H(C)$ and not $H^\bullet(C)$.
- ▶ Cohomology is regarded as a complex with 0 differentials

$$H(C) = \bigoplus_{i \in \mathbb{Z}} H^i(C)[-i] := \left(\cdots H^{-1}(C) \xrightarrow{0} H^0(C) \xrightarrow{0} H^1(C) \cdots \right).$$

Notation and conventions [Sta20]

- ▶ Only finite dimensional vector spaces over \mathbb{Q} for simplicity.
- ▶ (Cochain) complexes denoted with capital letters C and not C^\bullet .
- ▶ Cohomology again denoted just $H(C)$ and not $H^\bullet(C)$.
- ▶ Cohomology is regarded as a complex with 0 differentials

$$H(C) = \bigoplus_{i \in \mathbb{Z}} H^i(C)[-i] := \left(\cdots H^{-1}(C) \xrightarrow{0} H^0(C) \xrightarrow{0} H^1(C) \cdots \right).$$

- ▶ If it is clear to which C we refer, denote $H(C)$ simply by H .

Notation and conventions [Sta20]

- ▶ Only finite dimensional vector spaces over \mathbb{Q} for simplicity.
- ▶ (Cochain) complexes denoted with capital letters C and not C^\bullet .
- ▶ Cohomology again denoted just $H(C)$ and not $H^\bullet(C)$.
- ▶ Cohomology is regarded as a complex with 0 differentials

$$H(C) = \bigoplus_{i \in \mathbb{Z}} H^i(C)[-i] := \left(\cdots H^{-1}(C) \xrightarrow{0} H^0(C) \xrightarrow{0} H^1(C) \cdots \right).$$

- ▶ If it is clear to which C we refer, denote $H(C)$ simply by H .
- ▶ (Decreasing) filtrations are always finite for simplicity:

$$F^\bullet A: \quad 0 = F^n A \subseteq \cdots \subseteq F^0 A = A.$$

Notation and conventions [Sta20]

- ▶ Only finite dimensional vector spaces over \mathbb{Q} for simplicity.
- ▶ (Cochain) complexes denoted with capital letters C and not C^\bullet .
- ▶ Cohomology again denoted just $H(C)$ and not $H^\bullet(C)$.
- ▶ Cohomology is regarded as a complex with 0 differentials

$$H(C) = \bigoplus_{i \in \mathbb{Z}} H^i(C)[-i] := \left(\cdots H^{-1}(C) \xrightarrow{0} H^0(C) \xrightarrow{0} H^1(C) \cdots \right).$$


- ▶ If it is clear to which C we refer, denote $H(C)$ simply by H .
- ▶ (Decreasing) filtrations are always finite for simplicity:

$$F^\bullet A: \quad 0 = F^n A \subseteq \cdots \subseteq F^0 A = A.$$

- ▶ The graded object associated to a filtration $F^\bullet A$ is denoted

$$\mathrm{gr} A = \bigoplus_{p \in \mathbb{N}} \mathrm{gr}^p A := \bigoplus_{p \in \mathbb{N}} (F^p A / F^{p+1} A).$$

GOAL: compute cohomology with the help of a filtration



Filtration induced in cohomology

A filtration on the complex $F^\bullet C$ induces a filtration in cohomology

$$F^p H := \operatorname{im}(H(F^p C) \rightarrow H(C)) \subseteq H(C) = H,$$

where $H(F^p C) \rightarrow H(C)$ is induced by the inclusion map $F^p C \hookrightarrow C$.

BABY EXAMPLE: 2 step filtration on a length 1 complex

$$\begin{array}{ccccccc}
 C & & & & & & \\
 \parallel & & & & & & \\
 F^0 C & \cdots \rightarrow 0 \rightarrow C^0 \xrightarrow{d^0} C^1 \rightarrow 0 \rightarrow \cdots \\
 \cup \mid & & \cup \mid & & \cup \mid & & \\
 F^1 C & \cdots \rightarrow 0 \rightarrow F^1 C^0 \xrightarrow{d^0|_{F^1 C^0}} F^1 C^1 \rightarrow 0 \rightarrow \cdots \\
 \cup \mid & & \cup \mid & & \cup \mid & & \\
 F^2 C & \cdots \rightarrow 0 \rightarrow 0 \longrightarrow 0 \longrightarrow 0 \rightarrow \cdots \\
 \parallel & & & & & & \\
 0 & & & & & &
 \end{array}$$

BABY EXAMPLE: filtration induced in cohomology

We first look at the map induced by $F^1 C \hookrightarrow C$ in cohomology:

$$\begin{array}{ccc} H(F^1 C) & = & \ker(d^0|_{F^1 C}) [0] \oplus \left(\frac{F^1 C^1}{\operatorname{im}(d^0|_{F^1 C})} \right) [-1] \\ \downarrow & & \downarrow \\ H(C) & = & \ker(d^0) [0] \oplus \left(\frac{C^1}{\operatorname{im}(d^0)} \right) [-1] \end{array}$$

BABY EXAMPLE: filtration induced in cohomology

We first look at the map induced by $F^1 C \hookrightarrow C$ in cohomology:

$$\begin{array}{ccc} H(F^1 C) & = & \ker(d^0|_{F^1 C}) [0] \oplus \left(\frac{F^1 C^1}{\operatorname{im}(d^0|_{F^1 C})} \right) [-1] \\ \downarrow & & \downarrow \\ H(C) & = & \ker(d^0) [0] \oplus \left(\frac{C^1}{\operatorname{im}(d^0)} \right) [-1] \end{array}$$

The filtration on cohomology was given by its image, hence

$$F^1 H = \ker(d^0|_{F^1 C}) [0] \oplus \left(\frac{F^1 C^1 + \operatorname{im}(d^0)}{\operatorname{im}(d^0)} \right) [-1]$$

BABY EXAMPLE: graded cohomology pieces

Recall that $\mathrm{gr}^0 H := F^0 H / F^1 H = H / F^1 H$, hence (after \cong -theorem)

$$\mathrm{gr}^0 H = \left(\frac{\ker(d^0)}{\ker(d^0|_{F^1 C})} \right) [0] \oplus \left(\frac{C^1}{F^1 C^1 + \mathrm{im}(d^0)} \right) [-1]$$

BABY EXAMPLE: graded cohomology pieces


Recall that $\mathrm{gr}^0 H := F^0 H / F^1 H = H / F^1 H$, hence (after \cong -theorem)

$$\mathrm{gr}^0 H = \left(\frac{\ker(d^0)}{\ker(d^0|_{F^1 C})} \right) [0] \oplus \left(\frac{C^1}{F^1 C^1 + \mathrm{im}(d^0)} \right) [-1]$$


Similarly, $\mathrm{gr}^1 H = F^1 H / 0 = F^1 H$, thus

$$\mathrm{gr}^1 H = \ker(d^0|_{F^1 C}) [0] \oplus \left(\frac{F^1 C^1 + \mathrm{im}(d^0)}{\mathrm{im}(d^0)} \right) [-1]$$

BABY EXAMPLE: define $E_0^{p,q} := (\text{gr}^p C)^{p+q}$ and visualize in \mathbb{Z}^2




BABY EXAMPLE: define $E_0^{p,q} := (\operatorname{gr}^p C)^{p+q}$ and visualize in \mathbb{Z}^2




- Since $F^1C \subseteq C$ is a subcomplex, $d^0: C^0 \rightarrow C^1$ induces the differential $d_0^{0,0}$.

BABY EXAMPLE: define $E_0^{p,q} := (\operatorname{gr}^p C)^{p+q}$ and visualize in \mathbb{Z}^2



- ▶ Since $F^1 C \subseteq C$ is a subcomplex, $d^0: C^0 \rightarrow C^1$ induces the differential $d_0^{0,0}$.
- ▶ For the same reason, $d^0|_{F^1 C}$ has image in $F^1 C^1$, so it induces the differential $d_0^{1,-1}$.

BABY EXAMPLE: define $E_0^{p,q} := (\operatorname{gr}^p C)^{p+q}$ and visualize in \mathbb{Z}^2



- ▶ Since $F^1 C \subseteq C$ is a subcomplex, $d^0: C^0 \rightarrow C^1$ induces the differential $d_0^{0,0}$.
- ▶ For the same reason, $d^0|_{F^1 C}$ has image in $F^1 C^1$, so it induces the differential $d_0^{1,-1}$.
- ▶ In general, since $F^{p+1} C \subseteq F^p C$ is a subcomplex, the original differentials induce the vertical differentials.

BABY EXAMPLE: compute cohomology of the columns $\mathrm{gr}^p C$

From the $p = 0$ column in the “ E_0 -page” we compute

$$H(\mathrm{gr}^0 C) = \left(\frac{\ker(d^0) + (d^0)^{-1}(F^1 C^1)}{F^1 C^0} \right) [0] \oplus \left(\frac{C^1}{F^1 C^1 + \mathrm{im}(d^0)} \right) [-1]$$

BABY EXAMPLE: compute cohomology of the columns $\text{gr}^p C$

From the $p = 0$ column in the “ E_0 -page” we compute

$$H(\text{gr}^0 C) = \left(\frac{\ker(d^0) + (d^0)^{-1}(F^1 C^1)}{F^1 C^0} \right) [0] \oplus \left(\frac{C^1}{F^1 C^1 + \text{im}(d^0)} \right) [-1]$$

And from the $p = 1$ column in the “ E_0 -page” we compute

$$H(\text{gr}^1 C) = \ker(d^0|_{F^1 C}) [0] \oplus \left(\frac{F^1 C^1}{\text{im}(d^0|_{F^1 C})} \right) [-1]$$

BABY EXAMPLE: 1st approximation to the $\text{gr}^0 H$ part

We compare $\text{gr}^0 H$ (left column) to $H(\text{gr}^0 C)$ (right column):

$$\left(\frac{\ker(d^0)}{\ker(d^0|_{F^1 C})} \right) [0]$$

$$\oplus$$

$$\left(\frac{C^1}{F^1 C^1 + \text{im}(d^0)} \right) [-1]$$

$$\left(\frac{\ker(d^0) + (d^0)^{-1}(F^1 C^1)}{F^1 C^0} \right) [0]$$

$$\oplus$$

$$\left(\frac{C^1}{F^1 C^1 + \text{im}(d^0)} \right) [-1]$$

BABY EXAMPLE: 1st approximation to the $\mathrm{gr}^0 H$ part

We compare $\mathrm{gr}^0 H$ (left column) to $H(\mathrm{gr}^0 C)$ (right column):

$$\left(\frac{\ker(d^0)}{\ker(d^0|_{F^1 C})} \right) [0]$$

$$\oplus$$

$$\left(\frac{C^1}{F^1 C^1 + \mathrm{im}(d^0)} \right) [-1]$$

$$\left(\frac{\ker(d^0) + (d^0)^{-1}(F^1 C^1)}{F^1 C^0} \right) [0]$$

$$\oplus$$

$$\left(\frac{C^1}{F^1 C^1 + \mathrm{im}(d^0)} \right) [-1]$$

- $H^1(\mathrm{gr}^0 C)$ does compute $\mathrm{gr}^0 H^1$ (bottom row),

BABY EXAMPLE: 1st approximation to the $\text{gr}^0 H$ part

We compare $\text{gr}^0 H$ (left column) to $H(\text{gr}^0 C)$ (right column):

$$\left(\frac{\ker(d^0)}{\ker(d^0|_{F^1 C})} \right) [0]$$

$$\oplus$$

$$\left(\frac{C^1}{F^1 C^1 + \text{im}(d^0)} \right) [-1]$$

$$\left(\frac{\ker(d^0) + (d^0)^{-1}(F^1 C^1)}{F^1 C^0} \right) [0]$$

$$\oplus$$

$$\left(\frac{C^1}{F^1 C^1 + \text{im}(d^0)} \right) [-1]$$

- ▶ $H^1(\text{gr}^0 C)$ does compute $\text{gr}^0 H^1$ (bottom row),
- ▶ but $H^0(\text{gr}^0 C)$ is not isomorphic to $\text{gr}^0 H^0$ (top row).

BABY EXAMPLE: 1st approximation to the $\text{gr}^1 H$ part

We compare now $\text{gr}^1 H$ (left column) to $H(\text{gr}^1 C)$ (right column):

$$\ker (d^0|_{F^1 C}) [0]$$

$$\oplus$$

$$\left(\frac{F^1 C^1 + \text{im}(d^0)}{\text{im}(d^0)} \right) [-1]$$

$$\ker (d^0|_{F^1 C}) [0]$$

$$\oplus$$

$$\left(\frac{F^1 C^1}{\text{im}(d^0|_{F^1 C})} \right) [-1]$$

BABY EXAMPLE: 1st approximation to the $\mathrm{gr}^1 H$ part

We compare now $\mathrm{gr}^1 H$ (left column) to $H(\mathrm{gr}^1 C)$ (right column):

$$\ker(d^0|_{F^1 C}) [0]$$

$$\oplus$$

$$\left(\frac{F^1 C^1 + \mathrm{im}(d^0)}{\mathrm{im}(d^0)} \right) [-1]$$

$$\ker(d^0|_{F^1 C}) [0]$$

$$\oplus$$

$$\left(\frac{F^1 C^1}{\mathrm{im}(d^0|_{F^1 C})} \right) [-1]$$

- In this case $H^0(\mathrm{gr}^1 C)$ does compute $\mathrm{gr}^1 H^0$ (top row),

BABY EXAMPLE: 1st approximation to the $\mathrm{gr}^1 H$ part

We compare now $\mathrm{gr}^1 H$ (left column) to $H(\mathrm{gr}^1 C)$ (right column):

$$\ker(d^0|_{F^1 C}) [0]$$

$$\oplus$$

$$\left(\frac{F^1 C^1 + \mathrm{im}(d^0)}{\mathrm{im}(d^0)} \right) [-1]$$


$$\ker(d^0|_{F^1 C}) [0]$$

$$\oplus$$


$$\left(\frac{F^1 C^1}{\mathrm{im}(d^0|_{F^1 C})} \right) [-1]$$

- ▶ In this case $H^0(\mathrm{gr}^1 C)$ does compute $\mathrm{gr}^1 H^0$ (top row),
- ▶ but $H^1(\mathrm{gr}^1 C)$ is not isomorphic to $\mathrm{gr}^1 H^1$ (bottom row).

BABY EXAMPLE: define $E_1^{p,q} := H^{p+q}(\text{gr}^p C)$ and plot in \mathbb{Z}^2




BABY EXAMPLE: define $E_1^{p,q} := H^{p+q}(\text{gr}^p C)$ and plot in \mathbb{Z}^2



- The original differential d^0 induces the differential $d_1^{0,0}$.

BABY EXAMPLE: define $E_1^{p,q} := H^{p+q}(\text{gr}^p C)$ and plot in \mathbb{Z}^2



- ▶ The original differential d^0 induces the differential $d_1^{0,0}$.
- ▶ More generally, if we had started from a longer complex, the original differentials would induce the horizontal differentials.

BABY EXAMPLE: 2nd approximation to $\mathrm{gr} H$

- ▶ We obtained the 1st approximation by taking cohomologies on the “ E_0 -page”.

BABY EXAMPLE: 2nd approximation to $\operatorname{gr} H$

- ▶ We obtained the 1st approximation by taking cohomologies on the “ E_0 -page”.
- ▶ These 1st approximations were then organized into the “ E_1 -page” in such a way that $\operatorname{gr}^p H^n$ would correspond to $E_1^{p, n-p}$.

BABY EXAMPLE: 2^{nd} approximation to $\text{gr } H$

- ▶ We obtained the 1^{st} approximation by taking cohomologies on the “ E_0 -page”.
- ▶ These 1^{st} approximations were then organized into the “ E_1 -page” in such a way that $\text{gr}^p H^n$ would correspond to $E_1^{p,n-p}$.
- ▶ For example, we have seen that
 - ▶ $\text{gr}^0 H^1 = E_1^{0,1}$, although $\text{gr}^0 H^0 \neq E_1^{0,0}$;

BABY EXAMPLE: 2nd approximation to $\mathrm{gr} H$

- ▶ We obtained the 1st approximation by taking cohomologies on the “ E_0 -page”.
- ▶ These 1st approximations were then organized into the “ E_1 -page” in such a way that $\mathrm{gr}^p H^n$ would correspond to $E_1^{p,n-p}$.
- ▶ For example, we have seen that
 - ▶ $\mathrm{gr}^0 H^1 = E_1^{0,1}$, although $\mathrm{gr}^0 H^0 \neq E_1^{0,0}$;
 - ▶ $\mathrm{gr}^1 H^0 = E_1^{1,-1}$, although $\mathrm{gr}^1 H^1 \neq E_1^{1,0}$.

BABY EXAMPLE: 2nd approximation to $\text{gr } H$

- ▶ We obtained the 1st approximation by taking cohomologies on the “ E_0 -page”.
- ▶ These 1st approximations were then organized into the “ E_1 -page” in such a way that $\text{gr}^p H^n$ would correspond to $E_1^{p,n-p}$.
- ▶ For example, we have seen that
 - ▶ $\text{gr}^0 H^1 = E_1^{0,1}$, although $\text{gr}^0 H^0 \neq E_1^{0,0}$;
 - ▶ $\text{gr}^1 H^0 = E_1^{1,-1}$, although $\text{gr}^1 H^1 \neq E_1^{1,0}$.
- ▶ Continuing with this way to arrange things, we define the “ E_2 -page” by taking cohomologies at each point of the “ E_1 -page”, that is

$$E_2^{p,q} := H^p(E_1^{\bullet,q}),$$

so that $\text{gr}^p H^n$ would again correspond to $E_2^{p,n-p}$.

BABY EXAMPLE: 2^{nd} approximation to the $\text{gr}^0 H$ part

We compare then $\text{gr}^0 H$ (left column) to $E_2^{0,\bullet}$ (right column):

$$\left(\frac{\ker(d^0)}{\ker(d^0|_{F^1 C})} \right) [0]$$

$$\oplus$$

$$\left(\frac{C^1}{F^1 C^1 + \text{im}(d^0)} \right) [-1]$$

$$\left(\frac{\ker(d^0) + F^1 C^0}{F^1 C^0} \right) [0]$$

$$\oplus$$

$$\left(\frac{C^1}{F^1 C^1 + \text{im}(d^0)} \right) [-1]$$

BABY EXAMPLE: 2^{nd} approximation to the $\text{gr}^0 H$ part

We compare then $\text{gr}^0 H$ (left column) to $E_2^{0,\bullet}$ (right column):

$$\left(\frac{\ker(d^0)}{\ker(d^0|_{F^1 C})} \right) [0]$$

$$\oplus$$

$$\left(\frac{C^1}{F^1 C^1 + \text{im}(d^0)} \right) [-1]$$

$$\left(\frac{\ker(d^0) + F^1 C^0}{F^1 C^0} \right) [0]$$

$$\oplus$$

$$\left(\frac{C^1}{F^1 C^1 + \text{im}(d^0)} \right) [-1]$$

After applying an \cong -theorem we see that the two agree!

BABY EXAMPLE: 2^{nd} approximation to the $\text{gr}^1 H$ part

We compare now $\text{gr}^1 H$ (left column) to $E_2^{1,\bullet-1}$ (right column):

$$\ker (d^0|_{F^1 C}) [0]$$

$$\oplus$$

$$\left(\frac{F^1 C^1 + \text{im}(d^0)}{\text{im}(d^0)} \right) [-1]$$

$$\ker (d^0|_{F^1 C}) [0]$$

$$\oplus$$

$$\left(\frac{F^1 C^1}{\text{im}(d^0) \cap F^1 C^1} \right) [-1]$$


BABY EXAMPLE: 2^{nd} approximation to the $\text{gr}^1 H$ part

We compare now $\text{gr}^1 H$ (left column) to $E_2^{1,\bullet-1}$ (right column):


$$\begin{array}{ccc} \ker (d^0|_{F^1 C}) [0] & & \ker (d^0|_{F^1 C}) [0] \\ \oplus & & \oplus \\ \left(\frac{F^1 C^1 + \text{im}(d^0)}{\text{im}(d^0)} \right) [-1] & & \left(\frac{F^1 C^1}{\text{im}(d^0) \cap F^1 C^1} \right) [-1] \end{array}$$

After applying an \cong -theorem we see that the two sides agree again!

BABY EXAMPLE: reading off the result from the “ E_2 -page”




BABY EXAMPLE: reading off the result from the “ E_2 -page”




► $\text{gr}^0 H \cong E_2^{0,0}[0] \oplus E_2^{0,1}[-1];$

BABY EXAMPLE: reading off the result from the “ E_2 -page”



- ▶ $\text{gr}^0 H \cong E_2^{0,0}[0] \oplus E_2^{0,1}[-1];$
- ▶ $\text{gr}^1 H \cong E_2^{1,-1}[0] \oplus E_2^{1,0}[-1].$

BABY EXAMPLE: reading off the result from the “ E_2 -page”



- $$\begin{aligned} \blacktriangleright \operatorname{gr}^0 H &\cong E_2^{0,0}[0] \oplus E_2^{0,1}[-1]; \\ \blacktriangleright \operatorname{gr}^1 H &\cong E_2^{1,-1}[0] \oplus E_2^{1,0}[-1]. \end{aligned} \quad \begin{cases} H^0 \cong \operatorname{gr}^0 H^0 \oplus \operatorname{gr}^1 H^0 \cong E_2^{0,0} \oplus E_2^{1,-1} \\ H^1 \cong \operatorname{gr}^1 H^0 \oplus \operatorname{gr}^1 H^1 \cong E_2^{0,1} \oplus E_2^{1,0} \end{cases}$$

DEFINITION [Sta20, Tag 011N]

A *spectral sequence* in an abelian category \mathcal{A} is given by a system $(E_r, d_r)_{r \in \mathbb{N}}$ such that for all $r \in \mathbb{N}$ we have:

DEFINITION [Sta20, Tag 011N]

A *spectral sequence* in an abelian category \mathcal{A} is given by a system $(E_r, d_r)_{r \in \mathbb{N}}$ such that for all $r \in \mathbb{N}$ we have:

- ▶ E_r is an object of \mathcal{A} ;

DEFINITION [Sta20, Tag 011N]

A *spectral sequence* in an abelian category \mathcal{A} is given by a system $(E_r, d_r)_{r \in \mathbb{N}}$ such that for all $r \in \mathbb{N}$ we have:

- ▶ E_r is an object of \mathcal{A} ;
- ▶ $d_r: E_r \rightarrow E_r$ is a morphism such that $d_r \circ d_r = 0$;

DEFINITION [Sta20, Tag 011N]

A *spectral sequence* in an abelian category \mathcal{A} is given by a system $(E_r, d_r)_{r \in \mathbb{N}}$ such that for all $r \in \mathbb{N}$ we have:

- ▶ E_r is an object of \mathcal{A} ;
- ▶ $d_r: E_r \rightarrow E_r$ is a morphism such that $d_r \circ d_r = 0$;
- ▶ and $E_{r+1} = \ker(d_r)/\operatorname{im}(d_r)$.

EXISTENCE [Sta20, Tag 012M]

Let $F^\bullet C$ be a filtered complex in an abelian category \mathcal{A} .

EXISTENCE [Sta20, Tag 012M]

Let $F^\bullet C$ be a filtered complex in an abelian category \mathcal{A} . Then there is a spectral sequence $(E_r, d_r)_{r \in \mathbb{N}}$ of bigraded objects in \mathcal{A} such that d_r has bidegree $(r, -r + 1)$.

EXISTENCE [Sta20, Tag 012M]

Let $F^\bullet C$ be a filtered complex in an abelian category \mathcal{A} . Then there is a spectral sequence $(E_r, d_r)_{r \in \mathbb{N}}$ of bigraded objects in \mathcal{A} such that d_r has bidegree $(r, -r + 1)$. More explicitly, the “ E_0 -page” is given by

- ▶ $E_0^{p,q} = \operatorname{gr}^p C^{p+q},$
- ▶ $d_0^{p,q} = \operatorname{gr}^p(d^{p+q}): E_0^{p,q} \rightarrow E_0^{p,q+1};$

EXISTENCE [Sta20, Tag 012M]

Let $F^\bullet C$ be a filtered complex in an abelian category \mathcal{A} . Then there is a spectral sequence $(E_r, d_r)_{r \in \mathbb{N}}$ of bigraded objects in \mathcal{A} such that d_r has bidegree $(r, -r+1)$. More explicitly, the “ E_0 -page” is given by

- ▶ $E_0^{p,q} = \operatorname{gr}^p C^{p+q},$
- ▶ $d_0^{p,q} = \operatorname{gr}^p(d^{p+q}): E_0^{p,q} \rightarrow E_0^{p,q+1};$

and the “ E_r -page” is given by

- ▶ $E_r^{p,q} = (F^p C^{p+q} \cap d^{-1}(F^{p+r} C^{p+q+1})) / (F^p C^{p+q} \cap d(F^{p-r} C^{p+q-1})),$
- ▶ $d_r^{p,q}: E_r^{p,q} \rightarrow E_r^{p+r, q-r+1}$ induced by $d^{p+q}: F^p C^{p+q} \rightarrow F^{p+r} C^{p+r+q-r+1}.$

EXISTENCE [Sta20, Tag 012M]

Let $F^\bullet C$ be a filtered complex in an abelian category \mathcal{A} . Then there is a spectral sequence $(E_r, d_r)_{r \in \mathbb{N}}$ of bigraded objects in \mathcal{A} such that d_r has bidegree $(r, -r+1)$. More explicitly, the “ E_0 -page” is given by

- ▶ $E_0^{p,q} = \operatorname{gr}^p C^{p+q},$
- ▶ $d_0^{p,q} = \operatorname{gr}^p(d^{p+q}): E_0^{p,q} \rightarrow E_0^{p,q+1};$

and the “ E_r -page” is given by

- ▶ $E_r^{p,q} = (F^p C^{p+q} \cap d^{-1}(F^{p+r} C^{p+q+1})) / (F^p C^{p+q} \cap d(F^{p-r} C^{p+q-1})),$
- ▶ $d_r^{p,q}: E_r^{p,q} \rightarrow E_r^{p+r, q-r+1}$ induced by
 $d^{p+q}: F^p C^{p+q} \rightarrow F^{p+r} C^{p+r+q-r+1}.$

If the filtration induced on each C^n was finite, then letting $r \rightarrow \infty$ we would obtain the *limit* E_∞ of the spectral sequence, given by

$$E_\infty^{p,q} = (F^p C^{p+q} \cap \ker(d)) / (F^p C^{p+q} \cap \operatorname{im}(d)).$$

CONVERGENCE: $E_r^{p,q} \Rightarrow H^{p+q}(C)$ [McC01, Theorem 2.6]

Let $F^\bullet C$ be a filtered complex in an abelian category \mathcal{A} and assume that the filtration induced on each C^n is finite. Then:

CONVERGENCE: $E_r^{p,q} \Rightarrow H^{p+q}(C)$ [McC01, Theorem 2.6]

Let $F^\bullet C$ be a filtered complex in an abelian category \mathcal{A} and assume that the filtration induced on each C^n is finite. Then:

- ▶ The induced filtration $F^\bullet H$ is finite.
- ▶ For each $p, q \in \mathbb{Z}$ there exists $r_0 \in \mathbb{N}$ such that

$$E_\infty^{p,q} = E_r^{p,q} \text{ for all } r \geq r_0.$$

- ▶ The spectral sequence *converges* to $F^\bullet H$, i.e.

$$E_\infty^{p,q} \cong \operatorname{gr}^p H^{p+q} \text{ for all } p, q \in \mathbb{Z}.$$

In particular, modulo extension problems (e.g. if we are working with vector spaces), we can recover H as

$$H^n(C) \cong \bigoplus_{p \in \mathbb{Z}} E_\infty^{p, n-p} \text{ for all } n \in \mathbb{Z}.$$

APPLICATIONS of spectral sequences from filtrations

- ▶ Cellular homology and singular homology of a CW complex agree [[McC01](#), Theorem 4.13].

APPLICATIONS of spectral sequences from filtrations

- ▶ Cellular homology and singular homology of a CW complex agree [[McC01](#), Theorem 4.13].
- ▶ “Cheap Hodge decomposition” [[Voi02](#), Remark 8.29].

APPLICATIONS of spectral sequences from filtrations

- ▶ Cellular homology and singular homology of a CW complex agree [[McC01](#), Theorem 4.13].
- ▶ “Cheap Hodge decomposition” [[Voi02](#), Remark 8.29].
- ▶ Two different ways to compute the cohomology of the total complex of a double complex.

APPLICATIONS of spectral sequences from filtrations

- ▶ Cellular homology and singular homology of a CW complex agree [[McC01](#), Theorem 4.13].
- ▶ “Cheap Hodge decomposition” [[Voi02](#), Remark 8.29].
- ▶ Two different ways to compute the cohomology of the total complex of a double complex.

To avoid getting too much into specific topics, we will focus on the third application. We will do this again with an example.

SNAKE LEMMA via spectral sequences

References



Timothy Y. Chow.

You could have invented spectral sequences.

Notices Amer. Math. Soc., 53(1):15–19, 2006.

1



John McCleary.

A user's guide to spectral sequences, volume 58 of *Cambridge Studies in Advanced Mathematics*.

Cambridge University Press, Cambridge, second edition, 2001.

(document), 22



The Stacks project authors.

The stacks project.

<https://stacks.math.columbia.edu>, 2020.

(document)



Claire Voisin.

Hodge theory and complex algebraic geometry. I, volume 76 of *Cambridge Studies in Advanced Mathematics*.

Cambridge University Press, Cambridge, 2002.