

# Basic Notions — Spectral Sequences

Pedro Núñez

University of Freiburg

10th December 2020

“You could have invented spectral sequences”<sup>1</sup>



---

<sup>1</sup>Title of the expository article [Cho06]

Notation gets ugly real quick



# Notation usually following the Stacks project [Sta20]

- ▶ Only finite dimensional vector spaces over  $\mathbb{Q}$  for simplicity.

## Notation usually following the Stacks project [Sta20]

- ▶ Only finite dimensional vector spaces over  $\mathbb{Q}$  for simplicity.
- ▶ (Cochain) complexes denoted with capital letters  $C$  and not  $C^\bullet$ .

## Notation usually following the Stacks project [Sta20]

- ▶ Only finite dimensional vector spaces over  $\mathbb{Q}$  for simplicity.
- ▶ (Cochain) complexes denoted with capital letters  $C$  and not  $C^\bullet$ .
- ▶ Cohomology again denoted just  $H(C)$  and not  $H^\bullet(C)$ .

## Notation usually following the Stacks project [Sta20]

- ▶ Only finite dimensional vector spaces over  $\mathbb{Q}$  for simplicity.
- ▶ (Cochain) complexes denoted with capital letters  $C$  and not  $C^\bullet$ .
- ▶ Cohomology again denoted just  $H(C)$  and not  $H^\bullet(C)$ .
- ▶ Cohomology is regarded as a complex with 0 differentials

$$H(C) = \bigoplus_{i \in \mathbb{Z}} H^i(C)[-i] := \left( \cdots H^{-1}(C) \xrightarrow{0} H^0(C) \xrightarrow{0} H^1(C) \cdots \right).$$

## Notation usually following the Stacks project [Sta20]

- ▶ Only finite dimensional vector spaces over  $\mathbb{Q}$  for simplicity.
- ▶ (Cochain) complexes denoted with capital letters  $C$  and not  $C^\bullet$ .
- ▶ Cohomology again denoted just  $H(C)$  and not  $H^\bullet(C)$ .
- ▶ Cohomology is regarded as a complex with 0 differentials

$$H(C) = \bigoplus_{i \in \mathbb{Z}} H^i(C)[-i] := \left( \cdots H^{-1}(C) \xrightarrow{0} H^0(C) \xrightarrow{0} H^1(C) \cdots \right).$$

- ▶ If it is clear to which  $C$  we refer, denote  $H(C)$  simply by  $H$ .



## Notation usually following the Stacks project [Sta20]

- ▶ Only finite dimensional vector spaces over  $\mathbb{Q}$  for simplicity.
- ▶ (Cochain) complexes denoted with capital letters  $C$  and not  $C^\bullet$ .
- ▶ Cohomology again denoted just  $H(C)$  and not  $H^\bullet(C)$ .
- ▶ Cohomology is regarded as a complex with 0 differentials

$$H(C) = \bigoplus_{i \in \mathbb{Z}} H^i(C)[-i] := \left( \cdots H^{-1}(C) \xrightarrow{0} H^0(C) \xrightarrow{0} H^1(C) \cdots \right).$$

- ▶ If it is clear to which  $C$  we refer, denote  $H(C)$  simply by  $H$ .
- ▶ (Decreasing) filtrations are always finite for simplicity:

$$F^\bullet A: \quad 0 = F^n A \subseteq \cdots \subseteq F^0 A = A.$$

## Notation usually following the Stacks project [Sta20]

- ▶ Only finite dimensional vector spaces over  $\mathbb{Q}$  for simplicity.
- ▶ (Cochain) complexes denoted with capital letters  $C$  and not  $C^\bullet$ .
- ▶ Cohomology again denoted just  $H(C)$  and not  $H^\bullet(C)$ .
- ▶ Cohomology is regarded as a complex with 0 differentials

$$H(C) = \bigoplus_{i \in \mathbb{Z}} H^i(C)[-i] := \left( \cdots H^{-1}(C) \xrightarrow{0} H^0(C) \xrightarrow{0} H^1(C) \cdots \right).$$


- ▶ If it is clear to which  $C$  we refer, denote  $H(C)$  simply by  $H$ .
- ▶ (Decreasing) filtrations are always finite for simplicity:

$$F^\bullet A: \quad 0 = F^n A \subseteq \cdots \subseteq F^0 A = A.$$

- ▶ The graded object associated to a filtration  $F^\bullet A$  is denoted

$$\mathrm{gr} A = \bigoplus_{p \in \mathbb{N}} \mathrm{gr}^p A := \bigoplus_{p \in \mathbb{N}} (F^p A / F^{p+1} A).$$

# Goal: compute cohomology with the help of a filtration



# Filtration induced in cohomology

A filtration on the complex  $F^\bullet C$  induces a filtration in cohomology

$$F^p H := \operatorname{im}(H(F^p C) \rightarrow H(C)) \subseteq H(C) = H,$$

where  $H(F^p C) \rightarrow H(C)$  is induced by the inclusion map  $F^p C \hookrightarrow C$ .

## Baby example: 2 step filtration on a length 1 complex

$$\begin{array}{ccccccc}
 C = F^0 C & \cdots \rightarrow & 0 \rightarrow & C^0 & \xrightarrow{d^0} & C^1 & \rightarrow 0 \rightarrow \cdots \\
 \cup & & & \cup & & \cup & \\
 F^1 C & \cdots \rightarrow & 0 \rightarrow & F^1 C^0 & \xrightarrow{d^0|_{F^1 C^0}} & F^1 C^1 & \rightarrow 0 \rightarrow \cdots \\
 \cup & & & \cup & & \cup & \\
 0 = F^2 C & \cdots \rightarrow & 0 \rightarrow & 0 & \longrightarrow & 0 & \rightarrow 0 \rightarrow \cdots
 \end{array}$$

## Baby example: filtration induced in cohomology

$$\begin{array}{ccc} H(F^1 C) & = & \ker(d^0|_{F^1 C}) [0] \oplus \left( \frac{F^1 C^1}{\operatorname{im}(d^0|_{F^1 C})} \right) [-1] \\ \downarrow & & \downarrow \\ H(C) & = & \ker(d^0) [0] \oplus \left( \frac{C^1}{\operatorname{im}(d^0)} \right) [-1] \end{array}$$

## Baby example: filtration induced in cohomology

$$\begin{array}{ccc} H(F^1 C) & = & \ker(d^0|_{F^1 C}) [0] \oplus \left( \frac{F^1 C^1}{\operatorname{im}(d^0|_{F^1 C})} \right) [-1] \\ \downarrow & & \downarrow \\ H(C) & = & \ker(d^0) [0] \oplus \left( \frac{C^1}{\operatorname{im}(d^0)} \right) [-1] \end{array}$$

$$\Rightarrow F^1 H = \ker(d^0|_{F^1 C}) [0] \oplus \left( \frac{F^1 C^1 + \operatorname{im}(d^0)}{\operatorname{im}(d^0)} \right) [-1]$$

# References



Timothy Y. Chow.

You could have invented spectral sequences.

*Notices Amer. Math. Soc.*, 53(1):15–19, 2006.

1



The Stacks project authors.

The stacks project.

<https://stacks.math.columbia.edu>, 2020.

(document)