

Basic Notions — Spectral Sequences

Pedro Núñez

University of Freiburg

10th December 2020

“You could have invented spectral sequences”¹



¹Title of the expository article [Cho06]

Notation gets ugly real quick



Notation usually following the Stacks project [Sta20]

- ▶ Only finite dimensional vector spaces over \mathbb{Q} for simplicity.

Notation usually following the Stacks project [Sta20]

- ▶ Only finite dimensional vector spaces over \mathbb{Q} for simplicity.
- ▶ (Cochain) complexes denoted with capital letters C and not C^\bullet .

Notation usually following the Stacks project [Sta20]

- ▶ Only finite dimensional vector spaces over \mathbb{Q} for simplicity.
- ▶ (Cochain) complexes denoted with capital letters C and not C^\bullet .
- ▶ Cohomology again denoted just $H(C)$ and not $H^\bullet(C)$.

Notation usually following the Stacks project [Sta20]

- ▶ Only finite dimensional vector spaces over \mathbb{Q} for simplicity.
- ▶ (Cochain) complexes denoted with capital letters C and not C^\bullet .
- ▶ Cohomology again denoted just $H(C)$ and not $H^\bullet(C)$.
- ▶ Cohomology is regarded as a complex with 0 differentials

$$H(C) = \bigoplus_{i \in \mathbb{Z}} H^i(C)[-i] := \left(\cdots H^{-1}(C) \xrightarrow{0} H^0(C) \xrightarrow{0} H^1(C) \cdots \right).$$

Notation usually following the Stacks project [Sta20]

- ▶ Only finite dimensional vector spaces over \mathbb{Q} for simplicity.
- ▶ (Cochain) complexes denoted with capital letters C and not C^\bullet .
- ▶ Cohomology again denoted just $H(C)$ and not $H^\bullet(C)$.
- ▶ Cohomology is regarded as a complex with 0 differentials

$$H(C) = \bigoplus_{i \in \mathbb{Z}} H^i(C)[-i] := \left(\cdots H^{-1}(C) \xrightarrow{0} H^0(C) \xrightarrow{0} H^1(C) \cdots \right).$$

- ▶ If it is clear to which C we refer, denote $H(C)$ simply by H .

Notation usually following the Stacks project [Sta20]

- ▶ Only finite dimensional vector spaces over \mathbb{Q} for simplicity.
- ▶ (Cochain) complexes denoted with capital letters C and not C^\bullet .
- ▶ Cohomology again denoted just $H(C)$ and not $H^\bullet(C)$.
- ▶ Cohomology is regarded as a complex with 0 differentials

$$H(C) = \bigoplus_{i \in \mathbb{Z}} H^i(C)[-i] := \left(\cdots H^{-1}(C) \xrightarrow{0} H^0(C) \xrightarrow{0} H^1(C) \cdots \right).$$

- ▶ If it is clear to which C we refer, denote $H(C)$ simply by H .
- ▶ (Decreasing) filtrations are always finite for simplicity:

$$F^\bullet A: \quad 0 = F^n A \subseteq \cdots \subseteq F^0 A = A.$$

Notation usually following the Stacks project [Sta20]

- ▶ Only finite dimensional vector spaces over \mathbb{Q} for simplicity.
- ▶ (Cochain) complexes denoted with capital letters C and not C^\bullet .
- ▶ Cohomology again denoted just $H(C)$ and not $H^\bullet(C)$.
- ▶ Cohomology is regarded as a complex with 0 differentials

$$H(C) = \bigoplus_{i \in \mathbb{Z}} H^i(C)[-i] := \left(\cdots H^{-1}(C) \xrightarrow{0} H^0(C) \xrightarrow{0} H^1(C) \cdots \right).$$


- ▶ If it is clear to which C we refer, denote $H(C)$ simply by H .
- ▶ (Decreasing) filtrations are always finite for simplicity:

$$F^\bullet A: \quad 0 = F^n A \subseteq \cdots \subseteq F^0 A = A.$$

- ▶ The graded object associated to a filtration $F^\bullet A$ is denoted

$$\mathrm{gr} A = \bigoplus_{p \in \mathbb{N}} \mathrm{gr}^p A := \bigoplus_{p \in \mathbb{N}} (F^p A / F^{p+1} A).$$

Goal: compute cohomology with the help of a filtration



Starting point: a filtration on our complex

- If we do not have any extra information about C^\bullet , there is not much more we can do besides applying the definition

$$H^\bullet = \ker(d^0)[0] \oplus \frac{\ker(d^1)}{\operatorname{im}(d^0)}[-1] \oplus \frac{C^2}{\operatorname{im}(d^1)}[-2].$$

Starting point: a filtration on our complex

- ▶ If we do not have any extra information about C^\bullet , there is not much more we can do besides applying the definition

$$H^\bullet = \ker(d^0)[0] \oplus \frac{\ker(d^1)}{\operatorname{im}(d^0)}[-1] \oplus \frac{C^2}{\operatorname{im}(d^1)}[-2].$$

- ▶ But C^\bullet has often some useful filtration $F^\bullet C^\bullet$, say

$$\begin{array}{ccccccc} C^\bullet = F^0 C^\bullet & \cdots 0 & \longrightarrow & C^0 & \xrightarrow{d^0} & C^1 & \xrightarrow{d^1} & C^2 & \longrightarrow & 0 \cdots \\ \cup & & & \cup & & \cup & & \cup & & \\ F^1 C^\bullet & \cdots 0 & \longrightarrow & F^1 C^0 & \longrightarrow & F^1 C^1 & \longrightarrow & F^1 C^2 & \longrightarrow & 0 \cdots \\ \cup & & & \cup & & \cup & & \cup & & \\ 0 = F^2 C^\bullet & \cdots 0 & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & 0 \cdots \end{array}$$

How could a filtration on C^\bullet potentially help?

How could a filtration on C^\bullet potentially help?

1. $0 \subseteq F^1 C^\bullet \subseteq C^\bullet$ induces a filtration $0 \subseteq F^1 H^\bullet \subseteq H^\bullet$ given by

$$F^1 H^i := \operatorname{im} (H^i(F^1 C^\bullet) \rightarrow H^i(C^\bullet)) \text{ for all } i \in \{0, 1, 2\}.$$

How could a filtration on C^\bullet potentially help?

1. $0 \subseteq F^1 C^\bullet \subseteq C^\bullet$ induces a filtration $0 \subseteq F^1 H^\bullet \subseteq H^\bullet$ given by

$$F^1 H^i := \operatorname{im} (H^i(F^1 C^\bullet) \rightarrow H^i(C^\bullet)) \text{ for all } i \in \{0, 1, 2\}.$$

2. We can then recover H^i as the associated graded vector space

$$\operatorname{gr}(F^\bullet H^i) = F^1 H^i \oplus (H^i / F^1 H^i) \cong H^i \text{ for all } i \in \{0, 1, 2\}.$$

How could a filtration on C^\bullet potentially help?

1. $0 \subseteq F^1 C^\bullet \subseteq C^\bullet$ induces a filtration $0 \subseteq F^1 H^\bullet \subseteq H^\bullet$ given by

$$F^1 H^i := \operatorname{im} (H^i(F^1 C^\bullet) \rightarrow H^i(C^\bullet)) \text{ for all } i \in \{0, 1, 2\}.$$

2. We can then recover H^i as the associated graded vector space

$$\operatorname{gr}(F^\bullet H^i) = F^1 H^i \oplus (H^i / F^1 H^i) \cong H^i \text{ for all } i \in \{0, 1, 2\}.$$

3. $F^\bullet C^\bullet$ being a filtered complex means that $F^1 C^\bullet \subseteq C^\bullet$ is a subcomplex, so that $F^1 C^\bullet$ and $C^\bullet / F^1 C^\bullet$ are complexes as well.

How could a filtration on C^\bullet potentially help?

1. $0 \subseteq F^1 C^\bullet \subseteq C^\bullet$ induces a filtration $0 \subseteq F^1 H^\bullet \subseteq H^\bullet$ given by

$$F^1 H^i := \operatorname{im} (H^i(F^1 C^\bullet) \rightarrow H^i(C^\bullet)) \text{ for all } i \in \{0, 1, 2\}.$$

2. We can then recover H^i as the associated graded vector space

$$\operatorname{gr}(F^\bullet H^i) = F^1 H^i \oplus (H^i / F^1 H^i) \cong H^i \text{ for all } i \in \{0, 1, 2\}.$$

3. $F^\bullet C^\bullet$ being a filtered complex means that $F^1 C^\bullet \subseteq C^\bullet$ is a subcomplex, so that $F^1 C^\bullet$ and $C^\bullet / F^1 C^\bullet$ are complexes as well.

- **Q1:** Can we compute the cohomologies of $F^1 C^\bullet$ and $C^\bullet / F^1 C^\bullet$?

How could a filtration on C^\bullet potentially help?

1. $0 \subseteq F^1 C^\bullet \subseteq C^\bullet$ induces a filtration $0 \subseteq F^1 H^\bullet \subseteq H^\bullet$ given by

$$F^1 H^i := \operatorname{im} (H^i(F^1 C^\bullet) \rightarrow H^i(C^\bullet)) \text{ for all } i \in \{0, 1, 2\}.$$

2. We can then recover H^i as the associated graded vector space

$$\operatorname{gr}(F^\bullet H^i) = F^1 H^i \oplus (H^i / F^1 H^i) \cong H^i \text{ for all } i \in \{0, 1, 2\}.$$

3. $F^\bullet C^\bullet$ being a filtered complex means that $F^1 C^\bullet \subseteq C^\bullet$ is a subcomplex, so that $F^1 C^\bullet$ and $C^\bullet / F^1 C^\bullet$ are complexes as well.

- **Q1:** Can we compute the cohomologies of $F^1 C^\bullet$ and $C^\bullet / F^1 C^\bullet$?
- **Q2:** Are these cohomologies related to $F^1 H^\bullet$ and $H^\bullet / F^1 H^\bullet$?

Possible scenario I: the good

- ▶ Suppose $C^\bullet = C_1^\bullet \oplus C_2^\bullet$ with $H^\bullet(C_1^\bullet)$ and $H^\bullet(C_2^\bullet)$ computable.

Possible scenario I: the good

- ▶ Suppose $C^\bullet = C_1^\bullet \oplus C_2^\bullet$ with $H^\bullet(C_1^\bullet)$ and $H^\bullet(C_2^\bullet)$ computable.
- ▶ Then we are done, because $H^\bullet(C^\bullet) = H^\bullet(C_1^\bullet) \oplus H^\bullet(C_2^\bullet)$.

Possible scenario I: the good

- ▶ Suppose $C^\bullet = C_1^\bullet \oplus C_2^\bullet$ with $H^\bullet(C_1^\bullet)$ and $H^\bullet(C_2^\bullet)$ computable.
- ▶ Then we are done, because $H^\bullet(C^\bullet) = H^\bullet(C_1^\bullet) \oplus H^\bullet(C_2^\bullet)$.
- ▶ To rephrase this with filtrations we take

$$F^1 C^\bullet := C_1^\bullet \quad (\Rightarrow \quad C^\bullet / F^1 C^\bullet = C_2^\bullet.)$$

Possible scenario I: the good

- ▶ Suppose $C^\bullet = C_1^\bullet \oplus C_2^\bullet$ with $H^\bullet(C_1^\bullet)$ and $H^\bullet(C_2^\bullet)$ computable.
- ▶ Then we are done, because $H^\bullet(C^\bullet) = H^\bullet(C_1^\bullet) \oplus H^\bullet(C_2^\bullet)$.
- ▶ To rephrase this with filtrations we take

$$F^1 C^\bullet := C_1^\bullet \quad (\Rightarrow \quad C^\bullet / F^1 C^\bullet = C_2^\bullet.)$$

- ▶ Then we have

$$F^1 H^\bullet := \text{im}(H^\bullet(F^1 C^\bullet) \rightarrow H^\bullet(C^\bullet)) = H^\bullet(C_1^\bullet) = H^\bullet(F^1 C^\bullet)$$

and similarly $H^\bullet / F^1 H^\bullet = H^\bullet(C_2^\bullet) = H^\bullet(C^\bullet / F^1 C^\bullet)$.

Possible scenario I: the good

- ▶ Suppose $C^\bullet = C_1^\bullet \oplus C_2^\bullet$ with $H^\bullet(C_1^\bullet)$ and $H^\bullet(C_2^\bullet)$ computable.
- ▶ Then we are done, because $H^\bullet(C^\bullet) = H^\bullet(C_1^\bullet) \oplus H^\bullet(C_2^\bullet)$.
- ▶ To rephrase this with filtrations we take


$$F^1 C^\bullet := C_1^\bullet \quad (\Rightarrow \quad C^\bullet / F^1 C^\bullet = C_2^\bullet.)$$

- ▶ Then we have

$$F^1 H^\bullet := \text{im}(H^\bullet(F^1 C^\bullet) \rightarrow H^\bullet(C^\bullet)) = H^\bullet(C_1^\bullet) = H^\bullet(F^1 C^\bullet)$$

and similarly $H^\bullet / F^1 H^\bullet = H^\bullet(C_2^\bullet) = H^\bullet(C^\bullet / F^1 C^\bullet)$.

So in this case we can carry out our naive strategy of computing each graded piece of the induced filtration in cohomology separately:



Possible scenario II: the bad

Suppose we know nothing about C^\bullet . The *stupid filtration*

$$\begin{cases} F^1 C^\bullet := \cdots \rightarrow 0 \rightarrow 0 \rightarrow C^1 \xrightarrow{d^1} C^2 \rightarrow 0 \rightarrow \cdots \\ F^2 C^\bullet := \cdots \rightarrow 0 \rightarrow 0 \rightarrow 0 \rightarrow C^2 \rightarrow 0 \rightarrow \cdots \end{cases}$$

induces the cohomology filtration

$$\begin{cases} F^1 H^\bullet := \operatorname{im}(H^\bullet(F^1 C^\bullet) \rightarrow H^\bullet(C^\bullet)) = H^1(C^\bullet)[-1] \oplus H^2(C^\bullet)[-2] \\ F^2 H^\bullet := \operatorname{im}(H^\bullet(F^2 C^\bullet) \rightarrow H^\bullet(C^\bullet)) = H^2(C^\bullet)[-2] \end{cases}$$

whose graded pieces are $H^0(C^\bullet)[0]$, $H^1(C^\bullet)[-1]$ and $H^2(C^\bullet)[-2]$.
So in this case computing $\operatorname{gr}^i H^\bullet(C^\bullet)$ is as hard as computing $H^i(C^\bullet)$ and we just wasted our time with this detour.

Possible scenario III: the ugly

Suppose now that we are somewhere in between the good and the bad scenarios. Namely, assume that we have some filtration $0 \subsetneq F^1 C^\bullet \subsetneq C^\bullet$ which does not correspond to a direct sum decomposition but such that we can compute the graded pieces $\operatorname{gr}^i(H^\bullet(C^\bullet))$ of the filtration induced in cohomology.

References



Timothy Y. Chow.

You could have invented spectral sequences.

Notices Amer. Math. Soc., 53(1):15–19, 2006.

1



The Stacks project authors.

The stacks project.

<https://stacks.math.columbia.edu>, 2020.

(document)