

# Seesaw Principle and Theorem of the Cube

Remarks on Section I.5 of Milne's *Abelian Varieties*

University of Freiburg

26th May 2020

# Seesaw Principle — Idea

## Theorem (Weil)

*A limit of trivial line bundles on a complete variety is again trivial.*

# Seesaw Principle — Idea

## Theorem (Weil)

*A limit of trivial line bundles on a complete variety is again trivial.*

↓            making this idea more precise

## Theorem (Statement for varieties over $k = \bar{k}$ )

$V$  complete,  $T$  any (“parameter space”),  $\mathcal{L} \in \text{Pic}(V \times T)$ . Then

$$Z := \{t \in T \mid \mathcal{L}_t \text{ is trivial}\} \hookrightarrow T$$

*is closed and  $\mathcal{L}|_{V \times Z}$  is the pullback of a line bundle on  $Z$ .*

# Seesaw Principle — Idea

## Theorem (Weil)

*A limit of trivial line bundles on a complete variety is again trivial.*

↓            making this idea more precise

## Theorem (Statement for varieties over $k = \bar{k}$ )

$V$  complete,  $T$  any (“parameter space”),  $\mathcal{L} \in \text{Pic}(V \times T)$ . Then

$$Z := \{t \in T \mid \mathcal{L}_t \text{ is trivial}\} \hookrightarrow T$$

*is closed and  $\mathcal{L}|_{V \times Z}$  is the pullback of a line bundle on  $Z$ .*

This is the conclusion of the subsection on the seesaw principle in [Mil08], which can be read independently. This statement will be used to **reduce** the Theorem of the Cube to an easier case.

# Seesaw Principle — Generalisation

Theorem ([Bha17, Theorem 6.3] or [Fra18, Theorem 5])

*Let  $f: X \rightarrow S$  be a proper flat morphism of (locally) noetherian schemes with geometrically integral fibres and let  $\mathcal{L} \in X$ . Then*

$$Z := \{s \in S \mid \mathcal{L}_s \text{ is trivial}\} \hookrightarrow S$$

*and  $\mathcal{L}|_{f^{-1}(Z_{\text{red}})} = f^* \mathcal{M}_0$  for some  $\mathcal{M}_0 \in \text{Pic}(Z_{\text{red}})$ . Moreover, there exists a (unique) closed subscheme structure on  $Z$  such that  $\mathcal{L}|_{f^{-1}(Z)} = f^* \mathcal{M}$  for some  $\mathcal{M} \in \text{Pic}(Z)$  and  $Z$  is universal amongst all (locally) noetherian  $S$ -schemes with this property.*

# Seesaw Principle — Generalisation

Theorem ([Bha17, Theorem 6.3] or [Fra18, Theorem 5])


*Let  $f: X \rightarrow S$  be a proper flat morphism of (locally) noetherian schemes with geometrically integral fibres and let  $\mathcal{L} \in X$ . Then*

$$Z := \{s \in S \mid \mathcal{L}_s \text{ is trivial}\} \hookrightarrow S$$


*and  $\mathcal{L}|_{f^{-1}(Z_{\text{red}})} = f^* \mathcal{M}_0$  for some  $\mathcal{M}_0 \in \text{Pic}(Z_{\text{red}})$ . Moreover, there exists a (unique) closed subscheme structure on  $Z$  such that  $\mathcal{L}|_{f^{-1}(Z)} = f^* \mathcal{M}$  for some  $\mathcal{M} \in \text{Pic}(Z)$  and  $Z$  is universal amongst all (locally) noetherian  $S$ -schemes with this property.*

Brian Conrad proves this using existence and separatedness of  $\text{Pic}_{X/S}$  [Con15, Theorem 3.1.1]:  $\mathcal{O}_X$  gives us  $[\mathcal{O}_X]: S \rightarrow \text{Pic}_{X/S}$ , whose image we identify with  $S$ , and then  $Z = [\mathcal{L}](S) \cap [\mathcal{O}_X](S)$ . Conversely, assuming  $\text{Pic}_{X/S}$  exists, the seesaw theorem implies that it is separated [Fra18, Remark 2.4.2] (automatic if  $S = k$ ).

## Seesaw Principle — Particular case in a picture




## Seesaw Principle — Particular case in a picture



We have some  $\mathcal{L} \in \text{Pic}(X \times Y)$ . We can think of  $\mathcal{L}$  as a weight distribution on the two extremes  $X$  and  $Y$  of this seesaw: the more points  $x \in X$  such that  $\mathcal{L}|_{\{x\} \times Y}$  is trivial, the heavier  $\mathcal{L}$  sits on  $X$ .



## Seesaw Principle — Particular case in a picture



We have some  $\mathcal{L} \in \text{Pic}(X \times Y)$ . We can think of  $\mathcal{L}$  as a weight distribution on the two extremes  $X$  and  $Y$  of this seesaw: the more points  $x \in X$  such that  $\mathcal{L}|_{\{x\} \times Y}$  is trivial, the heavier  $\mathcal{L}$  sits on  $X$ . In this case, let's say, we have  $\mathcal{L}|_{X \times \{y\}}$  trivial for all  $y \in Y$ . Then  $\mathcal{L}$  sits with full weight on  $Y$ , so the seesaw principle tells us that there must be some line bundle on  $Y$  from which  $\mathcal{L}$  is the pullback.


# Theorem of the Cube — Statement

Theorem (Statement for varieties over  $k = \bar{k}$ )

$X, Y, Z$  complete and  $\mathcal{L} \in \text{Pic}(X \times Y \times Z)$  such that

$$\mathcal{L}|_{X \times Y \times \{z_0\}}, \mathcal{L}|_{X \times \{y_0\} \times Z} \text{ and } \mathcal{L}|_{\{x_0\} \times Y \times Z}$$

are trivial. Then  $\mathcal{L}$  is trivial.



If the restriction of  $\mathcal{L} \in \text{Pic}(\text{Cube})$  to each face  $XY$ ,  $YZ$  and  $XZ$  is trivial, then  $\mathcal{L}$  is trivial.

# Theorem of the Cube — Generalisation

# References



Bhargav Bhatt.

Math 731: Topics in Algebraic Geometry I – Abelian Varieties, 2017.

Lecture notes by Matt Stevenson, available at [umich.edu/~stevmatt](https://umich.edu/~stevmatt).



Brian Conrad.

Abelian Varieties, 2015.

Lecture notes by Tony Feng, available at [math.stanford.edu/~conrad](https://math.stanford.edu/~conrad).



Jens Franke.

Jacobians of Curves, 2018.

Lecture notes by Ferdinand Wagner, available at [github.com/Nicholas42/AlgebraFranke](https://github.com/Nicholas42/AlgebraFranke).



James S. Milne.

Abelian Varieties (v2.00), 2008.

Available at [jmilne.org/math](https://jmilne.org/math).