

HIGGS BUNDLES — EXISTENCE OF SOLUTIONS

PEDRO NÚÑEZ

ABSTRACT. In this talk we introduce the stability condition for Higgs bundles and prove the Hitchin–Kobayashi correspondence. The main result is [Hit87, Theorem 4.3]. Relevant literature is [Hit87, §3 and §4] and [Wen14, §2 and §3]. Maybe we will also use [Wen16] every now and then.

This talk is related to Tanuj’s talk on *Stable vector bundles*, for which the main reference is [Kob87]. Therefore we will also use [Kob87] as a main reference for generalities on complex vector bundles.

CONTENTS

Notation and conventions	1
1. Self-duality	2
Appendix A. Principal bundles on smooth manifolds	2
References	3

NOTATION AND CONVENTIONS

- **Top** is the category of topological spaces.
- **Diff** is the category of smooth manifolds.
- If \mathbf{C} is a category and $X \in \mathbf{C}$ is an object, then \mathbf{C}/X denotes the category of objects of \mathbf{C} over X , i.e. the category whose objects are morphisms $f: Y \rightarrow X$ in \mathbf{C} and whose morphisms are commutative triangles in \mathbf{C} of the form

$$\begin{array}{ccc} Y_1 & \xrightarrow{f} & Y_2 \\ & \searrow f_1 & \swarrow f_2 \\ & X & \end{array}$$

We will often talk about Y instead of $f: Y \rightarrow X$, leaving this structure morphism implicit.

Date: 6th June 2020.

Supported by the DFG-Graduiertenkolleg GK1821 “Cohomological Methods in Geometry” at the University of Freiburg.

- Let \mathbf{C} be a category which has **Top** as an underlying category, e.g. **Diff**. Let $X \in \mathbf{C}$ and $f \in \mathbf{C}/X$ and let \mathbf{P} be a property of morphisms in \mathbf{C} . We will say that f has some property \mathbf{P} *locally on* X if every point $x \in X$ has an open neighbourhood $x \in U \subseteq X$ in X such that the morphism $f|_{f^{-1}(U)}: f^{-1}(U) \rightarrow U$ has the property \mathbf{P} .
- The category of group objects in a category \mathbf{C} will be denoted by **CGrp**.
- Let \mathbf{C} be a category and let $X \in \mathbf{C}$ and $G \in \mathbf{CGrp}$. A *left action* of G on X , denoted $G \curvearrowright X$, is a morphism

$$\rho: G \times X \rightarrow X$$

such that the following diagrams commute:

$$\begin{array}{ccc} G \times G \times X & \xrightarrow{m \times \text{id}_X} & G \times X \\ \text{id}_G \times \rho \downarrow & & \downarrow \rho \\ G \times X & \xrightarrow{\rho} & X \end{array} \quad \begin{array}{ccc} * \times X & \xrightarrow{e \times \text{id}_X} & G \times X \\ & \searrow & \downarrow \rho \\ & & X \end{array}$$

We can similarly define right actions.

- Let \mathbf{C} be a category. Let $X \in \mathbf{C}$ and $G \in \mathbf{CGrp}$. Then we say that X is a G -object of \mathbf{C} . A morphism of G -objects of \mathbf{C} is a G -equivariant morphism $f: X_1 \rightarrow X_2$, meaning that the following diagram commutes:

$$\begin{array}{ccc} G \times X_1 & \xrightarrow{\text{id}_G \times f} & G \times X_2 \\ \rho_1 \downarrow & & \downarrow \rho_2 \\ X_1 & \xrightarrow{f} & X_2 \end{array}$$

The category of G -objects of \mathbf{C} is denoted $G\text{-}\mathbf{C}$.

1. SELF-DUALITY

We consider \mathbb{R}^4 with its standard smooth structure [Lee13, Example 1.22].

APPENDIX A. PRINCIPAL BUNDLES ON SMOOTH MANIFOLDS

In this appendix we recall the basics of principal G -bundles on smooth manifolds, where G is a Lie group.

Definition A.1 (Lie group). A *Lie group* is a group object in the category **Diff** of smooth manifolds.

Remark A.2. $G \in \mathbf{DiffGrp}$ if and only if its underlying set is equipped with a group structure such that the map $G \times G \rightarrow G$ given by $(g, h) \rightarrow gh^{-1}$ is smooth [Lee13, Proposition 7.1].

Let $M \in \mathbf{Diff}$ and $G \in \mathbf{DiffGrp}$. Then the projection $\pi: M \times G \rightarrow M$ has some nice properties, namely:

- $G \curvearrowright M \times G$ smoothly and fibrewise via $(x, g) \cdot h \mapsto (x, gh)$.
- For all $x \in M$, $G \curvearrowright \pi^{-1}(x)$ induces $G \cong \{x\} \times G \cong \pi^{-1}(x)$.

The smooth manifold $M \times G$ over M equipped with this right fibrewise action is called the *trivial principal G -bundle* on M . We can encode all this structure by saying that

$$M \times G \in (G\text{-}\mathbf{Diff})/M,$$

where we consider M with the trivial G -action.

Definition A.3 (Principal bundle). Let $M \in \mathbf{Diff}$ and $G \in \mathbf{DiffGrp}$. Consider $M \in G\text{-}\mathbf{Diff}$ with the trivial action. A *principal G -bundle* on M is an object $P \in (G\text{-}\mathbf{Diff})/M$ which is trivial locally on M .

Example A.4. Let $M \in \mathbf{Diff}$ and $G := \mathrm{GL}(n, \mathbb{R}) \in \mathbf{DiffGrp}$. Then the *frame bundle* of M , denoted $\mathrm{GL}(M)$, is the principal G -bundle whose fibra over $x \in M$ is the set of all frames for the tangent space $T_x M$.

Example A.5. Let $G \in \mathbf{Diff}$ and $H \subseteq G$ a closed subgroup. Then G is a principal H -bundle over the left coset space G/H .

Some nice properties in the topological category, which probably extend to the smooth category (check!):

- Proposition A.6.**
- i) Any morphism of principal G -bundles is an isomorphism.*
 - ii) A principal G -bundle is trivial if and only if it admits a section, where trivial means isomorphic to a trivial principal G -bundle.*

REFERENCES

- [Hit87] N. J. Hitchin. The self-duality equations on a Riemann surface. *Proc. London Math. Soc.* (3), 55(1):59–126, 1987. [↑ 1](#)
- [Kob87] Shoshichi Kobayashi. *Differential geometry of complex vector bundles*, volume 15 of *Publications of the Mathematical Society of Japan*. Princeton University Press, Princeton, NJ; Princeton University Press, Princeton, NJ, 1987. Kanô Memorial Lectures, 5. [↑ 1](#)
- [Lee13] John M. Lee. *Introduction to smooth manifolds*, volume 218 of *Graduate Texts in Mathematics*. Springer, New York, second edition, 2013. [↑ 2](#)
- [Wen14] Richard A. Wentworth. Higgs bundles and local systems on riemann surfaces, 2014. [↑ 1](#)
- [Wen16] Richard Wentworth. *Higgs Bundles and Local Systems on Riemann Surfaces*, pages 165–219. Springer International Publishing, Cham, 2016. [↑ 1](#)

PEDRO NÚÑEZ, MATHEMATISCHES INSTITUT, ALBERT-LUDWIGS-UNIVERSITÄT
FREIBURG, ERNST-ZERMELO-STRASSE 1, 79104 FREIBURG IM BREISGAU, GER-
MANY

Email address: pedro.nunez@math.uni-freiburg.de

URL: <https://home.mathematik.uni-freiburg.de/nunez>