## **SHORT NOTES ON...**

## Contents

1.	Example section	1
Refe	erences	1

## 1. Example section

Hirzebruch–Riemann–Roch [Har77, Thm. A.4.1] states that for a locally free sheaf  $\mathscr E$  of rank r on a nonsingular projective variety X of dimension n,

$$\chi(\mathscr{E}) = \deg(\operatorname{ch}(\mathscr{E}).\operatorname{td}(\mathscr{T}))_n,$$

where ( )<sub>n</sub> denotes the component of degree n in  $A(X) \otimes \mathbb{Q}$ .

## REFERENCES

[Har77] Robin Hartshorne. *Algebraic geometry*. Graduate Texts in Mathematics, No. 52. Springer-Verlag, New York-Heidelberg, 1977, pp. xvi+496. ISBN: 0-387-90244-9.

Pedro Núñez

Albert-Ludwigs-Universität Freiburg, Mathematisches Institut Ernst-Zermelo-Strasse 1, 79104 Freiburg im Breisgau (Germany) *Email address*: pedro.nunez@math.uni-freiburg.de *Homepage*: https://home.mathematik.uni-freiburg.de/nunez

I would like to thank the DFG-Graduiertenkolleg GK1821 "Cohomological Methods in Geometry" for their support!

Date: 19th March 2021.