

## SOLUTIONS TO EXERCISES

The exercises below are taken from [Har77, Chapter IV].

*Exercise 1.1.* Let  $X$  be a curve, and let  $P \in X$  be a point. Then there exists a nonconstant rational function  $f \in K(X)$ , which is regular everywhere except at  $P$ .

*Solution.* We need to show that  $l(nP) > 1$  for some non-zero  $n \in \mathbb{N}$ . Indeed, Riemann–Roch implies that this holds for all large enough  $n \in \mathbb{N}$ , as explained in [Har77, Remark 1.3.2]. ■

## REFERENCES

- [Har77] Robin Hartshorne. *Algebraic geometry*. Springer-Verlag, New York-Heidelberg, 1977. Graduate Texts in Mathematics, No. 52.

PEDRO NÚÑEZ

ALBERT-LUDWIGS-UNIVERSITÄT FREIBURG, MATHEMATISCHES INSTITUT  
ERNST-ZERMELO-STRASSE 1, 79104 FREIBURG IM BREISGAU (GERMANY)

*Email address:* [pedro.nunez@math.uni-freiburg.de](mailto:pedro.nunez@math.uni-freiburg.de)

*Homepage:* <https://home.mathematik.uni-freiburg.de/nunez>

I would like to thank the DFG-Graduiertenkolleg GK1821 “Cohomological Methods in Geometry” for their support!

---

*Date:* 28th April 2021.