

SOLUTIONS TO EXERCISES

The exercises below are taken from [Har77, Chapter IV].

Exercise 1.1. Let X be a curve, and let $P \in X$ be a point. Then there exists a nonconstant rational function $f \in K(X)$, which is regular everywhere except at P .

Solution. We need to show that $l(nP) > 1$ for some non-zero $n \in \mathbb{N}$. Indeed, Riemann–Roch implies that this holds for all large enough $n \in \mathbb{N}$, as explained in [Har77, Remark 1.3.2]. ■

REFERENCES

- [Har77] Robin Hartshorne. *Algebraic geometry*. Springer-Verlag, New York-Heidelberg, 1977. Graduate Texts in Mathematics, No. 52.

PEDRO NÚÑEZ

ALBERT-LUDWIGS-UNIVERSITÄT FREIBURG, MATHEMATISCHES INSTITUT
ERNST-ZERMELO-STRASSE 1, 79104 FREIBURG IM BREISGAU (GERMANY)

Email address: pedro.nunez@math.uni-freiburg.de

Homepage: <https://home.mathematik.uni-freiburg.de/nunez>

I would like to thank the DFG-Graduiertenkolleg GK1821 “Cohomological Methods in Geometry” for their support!

Date: 28th April 2021.