

Torelli's Theorem

Remarks on Sections III.12–13 of Milne's *Abelian Varieties*

University of Freiburg

28th July 2020

Assumptions, notation and recollections

- k is an algebraically closed field.

Assumptions, notation and recollections

- k is an algebraically closed field.
- C is a smooth complete curve over k of genus $g \geq 2$.

Assumptions, notation and recollections

- k is an algebraically closed field.
- C is a smooth complete curve over k of genus $g \geq 2$.
- J is its Jacobian, which is an abelian variety representing

$$T \mapsto \{\mathcal{L} \in \text{Pic}(C \times T) \mid \deg(\mathcal{L}_t) = 0 \text{ for all } t \in T\} / \text{Pic}(T).$$

Assumptions, notation and recollections

- k is an algebraically closed field.
- C is a smooth complete curve over k of genus $g \geq 2$.
- J is its Jacobian, which is an abelian variety representing

$$T \mapsto \{\mathcal{L} \in \text{Pic}(C \times T) \mid \deg(\mathcal{L}_t) = 0 \text{ for all } t \in T\} / \text{Pic}(T).$$

- In particular, we may and will identify $J(k) = \text{Pic}^0(C)$.

Assumptions, notation and recollections

- k is an algebraically closed field.
- C is a smooth complete curve over k of genus $g \geq 2$.
- J is its Jacobian, which is an abelian variety representing

$$T \mapsto \{\mathcal{L} \in \text{Pic}(C \times T) \mid \deg(\mathcal{L}_t) = 0 \text{ for all } t \in T\} / \text{Pic}(T).$$

- In particular, we may and will identify $J(k) = \text{Pic}^0(C)$.
- Fix $P \in C(k)$ once and for all. Then we get a canonical map

$$f: C \rightarrow J, \quad Q \mapsto [Q - P].$$

Assumptions, notation and recollections

- k is an algebraically closed field.
- C is a smooth complete curve over k of genus $g \geq 2$.
- J is its Jacobian, which is an abelian variety representing

$$T \mapsto \{\mathcal{L} \in \text{Pic}(C \times T) \mid \deg(\mathcal{L}_t) = 0 \text{ for all } t \in T\} / \text{Pic}(T).$$

- In particular, we may and will identify $J(k) = \text{Pic}^0(C)$.
- Fix $P \in C(k)$ once and for all. Then we get a canonical map

$$f: C \rightarrow J, \quad Q \mapsto [Q - P].$$

- For all $1 \leq r \leq g$ we get an induced map

$$f: C^{(r)} \rightarrow J, \quad P_1 \cdot \dots \cdot P_r \mapsto [P_1 + \dots + P_r - rP]$$


birational onto its image $W^r \subseteq J$, which is a closed subvariety.

Canonical polarization

- For $r = g - 1$ we get a divisor $\Theta = W^{g-1} \subseteq J$, the image of $f: C^{(g-1)} \rightarrow J, \quad P_1 \cdot \dots \cdot P_{g-1} \mapsto [P_1 + \dots + P_{g-1} - (g-1)P].$


Canonical polarization

- For $r = g - 1$ we get a divisor $\Theta = W^{g-1} \subseteq J$, the image of $f: C^{(g-1)} \rightarrow J$, $P_1 \cdots P_{g-1} \mapsto [P_1 + \cdots + P_{g-1} - (g-1)P]$.



Canonical polarization

- For $r = g - 1$ we get a divisor $\Theta = W^{g-1} \subseteq J$, the image of $f: C^{(g-1)} \rightarrow J$, $P_1 \cdots P_{g-1} \mapsto [P_1 + \cdots + P_{g-1} - (g-1)P]$.



- The induced $\lambda: J \rightarrow J^\vee$ is an isomorphism, so Θ gives us a principal polarization of J called the *canonical polarization*.

Statement — Existence

Let C and C' be curves as before and let $\beta: (J, \lambda) \xrightarrow{\sim} (J', \lambda')$ be an isomorphism such that $\lambda' \circ \beta = \beta^\vee \circ \lambda$.

Statement — Existence

Let C and C' be curves as before and let $\beta: (J, \lambda) \xrightarrow{\sim} (J', \lambda')$ be an isomorphism such that $\lambda' \circ \beta = \beta^\vee \circ \lambda$.

Then there exists an isomorphism $\alpha: C \xrightarrow{\sim} C'$ such that

$$\begin{array}{ccc} C & \xrightarrow{f} & J \\ \alpha \downarrow & & \downarrow \beta \\ C' & \xrightarrow{f'} & J' \end{array}$$

commutes up to a sign and translation by some $c \in J'(k)$.

Statement — Uniqueness

A curve C as before is called *hyperelliptic* if there is a $2 : 1$ branched covering $\gamma: C \rightarrow \mathbb{P}^1$.

Statement — Uniqueness

A curve C as before is called *hyperelliptic* if there is a $2 : 1$ branched covering $\gamma: C \rightarrow \mathbb{P}^1$. From Hartshorne's exercises:

- A curve of genus 2 is always hyperelliptic.
- There are hyperelliptic curves of any genus $g \geq 2$.
- A plane curve of degree 4 (thus $g = 3$) is not hyperelliptic.

Statement — Uniqueness

A curve C as before is called *hyperelliptic* if there is a $2 : 1$ branched covering $\gamma: C \rightarrow \mathbb{P}^1$. From Hartshorne's exercises:

- A curve of genus 2 is always hyperelliptic.
- There are hyperelliptic curves of any genus $g \geq 2$.
- A plane curve of degree 4 (thus $g = 3$) is not hyperelliptic.

We can now state uniqueness distinguishing two cases:

- If C is not hyperelliptic, then α , the sign and the element $c \in J'(k)$ by which we have to translate are uniquely determined by β , P and P' .
- If C is hyperelliptic, then the sign can be chosen arbitrarily, and α and c are uniquely determined by β , P , P' and the chosen sign.

Thanks for your attention! Here are some references: