

Torelli's Theorem

Remarks on Sections III.12–13 of Milne's *Abelian Varieties*

University of Freiburg

28th July 2020

Assumptions, notation and recollections

- k is an algebraically closed field.

Assumptions, notation and recollections

- k is an algebraically closed field.
- C is a smooth complete curve over k of genus $g \geq 2$.

Assumptions, notation and recollections

- k is an algebraically closed field.
- C is a smooth complete curve over k of genus $g \geq 2$.
- J is its Jacobian, which is an abelian variety representing

$$T \mapsto \{\mathcal{L} \in \text{Pic}(C \times T) \mid \deg(\mathcal{L}_t) = 0 \text{ for all } t \in T\} / \text{Pic}(T).$$

Assumptions, notation and recollections

- k is an algebraically closed field.
- C is a smooth complete curve over k of genus $g \geq 2$.
- J is its Jacobian, which is an abelian variety representing

$$T \mapsto \{ \mathcal{L} \in \text{Pic}(C \times T) \mid \deg(\mathcal{L}_t) = 0 \text{ for all } t \in T \} / \text{Pic}(T).$$

- In particular, we may and will identify $J(k) = \text{Pic}^0(C)$.

Assumptions, notation and recollections

- k is an algebraically closed field.
- C is a smooth complete curve over k of genus $g \geq 2$.
- J is its Jacobian, which is an abelian variety representing

$$T \mapsto \{\mathcal{L} \in \text{Pic}(C \times T) \mid \deg(\mathcal{L}_t) = 0 \text{ for all } t \in T\} / \text{Pic}(T).$$

- In particular, we may and will identify $J(k) = \text{Pic}^0(C)$.
- Fix $P \in C(k)$ once and for all. Then we get a canonical map

$$f: C \rightarrow J, \quad Q \mapsto [Q - P].$$

Assumptions, notation and recollections

- k is an algebraically closed field.
- C is a smooth complete curve over k of genus $g \geq 2$.
- J is its Jacobian, which is an abelian variety representing

$$T \mapsto \{\mathcal{L} \in \text{Pic}(C \times T) \mid \deg(\mathcal{L}_t) = 0 \text{ for all } t \in T\} / \text{Pic}(T).$$

- In particular, we may and will identify $J(k) = \text{Pic}^0(C)$.
- Fix $P \in C(k)$ once and for all. Then we get a canonical map

$$f: C \rightarrow J, \quad Q \mapsto [Q - P].$$

- For all $1 \leq r \leq g$ we get an induced map

$$f: C^{(r)} \rightarrow J, \quad P_1 \cdot \dots \cdot P_r \mapsto [P_1 + \dots + P_r - rP]$$

birational onto its image $W^r \subseteq J$, which is a closed subvariety.

Canonical polarization


- For $r = g - 1$ we get a divisor $\Theta = W^{g-1} \subseteq J$, the image of

$$f: C^{(g-1)} \rightarrow J, \quad P_1 \cdot \dots \cdot P_{g-1} \mapsto [P_1 + \dots + P_{g-1} - (g-1)P].$$

Canonical polarization

- For $r = g - 1$ we get a divisor $\Theta = W^{g-1} \subseteq J$, the image of


$$f: C^{(g-1)} \rightarrow J, \quad P_1 \cdot \dots \cdot P_{g-1} \mapsto [P_1 + \dots + P_{g-1} - (g-1)P].$$



Canonical polarization

- For $r = g - 1$ we get a divisor $\Theta = W^{g-1} \subseteq J$, the image of

$$f: C^{(g-1)} \rightarrow J, \quad P_1 \cdot \dots \cdot P_{g-1} \mapsto [P_1 + \dots + P_{g-1} - (g-1)P].$$



- The induced $\lambda: J \rightarrow J^\vee$ is an isomorphism, so Θ gives us a principal polarization of J called the *canonical polarization*.

Statement — Existence

Let C and C' be curves as before and let $\beta: (J, \lambda) \xrightarrow{\sim} (J', \lambda')$ be an isomorphism such that $\lambda' \circ \beta = \beta^\vee \circ \lambda$.

Statement — Existence

Let C and C' be curves as before and let $\beta: (J, \lambda) \xrightarrow{\sim} (J', \lambda')$ be an isomorphism such that $\lambda' \circ \beta = \beta^\vee \circ \lambda$.

Then there exists an isomorphism $\alpha: C \xrightarrow{\sim} C'$ such that

$$\begin{array}{ccc} C & \xrightarrow{f} & J \\ \alpha \downarrow & & \downarrow \beta \\ C' & \xrightarrow{f'} & J' \end{array}$$

commutes up to a sign and a translation by some $c \in J'(k)$.

Statement — Uniqueness

A curve C as before is called *hyperelliptic* if there is a (unique) $2 : 1$ branched covering $\pi : C \rightarrow \mathbb{P}^1$.

Statement — Uniqueness

A curve C as before is called *hyperelliptic* if there is a (unique) $2 : 1$ branched covering $\pi : C \rightarrow \mathbb{P}^1$. From Hartshorne's exercises:

- A curve of genus 2 is always hyperelliptic.
- There are hyperelliptic curves of any genus $g \geq 2$.
- A plane curve of degree 4 (thus $g = 3$) is not hyperelliptic.

Statement — Uniqueness

A curve C as before is called *hyperelliptic* if there is a (unique) $2 : 1$ branched covering $\pi : C \rightarrow \mathbb{P}^1$. From Hartshorne's exercises:

- A curve of genus 2 is always hyperelliptic.
- There are hyperelliptic curves of any genus $g \geq 2$.
- A plane curve of degree 4 (thus $g = 3$) is not hyperelliptic.

We can now state uniqueness distinguishing two cases:

- If C is not hyperelliptic, then the sign, α and c are uniquely determined by β , P and P' .
- If C is hyperelliptic, then the sign can be chosen arbitrarily, and α and c are uniquely determined by β , P , P' and the chosen sign.

Proof — Uniqueness (modulo case distinctions for signs)

- Suppose α_1, α_2, c_1 and c_2 were such that $f' \circ \alpha_i = \beta \circ f + c_i$.

Proof — Uniqueness (modulo case distinctions for signs)

- Suppose α_1, α_2, c_1 and c_2 were such that $f' \circ \alpha_i = \beta \circ f + c_i$.
- Then $f' \circ (\alpha_1 - \alpha_2): C \rightarrow J'$ is constant, so $\alpha_1 - \alpha_2$ sends every pair of points in C to the same fibre of $f': C' \rightarrow J'$.

Proof — Uniqueness (modulo case distinctions for signs)

- Suppose α_1, α_2, c_1 and c_2 were such that $f' \circ \alpha_i = \beta \circ f + c_i$.
- Then $f' \circ (\alpha_1 - \alpha_2): C \rightarrow J'$ is constant, so $\alpha_1 - \alpha_2$ sends every pair of points in C to the same fibre of $f': C' \rightarrow J'$.
- Hence $(\alpha_1 - \alpha_2)(Q_1) \sim (\alpha_1 - \alpha_2)(Q_2)$ and

$$(*) \quad \alpha_1(Q_1) + \alpha_2(Q_2) \sim \alpha_2(Q_1) + \alpha_1(Q_2) \quad (\forall Q_1, Q_2).$$

Proof — Uniqueness (modulo case distinctions for signs)

- Suppose α_1, α_2, c_1 and c_2 were such that $f' \circ \alpha_i = \beta \circ f + c_i$.
- Then $f' \circ (\alpha_1 - \alpha_2): C \rightarrow J'$ is constant, so $\alpha_1 - \alpha_2$ sends every pair of points in C to the same fibre of $f': C' \rightarrow J'$.
- Hence $(\alpha_1 - \alpha_2)(Q_1) \sim (\alpha_1 - \alpha_2)(Q_2)$ and

$$(*) \quad \alpha_1(Q_1) + \alpha_2(Q_2) \sim \alpha_2(Q_1) + \alpha_1(Q_2) \quad (\forall Q_1, Q_2).$$

- Suppose $\alpha_1 \neq \alpha_2$. Then $\exists Q_1 \in C(k)$ s.t. $\alpha_1(Q_1) \neq \alpha_2(Q_1)$. Since α_1 is an isomorphism, there are also plenty $Q_2 \in C(k)$ s.t. $\alpha_1(Q_1) \neq \alpha_1(Q_2)$, hence $\alpha_1(Q_1) \notin \{\alpha_2(Q_1), \alpha_1(Q_2)\}$.

Proof — Uniqueness (modulo case distinctions for signs)

- Suppose α_1, α_2, c_1 and c_2 were such that $f' \circ \alpha_i = \beta \circ f + c_i$.
- Then $f' \circ (\alpha_1 - \alpha_2): C \rightarrow J'$ is constant, so $\alpha_1 - \alpha_2$ sends every pair of points in C to the same fibre of $f': C' \rightarrow J'$.
- Hence $(\alpha_1 - \alpha_2)(Q_1) \sim (\alpha_1 - \alpha_2)(Q_2)$ and

$$(*) \quad \alpha_1(Q_1) + \alpha_2(Q_2) \sim \alpha_2(Q_1) + \alpha_1(Q_2) \quad (\forall Q_1, Q_2).$$

- Suppose $\alpha_1 \neq \alpha_2$. Then $\exists Q_1 \in C(k)$ s.t. $\alpha_1(Q_1) \neq \alpha_2(Q_1)$. Since α_1 is an isomorphism, there are also plenty $Q_2 \in C(k)$ s.t. $\alpha_1(Q_1) \neq \alpha_1(Q_2)$, hence $\alpha_1(Q_1) \notin \{\alpha_2(Q_1), \alpha_1(Q_2)\}$.
- So the degree 2 linear system given by $(*)$ contains at least two divisors, which implies that it is of dimension at least 1.

Proof — Uniqueness (modulo case distinctions for signs)

- Suppose α_1, α_2, c_1 and c_2 were such that $f' \circ \alpha_i = \beta \circ f + c_i$.
- Then $f' \circ (\alpha_1 - \alpha_2): C \rightarrow J'$ is constant, so $\alpha_1 - \alpha_2$ sends every pair of points in C to the same fibre of $f': C' \rightarrow J'$.
- Hence $(\alpha_1 - \alpha_2)(Q_1) \sim (\alpha_1 - \alpha_2)(Q_2)$ and

$$(*) \quad \alpha_1(Q_1) + \alpha_2(Q_2) \sim \alpha_2(Q_1) + \alpha_1(Q_2) \quad (\forall Q_1, Q_2).$$

- Suppose $\alpha_1 \neq \alpha_2$. Then $\exists Q_1 \in C(k)$ s.t. $\alpha_1(Q_1) \neq \alpha_2(Q_1)$. Since α_1 is an isomorphism, there are also plenty $Q_2 \in C(k)$ s.t. $\alpha_1(Q_1) \neq \alpha_1(Q_2)$, hence $\alpha_1(Q_1) \notin \{\alpha_2(Q_1), \alpha_1(Q_2)\}$.
- So the degree 2 linear system given by $(*)$ contains at least two divisors, which implies that it is of dimension at least 1.
- Varying Q_1 we obtain more such linear systems, and curves of general type can have at most one such linear system.

Proof — Uniqueness (modulo case distinctions for signs)

- Suppose α_1, α_2, c_1 and c_2 were such that $f' \circ \alpha_i = \beta \circ f + c_i$.
- Then $f' \circ (\alpha_1 - \alpha_2): C \rightarrow J'$ is constant, so $\alpha_1 - \alpha_2$ sends every pair of points in C to the same fibre of $f': C' \rightarrow J'$.
- Hence $(\alpha_1 - \alpha_2)(Q_1) \sim (\alpha_1 - \alpha_2)(Q_2)$ and

$$(*) \quad \alpha_1(Q_1) + \alpha_2(Q_2) \sim \alpha_2(Q_1) + \alpha_1(Q_2) \quad (\forall Q_1, Q_2).$$

- Suppose $\alpha_1 \neq \alpha_2$. Then $\exists Q_1 \in C(k)$ s.t. $\alpha_1(Q_1) \neq \alpha_2(Q_1)$. Since α_1 is an isomorphism, there are also plenty $Q_2 \in C(k)$ s.t. $\alpha_1(Q_1) \neq \alpha_1(Q_2)$, hence $\alpha_1(Q_1) \notin \{\alpha_2(Q_1), \alpha_1(Q_2)\}$.
- So the degree 2 linear system given by $(*)$ contains at least two divisors, which implies that it is of dimension at least 1.
- Varying Q_1 we obtain more such linear systems, and curves of general type can have at most one such linear system.
- This contradiction shows that $\alpha_1 = \alpha_2$, thus $c_1 = c_2$.

Corollary — Torelli over a perfect field $F \subseteq k$

If C , C' and β are defined over F , then α is defined over F as well. In particular, $C \cong C'$ over F .

Corollary — Torelli over a perfect field $F \subseteq k$

If C , C' and β are defined over F , then α is defined over F as well. In particular, $C \cong C'$ over F .

Sketch of proof (choose a sign if C hyperelliptic):

(1) α is characterized by: $\exists c \in J(k)$ s.t. $f^{P'} \circ \alpha = \pm \beta \circ f^P + c$.

Corollary — Torelli over a perfect field $F \subseteq k$

If C , C' and β are defined over F , then α is defined over F as well. In particular, $C \cong C'$ over F .

Sketch of proof (choose a sign if C hyperelliptic):

- (1) α is characterized by: $\exists c \in J(k)$ s.t. $f^{P'} \circ \alpha = \pm \beta \circ f^P + c$.
- (2) Replacing P by Q we get $f^Q = f^P + d$, resp. for $(-)'$. Hence

$$f^{Q'} \circ \alpha = f^{P'} \circ \alpha + d' = \pm \beta \circ f^P + c + d' = \pm \beta \circ f^Q \mp \beta(d) + c + d'.$$

Corollary — Torelli over a perfect field $F \subseteq k$

If C , C' and β are defined over F , then α is defined over F as well. In particular, $C \cong C'$ over F .

Sketch of proof (choose a sign if C hyperelliptic):

- (1) α is characterized by: $\exists c \in J(k)$ s.t. $f^{P'} \circ \alpha = \pm \beta \circ f^P + c$.
- (2) Replacing P by Q we get $f^Q = f^P + d$, resp. for $(-)'$. Hence

$$f^{Q'} \circ \alpha = f^{P'} \circ \alpha + d' = \pm \beta \circ f^P + c + d' = \pm \beta \circ f^Q \mp \beta(d) + c + d'.$$

- (3) Hence α is independent of the chosen k -points P and P' .

Corollary — Torelli over a perfect field $F \subseteq k$

If C , C' and β are defined over F , then α is defined over F as well. In particular, $C \cong C'$ over F .

Sketch of proof (choose a sign if C hyperelliptic):

- (1) α is characterized by: $\exists c \in J(k)$ s.t. $f^{P'} \circ \alpha = \pm \beta \circ f^P + c$.
- (2) Replacing P by Q we get $f^Q = f^P + d$, resp. for $(-)'$. Hence

$$f^{Q'} \circ \alpha = f^{P'} \circ \alpha + d' = \pm \beta \circ f^P + c + d' = \pm \beta \circ f^Q \mp \beta(d) + c + d'.$$


- (3) Hence α is independent of the chosen k -points P and P' .
- (4) For $\sigma \in \text{Gal}(k/F)$ we have $\sigma f^P = f^{\sigma P}$, resp. for $(-)'$. Hence

$$f^{\sigma P'} \circ \sigma \alpha = \sigma f^{P'} \circ \sigma \alpha = \pm \sigma \beta \circ \sigma f^P + \sigma c = \pm \beta \circ f^{\sigma P} + \sigma c.$$

Point (3) implies then $\sigma \alpha = \alpha$.


Idea of the existence proof for $g = 2$

Suppose $g = 2$ as in the previous picture. If we identify J with J' via β , the fact that β is a polarized isomorphism guarantees the following situation:




Idea of the existence proof for $g > 2$

If $g > 2$, the fact that β is a polarized isomorphism guarantees a priori only that Θ' is a translation Θ_a of Θ by some $a \in J(k)$:



Idea of the existence proof for $g > 2$

If $g > 2$, the fact that β is a polarized isomorphism guarantees a priori only that Θ' is a translation Θ_a of Θ by some $a \in J(k)$:



Is $V_a^1 = W_b^1$ for some $b \in J(k)$?

[Yes! Modulo replacing W^1 by its image under $x \mapsto -x$.]

Thanks for listening!