

TITLE

PEDRO NÚÑEZ

CONTENTS

1. First section	1
References	1

1. FIRST SECTION

Let \mathcal{F} be a coherent sheaf on X . Then

$$\chi(\mathcal{F}) = \sum_{i=1}^{\dim X} (-1)^i h^i(X, \mathcal{F}).$$

Definition 1.1. First definition!

Fact. *This is an unnumbered fact.*

Theorem A (Custom numbered theorem). *This theorem has custom numbering.*

References included in the .bib file by default are [Har77] and [Sta20].

REFERENCES

- [Har77] Robin Hartshorne. *Algebraic geometry*. Springer-Verlag, New York-Heidelberg, 1977. Graduate Texts in Mathematics, No. 52. ↑ 1
- [Sta20] The Stacks project authors. The stacks project. <https://stacks.math.columbia.edu>, 2020. ↑ 1

PEDRO NÚÑEZ

ALBERT-LUDWIGS-UNIVERSITÄT FREIBURG, MATHEMATISCHES INSTITUT
ERNST-ZERMELO-STRASSE 1, 79104 FREIBURG IM BREISGAU (GERMANY)

Email address: pedro.nunez@math.uni-freiburg.de

Homepage: <https://home.mathematik.uni-freiburg.de/nunez>

Date: 4th September 2020.

The author gratefully acknowledges support by the DFG-Graduiertenkolleg GK1821 “Cohomological Methods in Geometry” at the University of Freiburg.