

Derived Categories and Birational Geometry

Pedro Núñez


National Taiwan University

September 22, 2025

Mathematics Seminar at CUNEF University


Outline

- 1 Introduction
 - Birational Geometry
 - Derived Categories
- 2 Relation between Derived Categories and Birational Geometry
 - Relation between them: Kawamata's DK hypothesis
 - Indecomposability conjecture and known results
- 3 Previous and current work on indecomposability
 - Hyperelliptic varieties
(joint work with Pieter Belmans and Andreas Demleitner)
 - Threefolds on the Noether Line
(joint work in progress with Jungkai Chen)




Outline

- 1 Introduction
 - Birational Geometry
 - Derived Categories
- 2 Relation between Derived Categories and Birational Geometry
 - Relation between them: Kawamata's DK hypothesis
 - Indecomposability conjecture and known results
- 3 Previous and current work on indecomposability
 - Hyperelliptic varieties
(joint work with Pieter Belmans and Andreas Demleitner)
 - Threefolds on the Noether Line
(joint work in progress with Jungkai Chen)




A set of small navigation icons typically found in Beamer presentations, including symbols for back, forward, search, and other slide controls.

Projective algebraic varieties




$$\{(x, y) \in \mathbb{A}^2 \mid y = x^2 - 1\}$$




$$\{[x : y : z] \in \mathbb{P}^2 \mid yz = x^2 - z^2\}$$

- Work over \mathbb{C} , because we want FTA, Bézout's theorem, etc.

Projective algebraic varieties




$$\{(x, y) \in \mathbb{A}^2 \mid y = x^2 - 1\}$$




$$\{[x : y : z] \in \mathbb{P}^2 \mid yz = x^2 - z^2\}$$

- Work over \mathbb{C} , because we want FTA, Bézout's theorem, etc.
- Zariski topology: closed subsets are zero loci of polynomials.

Projective algebraic varieties




$$\{(x, y) \in \mathbb{A}^2 \mid y = x^2 - 1\}$$




$$\{[x : y : z] \in \mathbb{P}^2 \mid yz = x^2 - z^2\}$$

- Work over \mathbb{C} , because we want FTA, Bézout's theorem, etc.
- Zariski topology: closed subsets are zero loci of polynomials.
- Work with projective varieties, because we want compactness.

Projective algebraic varieties



$$\{(x, y) \in \mathbb{A}^2 \mid y = x^2 - 1\}$$



$$\{[x : y : z] \in \mathbb{P}^2 \mid yz = x^2 - z^2\}$$

- Work over \mathbb{C} , because we want FTA, Bézout's theorem, etc.
- Zariski topology: closed subsets are zero loci of polynomials.
- Work with projective varieties, because we want compactness.
- We assume varieties to be irreducible.

Birational equivalence

- A morphism of algebraic varieties is a continuous map which locally looks polynomial. So $x \mapsto x^2$ is, but $x \mapsto e^x$ isn't.

Birational equivalence

- A morphism of algebraic varieties is a continuous map which locally looks polynomial. So $x \mapsto x^2$ is, but $x \mapsto e^x$ isn't.
- An isomorphism of algebraic varieties is a morphism which is invertible (the inverse should be algebraic as well).

Birational equivalence

- A morphism of algebraic varieties is a continuous map which locally looks polynomial. So $x \mapsto x^2$ is, but $x \mapsto e^x$ isn't.
- An isomorphism of algebraic varieties is a morphism which is invertible (the inverse should be algebraic as well).
- **Goal:** Classify varieties up to isomorphism. (Too hard!)

Birational equivalence

- A morphism of algebraic varieties is a continuous map which locally looks polynomial. So $x \mapsto x^2$ is, but $x \mapsto e^x$ isn't.
- An isomorphism of algebraic varieties is a morphism which is invertible (the inverse should be algebraic as well).
- **Goal:** Classify varieties up to isomorphism. (Too hard!)
- Two varieties are *birationally equivalent* if they contain isomorphic dense open subsets, i.e., they are the same except possibly over some (lower-dimensional) proper closed subset.

Birational equivalence

- A morphism of algebraic varieties is a continuous map which locally looks polynomial. So $x \mapsto x^2$ is, but $x \mapsto e^x$ isn't.
- An isomorphism of algebraic varieties is a morphism which is invertible (the inverse should be algebraic as well).
- **Goal:** Classify varieties up to isomorphism. (Too hard!)
- Two varieties are *birationally equivalent* if they contain isomorphic dense open subsets, i.e., they are the same except possibly over some (lower-dimensional) proper closed subset.
- Isomorphic varieties are birationally equivalent, but not vice-versa. (Example: blow-up.)

Birational equivalence

- A morphism of algebraic varieties is a continuous map which locally looks polynomial. So $x \mapsto x^2$ is, but $x \mapsto e^x$ isn't.
- An isomorphism of algebraic varieties is a morphism which is invertible (the inverse should be algebraic as well).
- **Goal:** Classify varieties up to isomorphism. (Too hard!)
- Two varieties are *birationally equivalent* if they contain isomorphic dense open subsets, i.e., they are the same except possibly over some (lower-dimensional) proper closed subset.
- Isomorphic varieties are birationally equivalent, but not vice-versa. (Example: blow-up.)
- **Intermediate goal:** Classify up to birational equivalence.

Canonical line bundle/divisor

- If X is an n -dimensional smooth variety, its *canonical line bundle* is

$\omega_X := \bigwedge^n \Omega_X$, where Ω_X is its (holomorphic) cotangent bundle.

Canonical line bundle/divisor

- If X is an n -dimensional smooth variety, its *canonical line bundle* is

$\omega_X := \bigwedge^n \Omega_X$, where Ω_X is its (holomorphic) cotangent bundle.

- A divisor is a \mathbb{Z} -linear combination of codimension 1 subvarieties. E.g., on a curve, \mathbb{Z} -linear combination of points.

Canonical line bundle/divisor

- If X is an n -dimensional smooth variety, its *canonical line bundle* is

$$\omega_X := \bigwedge^n \Omega_X, \text{ where } \Omega_X \text{ is its (holomorphic) cotangent bundle.}$$

- A divisor is a \mathbb{Z} -linear combination of codimension 1 subvarieties. E.g., on a curve, \mathbb{Z} -linear combination of points.
- The *canonical divisor* K_X is the divisor of zeros and poles of any non-zero rational section of ω_X .

Canonical line bundle/divisor

- If X is an n -dimensional smooth variety, its *canonical line bundle* is

$\omega_X := \bigwedge^n \Omega_X$, where Ω_X is its (holomorphic) cotangent bundle.

- A divisor is a \mathbb{Z} -linear combination of codimension 1 subvarieties. E.g., on a curve, \mathbb{Z} -linear combination of points.
- The *canonical divisor* K_X is the divisor of zeros and poles of any non-zero rational section of ω_X .

Example: On $\mathbb{P}^1 = \{[x_0 : x_1] \mid (x_0, x_1) \neq (0, 0)\}$ we have coordinates $x := x_1/x_0$ when $x_0 \neq 0$ and $y := x_0/x_1$ when $x_1 \neq 0$. The rational differential form $dx = d(y^{-1}) = -y^{-2}dy$ has a pole of order 2 at the point $H := \{x_0 = 0\} = \{y = 0\}$, hence $K_{\mathbb{P}^1} = -2H$.

Minimal Model Program (MMP)

- To classify up to birational equivalence, we want to pick a single representative X' in the birational equivalence class $[X]$.

Minimal Model Program (MMP)

- To classify up to birational equivalence, we want to pick a single representative X' in the birational equivalence class $[X]$.
- If $\pi: \tilde{X} \rightarrow X$ is a blow-up, $\tilde{X} \sim_{\text{bir}} X$. Among them, X is simpler.

Minimal Model Program (MMP)

- To classify up to birational equivalence, we want to pick a single representative X' in the birational equivalence class $[X]$.
- If $\pi: \tilde{X} \rightarrow X$ is a blow-up, $\tilde{X} \sim_{\text{bir}} X$. Among them, X is simpler.
- We have the relation $K_{\tilde{S}} = \pi^* K_S + E$, and $K_{\tilde{S}} \cdot E = -1$.

Minimal Model Program (MMP)

- To classify up to birational equivalence, we want to pick a single representative X' in the birational equivalence class $[X]$.
- If $\pi: \tilde{X} \rightarrow X$ is a blow-up, $\tilde{X} \sim_{\text{bir}} X$. Among them, X is simpler.
- We have the relation $K_{\tilde{S}} = \pi^* K_S + E$, and $K_{\tilde{S}} \cdot E = -1$.
- **MMP's idea:** Look for curves C such that $K_X \cdot C < 0$. If you find one, you can contract it (Castelnuovo/Mori). Then repeat.

Minimal Model Program (MMP)

- To classify up to birational equivalence, we want to pick a single representative X' in the birational equivalence class $[X]$.
- If $\pi: \tilde{X} \rightarrow X$ is a blow-up, $\tilde{X} \sim_{\text{bir}} X$. Among them, X is simpler.
- We have the relation $K_{\tilde{S}} = \pi^* K_S + E$, and $K_{\tilde{S}} \cdot E = -1$.
- **MMP's idea:** Look for curves C such that $K_X \cdot C < 0$. If you find one, you can contract it (Castelnuovo/Mori). Then repeat.
- Conjecturally, this process terminates. The variety we are left with at the end is our chosen representative.

Minimal Model Program (MMP)


- To classify up to birational equivalence, we want to pick a single representative X' in the birational equivalence class $[X]$.
- If $\pi: \tilde{X} \rightarrow X$ is a blow-up, $\tilde{X} \sim_{\text{bir}} X$. Among them, X is simpler.
- We have the relation $K_{\tilde{S}} = \pi^* K_S + E$, and $K_{\tilde{S}} \cdot E = -1$.
- **MMP's idea:** Look for curves C such that $K_X \cdot C < 0$. If you find one, you can contract it (Castelnuovo/Mori). Then repeat.
- Conjecturally, this process terminates. The variety we are left with at the end is our chosen representative.

Definition: A projective variety X is called *minimal* when K_X has non-negative intersection with every irreducible curve in X .

Outline

1 Introduction

- Birational Geometry
- Derived Categories



From singular cohomology to sheaf cohomology


From sheaf cohomology to derived categories

Semiorthogonal decompositions (SOD)

- The category $D^b(X)$ has a natural *triangulated structure*.
- An *orthogonal decomposition* of $D^b(X)$ would consist of
 - triangulated subcategories $\mathcal{A}, \mathcal{B} \subseteq D^b(X)$;
 - such that $\text{Hom}(\mathcal{A}, \mathcal{B}) = \text{Hom}(\mathcal{B}, \mathcal{A}) = 0$, i.e.,
 $\text{Hom}(a, b) = \text{Hom}(b, a) = 0$ for all $a \in \mathcal{A}$ and all $b \in \mathcal{B}$;
 - and such that the smallest triangulated subcategory of $D^b(X)$ containing both of them is $D^b(X)$ itself.
- **Fact:** If X is connected, then $D^b(X)$ does not admit any orthogonal decomposition. (Bridgeland '99.)
- A *semiorthogonal decomposition* is the same thing, but without requiring $\text{Hom}(\mathcal{A}, \mathcal{B}) = 0$, only requiring $\text{Hom}(\mathcal{B}, \mathcal{A}) = 0$. It is denoted

$$D^b(X) = \langle \mathcal{A}, \mathcal{B} \rangle.$$

Outline




1 Introduction

- Birational Geometry
- Derived Categories

2 Relation between Derived Categories and Birational Geometry

- Relation between them: Kawamata's DK hypothesis
- Indecomposability conjecture and known results

Kawamata's DK hypothesis



Example: blow-up

$$\begin{array}{ccc}
 E & \xhookrightarrow{i} & \tilde{S} \\
 \pi \downarrow & & \downarrow q \\
 \{p\} & \xhookrightarrow{j} & S
 \end{array}$$

$$q: \tilde{S} = \mathrm{Bl}_p(S) \rightarrow S$$

$$\Leftrightarrow$$

$$\mathrm{D}^b(\tilde{S}) = \langle \mathrm{D}^b(p), \mathrm{D}^b(S) \rangle$$

$$\updownarrow$$

$$\updownarrow$$

$$\begin{aligned}
 K_{\tilde{S}} &= q^* K_S + E \\
 (K_{\tilde{S}} &\geq K_S)
 \end{aligned}$$


$$\Leftrightarrow$$

$$\begin{aligned}
 &q^*(\mathrm{D}^b(S)) \\
 &i_*(\mathcal{O}_E(-E) \otimes \pi^* \mathrm{D}^b(p))
 \end{aligned}$$

Outline

1 Introduction

- Birational Geometry
- Derived Categories



2 Relation between Derived Categories and Birational Geometry

- Relation between them: Kawamata's DK hypothesis
- Indecomposability conjecture and known results

Indecomposability conjecture

The previous discussion suggests that

minimal varieties \Leftrightarrow indecomposable derived categories.

Indecomposability conjecture

The previous discussion suggests that

minimal varieties \Leftrightarrow indecomposable derived categories.

This is not strictly true, but the following is a folklore conjecture:

Conjecture

Let X be a minimal smooth projective variety with $p_g > 0$.
Then $D^b(X)$ is indecomposable.

Main known results on indecomposability

- Bridgeland '99: Calaby–Yau varieties have indecomposable derived categories.

Main known results on indecomposability

- Bridgeland '99: Calaby–Yau varieties have indecomposable derived categories.
- Kawatani–Okawa '18: the base locus of the canonical linear system controls indecomposability.

Main known results on indecomposability

- Bridgeland '99: Calaby–Yau varieties have indecomposable derived categories.
- Kawatani–Okawa '18: the base locus of the canonical linear system controls indecomposability.
- Pirozhkov '23: stronger notion of indecomposability (NSSI); examples are finite covers of abelian varieties and varieties fibered in NSSI varieties over NSSI bases.

Main known results on indecomposability

- Bridgeland '99: Calaby–Yau varieties have indecomposable derived categories.
- Kawatani–Okawa '18: the base locus of the canonical linear system controls indecomposability.
- Pirozhkov '23: stronger notion of indecomposability (NSSI); examples are finite covers of abelian varieties and varieties fibered in NSSI varieties over NSSI bases.

Theorem (Kawatani–Okawa '18, Okawa '23, Pirozhkov '25, ...)

A minimal smooth projective surface has indecomposable derived category if and only if $(p_g, q) \neq (0, 0)$.

◀ ◻ ▶ ◀ ◻ ▶ ◀ ≡ ▶ ◀ ≡ ▶ ≡ ↺ 🔍 ↻

Hyperelliptic varieties: definition

- A *hyperelliptic variety* $X = A/G$ is the quotient of an abelian variety A by a finite group of automorphisms $G \subseteq \text{Aut}(A)$ acting freely and without translations on A .

Hyperelliptic varieties: definition

- A *hyperelliptic variety* $X = A/G$ is the quotient of an abelian variety A by a finite group of automorphisms $G \subseteq \operatorname{Aut}(A)$ acting freely and without translations on A .
- It follows that they are smooth projective minimal varieties with torsion canonical divisor, i.e., $mK_X \sim 0$ for some $m \in \mathbb{Z}_{>0}$.

Hyperelliptic varieties: definition

- A *hyperelliptic variety* $X = A/G$ is the quotient of an abelian variety A by a finite group of automorphisms $G \subseteq \operatorname{Aut}(A)$ acting freely and without translations on A .
- It follows that they are smooth projective minimal varieties with torsion canonical divisor, i.e., $mK_X \sim 0$ for some $m \in \mathbb{Z}_{>0}$.
- Equivalently, they are smooth projective varieties which are not abelian but admit an abelian variety as a finite étale cover.

Hyperelliptic varieties: definition

- A *hyperelliptic variety* $X = A/G$ is the quotient of an abelian variety A by a finite group of automorphisms $G \subseteq \operatorname{Aut}(A)$ acting freely and without translations on A .
- It follows that they are smooth projective minimal varieties with torsion canonical divisor, i.e., $mK_X \sim 0$ for some $m \in \mathbb{Z}_{>0}$.
- Equivalently, they are smooth projective varieties which are not abelian but admit an abelian variety as a finite étale cover.
- 1-dimensional hyperelliptic varieties do not exist, and 2-dimensional hyperelliptic varieties are bielliptic surfaces.

Hyperelliptic varieties: conjecture and main result

Conjecture

Let X be a hyperelliptic variety. Then $D^b(X)$ is indecomposable.

Hyperelliptic varieties: conjecture and main result

Conjecture

Let X be a hyperelliptic variety. Then $D^b(X)$ is indecomposable.

The *irregularity* of X is $q_X := h^1(X, \mathcal{O}_X)$ ($< \dim X$ if X hyperelliptic).

Theorem

The conjecture holds in the following cases:

1. X is cyclic, i.e., $X = A/G$ with G cyclic.
2. X has irregularity $q_X = \dim X - 2$ or $\dim X - 1$.
3. The fiber(s) of the Albanese morphism of X have trivial canonical bundle.

In particular, the conjecture holds if $\dim X \leq 3$.

Main approach: Albanese morphism + induction

- The Albanese morphism is a universal morphism into an abelian variety $\mathrm{alb}_X: X \rightarrow \mathrm{Alb}(X)$.
- By [Kawamata '85], if X is hyperelliptic, then the Albanese morphism is an étale fiber bundle with smooth connected fibers.

Main approach: Albanese morphism + induction

- The Albanese morphism is a universal morphism into an abelian variety $\mathrm{alb}_X: X \rightarrow \mathrm{Alb}(X)$.
- By [Kawamata '85], if X is hyperelliptic, then the Albanese morphism is an étale fiber bundle with smooth connected fibers.
- In our paper we show that the fibers are either abelian varieties or hyperelliptic varieties again.

Main approach: Albanese morphism + induction

- The Albanese morphism is a universal morphism into an abelian variety $\mathrm{alb}_X: X \rightarrow \mathrm{Alb}(X)$.
- By [Kawamata '85], if X is hyperelliptic, then the Albanese morphism is an étale fiber bundle with smooth connected fibers.
- In our paper we show that the fibers are either abelian varieties or hyperelliptic varieties again.
- Combining this with [Pirozhkov '23] and induction on the dimension, we can deduce indecomposability in the first two cases of the theorem.

- Birational Geometry
- Derived Categories

- Hyperelliptic varieties
(joint work with Pieter Belmans and Andreas Demleitner)
- Threefolds on the Noether Line
(joint work in progress with Jungkai Chen)

Threefolds on the Noether Line: definition

Two key birational invariants of a projective variety X are its geometric genus $p_g(X) := h^0(X, \omega_X)$ and its canonical volume

$$\mathrm{vol}(X) := \lim_{m \rightarrow \infty} \frac{h^0(X, \omega_X^{\otimes m})}{m^n / n!}, \quad \text{where } n := \dim(X).$$

Threefolds on the Noether Line: definition

Two key birational invariants of a projective variety X are its geometric genus $p_g(X) := h^0(X, \omega_X)$ and its canonical volume

$$\mathrm{vol}(X) := \lim_{m \rightarrow \infty} \frac{h^0(X, \omega_X^{\otimes m})}{m^n / n!}, \quad \text{where } n := \dim(X).$$

By work of Jungkai Chen, Meng Chen and Chen Jiang ('20), and others, we know that projective threefolds of general type satisfy

$$\mathrm{vol}(X) \geq \frac{4}{3} p_g(X) - \frac{10}{3} \quad (\text{Noether Inequality}).$$

Threefolds on the Noether Line: definition

Two key birational invariants of a projective variety X are its geometric genus $p_g(X) := h^0(X, \omega_X)$ and its canonical volume


$$\mathrm{vol}(X) := \lim_{m \rightarrow \infty} \frac{h^0(X, \omega_X^{\otimes m})}{m^n / n!}, \quad \text{where } n := \dim(X).$$

By work of Jungkai Chen, Meng Chen and Chen Jiang ('20), and others, we know that projective threefolds of general type satisfy

$$\mathrm{vol}(X) \geq \frac{4}{3} p_g(X) - \frac{10}{3} \quad (\text{Noether Inequality}).$$

Definition: A projective threefold of general type is said to be *on the Noether line* if equality holds above.

Kobayashi's construction ('92)



Threefolds on the Noether Line: current result

Theorem

Let X be a *general*^{*} minimal smooth projective threefold on the first^{**} Noether Line. Then $D^b(X)$ is indecomposable.

Threefolds on the Noether Line: current result


Theorem

Let X be a *general*^{*} minimal smooth projective threefold on the first^{**} Noether Line. Then $D^b(X)$ is indecomposable.

- * The moduli space of such threefolds has several irreducible components, and this statement applies to one of the top-dimensional irreducible components.
- ** There are three Noether Lines, and threefolds on the second and third Noether Lines are necessarily singular.

A set of small navigation icons typically found in Beamer presentations, including symbols for back, forward, search, and other slide controls.

Current work in progress: singular cases



Thanks for your attention!