

# Lecture slides with L<sup>A</sup>T<sub>E</sub>X

A user friendly template

**Pedro Fonseca**



**ISEG Lisbon School of Economics & Management**

April 24, 2020

# Contents



- 1 Introduction
- 2 Some complicated formulas
- 3 Code chunks
- 4 Conclusion
- 5 References

# Before we start



You should remember the following topics from last lecture:

- Interesting topic 1
  - ▶ Interesting sub topic 1
  - ▶ Interesting sub topic 2
  - ▶ Interesting sub topic 3
- Interesting topic 2
- Interesting topic 3

# Before we start




In this lecture you will see:

- a plot
- some formulas
- some code
- some citations

# including pictures



This picture is in the “figures” folder:



# Mathematical expressions



- Latex is very useful for expressions.
- Which is nice, since this lecture has lots of them!
- Lets see some Bayesian model selection formulas.

# Multinomial $\wedge$ Dirichlet Model



- Likelihood:  $\mathbf{x}|\boldsymbol{\theta} \sim \mathcal{M}_k(N, \boldsymbol{\theta})$
- Hypotheses:  $H_0 : \boldsymbol{\theta} = \boldsymbol{\theta}_0$  vs  $H_1 : \boldsymbol{\theta} \neq \boldsymbol{\theta}_0$
- Prior:  $\pi(\boldsymbol{\theta}|H_0) = \mathbf{1}_{\boldsymbol{\theta}_0}(\boldsymbol{\theta}_0)$  and  $\boldsymbol{\theta}|H_1, \boldsymbol{\alpha} \sim \text{Dir}_k(\boldsymbol{\alpha})$
- Marginal density:  $m_i(\mathbf{x}) = \int_{\Theta_i} f(\mathbf{x}|\boldsymbol{\theta})\pi(\boldsymbol{\theta}|H_i) d\boldsymbol{\theta}, \quad i = 0, 1$

$$\blacktriangleright m_0(\mathbf{x}) = \frac{N!}{\prod_{i=1}^{k+1} x_i!} \prod_{i=1}^{k+1} \theta_{0i}^{x_i}$$

$$\blacktriangleright m_1(\mathbf{x}) = \frac{N!}{\prod_{i=1}^{k+1} x_i!} \frac{B(\boldsymbol{\alpha} + \mathbf{x})}{B(\boldsymbol{\alpha})}$$

- Bayes factor:

$$B_{01}(\mathbf{x}) = \frac{m_0(\mathbf{x})}{m_1(\mathbf{x})} = \frac{\prod_{i=1}^{k+1} (\theta_{0i}^{x_i}) \prod_{i=1}^{k+1} [\Gamma(\alpha_i)] \Gamma[\sum_{i=1}^{k+1} (\alpha_i + x_i)]}{\Gamma(\sum_{i=1}^{k+1} \alpha_i) \prod_{i=1}^{k+1} \Gamma(\alpha_i + x_i)}$$

# Binomial $\wedge$ Beta Model



- Likelihood:  $x \sim \text{Bin}(N, \theta)$
- Hypotheses:  $H_0 : \theta = \theta_0$  vs  $H_1 : \theta \neq \theta_0$
- Prior:  $\pi(\theta|H_0) = 1_{\theta_0}(\theta)$  and  $\theta|H_1 \sim \text{Beta}(a,b)$
- Marginal density:  $m_i(x) = \int_0^1 f(x|\theta)\pi(\theta|H_i)d\theta, i = 0, 1$ 
  - ▶  $m_0(x) = \binom{N}{x} \theta_0^x (1 - \theta_0)^{N-x}$
  - ▶  $m_1(x) = \binom{N}{x} \frac{B(x+a, n-x+b)}{B(a,b)}$
- Bayes factor:  $B_{01}(x) = \frac{m_0(x)}{m_1(x)} = \frac{\theta_0(1-\theta_0)^{N-x} \Gamma(a) \Gamma(b) \Gamma(n+a+b)}{\Gamma(a+b) \Gamma(n+a-x) \Gamma(x+a)}$



# Want to know more?



- To learn about Bayes factors see Kass and Raftery, 1995.
- For an interesting application of the Multinomial  $\wedge$  Dirichlet model see Pericchi and Torres (2011).

## Some python code



This lecture also has some examples with code. Lets print the Fibonacci sequence:

```
nterms = int(input("How many terms? "))
n1, n2 = 0, 1
count = 0
print("Fibonacci sequence:")
while count < nterms:
    print(n1)
    nth = n1 + n2
    n1 = n2
    n2 = nth
    count += 1
}
```

# Now some R code



There is also a verbatim environment with background color.

```
getmode <- function(v) {  
  
  uniqv <- unique(v)  
  
  uniqv[which.max(tabulate(match(v, uniqv)))]  
  
}
```

With this R code you can build a function that calculates the mode.

# summary



- we talked about

# References



- Kass, RE and AE Raftery (1995). Bayes factors. *Journal of the American Statistical Association* **90**(430), 773–795.
- Pericchi, L and D Torres (2011). Quick anomaly detection by the Newcomb—Benford law, with applications to electoral processes data from the USA, Puerto Rico and Venezuela. *Statistical Science* **26**(4), 502–516.