

UNIVERSIDADE DE BRASÍLIA
FACULDADE DE ECONOMIA, ADMINISTRAÇÃO E CONTABILIDADE
DEPARTAMENTO DE ECONOMIA
PROGRAMA DE PÓS-GRADUAÇÃO EM ECONOMIA

High Dimensional Models for Forecast Power Electricity Consumption

Pedro Caiua Campelo Albuquerque

Orientador: Marina Rossi

Brasília
2019

Resumo

Resumo do trabalho.

Key-words: Power Electricity. Machine Learning. Forecast.

Abstract

This present work uses machine learning models to predict power electricity consumption, for 6 different time horizons. We observed that the machine learning models have superior results than the original econometric models, by removing the overfitting present in these models. The best predictions was from Random Forest model, that managed to keep up with the trend and the seasonality of the different time horizons. With this model, we are able to capture the trend of power electricity consumption and make inferences about economic activity, due to the positive relationship between the two variables. Our work further concludes that lagged consumption values are extremely important for very short-term forecasting (1 and 7 days) and calendar and meteorological variables become more important for forecasting as it increases the forecast horizon. Price and economic variables are more relevant for 7 and 15 days ahead forecast.

Key-words: Power Electricity. Machine Learning. Forecast.

Contents

1	Introduction	5
2	Econometric Models	7
2.1	Model Structure	7
2.2	Models	8
2.2.1	ARIMA	8
2.2.2	Random Walk	9
2.2.3	Lasso	9
2.2.4	Lars	9
2.2.5	Lasso Lars	10
2.2.6	Ridge	10
2.2.7	Elastic Net	11
2.2.8	Random Forest	11
3	Data	12
3.1	Power Electricity Consumption	14
3.2	Calendar Variables	16
3.3	Meteorological Variables	16
3.4	Price of Electric Energy	18
3.5	Economic Variables	19
4	Results	21
4.1	Very Short-Term Forecast Group	23
4.1.1	1 day	23
4.1.2	7 days	24
4.2	Short-Term Forecast Group	25
4.2.1	15 days	26
4.2.2	30 days	27
4.3	Medium-Term Forecast Group	27
4.3.1	60 days	28
4.3.2	90 days	29

5 Conclusion	32
---------------------	-----------

Bibliography	34
---------------------	-----------

1 Introduction

[Colocar motivacao do mundo real de pqp prever consumo de energia elétrica]

[O que eu estou fazeno aqui, depois segue uma breve introducao]

[falar um pouco do Medeiros aqui] Maza and Villaverde (2008) review the bibliography to understand the causality of electric energy and economic growth. The relationship can be divided into 4 categories: causality from electricity consumption to GDP growth, causality from GDP growth to electricity consumption, bidirectional causality and no causality at all. Literature does not necessarily converge to a specific result. Although a vast majority finds a relation between these two variables, the direction of this causality is not so clear. The authors argue that the newer and better controlled models point causality from electricity consumption to GDP growth.

(??) Still, most of the electricity consumption today is consumed by industrial and commercial activities. Therefore, the main objective of this work is to forecast electricity consumption in the short term and in the medium term, and thus, be able to measure a tendency for economic activity. With GDP released quarterly, we want to structure the forecast models of electricity consumption to capture the trend of economic activity.

[There are] Huang and Shih (2003) and Almeshaiei and Hassan Soltan (2011) uses just lagged demand to forecast short-term of energy load via ARMA Model. Rodrigues *et. al.* (2014) also uses just past values to forecast the daily and hourly energy consumption via Artificial Neural Networks. Lebotsa *et. al.* (2018) and Fan and Hyndman (2012) includes calendar and meteorological variables to forecast PEC.

Hence, its a fact that the variable selection for power electricity consumption forecast is not trivial. So, we prefer to build a high dimensional database that includes all the main variables and let the machine learning models choose the most relevant variables for each horizon prediction.

We construct a high dimensional database with more than 1500 variables, including lagged demand, calendar variables, meteorological variables, price of electric energy

variables and economic variables. We use as many traditional models as the machine learning models, expecting better performance for the machine learning models, by removing the overfitting present in traditional econometric models. We run our models for 6 different forecast horizons.

We also want to analyze which variables are most relevant for each forecast horizon, and thus, enable an improvement in the future variable selection to predict the power electricity consumption.

In Section 2, we present our model – describing in-depth the structure and all models used to forecast power electricity consumption. In Section 3, we exhibit our data set and analyze all sets of variables used. In Section 4, we show our general results and examine the best models. Also, we observe which variable are more importantes to predict PEC in different forecasts horizons. Lastly, in Section 5, we present a brief conclusion concerning our findings.

2 Econometric Models

2.1 Model Structure

This section delineates the empirical methods used in this paper for forecasting future power electricity consumption (PEC). As we said, electricity consumption is a high seasonal variable, so we consider that the forecast of PEC h , periods ahead is modeled as a function of blocks of predictors in different lag times. Accordingly, we used three lags of the candidate variables divided into five groups aligned with the literature of economic activity and power electricity forecast. The models were estimated using a rolling window of 546 observations.

The Brazilian GDP is released quarterly, hence the main idea of this paper, is to measure economic activity in the period that the data is not released yet. Finally, we estimated the models for 6 different forecast horizons, 1, 7, 15, 30, 60 and 90 days ahead. Moreover, we have divided the forecast horizons into two groups: (i) Short-Term Forecast Group (STFG), which contains 1, 7 and 15 days ahead and (ii) Medium-Term Forecast Group (MTFG), which contains 30, 60 and 90 days ahead. Our estimated equation is given by:

$$y_{t+h} = \alpha_0 + \sum_{i=0}^2 \gamma_i y_{t-(h*i)} + \sum_{i=0}^2 \beta_i X_{t-(h*i)} + u_{t+h}$$

where y_t is the power electricity consumption (KM/h) in Brazil at time t , α_0 is a constant term, X_t is a matrix containing all candidate variables, $\mathbf{B} = [\gamma, \beta]$ is a vector with all the linear parameters and u_t is an error term.

The lags structure means that if we make a forecast for $t + h$ periods ahead, we would use the variables from t , $t - h$ and $t - 2h$. That is, if we want a 7 day ahead forecast, we would use data from today and from past 7 and 14 days. This structure was the first hyper-parameters defined. We had tested for several other lags structures and this one had the minor forecast error. Which is quite intuitive, because PEC is a seasonal variable

and if we define a lag structure very close, like 1, 2 and 3 days past, we would lost forecast accuracy for picking close days.

Our explanatory variables are divided into five sets aligned to the literature of economic activity and power electricity consumption forecast. The first set, represented by the vector yt_h , are the lagged demand of power electricity. The next four sets are in the X_t vector, that is, $x_t^1, x_t^2, x_t^3, x_t^4 \in X_t$. These sets are, respectively, calendar variables, meteorological variables, price of electric energy and economic variables. Section 3 will better explain each set of variable.

2.2 Models

This section exhibit and explain all the models used to predict PEC, with the structure showed in last section. Shyh-Jier Huang *et al.* (2003*) forecast short-term load using of autoregressive moving average (ARMA). Medeiros *et al.* (2017) use high-dimensional and machine learning models to forecast inflation in real-time, and concluded that these models perform better with big data than traditional models*. Following this line, we use several models to forecast PEC and analyse which perform better. Our initial predict models, ARIMA and Random Walk, use only lagged consumption variables. Then, we begin to introduce the explanatory variables and to predict with OLS. After that, we use some machine learning models, such as Lasso, Lars, Lasso Lars, Ridge, Elastic Net and Random Forest. All models are explained below.

2.2.1 ARIMA

Autoregressive integrated moving average (ARIMA) uses just lagged demand of the variable of interest. The ARIMA (p,i,q) equation can be represented by

$$y_t^* = \gamma + \Phi_0 + \Phi_1 y_{t-1}^* + \cdots + \Phi_p y_{t-p}^* + u_t + \theta_1 u_{t-1} + \cdots + \theta_q u_{t-q}$$

where $y_t^* = y_t - y_{t-1}$, because our serial data is non-stationary and the first difference is.

2.2.2 Random Walk

Our random walk model (without drift) represents the last value we have available for forecast. That is, the best value for forecast k periods ahead is the value available today. The model can be represented by

$$y_t = y_{t-k} + u_t$$

This model is a good measure of sensitivity, because it is reasonable to accept, at least in the short term, that the last available value is a good prediction. So, if other models present better results than the random walk, it can be indicative of a good predictor.

2.2.3 Lasso

[Ref] Working with high dimensional data, *shrinkage* methods form a prosperous alternative to traditional models. The main idea is to shrink irrelevant variable to zero.

As well as the OLS, the least absolute shrinkage and selection operator (LASSO), also minimizes the MSE. However, lasso has a shrinkage coefficient, λ , that forces irrelevant variables to zero. Their parameters are determined by

$$\hat{\beta} = \underset{\hat{\beta}}{\operatorname{argmin}} \left[\|Y - X\beta\|_2^2 + \lambda \|\beta\|_1 \right]$$

LASSO can be seen as a generical OLS model, because if $\lambda = 0$, the parameters $\hat{\beta}$ and $\hat{\beta}_{OLS}$ are the same. Still, if the coefficient of shrinkage are equal to some $\lambda_c > 0$, all β_i will be equal to zero. As well, $0 = \lambda_{ols} \leq \lambda \leq \lambda_c$.

2.2.4 Lars

Least angle regression (LARS) is also a good model to use when working with high dimensional data. The main idea of his algorithm is to start with all β equal to zero and to increase the coefficient of the most correlated x_i with y . This growth stops when finds

some x_j that own as much correlation with the residual. Again, the model rises (β_i, β_j) in their joint least squares direction, until other predictor x_k , and so on.

Efrom *et. al.* (2004) concluded that the LARS algorithm and his variations works gracefully for the case where there are many more variables than observations. LARS is easy to modify to produce efficient algorithms, like LASSO LARS, and is useful in cross-validation or similar attempts.

2.2.5 Lasso Lars

Efrom *et. al.* (2004) proof that a simple modification of the LARS algorithm can result in the simulation of all LASSO models for all possible values of λ . So, in the traditional LARS method, if a coefficient changes the signal, the direction remains the same. However, in the LASSO model, when a coefficient reaches the value 0, it is discarded from the set of active variables. For that reason, LASSO LARS algorithm make a simple modification in the original LARS: if any coefficient becomes zero this variable is discarded and the model recalculates the search direction.

2.2.6 Ridge

As well as LASSO, Ridge Regression, is a generic version of OLS that also has a shrink coefficient λ . However, the difference is that the Ridge's restriction is squared and their parameters are determined below

$$\hat{\beta} = \underset{\hat{\beta}}{\operatorname{argmin}} \left[\|Y - X\beta\|_2^2 + \lambda \|\beta\|_2^2 \right]$$

By adding a squared constraints, the loss function is strictly convex, and therefore has a unique minimum. Consequently. Ridge algorithm make it harder to zero the coefficients and can't eliminate completely some irrelevant variables. This can result in several variables with close values to zero and a lower interpretability of the results.

2.2.7 Elastic Net

Elastic Net is also a regularized regression model that combines the constraints of Lasso and Ridge Regression. Their parameters are given by

$$\hat{\beta} = \underset{\hat{\beta}}{\operatorname{argmin}} \left[\|Y - X\beta\|_2^2 + \lambda_1 \|\beta\|_1 + \lambda_2 \|\beta\|_2^2 \right]$$

Elastic Net becomes a general case of LASSO and Ridge Regression. And so, as in Ridge model, the Elastic Net method makes the loss function strictly convex, forcing it to have a unique minimum.

2.2.8 Random Forest

[falar que é uma classe diferente Regularization vs Ensemble] Random Forest is a method of machine learning for classification and regression of variables. This model creates several random decision trees in the training period and mixes them to obtain the most accurate prediction possible, resulting in the node of the variables that appear most frequently

Random Forest seeks to reduce the overfitting of traditional models when making a bagging (or Botstrap), combining learning models. By making this average of the various decision nodes possible, the model decreases its variance in exchange for an increase in bias.

3 Data

Power Electricity Consumption (PEC) is measured using the hourly energy load by subsystem (KW/h) released by the National Electricity System Operator (ONS). The initial date from our dataset is February 1, 2017, when the hourly data of electricity consumption started to be disclosed, and ends July 31, 2018, totaling one and a half years of hourly data of PEC. However, the time lag analyzed is daily, due to the other variables, totaling 546 days observed.

We used February 2017 to April 2018 for the training set. The holdout set is from May 2018 to July 2018. We split the training set in two to choose the hyper-parameters of all models for each forecast horizon. We run several hyper-parameters values for each model, such as λ from Lasso. The chosen parameters was the one with the lowest mean square error (MSE).

After the hyper-parameters selection, we run the models for the entire training set and obtain the models parameters. Next, we used β calculated in the training set in our holdout set and we obtain the predictions for all the models and forecasts horizons. To evaluate our models, we use the MSE. Thus, the model that presents the lowest MSE is considered the best because arrives closer to the real value of PEC.

Figure 1 shows how the total daily electricity consumption in Brazil has behaved, its trend, seasonal and residual. We observe PEC is a variable with a great seasonality. However, besides being a cyclical variable, this graph showed that the average consumption of electric energy showed a slight decrease in its average, during the analyzed period.

An important chronological fact that can be observed in Figure 1 is the effect of the truckers' strike. At the end of May 2018, truck drivers blocked roads across Brazil demanding a reduction in diesel prices that had risen more than 50% in the last 12 months. With trucks stalled, partially blocking roads, fuels were no longer delivered at several gas stations and other activities that expected raw materials and essential products, such as food, also ran out of supplies.




Figure 1 – Seasonal Plot




Figure 2 – PEC by Hour

As the exogenous shock negatively affected several layers of the economy, we observe in Figure 1 that the consumption of electric energy fell sharply. This cointegration between the two variables reinforces the great causal relation among economic activity and consumption of electric energy.

Figure 2 exhibits the behavior of the PEC by region. It is easy to observe that the consumption of electric energy has different conduct by region of Brazil and the division of the variable by region better the performances of the forecast models.

3.1 Power Electricity Consumption

Huang and Shih (2003) use just lagged demand to forecast short-term of energy load via ARMA Model. The authors also use a variation of this model including non-Gaussian process considerations and concluded that the performance of ARMA model is better ensured, improving the load forecast accuracy significantly. Almeshaiei and Hassan Soltan (2011) also forecasts electric power load using ARMA/ARIMA models. The authors propose a practical guide for forecasting future electricity consumption, arguing that every electric network needs special forecasting method because each country is indifferent in the factors that affect the electricity demand.

Figure 3 shows how the electric energy consumption is correlated with its lagged demand. We observe a great correlation with its value from the previous day. However, the correlation is even greater with the consumption of a week ago, corroborating with the cyclicity of the PEC. These figures clarify the importance of including lagged demand in forecast models.

Therefore, we selected the hourly energy load by subsystem (KW/h) to Brazil and by Region on the [National Electricity System Operator \(ONS\) website](http://sdro.ons.org.br/SDRO/DIARIO/index.htm)¹. In addition, we took the daily data of PEC divided by consumption class for each region in [Electric Energy National Agency \(ANEEL\) website](http://www.aneel.gov.br)². The set of consumption classes includes PEC

¹ [<http://sdro.ons.org.br/SDRO/DIARIO/index.htm>](http://sdro.ons.org.br/SDRO/DIARIO/index.htm)

² [<http://www.aneel.gov.br>](http://www.aneel.gov.br)




Figure 3 – Autocorrelation Plot

variables divided in groups such as industrial, comercial and services, own consumption, residential, rural, street lighting, public service and others.




Figure 4 – PEC by sector

This division by consumption class not only improves the performance of prediction models, but also is fundamental to understand where the greatest demand for electricity consumption is. The vast majority of PEC are divided into commercial and services,

industrial and residential consumption. The first two are directly linked to economic activity of the country, while the third is indirectly related. Figure 4 shows the trend of these variables, and we can see that the sum of the first two variables exceeds residential consumption, consequently, the largest share of Brazil's electricity consumption is for commercial and industrial activities. This fact reinforces the relation of economic activity and electric energy consumption.

With the division of the variables by hour and by consumption class for each region, a total of 185 past PEC variables has been detected.

3.2 Calendar Variables

Lebotsa *et. al.* (2018) included lagged demand, calendar variables and meteorological variables to construct a partially linear model to predict short term electricity demand. These variables and the use of nonlinear trend variables made the short term electricity demand of perform better.

Fan and Hyndman (2012) propose a new statistical method to forecast the short term electricity demand, allowing nonlinear and nonparametric terms within the regression framework. Their model inputs are calendar variables, lagged demand and temperature variables, and try to catch the complex nonlinear relationship between electricity demand and its drivers.

As we saw above, PEC is an extremely seasonal variable. At least in the short term, the inclusion of calendar variables controls part of this cyclicity. This group of variables includes day of the week, day of the month, month, season of the year, year and a dummy for holidays, totaling 6 calendar variables.

3.3 Meteorological Variables

In the same line of calendar variables, the meteorological variables serve to control a little of the seasonality of the consumption of electric energy. Thus, as seen in the two

studies cited above, it is highly recommendable to place this set of variables in the forecast models of electricity consumption in the short term.

These data were collected from the [National Institute of Meteorology \(INMET\) website](http://www.inmet.gov.br)³ that release the historical series by stations scattered throughout Brazil. This set of variables contains air nebulosity, atmospheric pressure (mbar), dry bulb temperature (oC), humidity bulb temperature (oC), relative humidity (%), wind direction and wind speed (m/s) disclosed for the hours 9, 15 and 21 for each station.

After aggregating the data by region and for brazil, we have a total of 180 meteorological variables for all regions and all hours analyzed by INMET.

The most important meteorological variable for the forecast of electric energy consumption is the temperature of the dry bulb, which is also a very seasonal variable with a large variance. Boldin and Wright (2015) proposes an adjusted temperature that controls both the seasonal effect and the variation of the normal temperature per year.

$$temp_adj_i = temp_i - \frac{1}{n} \sum_{i=1}^n temp_i$$

where $temp_i$ is the temperature on day i .




Figure 5 – Average Temperature

³ <http://www.inmet.gov.br>




Figure 6 – Adjusted Average Temperature

Figures 5 and 6 show how the adjusted variability loses part of its variance and seasonality. Besides that, comparing figures 2 and 6, it is possible to observe that in the times of higher temperature also occurs a greater consumption of electric energy in the day. This is a signal that the meteorological variables are important to predict PEC.

3.4 Price of Electric Energy

Since our interest variable is a demand variable, we include variables that measure the cost of electric energy. We pick the price set from Electric Energy National Agency (ANEEL) and its also divided by consumption class each region. Aggregating all variables by region and by consumption class, we have a total of 60 price variables. Yet, cost of electricity is a monthly variable and as our dataset is a daily variable, we set the same price for the whole month.

When comparing figures 2 and 7, we can observe that the negative relation between price and demand, in this case, is not so strong. The Northeast region, for example, has the lowest tariff and yet does not have the highest consumption of electricity. It's all about the region's structural capacity. Hence, this set of variables may be indicating that the price elasticity of demand of electricity might be low in some time horizons.




Figure 7 – Average Price

3.5 Economic Variables

Shazly (2013) build a dynamic econometric model of nonstationary heterogeneous panels including economic variables. Their models provides a satisfactory technique to analyze energy demand at a disaggregated sectoral level for policy purposes. Maza and Villaverde (2008) argues that existis a causality from electricity consumption to GDP growth.

How the main idea of this paper is measure economic activity with consumption of electric energy, we include a wide list of economic variables, some daily and other monthly. As in the price variables, we set the same value to economic variable for the whole month. We pick the economic variables from a database of Central Bank, and include several indices of price, traffic of vehicles, unemployment, wage, industrial production variables (consumer goods, capital goods, automobile), monetary variables (monetary base and currency issued), consults to SPC and Serasa, BNDES expenses and several others. This set is composed by a total of 79 economic variables.

Adding all the explanatory variables listed in this section, we have a total of 510 variables. As I explained in the previous section, these variables are divided into 3 blocks

of lags that depend on the forecast horizon, k . Thus, our model has 1530 variables and its estimated using a rolling window of 546 observations. We expect that our machine learning models select the most relevant variables for forecast PEC for each forecast horizon.

4 Results

In this session we show the results of the four-best performed models for each time horizon and make an analysis of these models and their parameters. First of all, we list our best models by analyzing their MSE. All variables were normalized before running the models, facilitating the comparison between them.

Model Rank	1 dia (MSE)	7 dias (MSE)	15 dias (MSE)	30 dias (MSE)	60 dias (MSE)	90 dias (MSE)
1st	Random Forest (35753.39)	Random Forest (66621.08)	Lasso Lars (75882.53)	Lasso (72545.07)	Elastic Net (80127.27)	Lasso Lars (84758.67)
2nd	Lasso Lars (48300.59)	Lars (68400.7)	Random Forest (79375.37)	Lasso Lars (74151.69)	Ridge (80876.71)	Lars (85059.85)
3rd	Lars (50234.63)	Ridge (68846.63)	Lars (86543.14)	Ridge (79611.33)	Lars (81952.96)	Random Forest (86887.45)
4th	Lasso (61572.13)	Lasso Lars (68898.78)	Ridge (93536.83)	Lars (108984.56)	Lasso Lars (91794.13)	Ridge (92784.78)

Table 1 – Models Results

Table 1 exhibiths the four models that presented the smallest prediction error (MSE) for each forecast horizon. These results corroborate to what we delineate in section 2. The machine learning models present much better perfomance than the tradicional ones, which ends up being overfitted and presents poor results in all time horizons analyzed. For all forecasts horizons, our four-best prediction models were always the machine learning ones.

Rank	Model		
	Very Short-Term	Short-Term	Medium-Term
1st	Random Forest	Lasso Lars	Lasso Lars
2nd	Lars	Lasso	Lars
3rd	Lasso Lars	Random Forest	Random Forest
4th	Ridge	Lars	Elastic Net
5th	Lasso	Ridge	Ridge

Table 2 – Models Rank

In table 2, we classified the five-best models for each forecast horizon group via analyzing the results presents in Table 1. As we observe in Table 1, the main models in both forecast horizon group are the regularized models that select better the set of relevant variables. For VSTFG, our best prediction model was Random Forest and for STFG and MTFG was Lasso Lars. Both models presented the lowest MSE in most of the forecast horizons analyzed for each group.

Parameters	1 Day				7 Days			
	Random Forest	Lasso Lars	Lars	Lasso	Random Forest	Lars	Ridge	Lasso Lars
Σ (PED Variables)	0.74	-203.06	-251.15	80.12	0.9	266.75	361.89	246.78
Σ (Calendar Variables)	0.2	-55575.01	-54343.69	-122161.95	0.01	116088.1	29363.31	117827.2
Σ (Meteorological Variables)	0.05	2615.98	2633.04	22853.45	0.08	25852.79	18584.8	30527.85
Σ (Price Variables)	0.01	-16332.11	-20921.88	-77237.63	0.01	-61841.9	-20269.3	-53178.2
Σ (Economic Variables)	0.02	1502.11	1450.26	5352.6	0.03	2758.27	3560.09	3226.2

Table 3 – VSTFG: Parameters Analysis

Parameters	15 Days				30 Days			
	Lasso Lars	Random Forest	Lars	Ridge	Lasso	Lasso Lars	Ridge	Lars
Σ (PED Variables)	51.24	0.43	37.55	129.07	58.31	-101.15	196.96	-87.53
Σ (Calendar Variables)	-50502.52	0.33	-45871.43	-36740.47	-177819.21	-154571.91	-80242.57	-80369.78
Σ (Meteorological Variables)	-12791.22	0.13	13409.61	-12422.43	56999.25	54648.9	57121.47	54308.08
Σ (Price Variables)	-14790.79	0.04	-7991.82	-18047.1	-18402.41	10405.24	3155.14	47408.99
Σ (Economic Variables)	-7523.73	0.09	-2645.92	-3731.25	-9692.37	-4650.39	-20018.51	-2452.44

Table 4 – STFG: Parameters Analysis

Parameters	60 Days				90 Days			
	Elastic Net	Ridge	Lars	Lasso Lars	Lasso Lars	Lars	Random Forest	Ridge
Σ (PED Variables)	-47.56	-47.82	17.01	-34.51	-165.46	78.88	0.18	-400.96
Σ (Calendar Variables)	-18938.03	-16871.96	21533.29	-8004.23	42786.93	42731.1	0.51	110.25
Σ (Meteorological Variables)	-20711.24	-19325.94	-8132.87	-38750.11	-9214.65	-4171.4	0.28	-17230.67
Σ (Price Variables)	-84044.65	-76144.52	-22053.61	-121277.64	-82185.74	-93558.98	0.04	-16851.77
Σ (Economic Variables)	-7660.18	-6923.76	-50566.99	-50581.41	-42507.03	-42128.09	0.01	-30923.86

Table 5 – MTFG: Parameters Analysis

To make a better analysis of the most important variables for each model, we sum the coefficients values for each set of variables described in the previous session. Tables 3, 4 and 5 shows this sum for the four-best models in each forecast horizon.

Since each model behave differently in relation to the valuation of their paremeters, comparing the sum among them does not give us much. Moreover, the set of variables has different size and comparison between them also makes no sense. What we did in this work was compare the parameters sum for the same model and the same set of variable in differents forecasts horizons. That is, we basically analyze how the coeficientes sum varied for each set in every time horizon and infer the importance of this set of variables for each one.

In order to clarify our analysis, we will compare here the sum of the coefficients from our mains models for each forecast horizon group. However, we also checked with the coefficients sum of the other models that also presented good results.

4.1 Very Short-Term Forecast Group

For the forecast horizons in this group, the main variables for forecast PEC are almost only lagged values. These variables are essential due to its high autocorrelation and low variance. Other variables have almost no relevance for the prediction in this group.

4.1.1 1 day

Power Electricity Consumption 1 Day Forecast




Figure 8 – 30 days forecast

As can be observed in figures 3 and 4, the electricity consumption is highly correlated with the consumption of the previous day. The LASSO, for example, zeroed all variables parameters except for the aggregate power electric consumption at 11 p.m. of the previous day.

All major models gives greater importance for lagged values of electricity consumption. For example, LARS assigns no value to the coefficients of the 60 electric energy price variables. This makes a lot of sense, since cost of energy consumption in that month will have little or no influence on the electricity consumption of the previous day of the same month.

Figure 8 shows that the forecasts were able to capture the trend and seasonality of electric power consumption. Forecasting 1 day ahead, we could predict until the fall in PEC during the unpredictable truckers' strike. This shows that the main models can absorb up to exogenous shock to forecast horizons of very short term. However, as we increase the forecast horizon, major models have a greater difficulty capturing this shock.

4.1.2 7 days

If energy consumption has a large autocorrelation with the value of the previous day, the autocorrelation with the value of 7 days ago is even higher, as can be seen at figures 3 and 4. Thus, Table 3 shows that the weight of the past variables of electric power consumption is even higher for this forecast horizon, comparing with day the forecast horizon of 1 day.

At the same time as the lagged values increased their importance, the relevance of the others sets of variables remained relatively constant, with a slight increase meteorological and economic variables coefficients sum. That is, the other explanatory variables still not very important for forecasting very short-term, in which lagged demand has a large autocorrelation.

Figure 9 showed that the predictions of the main models are able to follow the trend and seasonality of the series of electric energy consumption. However he had a greater difficulty capturing the exogenous shock that occurred in May 2018. It is worth mentioning that even with a high autocorrelation from lagged demand in this forecast horizon, our main models performed better than values of the previous week, exhibited in Random Walk model.

Power Electricity Consumption 7 Days Forecast




Figure 9 – 7 days forecast

4.2 Short-Term Forecast Group

Now, lagged demand set of variables loses its relevance for the following dates, as its autocorrelation decreases. Consequently, from the 15-day forecast, all other variables significantly expand their importance in the major models. Calendar and meteorological variables coefficients sum had a large increase for all forecast horizons of this group.

The results are intuitive because PEC is highly autocorrelated and as we walk with the forecast horizon, the other variables gain more importance, emphasizing calendar and meteorological variables, which capture better the seasonality of the series.

4.2.1 15 days

For a forecast of 15 days ahead, the correlation with the past demand values are still present, however, increasingly weak. Comparing with the 7-day forecast, it is observed that the sum of the lag variability parameters decreases for the main models. In Random Forest, our main model, this figure decline by more than half.

On the other hand, other variables began to gain more importance as we advance in the forecast horizon. In Random Forest, all the variable sets obtain a greater importance in the main forecast models, with a greater significance for the calendar variables set, which increased their coefficients sum by more than 30 times. Still, meteorological, price and economic variables also were relevant for forecasting energy consumption for 15 days ahead.

Power Eletricity Consumption 15 Days Forecast




Figure 10 – 15 days forecast

Figure 10 shows that, again, the forecast and our best can follow the tendency and cyclicity of the series of electric energy consumption, with a small difficulty to capture the exogenous shock and find the local maximum and minimum of the series.

4.2.2 30 days

Forecasting PEC 30 days ahead, we observe that the lagged demand coefficients sum increases slightly. After the great fall in the last forecast horizon, this set of variables stabilizes its relevance for the prediction of PEC.

On the other hand, the coefficients sum of the calendar and meteorological variables increases somewhat and confirms their importance for short and medium-term forecast models. Both groups are able to catch the seasonality of the series, which does not occur in very short forecast horizons.

In addition, the coefficients sum of price and economic variables didn't varied significantly, but but they remain important to predict demand for electric energy.

Figure 11 shows that the results of forecasts closely follow the original series of electricity consumption. Our main models are able to follow well the variations between local maximum and minimum of the analyzed series, excepting the exogenous shock that occurred in middle of 2018 that was totally unpredictable 1 month before.

4.3 Medium-Term Forecast Group

In STFG calendar and meteorological variables have proved to be essential in prediction models. Now, economic and price variables grows their relevance in all forecasts horizons of this group.

This result is also intuitive, from 1 month and ahead, the variables can absorb the high seasonality of the series increase their importance and stagnate until the end of this forecast group. Also, for 2 and 3 month forecasts, the variables that measure the economic environment and a bigger price elasticity of demand of power electricity show their greater

Power Electricity Consumption 30 Days Forecast




Figure 11 – 30 days forecast

relevance to forecast models.

4.3.1 60 days

Comparing the results of the forecast models from 60-days ahead to 30-days, we observe that the lagged demand coefficients sum increases modestly. That is, this set remained with its relevance constant for the next periods. Also, after a large increase, calendar and meteorological variables coefficients sum fall and had their stagnant relevance for the other forecast horizon, after being able to capture the great variance of the series.

On the other hand, economic and price variables had a huge increase in their coefficients sum. Economics variables set is important for capturing the hole macroeconomic environment. In addition, price variables rise of relevance shows greater price elasticity of

demand of power electricity for longer forecast horizons.

Power Electricity Consumption 60 Days Forecast




Figure 12 – 60 days forecast

So far, our forecasts have converged almost on values to the original consumption. Now the result are able to catch the trend and cyclicity of the original series. This can be observed in Figure 12. With an excess of Random Forest model, with an increase in the forecast horizon the forecasts of the models are more difficult to reach the local maximums and minimums of the series. This is almost a distribution convergence of our forecasts.

4.3.2 90 days

Forecasting PEC 90-days ahead, we observe that coefficients sum did not change significantly, except for a slight decrease in meteorological and price variables, and a modest increase in calendar variables.

Therefore, we infer that the relevance of variables varies more in smaller time horizons and when predicting PEC for 2 and 3 months ahead, they have already stagnated in their relevance.

Power Eletricity Consumption 90 Days Forecast




Figure 13 – 90 days forecast

As we observe in Figure 13, in the last forecast horizon, the main models have a greater difficulty to follow the series cyclicity, but still, they can follow its trend. Random Forest did not performed better due to the unpredictable truckers' strike, where the model predicted a increase of consumption. Random Forest, although not having the smallest MSE, manages to follow the trend and the seasonality of the series better than the other models, with the exception of the negative shock.

In this way, if we are making a prediction of 90 days ahead, Random Forest remains the model that best captures the trend and the cyclicity of the series. While the other

models can give the trend of electricity consumption for the next 3 months, which is good enough when we want to measure the level of future economic activity.

5 Conclusion

In this work, we build a model to predict power electricity consumption (PEC) for 6 different forecast horizon. These horizons are divided between very short term ¹, short term ² and medium term ³. With GDP released quarterly, we want to structure the forecast models of electricity consumption to capture the trend of economic activity.

We construct a high dimensional database with 5 main sets of variables: lagged demand, calendar variables, meteorological variables, cost of electric energy variables and economic variables. We make predictions for more than 10 different models for all forecast horizons, among them are ARIMA, Random Walk and machine learning models. We expect the machine learning models perform better than the usual traditional models because they can make a better selection of the relevant variables. The results showed that the five best models are based on machine learning models, especially Random Forest, which presented the lowest mean square error for almost all forecast horizons. This model was able to capture the trend and the seasonality of electricity consumption with good precision.

In addition, we analyze the variation of the sets of variables for all time horizons analyzed. We conclude that for 1-day forecast, lagged demands are the most relevant variables for predicting the consumption of electricity and it loses its importance in the models as we advance in the time horizon until its relevance is stagnated from the medium term horizons. Calendar variables were very relevant for the 7-day forward forecast and have their relevance shows a slight growth until the forecast horizon of 3 months. Yet, the meteorological variables were relevant for medium-term forecast horizons, that is, their relevance grows in the forecast for 30 days and stagnates for the following values.

Finally, price of the power electricity variables and economic variables had a similar trajectory. Both presented a great relevance for the models when forecasting 15 and 30 days ahead, however, they had a drop of relevance in the models for the following horizons

¹ Forecast horizons: 1 and 7 days ahead

² Forecast horizontes: 15 and 30 days ahead

³ Forecast horizontes: 60 and 90 days ahead

of forecast. Our work presents a new model, which uses high dimensional data and machine learning models to predict power electricity consumption. With the good results of the models, we were able to measure the trend of this consumption and infer about the trend of economic activity in the short term.

In addition, we structure which variables are most relevant to each time horizon, in the short and medium term, which may facilitate the variable selection for future works that predicts power electricity consumption.

Bibliography