

# Math4AI Lecture No.2

Tin Hadži Veljković  
t.hadziveljkovic@uva.nl

20 September 2024.

# Table of Contents

Statistical modeling and Bayesian statistics

Kronecker deltas & Summations

Vector Calculus: Revision

# Basic Probability Rules

- ▶ Probability Notation:

- ▶  $p(x)$ : Probability of event  $x$ .
- ▶  $p(x, y)$ : Joint probability of events  $x$  and  $y$ .
- ▶  $p(x|y)$ : Conditional probability of event  $x$  given event  $y$ .

# Basic Probability Rules

- ▶ Probability Notation:
  - ▶  $p(x)$ : Probability of event  $x$ .
  - ▶  $p(x, y)$ : Joint probability of events  $x$  and  $y$ .
  - ▶  $p(x|y)$ : Conditional probability of event  $x$  given event  $y$ .
- ▶ Chain Rule (Two Variables):

$$p(x, y) = p(x|y) \cdot p(y) = p(y|x) \cdot p(x)$$

# Basic Probability Rules

- ▶ Probability Notation:

- ▶  $p(x)$ : Probability of event  $x$ .
- ▶  $p(x, y)$ : Joint probability of events  $x$  and  $y$ .
- ▶  $p(x|y)$ : Conditional probability of event  $x$  given event  $y$ .

- ▶ Chain Rule (Two Variables):

$$p(x, y) = p(x|y) \cdot p(y) = p(y|x) \cdot p(x)$$

- ▶ Chain Rule (Three Variables):

$$p(x, y, z) = p(x|y, z) \cdot p(y, z) = p(x|y, z) \cdot p(y|z) \cdot p(z)$$

# Basic Probability Rules

- ▶ Probability Notation:

- ▶  $p(x)$ : Probability of event  $x$ .
- ▶  $p(x, y)$ : Joint probability of events  $x$  and  $y$ .
- ▶  $p(x|y)$ : Conditional probability of event  $x$  given event  $y$ .

- ▶ Chain Rule (Two Variables):

$$p(x, y) = p(x|y) \cdot p(y) = p(y|x) \cdot p(x)$$

- ▶ Chain Rule (Three Variables):

$$p(x, y, z) = p(x|y, z) \cdot p(y, z) = p(x|y, z) \cdot p(y|z) \cdot p(z)$$

- ▶ Marginal Probability (Discrete case):

$$p(x) = \sum_y p(x, y) = \sum_y p(x|y) \cdot p(y)$$

# Basic Probability Rules

- ▶ Probability Notation:

- ▶  $p(x)$ : Probability of event  $x$ .
- ▶  $p(x, y)$ : Joint probability of events  $x$  and  $y$ .
- ▶  $p(x|y)$ : Conditional probability of event  $x$  given event  $y$ .

- ▶ Chain Rule (Two Variables):

$$p(x, y) = p(x|y) \cdot p(y) = p(y|x) \cdot p(x)$$

- ▶ Chain Rule (Three Variables):

$$p(x, y, z) = p(x|y, z) \cdot p(y, z) = p(x|y, z) \cdot p(y|z) \cdot p(z)$$

- ▶ Marginal Probability (Discrete case):

$$p(x) = \sum_y p(x, y) = \sum_y p(x|y) \cdot p(y)$$

- ▶ Marginal Probability (Continuous case):

$$p(x) = \int p(x, y) dy = \int p(x|y) \cdot p(y) dy$$

# Statistical modeling

- ▶ *Idea*: Construct a model  $\mathcal{M}$  that describes inherent uncertainty/variability in the observed data  $\mathcal{D}$ , we wish to model the process that generated the data.



# Statistical modeling

- ▶ *Idea*: Construct a model  $\mathcal{M}$  that describes inherent uncertainty/variability in the observed data  $\mathcal{D}$ , we wish to model the process that generated the data.
- ▶ Given a model  $\mathcal{M}$  with its parameters  $\mathbf{w}$  and data  $\mathcal{D}$  we can write the *likelihood function*  $p(\mathcal{D}|\mathbf{w}, \mathcal{M})$ .

# Statistical modeling

- ▶ *Idea*: Construct a model  $\mathcal{M}$  that describes inherent uncertainty/variability in the observed data  $\mathcal{D}$ , we wish to model the process that generated the data.
- ▶ Given a model  $\mathcal{M}$  with its parameters  $\mathbf{w}$  and data  $\mathcal{D}$  we can write the *likelihood function*  $p(\mathcal{D}|\mathbf{w}, \mathcal{M})$ .
- ▶ *Point estimations*: finding specific optimal parameters  $\mathbf{w}^*$  that maximize/minimize some function.

# Statistical modeling

- ▶ *Idea*: Construct a model  $\mathcal{M}$  that describes inherent uncertainty/variability in the observed data  $\mathcal{D}$ , we wish to model the process that generated the data.
- ▶ Given a model  $\mathcal{M}$  with its parameters  $\mathbf{w}$  and data  $\mathcal{D}$  we can write the *likelihood function*  $p(\mathcal{D}|\mathbf{w}, \mathcal{M})$ .
- ▶ *Point estimations*: finding specific optimal parameters  $\mathbf{w}^*$  that maximize/minimize some function.
- ▶ For example, a *maximum likelihood* solution corresponds to the weights  $\mathbf{w}^*$  that maximize the likelihood (or log-likelihood).

# Bayesian Inference in Machine Learning

- ▶ Objective:
  - ▶ Let's assume we have data  $\mathcal{D}$  and a model  $\mathcal{M}$  with parameters  $\mathbf{w}$ .
  - ▶ The goal is to find the posterior distribution over model parameters  $\mathbf{w}$ .
  - ▶ **Disclaimer:** model parameters  $\mathbf{w} \neq$  model  $\mathcal{M}$ !

# Bayesian Inference in Machine Learning

- ▶ Objective:
  - ▶ Let's assume we have data  $\mathcal{D}$  and a model  $\mathcal{M}$  with parameters  $\mathbf{w}$ .
  - ▶ The goal is to find the posterior distribution over model parameters  $\mathbf{w}$ .
  - ▶ **Disclaimer:** model parameters  $\mathbf{w} \neq$  model  $\mathcal{M}$ !
- ▶ The posterior is given by:

$$p(\mathbf{w}|\mathcal{D}, \mathcal{M}) =$$

# Bayesian Inference in Machine Learning

- ▶ Objective:
  - ▶ Let's assume we have data  $\mathcal{D}$  and a model  $\mathcal{M}$  with parameters  $\mathbf{w}$ .
  - ▶ The goal is to find the posterior distribution over model parameters  $\mathbf{w}$ .
  - ▶ **Disclaimer:** model parameters  $\mathbf{w} \neq$  model  $\mathcal{M}$ !
- ▶ The posterior is given by:

$$p(\mathbf{w}|\mathcal{D}, \mathcal{M}) = \frac{\overbrace{p(\mathcal{D}|\mathbf{w}, \mathcal{M})}^{\text{Likelihood}} \cdot \overbrace{p(\mathbf{w}|\mathcal{M})}^{\text{Prior}}}{\underbrace{p(\mathcal{D}|\mathcal{M})}_{\text{Evidence}}}$$

# Bayesian Inference in Machine Learning

- ▶ Objective:
  - ▶ Let's assume we have data  $\mathcal{D}$  and a model  $\mathcal{M}$  with parameters  $\mathbf{w}$ .
  - ▶ The goal is to find the posterior distribution over model parameters  $\mathbf{w}$ .
  - ▶ **Disclaimer:** model parameters  $\mathbf{w} \neq$  model  $\mathcal{M}$ !
- ▶ The posterior is given by:

$$p(\mathbf{w}|\mathcal{D}, \mathcal{M}) = \frac{\overbrace{p(\mathcal{D}|\mathbf{w}, \mathcal{M})}^{\text{Likelihood}} \cdot \overbrace{p(\mathbf{w}|\mathcal{M})}^{\text{Prior}}}{\underbrace{p(\mathcal{D}|\mathcal{M})}_{\text{Evidence}}}$$

- ▶ The evidence is given by:

$$p(\mathcal{D}|\mathcal{M}) = \int p(\mathcal{D}|\mathbf{w}, \mathcal{M})p(\mathbf{w}|\mathcal{M}) d\mathbf{w}$$

# Bayesian Inference in Machine Learning

- ▶ Objective:
  - ▶ Let's assume we have data  $\mathcal{D}$  and a model  $\mathcal{M}$  with parameters  $\mathbf{w}$ .
  - ▶ The goal is to find the posterior distribution over model parameters  $\mathbf{w}$ .
  - ▶ **Disclaimer:** model parameters  $\mathbf{w} \neq$  model  $\mathcal{M}$ !
- ▶ The posterior is given by:

$$p(\mathbf{w}|\mathcal{D}, \mathcal{M}) = \frac{\overbrace{p(\mathcal{D}|\mathbf{w}, \mathcal{M})}^{\text{Likelihood}} \cdot \overbrace{p(\mathbf{w}|\mathcal{M})}^{\text{Prior}}}{\underbrace{p(\mathcal{D}|\mathcal{M})}_{\text{Evidence}}}$$

- ▶ The evidence is given by:

$$p(\mathcal{D}|\mathcal{M}) = \int p(\mathcal{D}|\mathbf{w}, \mathcal{M})p(\mathbf{w}|\mathcal{M}) d\mathbf{w}$$

- ▶ Okay... now what?



# MLE and MAP

- ▶ Evaluating evidence often isn't tractable.

# MLE and MAP

- ▶ Evaluating evidence often isn't tractable.
- ▶ Let's write the log-posterior:

$$\log p(\mathbf{w}|\mathcal{D}, \mathcal{M}) = \log p(\mathcal{D}|\mathbf{w}, \mathcal{M}) + \log p(\mathbf{w}|\mathcal{M}) - \log p(\mathcal{D}|\mathcal{M})$$

# MLE and MAP

- ▶ Evaluating evidence often isn't tractable.
- ▶ Let's write the log-posterior:

$$\log p(\mathbf{w}|\mathcal{D}, \mathcal{M}) = \log p(\mathcal{D}|\mathbf{w}, \mathcal{M}) + \log p(\mathbf{w}|\mathcal{M}) - \log p(\mathcal{D}|\mathcal{M})$$

- ▶ Finding the *maximum a posteriori* estimation has the form:

$$\frac{d \log p(\mathbf{w}|\mathcal{D}, \mathcal{M})}{d\mathbf{w}} = 0 \rightarrow \frac{d \log p(\mathcal{D}|\mathbf{w}, \mathcal{M})}{d\mathbf{w}} + \frac{d \log p(\mathbf{w}|\mathcal{M})}{d\mathbf{w}} = 0$$

# MLE and MAP

- ▶ Evaluating evidence often isn't tractable.
- ▶ Let's write the log-posterior:

$$\log p(\mathbf{w}|\mathcal{D}, \mathcal{M}) = \log p(\mathcal{D}|\mathbf{w}, \mathcal{M}) + \log p(\mathbf{w}|\mathcal{M}) - \log p(\mathcal{D}|\mathcal{M})$$

- ▶ Finding the *maximum a posteriori* estimation has the form:

$$\frac{d \log p(\mathbf{w}|\mathcal{D}, \mathcal{M})}{d\mathbf{w}} = 0 \rightarrow \frac{d \log p(\mathcal{D}|\mathbf{w}, \mathcal{M})}{d\mathbf{w}} + \frac{d \log p(\mathbf{w}|\mathcal{M})}{d\mathbf{w}} = 0$$

- ▶ Similarly to MLE, we would obtain specific optimal parameters  $\mathbf{w}^*$  that maximize the posterior.

# MLE and MAP

- ▶ Evaluating evidence often isn't tractable.
- ▶ Let's write the log-posterior:

$$\log p(\mathbf{w}|\mathcal{D}, \mathcal{M}) = \log p(\mathcal{D}|\mathbf{w}, \mathcal{M}) + \log p(\mathbf{w}|\mathcal{M}) - \log p(\mathcal{D}|\mathcal{M})$$

- ▶ Finding the *maximum a posteriori* estimation has the form:

$$\frac{d \log p(\mathbf{w}|\mathcal{D}, \mathcal{M})}{d\mathbf{w}} = 0 \rightarrow \frac{d \log p(\mathcal{D}|\mathbf{w}, \mathcal{M})}{d\mathbf{w}} + \frac{d \log p(\mathbf{w}|\mathcal{M})}{d\mathbf{w}} = 0$$

- ▶ Similarly to MLE, we would obtain specific optimal parameters  $\mathbf{w}^*$  that maximize the posterior.
- ▶ *Insight*: when are MLE and MAP solutions equal?

## Bayesian predictive distribution

- ▶ Let's assume we managed to obtain the posterior  $p(\mathbf{w}|\mathcal{D}, \mathcal{M})$  somehow. What can we do with it?

## Bayesian predictive distribution

- ▶ Let's assume we managed to obtain the posterior  $p(\mathbf{w}|\mathcal{D}, \mathcal{M})$  somehow. What can we do with it?
- ▶ Given new datapoint  $\mathbf{x}$ , the predictive distribution for the observation  $\mathbf{t}$  is given by:

$$p(\mathbf{t}|\mathbf{x}, \mathcal{D}, \mathcal{M}) = \int p(\mathbf{t}|\mathbf{x}, \mathbf{w}, \mathcal{M})p(\mathbf{w}|\mathcal{D}, \mathcal{M}) d\mathbf{w}$$

## Bayesian predictive distribution

- ▶ Let's assume we managed to obtain the posterior  $p(\mathbf{w}|\mathcal{D}, \mathcal{M})$  somehow. What can we do with it?
- ▶ Given new datapoint  $\mathbf{x}$ , the predictive distribution for the observation  $\mathbf{t}$  is given by:

$$p(\mathbf{t}|\mathbf{x}, \mathcal{D}, \mathcal{M}) = \int p(\mathbf{t}|\mathbf{x}, \mathbf{w}, \mathcal{M})p(\mathbf{w}|\mathcal{D}, \mathcal{M}) d\mathbf{w}$$

- ▶ What are the benefits of this approach?



## Bayesian predictive distribution

- ▶ Let's assume we managed to obtain the posterior  $p(\mathbf{w}|\mathcal{D}, \mathcal{M})$  somehow. What can we do with it?
- ▶ Given new datapoint  $\mathbf{x}$ , the predictive distribution for the observation  $\mathbf{t}$  is given by:

$$p(\mathbf{t}|\mathbf{x}, \mathcal{D}, \mathcal{M}) = \int p(\mathbf{t}|\mathbf{x}, \mathbf{w}, \mathcal{M})p(\mathbf{w}|\mathcal{D}, \mathcal{M}) d\mathbf{w}$$

- ▶ What are the benefits of this approach?
- ▶ The predictive distribution in case of a Gaussian distributions is given by:

$$p(\mathbf{t}|\mathbf{x}, \mathcal{D}, \mathcal{M}) = \mathcal{N}(\mathbf{t}|\mathbf{m}^T \phi(\mathbf{x}), \sigma_N^2(\mathbf{x})).$$

## Bayesian predictive distribution

- ▶ Let's assume we managed to obtain the posterior  $p(\mathbf{w}|\mathcal{D}, \mathcal{M})$  somehow. What can we do with it?
- ▶ Given new datapoint  $\mathbf{x}$ , the predictive distribution for the observation  $\mathbf{t}$  is given by:

$$p(\mathbf{t}|\mathbf{x}, \mathcal{D}, \mathcal{M}) = \int p(\mathbf{t}|\mathbf{x}, \mathbf{w}, \mathcal{M})p(\mathbf{w}|\mathcal{D}, \mathcal{M}) d\mathbf{w}$$





- ▶ What are the benefits of this approach?
- ▶ The predictive distribution in case of a Gaussian distributions is given by:

$$p(\mathbf{t}|\mathbf{x}, \mathcal{D}, \mathcal{M}) = \mathcal{N}(\mathbf{t}|\mathbf{m}^T \phi(\mathbf{x}), \sigma_N^2(\mathbf{x})).$$


- ▶ The variance is a function of datapoints  $\mathbf{x}$ :

$$\sigma_N^2(\mathbf{x}) = 1/\beta + \phi(\mathbf{x})^T \mathbf{S}_N \phi(\mathbf{x}).$$

# Visualizing uncertainty




## New predictions - point estimations




- ▶ Predictions using the likelihood  $\mathcal{L}$  with optimal weights  $\mathbf{w}^*$ .
- ▶ Selecting one specific instantiation of the model  $\mathcal{M}$ , no uncertainty in the model parameters.

## New predictions - Bayesian predictive distribution




- ▶ Obtaining the posterior distribution over model parameters  $\mathbf{w}$ .
- ▶ Predictions using the predictive distribution, without choosing a specific set of weights  $\mathbf{w}$ .
- ▶ The "average" prediction over all possible model instantiations.

## New predictions - combining models?



- Given the same data  $\mathcal{D}$ , can we combine different "worlds" (models)?



## Back to the evidence!

- ▶ The evidence was given by  $p(\mathcal{D}|\mathcal{M})$ .



## Back to the evidence!

- ▶ The evidence was given by  $p(\mathcal{D}|\mathcal{M})$ .
- ▶ More generally, for a model  $\mathcal{M}_i$ , the evidence is  $p(\mathcal{D}|\mathcal{M}_i)$ .

## Back to the evidence!

- ▶ The evidence was given by  $p(\mathcal{D}|\mathcal{M})$ .
- ▶ More generally, for a model  $\mathcal{M}_i$ , the evidence is  $p(\mathcal{D}|\mathcal{M}_i)$ .
- ▶ But what does it mean?

## Back to the evidence!

- ▶ The evidence was given by  $p(\mathcal{D}|\mathcal{M})$ .
- ▶ More generally, for a model  $\mathcal{M}_i$ , the evidence is  $p(\mathcal{D}|\mathcal{M}_i)$ .
- ▶ But what does it mean?
- ▶ Let's observe the following:

$$p(\mathcal{D}|\mathcal{M}_i) = \int p(\mathcal{D}|\mathbf{w}, \mathcal{M}_i)p(\mathbf{w}|\mathcal{M}_i) d\mathbf{w}$$

## Back to the evidence!

- ▶ The evidence was given by  $p(\mathcal{D}|\mathcal{M})$ .
- ▶ More generally, for a model  $\mathcal{M}_i$ , the evidence is  $p(\mathcal{D}|\mathcal{M}_i)$ .
- ▶ But what does it mean?
- ▶ Let's observe the following:

$$p(\mathcal{D}|\mathcal{M}_i) = \int p(\mathcal{D}|\mathbf{w}, \mathcal{M}_i)p(\mathbf{w}|\mathcal{M}_i) d\mathbf{w}$$

- ▶ The evidence is the measure of how well the model, as a whole, predicts the data.

## Back to the evidence!

- ▶ The evidence was given by  $p(\mathcal{D}|\mathcal{M})$ .
- ▶ More generally, for a model  $\mathcal{M}_i$ , the evidence is  $p(\mathcal{D}|\mathcal{M}_i)$ .
- ▶ But what does it mean?
- ▶ Let's observe the following:

$$p(\mathcal{D}|\mathcal{M}_i) = \int p(\mathcal{D}|\mathbf{w}, \mathcal{M}_i)p(\mathbf{w}|\mathcal{M}_i) d\mathbf{w}$$

- ▶ The evidence is the measure of how well the model, as a whole, predicts the data.
- ▶ We can optimize the parameters of a specific model, but can we optimize which **model** to choose?

## Bayes factor

- ▶ Assume we are given data  $\mathcal{D}$  and two models  $\mathcal{M}_1, \mathcal{M}_2$ .

## Bayes factor

- ▶ Assume we are given data  $\mathcal{D}$  and two models  $\mathcal{M}_1, \mathcal{M}_2$ .
- ▶ We calculate the evidence of models  $\mathcal{M}_1$  and  $\mathcal{M}_2$ .

## Bayes factor

- ▶ Assume we are given data  $\mathcal{D}$  and two models  $\mathcal{M}_1, \mathcal{M}_2$ .
- ▶ We calculate the evidence of models  $\mathcal{M}_1$  and  $\mathcal{M}_2$ .
- ▶ Their ratio  $K$  is called the *Bayes factor*:

$$K = \frac{p(\mathcal{D}|\mathcal{M}_1)}{p(\mathcal{D}|\mathcal{M}_2)}$$



## Bayes factor

- ▶ Assume we are given data  $\mathcal{D}$  and two models  $\mathcal{M}_1, \mathcal{M}_2$ .
- ▶ We calculate the evidence of models  $\mathcal{M}_1$  and  $\mathcal{M}_2$ .
- ▶ Their ratio  $K$  is called the *Bayes factor*:

$$K = \frac{p(\mathcal{D}|\mathcal{M}_1)}{p(\mathcal{D}|\mathcal{M}_2)}$$

- ▶ Robust, penalizes overfitting. How do we interpret the values?

$K$ Range	Strength of Evidence
1 to 3.2	Not worth more than a bare mention
3.2 to 10	Substantial
10 to 100	Strong
> 100	Decisive

Table: Strength of Evidence vs.  $K$

# Mixture distributions

- ▶ Let's think again of combining models!

# Mixture distributions

- ▶ Let's think again of combining models!
- ▶ Assume we have a set of models  $\{\mathcal{M}_i\}$ , where  $i = 1, \dots, L$  and data  $\mathcal{D}$ .

# Mixture distributions

- ▶ Let's think again of combining models!
- ▶ Assume we have a set of models  $\{\mathcal{M}_i\}$ , where  $i = 1, \dots, L$  and data  $\mathcal{D}$ .
- ▶ Let's find the *model posterior* distribution:

$$p(\mathcal{M}_i|\mathcal{D}) = \frac{\overbrace{p(\mathcal{D}|\mathcal{M}_i)}^{\text{Model evidence}} \cdot \overbrace{p(\mathcal{M}_i)}^{\text{Model prior}}}{\underbrace{p(\mathcal{D})}_{\text{Probability of the data}}}$$

# Mixture distributions


- ▶ Let's think again of combining models!
- ▶ Assume we have a set of models  $\{\mathcal{M}_i\}$ , where  $i = 1, \dots, L$  and data  $\mathcal{D}$ .
- ▶ Let's find the *model posterior* distribution:

$$p(\mathcal{M}_i|\mathcal{D}) = \frac{\overbrace{p(\mathcal{D}|\mathcal{M}_i)}^{\text{Model evidence}} \cdot \overbrace{p(\mathcal{M}_i)}^{\text{Model prior}}}{\underbrace{p(\mathcal{D})}_{\text{Probability of the data}}}$$

- ▶ A *mixture distribution* that uses all models is given by:

$$p(\mathbf{t}_{\text{new}}|\mathbf{x}_{\text{new}}, \mathcal{D}) = \sum_{i=1}^L p(\mathbf{t}_{\text{new}}|\mathbf{x}_{\text{new}}, \mathcal{M}_i, \mathcal{D})p(\mathcal{M}_i|\mathcal{D}).$$

## New predictions - combining models



- New predictions by averaging predictive distributions of individual models, weighted by their posterior probabilities.

# Kronecker delta

- Definition of the Kronecker delta:

$$\delta_{ij} = \begin{cases} 1, & \text{if } i = j, \\ 0, & \text{if } i \neq j. \end{cases}$$

# Kronecker delta

- Definition of the Kronecker delta:

$$\delta_{ij} = \begin{cases} 1, & \text{if } i = j, \\ 0, & \text{if } i \neq j. \end{cases}$$

- Let  $\mathbf{a} = (a_1, \dots, a_n)^T$ . Summing over the elements of this vector with  $\delta_{ij}$  yields:

$$\sum_{i=1}^n \delta_{ik} a_i = a_k$$



# Kronecker delta

- ▶ Similarly, if we have a matrix  $\mathbf{A} \in \mathbb{R}^{m \times n}$  with elements  $A_{ij}$ , and we sum over all elements of the matrix using one Kronecker delta, we get:

$$\sum_{i=1}^n \sum_{j=1}^m A_{ij} \delta_{ik} = \sum_{j=1}^m A_{kj}$$

# Kronecker delta

- ▶ Similarly, if we have a matrix  $\mathbf{A} \in \mathbb{R}^{m \times n}$  with elements  $A_{ij}$ , and we sum over all elements of the matrix using one Kronecker delta, we get:

$$\sum_{i=1}^n \sum_{j=1}^m A_{ij} \delta_{ik} = \sum_{j=1}^m A_{kj}$$

- ▶ Sometimes we will encounter summations that include multiple Kronecker delta symbols (both which include indices of the matrix). In this case, we would get:

$$\sum_{i=1}^n \sum_{j=1}^m A_{ij} \delta_{ik} \delta_{jl} = \sum_{j=1}^m A_{kj} \delta_{jl} = A_{kl}$$

# Kronecker delta

- ▶ Similarly, if we have a matrix  $\mathbf{A} \in \mathbb{R}^{m \times n}$  with elements  $A_{ij}$ , and we sum over all elements of the matrix using one Kronecker delta, we get:

$$\sum_{i=1}^n \sum_{j=1}^m A_{ij} \delta_{ik} = \sum_{j=1}^m A_{kj}$$

- ▶ Sometimes we will encounter summations that include multiple Kronecker delta symbols (both which include indices of the matrix). In this case, we would get:

$$\sum_{i=1}^n \sum_{j=1}^m A_{ij} \delta_{ik} \delta_{jl} = \sum_{j=1}^m A_{kj} \delta_{jl} = A_{kl}$$

- ▶ Kronecker delta acts as an index selector and eliminates sums over indices which are also part of the Kronecker delta symbols.

## Linear Algebra: General notes

- ▶ Matrix-vector multiplication:  $\mathbf{b} = \mathbf{X}\mathbf{a}$ . The  $i$ -th element  $b_i$  is given by:

$$b_i = \sum_k X_{ik} a_k$$

## Linear Algebra: General notes

- ▶ Matrix-vector multiplication:  $\mathbf{b} = \mathbf{X}\mathbf{a}$ . The  $i$ -th element  $b_i$  is given by:

$$b_i = \sum_k X_{ik} a_k$$

- ▶ Matrix-matrix multiplication:  $\mathbf{C} = \mathbf{A}\mathbf{B}$ . The  $ij$ -th element  $C_{ij}$  is given by:

$$C_{ij} = \sum_k A_{ik} B_{kj}$$

## Linear Algebra: General notes

- ▶ Matrix-vector multiplication:  $\mathbf{b} = \mathbf{X}\mathbf{a}$ . The  $i$ -th element  $b_i$  is given by:

$$b_i = \sum_k X_{ik} a_k$$

- ▶ Matrix-matrix multiplication:  $\mathbf{C} = \mathbf{AB}$ . The  $ij$ -th element  $C_{ij}$  is given by:

$$C_{ij} = \sum_k A_{ik} B_{kj}$$

- ▶ Dot product:  $\mathbf{a} \cdot \mathbf{b} = \mathbf{a}^T \mathbf{b}$ . The resulting scalar is given by:

$$\mathbf{a}^T \mathbf{b} = \sum_k a_k b_k$$

## Vector Calculus: General notes

- ▶ Suppose we have a vector valued function  $\mathbf{f}(\mathbf{X}) = \mathbf{Y}$ , where  $\mathbf{X} \in \mathbb{R}^n$ , and  $\mathbf{Y} \in \mathbb{R}^m$ . We can write this mapping as  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^m$

## Vector Calculus: General notes

- ▶ Suppose we have a vector valued function  $\mathbf{f}(\mathbf{X}) = \mathbf{Y}$ , where  $\mathbf{X} \in \mathbb{R}^n$ , and  $\mathbf{Y} \in \mathbb{R}^m$ . We can write this mapping as  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^m$
- ▶ The derivative  $\frac{d\mathbf{f}}{d\mathbf{x}}$  will have, by definition, shape  $\mathbb{R}^{m \times n}$ .



## Vector Calculus: General notes

- ▶ Suppose we have a vector valued function  $\mathbf{f}(\mathbf{X}) = \mathbf{Y}$ , where  $\mathbf{X} \in \mathbb{R}^n$ , and  $\mathbf{Y} \in \mathbb{R}^m$ . We can write this mapping as  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^m$
- ▶ The derivative  $\frac{d\mathbf{f}}{d\mathbf{x}}$  will have, by definition, shape  $\mathbb{R}^{m \times n}$ .
- ▶ For example, if the function  $f \in \mathbb{R}$  (a scalar), and the input is a vector  $\mathbf{x} \in \mathbb{R}^n$ , then the derivative will have the shape  $\mathbb{R}^{1 \times n}$  (i.e. the derivative is an  $n$ -dimensional row vector).

# Vector Calculus: An example from A to Z

- ▶ Suppose we have a function  $f(\mathbf{X}) = \mathbf{b}^T \mathbf{X}^T \mathbf{X} \mathbf{c}$

## Vector Calculus: An example from A to Z

- ▶ Suppose we have a function  $f(\mathbf{X}) = \mathbf{b}^T \mathbf{X}^T \mathbf{X} \mathbf{c}$
- ▶ We have  $f \in \mathbb{R}$ ,  $\mathbf{X} \in \mathbb{R}^{n \times m}$ , while  $\mathbf{b} \in \mathbb{R}^m$ , and  $\mathbf{c} \in \mathbb{R}^m$

## Vector Calculus: An example from A to Z

$$A \in \mathbb{R}^{n \times m}$$

$$B \in \mathbb{R}^{m \times k}$$

$$C = AB \in \mathbb{R}^{n \times k}$$

$$(n \times m) \cdot (m \times k) = n \times k$$

- ▶ Suppose we have a function  $f(\mathbf{X}) = \mathbf{b}^T \mathbf{X}^T \mathbf{X} \mathbf{c}$
- ▶ We have  $f \in \mathbb{R}$ ,  $\mathbf{X} \in \mathbb{R}^{n \times m}$ , while  $\mathbf{b} \in \mathbb{R}^m$ , and  $\mathbf{c} \in \mathbb{R}^m$
- ▶ What is the shape of the derivative  $\partial f(\mathbf{X}) / \partial \mathbf{X}$ ?

# Vector Calculus: An example from A to Z

- ▶ Let's calculate  $\partial f(\mathbf{X})/\partial \mathbf{X}$ .

## Vector Calculus: An example from A to Z

- ▶ Let's calculate  $\partial f(\mathbf{X})/\partial \mathbf{X}$ .
- ▶ Before we continue, let's expand the expression for the function  $f = \mathbf{b}^T \mathbf{X}^T \mathbf{X} \mathbf{c}$ .

## Vector Calculus: An example from A to Z

$$\mathbf{b}^T \mathbf{X}^T \mathbf{v}$$

- ▶ Let's calculate  $\partial f(\mathbf{X})/\partial \mathbf{X}$ .
- ▶ Before we continue, let's expand the expression for the function  $f = \mathbf{b}^T \mathbf{X}^T \mathbf{X} \mathbf{c}$ .
- ▶ First we will expand the term  $\mathbf{X} \mathbf{c}$ , which is a matrix-vector multiplication. We will call this new vector  $\mathbf{v} = \mathbf{X} \mathbf{c}$ . The  $l$ -th element of this vector is given by:

$$v_l = \sum_k X_{lk} c_k$$

## Vector Calculus: An example from A to Z

- ▶ Let's calculate  $\partial f(\mathbf{X})/\partial \mathbf{X}$ .
- ▶ Before we continue, let's expand the expression for the function  $f = \mathbf{b}^T \mathbf{X}^T \mathbf{X} \mathbf{c}$ .
- ▶ First we will expand the term  $\mathbf{X} \mathbf{c}$ , which is a matrix-vector multiplication. We will call this new vector  $\mathbf{v} = \mathbf{X} \mathbf{c}$ . The  $l$ -th element of this vector is given by:

$$v_l = \sum_k X_{lk} c_k$$

- ▶ Next, we see that we can write out the function  $f$  as  $\mathbf{b}^T \mathbf{X}^T \mathbf{v}$ .



## Vector Calculus: An example from A to Z

- ▶ Let's calculate  $\partial f(\mathbf{X})/\partial \mathbf{X}$ .
- ▶ Before we continue, let's expand the expression for the function  $f = \mathbf{b}^T \mathbf{X}^T \mathbf{X} \mathbf{c}$ .
- ▶ First we will expand the term  $\mathbf{X} \mathbf{c}$ , which is a matrix-vector multiplication. We will call this new vector  $\mathbf{v} = \mathbf{X} \mathbf{c}$ . The  $l$ -th element of this vector is given by:

$$v_l = \sum_k X_{lk} c_k$$

- ▶ Next, we see that we can write out the function  $f$  as  $\mathbf{b}^T \mathbf{X}^T \mathbf{v}$ .
- ▶ Using similar logic, we will write the matrix-vector multiplication  $\mathbf{X}^T \mathbf{v}$  as  $\mathbf{w}$ . Now, the  $m$ -th element of this vector is given by:

$$w_m = \sum_l X_{ml}^T v_l = \sum_l X_{lm} v_l = \sum_l \sum_k X_{lm} X_{lk} c_k$$

## Vector Calculus: An example from A to Z

- Using this substitution, we can see that the function  $f$  can be written as  $\mathbf{b}^T \mathbf{w}$ , which is just a dot product between the two vectors. Thus, we can write:

$$f = \mathbf{b}^T \mathbf{w} = \sum_m b_m w_m = \sum_l \sum_k \sum_m b_m X_{lm} X_{lk} c_k$$

## Vector Calculus: An example from A to Z

- ▶ Using this substitution, we can see that the function  $f$  can be written as  $\mathbf{b}^T \mathbf{w}$ , which is just a dot product between the two vectors. Thus, we can write:

$$f = \mathbf{b}^T \mathbf{w} = \sum_m b_m w_m = \sum_l \sum_k \sum_m b_m X_{lm} X_{lk} c_k$$

- ▶ The derivative will be a matrix, and we find the  $ij$ -th element of the derivative by taking the derivative wrt.  $ij$ -th element of the matrix  $\mathbf{X}$ :

$$\left[ \frac{df}{d\mathbf{X}} \right]_{ij} = \frac{\partial f}{\partial X_{ij}}$$

## Vector Calculus: An example from A to Z

- ▶ Using this substitution, we can see that the function  $f$  can be written as  $\mathbf{b}^T \mathbf{w}$ , which is just a dot product between the two vectors. Thus, we can write:

$$f = \mathbf{b}^T \mathbf{w} = \sum_m b_m w_m = \sum_l \sum_k \sum_m b_m X_{lm} X_{lk} c_k$$

- ▶ The derivative will be a matrix, and we find the  $ij$ -th element of the derivative by taking the derivative wrt.  $ij$ -th element of the matrix  $\mathbf{X}$ :

$$\left[ \frac{df}{d\mathbf{X}} \right]_{ij} = \frac{\partial f}{\partial X_{ij}}$$

- ▶ Let's take the derivative!

# Vector Calculus: An example from A to Z

- ▶ The derivative is given by:



$$\frac{\partial f}{\partial X_{ij}} = \frac{\partial}{\partial X_{ij}} \left( \sum_l \sum_k \sum_m b_m X_{lm} X_{lk} c_k \right)$$

# Vector Calculus: An example from A to Z

- ▶ The derivative is given by:



$$\begin{aligned}\frac{\partial f}{\partial X_{ij}} &= \frac{\partial}{\partial X_{ij}} \left( \sum_l \sum_k \sum_m b_m X_{lm} X_{lk} c_k \right) \\ &= \sum_l \sum_k \sum_m b_m \frac{\partial X_{lm}}{\partial X_{ij}} X_{lk} c_k + \sum_l \sum_k \sum_m b_m X_{lm} \frac{\partial X_{lk}}{\partial X_{ij}} c_k\end{aligned}$$

# Vector Calculus: An example from A to Z

- ▶ The derivative is given by:



$$\begin{aligned}\frac{\partial f}{\partial X_{ij}} &= \frac{\partial}{\partial X_{ij}} \left( \sum_l \sum_k \sum_m b_m X_{lm} X_{lk} c_k \right) \\ &= \sum_l \sum_k \sum_m b_m \frac{\partial X_{lm}}{\partial X_{ij}} X_{lk} c_k + \sum_l \sum_k \sum_m b_m X_{lm} \frac{\partial X_{lk}}{\partial X_{ij}} c_k \\ &= \sum_l \sum_k \sum_m b_m \delta_{li} \delta_{mj} X_{lk} c_k + \sum_l \sum_k \sum_m b_m X_{lm} \delta_{li} \delta_{kj} c_k\end{aligned}$$

# Vector Calculus: An example from A to Z

- ▶ The derivative is given by:



$$\begin{aligned}\frac{\partial f}{\partial X_{ij}} &= \frac{\partial}{\partial X_{ij}} \left( \sum_l \sum_k \sum_m b_m X_{lm} X_{lk} c_k \right) \\&= \sum_l \sum_k \sum_m b_m \frac{\partial X_{lm}}{\partial X_{ij}} X_{lk} c_k + \sum_l \sum_k \sum_m b_m X_{lm} \frac{\partial X_{lk}}{\partial X_{ij}} c_k \\&= \sum_l \sum_k \sum_m b_m \delta_{li} \delta_{mj} X_{lk} c_k + \sum_l \sum_k \sum_m b_m X_{lm} \delta_{li} \delta_{kj} c_k \\&= \sum_k b_j X_{ik} c_k + \sum_m X_{im} b_m c_j\end{aligned}$$



## Vector Calculus: An example from A to Z

- ▶ The derivative is given by:



$$\begin{aligned}\frac{\partial f}{\partial X_{ij}} &= \frac{\partial}{\partial X_{ij}} \left( \sum_l \sum_k \sum_m b_m X_{lm} X_{lk} c_k \right) \\&= \sum_l \sum_k \sum_m b_m \frac{\partial X_{lm}}{\partial X_{ij}} X_{lk} c_k + \sum_l \sum_k \sum_m b_m X_{lm} \frac{\partial X_{lk}}{\partial X_{ij}} c_k \\&= \sum_l \sum_k \sum_m b_m \delta_{li} \delta_{mj} X_{lk} c_k + \sum_l \sum_k \sum_m b_m X_{lm} \delta_{li} \delta_{kj} c_k \\&= \sum_k b_j X_{ik} c_k + \sum_m X_{im} b_m c_j \\&= \left( \sum_k X_{ik} c_k \right) b_j + \left( \sum_m X_{im} b_m \right) c_j\end{aligned}$$

## Vector Calculus: An example from A to Z

- ▶ Let's define new vectors  $\tilde{\mathbf{c}} = \mathbf{X}\mathbf{c}$  and  $\tilde{\mathbf{b}} = \mathbf{X}\mathbf{b}$ .

## Vector Calculus: An example from A to Z

- ▶ Let's define new vectors  $\tilde{\mathbf{c}} = \mathbf{X}\mathbf{c}$  and  $\tilde{\mathbf{b}} = \mathbf{X}\mathbf{b}$ .
- ▶ Then, their  $p$ -th element is given by  $\tilde{b}_p = \sum_r X_{pr}b_r$  and  $\tilde{c}_p = \sum_r X_{pr}c_r$  respectively.

## Vector Calculus: An example from A to Z

- ▶ Let's define new vectors  $\tilde{\mathbf{c}} = \mathbf{X}\mathbf{c}$  and  $\tilde{\mathbf{b}} = \mathbf{X}\mathbf{b}$ .
- ▶ Then, their  $p$ -th element is given by  $\tilde{b}_p = \sum_r X_{pr} b_r$  and  $\tilde{c}_p = \sum_r X_{pr} c_r$  respectively.
- ▶ We can recognize that in the previous expression!

$$\frac{\partial f}{\partial X_{ij}} = \left( \sum_k X_{ik} c_k \right) b_j + \left( \sum_m X_{im} b_m \right) c_j = \tilde{c}_i b_j + \tilde{b}_i c_j$$

## Vector Calculus: An example from A to Z

- ▶ Let's define new vectors  $\tilde{\mathbf{c}} = \mathbf{X}\mathbf{c}$  and  $\tilde{\mathbf{b}} = \mathbf{X}\mathbf{b}$ .
- ▶ Then, their  $p$ -th element is given by  $\tilde{b}_p = \sum_r X_{pr} b_r$  and  $\tilde{c}_p = \sum_r X_{pr} c_r$  respectively.
- ▶ We can recognize that in the previous expression!

$$\frac{\partial f}{\partial X_{ij}} = \left( \sum_k X_{ik} c_k \right) b_j + \left( \sum_m X_{im} b_m \right) c_j = \tilde{c}_i b_j + \tilde{b}_i c_j$$

- ▶ This is just the outer product, so we have:

$$\frac{df}{d\mathbf{X}} = \tilde{\mathbf{c}}\mathbf{b}^T + \tilde{\mathbf{b}}\mathbf{c}^T = \mathbf{X}\mathbf{c}\mathbf{b}^T + \mathbf{X}\mathbf{b}\mathbf{c}^T$$

## Vector Calculus: An example from A to Z

- ▶ Let's define new vectors  $\tilde{\mathbf{c}} = \mathbf{X}\mathbf{c}$  and  $\tilde{\mathbf{b}} = \mathbf{X}\mathbf{b}$ .
- ▶ Then, their  $p$ -th element is given by  $\tilde{b}_p = \sum_r X_{pr} b_r$  and  $\tilde{c}_p = \sum_r X_{pr} c_r$  respectively.
- ▶ We can recognize that in the previous expression!

$$D_{ij} = \frac{\partial f}{\partial X_{ij}} = \left( \sum_k X_{ik} c_k \right) b_j + \left( \sum_m X_{im} b_m \right) c_j = \tilde{c}_i b_j + \tilde{b}_i c_j$$

- ▶ This is just the outer product, so we have:

$$\frac{df}{d\mathbf{X}} = \tilde{\mathbf{c}}\mathbf{b}^T + \tilde{\mathbf{b}}\mathbf{c}^T = \mathbf{X}\mathbf{c}\mathbf{b}^T + \mathbf{X}\mathbf{b}\mathbf{c}^T$$

- ▶ Therefore the full derivative is given by:

$$\boxed{\frac{df}{d\mathbf{X}} = \mathbf{X} (\mathbf{c}\mathbf{b}^T + \mathbf{b}\mathbf{c}^T)}$$

Handwritten notes on the right side of the slide:

- $f \in \mathbb{R}$
- $\mathbf{X} \in \mathbb{R}^{n \times m}$
- $\frac{df}{d\mathbf{X}} \in \mathbb{R}^{1 \times (n \times m)}$
- $\mathbb{R}^{n \times m}$

Questions?

Ask me anything!