

1 **Running title:** Forbidden interactions
2 **Number of words:** ~910
3 **Number of tables:** 0
4 **Number of figures:** 1
5 **Number of references:** 7

6 **IN FOCUS**
7 **Natural history matters: how biological**
8 **constraints shape diversified interactions in**
9 **pollination networks**

10 **Pedro Jordano¹**

11 *1. Integrative Ecology Group, Estación Biológica de Doñana, Consejo Superior*
12 *de Investigaciones Científicas (EBD-CSIC), Avenida Americo Vespucio s/n,*
13 *E-41092 Sevilla, Spain*

14 *Correspondence author. jordano@ebd.csic.es

15 IN FOCUS: Sazatornil, F.D., Moré, M., Benitez-Vieyra, S., Cocucci, A.A.,
16 Kitching, I.J., Schlumpberger, B.O., Oliveira, P.E., Sazima, M. & Amorim,
17 F.W. (2016) Beyond neutral and forbidden links: morphological matches
18 and the assembly of mutualistic hawkmoth-plant networks. *Journal of An-*
19 *imal Ecology*, 00, 000–000. doi:10.1111/1365-2656.12509

20
21 **Species-specific traits and life-history characteristics constrain**
22 **the ways organisms interact in nature. For example, gape-limited**
23 **predators are constrained in the sizes of prey they can handle and**
24 **efficiently consume. When we consider the ubiquity of such con-**
25 **straints it is evident how hard it can be to be a generalist partner**
26 **in ecological interactions: a free living animal or plant can't simply**
27 **interact with every available partner it encounters. Some pairwise**
28 **interactions among coexisting species simply do not occur; they**
29 **are impossible to observe despite the fact that partners coexist in**
30 **the same place. Sazatornil *et al.* (1) explore the nature of such**
31 **constraints in the mutualisms among hawkmoths and the plants**
32 **they pollinate. In this iconic interaction, used by Darwin and Wal-**
33 **lace to vividly illustrate the power of natural selection in shaping**
34 **evolutionary change, both pollinators and plants are sharply con-**
35 **strained in their interaction modes and outcomes.**

36
37 **Keywords:** complex networks, forbidden links, long-tubed flowers, mutual-
38 **ism, pollination, Sphingidae**

39 Size-limited foragers show clear restrictions on the size of prey items they
40 can efficiently handle. In the case of plant-pollinator interactions, size un-
41 coupling between pollinator bodies and flower sizes or structure are specially
42 relevant in filtering out a range of potential partners (2). The idea, when
43 applied to the bizarre flowers of some plants pollinated by sphingid moths
44 (Lepidoptera: Sphingidae) (Fig. 1), was seminal in Darwinian evolutionary
45 theory to support the potential of natural selection in shaping adaptations
46 (3). Wallace (4) in his book, *Creation by law*, vividly uses the famous example
47 of the Malagasy orchid and its sphingid pollinator to refute the arguments
48 of the Duke of Argyll against natural selection and Darwinism:

49 "There is a Madagascar Orchis—the *Angræcum sesquipedale*—
50 with an immensely long and deep nectary. How did such an
51 extraordinary organ come to be developed? Mr. Darwin's [[p.
52 475]] explanation is this. The pollen of this flower can only be
53 removed by the proboscis of some very large moths trying to get
54 at the nectar at the bottom of the vessel. The moths with the
55 longest proboscis would do this most effectually; they would be
56 rewarded for their long noses by getting the most nectar; whilst
57 on the other hand, the flowers with the deepest nectaries would
58 be the best fertilized by the largest moths preferring them. Con-
59 sequently, the deepest nectaried Orchids and the longest nosed
60 moths would each confer on the other a great advantage in the
61 'battle of life.' This would tend to their respective perpetuation
62 and to the constant lengthening of nectar and noses."

63 Phenotypic fitting of corolla length and shape and the pollinators' feeding
64 apparatus and body sizes are important because the better the fit, the better
65 the consequences in terms of fitness outcomes for the interaction partners
66 (5). Yet the expectation of perfect trait matching across populations or com-
67 munities is too simplistic (7): "arms races" as initially suggested by Darwin
68 and Wallace are frequently asymmetric, originating pollinator shifts rather

69 than tight phenotypic trait matching (Fig. 2). Extensive local variation
70 in phenotypic mismatch exists in different plant-pollinator systems (e.g.,
71 2; 7; 8), with pollinator-mediated selection geographic mosaics of locally
72 coevolved partners.

73 Recent work by Sazatornil *et al.* (1) provides evidences that the types
74 of trait mismatching outlined in Fig. 2 limit the ranges of host plants for
75 sphingid pollinators, and ultimately shape their complex plant-pollinator net-
76 works. By using a comparative analysis of five different hawkmoth/flower
77 assemblages across four South American biotas (Atlantic rainforest and Cer-
78 rado in Brazil, Chaco montane dry woodland, and the ecotone between west-
79 ern Chaco woodland and Yungas montane rain forest in Argentina) they
80 tested the contributions of phenotypic matching to explain observed patterns
81 of moth-flower interactions.

82 How are these moth-flower interactions assembled? Sazatornil *et al.* (1)
83 first tested a neutral model, where interactions are independent of trait-
84 matching. Under this hypothesis distribution parameters (mean and stand-
85 ard deviation) must be the same for both distributions. They further tested
86 a Forbidden links hypothesis, where interactions occurred only if the hawk-
87 moth proboscis length (HPL) is equal to or greater than the effective length
88 of the flower (EFL). EFL is just the corolla tube length (as in Fig. 2 for long-
89 tubed and salverform corollas) or the stamen protrusion length in brush-type
90 and funnel-shape flowers (as in Fig. 1). Sazatornil *et al.* further tested the
91 morphological match hypothesis, where the probability of occurrence of an
92 interaction depends on the frequency of possible pairwise differences between
93 HPL and EFL, i.e., all possible pairwise HPL-EFL differences were weighted

94 by their respective interaction frequency.
 95 Th trait matching between HPL and EFL is crucial in this type of in-
 96 teraction and determines its outcome in terms of fitness for both partners.
 97 Nilsson (5) demonstrated experimentally that shortening the nectary tube
 98 of long-spurred corollas decreased both seed set and pollinia removal for
 99 *Platanthera* orchids. Further experimental evidence has been provided for
 100 long-tongued nemestrinid flies pollinating long-tubed irises in South Africa,
 101 where increased mismatch decreases both plant fitness and the nectar extrac-
 102 tion efficiency of the pollinators (6). Sazatornil *et al.* extend those results
 103 to the scale of the whole moth-plant assemblage and demonstrate that trait
 104 matching successfully predicts the diversity of interactions recorded. In-
 105 terestingly enough, the interaction patterns in two local assemblages from
 106 ecotone areas of the Chaco woodland-Yungas montane rain forest transition
 107 are better fitted by a neutral model where pairwise interactions are driven by
 108 probability of interspecific encounter. Yet Sazatornil *et al.* did not include
 109 the morphological difference for parameter estimation when interactions were
 110 not recorded. Thus the test of the mismatch hypothesis implicitly includes
 111 forbidden links effects: a full mismatch of corolla tube/proboscis lengths ac-
 112 tually means a forbidden link. Furthermore, a fraction of unobserved inter-
 113 actions was likely caused by phenological uncoupling between flowering and
 114 hawkmoth activity phenophases (9; 1). In any case the mismatch hypothesis
 115 somehow captures the fact that a fraction of the unobserved interactions in
 116 these hawkmoth/flower assemblages is due to extreme phenotypic mismatch
 117 (i.e., size-related forbidden links) (1).

118 Sazatornil *et al.* approach would be most useful for proper tests of coe-

119 evolutionary hypotheses in hawkmoth/flower assemblages (and plant-animal
120 mutualisms in general): assessing match/mismatch patterns for every pos-
121 sible pairwise interaction among partners within complex webs of interaction
122 where multiple life-history attributes may contribute biological reasons to
123 expect forbidden links. The morphological match hypothesis is not the only
124 mechanism to explain patterns of hawkmoth–plant interactions, where other
125 life-history limitations may operate generating forbidden links, e.g., phenolo-
126 gical mismatches (for example in the case of long-distance or elevational mi-
127 gratory hawkmoths), energetic constraints due to nectar availability/foraging
128 costs balances, etc.

129 **Acknowledgments**

130 My work was funded by a Severo-Ochoa Excellence Grant (SEV2012-0262)
131 from the Spanish Ministerio de Economía y Competitividad (MINECO),
132 and RNM-5731 from the Junta de Andalucía. Andrea Cocucci generously
133 provided material for Fig. 1 and insightful discussions on sphingids and
134 long-tubed flowers.

References

- [1] Sazatornil, F.D., Moré, M., Benitez-Vieyra, S., Cocucci, A.A., Kitching, I.J., Schlumpberger, B.O., Oliveira, P.E., Sazima, M. & Amorim, F.W. (2016) Beyond neutral and forbidden links: morphological matches and the assembly of mutualistic hawkmoth-plant networks. *Journal of Animal Ecology*, 00, 000–000.
- [2] Cocucci, A.A., Moré, M. & Sérsic, A.N. (2009) Restricciones mecánicas en las interacciones planta-polinizador: estudio de casos en plantas polinizadas por esfíngidos. *Interacciones planta—animal y la conservación de la biodiversidad* (eds R. Medel, R. Zamora, M. Aizen & R. Dirzo), pp. 43–59. CYTED, Madrid.
- [3] Arditti, J., Elliott, J., Kitching, I.J. & Wasserthal, L.T. (2012) “Good Heavens what insect can suck it”—Charles Darwin, *Angraecum sesquipedale* and *Xanthopan morgani praedicta*. *Botanical Journal of the Linnean Society*, 169, 403–432.
- [4] Wallace, A.R. (1867) Creation by law. *The Quarterly Journal of Science*, 4, 471–488.
- [5] Nilsson, L.A. (1988) The evolution of flowers with deep corolla tubes. *Nature*, 334, 147–149.
- [6] Pauw, A., Stofberg, J. & Waterman, R.J. (2009) Flies and flowers in Darwin’s race. *Evolution*, 63, 268–279.
- [7] Anderson, B., Terblanche, J.S. & Ellis, A.G. (2010) Predictable patterns of trait mismatches between interacting plants and insects. *BMC Evolutionary Biology*, 10, 204.
- [8] Moré, M., Amorim, F.W., Benitez-Vieyra, S., Medina, A.M., Sazima, M. & Cocucci, A.A. (2012) Armament imbalances: match and mismatch in plant-pollinator traits of highly specialized long-spurred orchids (ed M. Heil). *PLoS ONE*, 7, e41878.
- [9] Bascompte, J. & Jordano, P. (2014) *Mutualistic Networks*. Princeton University Press, Princeton, NJ.

Pedro Jordano

*Integrative Ecology Group, Estación Biológica de Doñana,
Consejo Superior de Investigaciones Científicas (EBD-CSIC),
Avenida Americo Vespucio s/n,
E-41092 Sevilla, Spain*

Figures

Fig. 1 Morphological mismatches set important biological constraints for size-limited foragers, including e.g., predators, pollinators, and frugivores. In plant-animal mutualisms, a morphological mismatch between partners sets size limits that filter out a range of phenotypes that otherwise could eventually interact. Other reasons for forbidden links include, e.g., phenological differences (9). Thus, a number of the potential interactions that could take place in a given mutualistic assemblage simply cannot occur because of biological reasons: these are forbidden interactions. Photo: Andrea Cocucci. An sphingid moth, *Agrius cingulata*, visiting a flower of *Bauhinia mollis* (Fabaceae), Las Yungas, Argentina.



Fig. 2 The mechanistic basis of morphological mismatches in hawkmoth-flower interactions. Modified and redrawn from (3).

