

# What is long-distance dispersal? And a taxonomy of dispersal events

MS JEcol-2016-0422.R1

Pedro Jordano<sup>†</sup>

Sevilla, September 2, 2016

<sup>†</sup> Integrative Ecology Group, Estación Biológica de Doñana, CSIC, Avda. Americo Vespucio, s/n, Isla de La Cartuja E-41092 Sevilla, Spain.

*Corresponding author:* Pedro Jordano. Integrative Ecology Group, Estación Biológica de Doñana, CSIC, Avda. americo Vespucio, s/n, E-41092 Sevilla, Spain. Email address: jordano@ebd.csic.es

**Key words:** dispersal, frugivory, plant-animal interactions, pollination, seed dispersal

**Manuscript information:** \*\* Words; \*\* Chars; \*\* Pages, \* Figures; \* Tables.

## Abstract

1 Dispersal is a key individual-based process influencing many life-history attributes,  
2 scaling up to population-level properties (e.g., metapopulation connectivity). A  
3 persistent challenge in dispersal ecology has been the robust characterization of  
4 dispersal functions (kernels), a fundamental tool to predict how dispersal processes  
5 respond under global change scenarios. Especially the rightmost tail of these func-  
6 tions, i.e. the long-distance dispersal (LDD) events, are difficult to characterize  
7 empirically and to model in realistic ways. But, when is it a LDD event? In the  
8 specific case of plants, dispersal has three basic components: 1) a distinct (sessile)  
9 source, the maternal plant producing the fruits or the paternal tree acting as a  
10 source of pollen; 2) a distance component between source and target locations;  
11 and 3) a vector actually performing the movement entailing the dispersal event.  
12 Here we discuss operative definitions of LDD based on their intrinsic properties: 1)  
13 events crossing geographic boundaries among stands; and 2) events contributing  
14 to effective gene flow and propagule migration. Strict-sense long distance disper-  
15 sal involves movement both outside the stand geographic limits and outside the  
16 genetic neighborhood area of individuals. Combinations of propagule movements  
17 within/outside these two spatial reference frames results in four distinct modes  
18 of LDD. Beyond traditional statistical approaches to characterize distributions,  
19 Extreme Value Analysis (EVA) can be used to properly and explicitly evaluate  
20 the properties of frequency and extent of LDD events. We discuss conditions  
21 where global change scenarios truncate dispersal processes, leading to the loss of  
22 key dispersal services in natural populations. Proper characterization of the LDD

<sup>23</sup> events helps to assess, for example, how the ongoing defaunation of large-bodied  
<sup>24</sup> frugivores pervasively entails the loss of crucial LDD functions.

## 25 Introduction

26 Dispersal is a key individual-based process influencing many life-history attributes  
27 and scaling up to population-level properties (e.g., metapopulation connectivity,  
28 Cousens *et al.* 2008). In the specific case of plants, largely sessile organisms,  
29 dispersal has three basic components: 1) a distinct (sessile) source, the maternal  
30 plant producing the fruits or the paternal tree acting as a source of pollen; 2) a  
31 distance component between source and target locations; and 3) a vector actually  
32 performing the movement entailing the dispersal event. While realized dispersal  
33 also depends upon stages subsequent to dissemination (e.g., successful germination  
34 and seedling establishment) Schupp (1995), the three previous components fully  
35 characterize the dispersal process per se. Therefore, plant movement differs in  
36 important natural history details from animal dispersal, yet both can be assessed  
37 within a common conceptual framework (e.g., Nathan, 2006). Characteristically,  
38 animal-assisted plant dispersal has three distinct, highly integrated, components  
39 missing in the process of animal dispersal: the properties of the source (parental)  
40 plant, that mediate in the foraging of the animal vector (pollinator or frugivore),  
41 the intrinsic properties of the propagule, and the functional characteristics of the  
42 animal vector who performs the movement (Nathan *et al.*, 2008a).

43 The movement of pollen and seeds by animals and its consequences have intrigued  
44 population geneticists and field ecologists since the infancy of both research disci-  
45 plines. Each has generated an impressive body of theoretical and empirical research  
46 through the past decades, yet advances have long been co-existing in ‘parallel

47 worlds' and the great synergistic potential of population genetics and demography  
48 for the study of plant dispersal by animals remains little explored. Knowledge  
49 gaps still having the imprint of this conceptual disconnection include the idea of  
50 long distance dispersal, and the paradoxes of forest fragmentation effects on ge-  
51 netic diversity (Kramer *et al.*, 2008), survival and persistence of relict tree species  
52 (Hampe & Jump, 2011), rapid post-Pleistocene recolonization of vast continental  
53 areas in response to climate modification (Clark *et al.*, 1998; Clark, 1998), among  
54 other persisting issues. This conceptual isolation has been exacerbated by techni-  
55 cal difficulties for the robust characterization of dispersal events, especially those  
56 involving movement over long-distances (long-distance dispersal, LDD). LDD is a  
57 characteristically extreme event of propagule movement in any plant population,  
58 typically occurring with an extremely low probability but potentially reaching  
59 an extremely long distance. Some progress has recently been made through the  
60 fast-paced implementation of molecular tools in ecological research labs and the  
61 availability of cutting-edge technology for biotelemetry applications. But much of  
62 the population geneticist and ecologist communities remains unaware of the state  
63 of the art in each other and likely under-appreciates their potential to validate  
64 and enrich dispersal studies (Jones & Muller-Landau, 2008). In particular, LDD  
65 events remain difficult to assess, both technically- with serious methodological  
66 problems for its reliable estimation- and conceptually. Our aim here is to review  
67 the LDD concept with a specific emphasis on dispersal of plant propagules (seeds  
68 and pollen), providing an extended definition that might be helpful in the robust  
69 quantification of LDD events.

Two main conceptual approaches have been used to assess dispersal (Fig. 1). The “forward” approach attempts to track the dispersal events away from the known sources, e.g., by tracking the movement patterns of frugivores as they leave fruiting plants after feeding (Fig. 1A). This is the main approach used in the movement ecology framework (Nathan *et al.*, 2008a), with extensive application to animal movement based on the use of advanced biotelemetry. The “backward” approach attempts to reconstruct the most likely source of a dispersed propagule by inferring the sources given the propagule delivery pattern, the fecundity of potential sources, and the dispersal function (Fig. 1B), i.e., by using an inverse modeling approach. The main technical challenge in Fig. 1A is to sample enough dispersal events away from the source to be able to fully characterize the tail (LDD events) of the dispersal function. In Fig. 1B, the main challenge is to have a robust sampling scheme with propagule collectors (e.g., seed traps) and a good characterization of the potential sources to derive robust estimates of the actual sources. Both approaches are limited logically by the difficulties to sample the vast areas required to assess LDD events from the focal source population.

No explicit definition of what constitutes an LDD event exists. Previous approaches (e.g., Nathan, 2006; Schurr *et al.*, 2009) include both absolute and proportional definitions to characterize LDD events. This means providing information about the absolute distances moved by a given percentile of the events and/or providing data on the proportion of events exceeding a given distance threshold (Nathan *et al.*, 2008b). The exact proportional or absolute thresholds selected remain arbitrary, as no reference spatial frame is provided within the

93 definition of LDD. This leaves the consideration of LDD as an extreme form of  
94 context-dependent phenomenon, strongly dependent upon the scale of the biological  
95 process studied (Kinlan *et al.*, 2005) and of the specific organism considered.  
96 For example, Kinlan *et al.* (2005) used a spatial reference frame to characterize  
97 LDD events of marine organisms, where sedentary adults and larvae differ enormously  
98 in the spatial scales of their dispersal (D'Aloia *et al.*, 2013). Therefore,  
99 any measure of extent and reach of LDD events requires reference to an explicit  
100 spatial frame or "local" scale (Kinlan *et al.*, 2005).

101 We aim at providing a general framework for the quantitative analysis of LDD  
102 events so that estimates of their frequency and extent could be comparable across  
103 different study systems. We argue that both demographic and genetic elements  
104 are needed for this framework, most likely requiring a combination of field-based  
105 movement data and genetic analyses. These elements can be overlaid on previous  
106 definitions based on absolute and proportional characterizations of LDD. We start  
107 with a definition of LDD events within a spatially-explicit mechanistic framework  
108 allowing an unambiguous meaning for setting long-distance thresholds. We then  
109 use a case study to assess differential contributions of animal frugivores performing  
110 LDD.

111 Long-distance dispersal is currently one of the most debated topics in dispersal  
112 ecology; it defines the connectedness within the network of local populations and  
113 the possibilities for range expansion and successful colonization events. We propose  
114 a first demogenetically-based, operational definition of what a long-distance  
115 dispersal event actually is, and review existing empirical literature on distance

116 thresholds from population and genetic perspectives. We also show how molecular  
117 tools have been used to identify the respective contributions of different animal  
118 species to the LDD portion of dispersal kernels of pollen and seeds by setting  
119 empirically-derived distance thresholds. Finally, we highlight potential applica-  
120 tions of molecular markers beyond the quantification of just the dispersal distances  
121 that prevails in current studies, e.g., experimental approaches to assess dispersal  
122 limitation and Janzen-Connell effects.

123 **LDD within a demo-genetic perspective: a taxonomy of dis-  
124 persal events**

125 Here we propose an explicit definition of LDD and what constitutes a LDD event.  
126 Previous definitions of dispersal patterns emphasized only their distance compo-  
127 nents and characterized LDD events basically in terms of geographic distance be-  
128 tween a dispersed propagule (or an established early seedling) and its most likely  
129 maternal or paternal (in case of pollen) source. Absolute and proportional defini-  
130 tions for the LDD events have been proposed depending on arbitrary thresholds  
131 of either the distance beyond which a dispersal event is LDD or the proportion of  
132 events occurring beyond a specific distance (Nathan, 2005; Nathan *et al.*, 2008b).  
133 Thus, two key biological aspects of LDD events involve the transport of propag-  
134 ules outside a reference area: moving away from the source stand or population,  
135 and moving away from the area where relatives stand (Kinlan *et al.*, 2005). These  
136 two movements do not necessarily concur: a propagule may move over a very

long distance yet still be disseminated within the reach of the neighborhood where parental individuals mate. Within a demo-genetic framework it is easy to envision a combination of situations concerning the spatial scale of the dispersal processes (Table 1) and unambiguously define different types of LDD events. The idea that dispersal occurs in reference to these two spatial reference frames, i.e., the population or stand and the genetic neighborhood area, is motivated by the fact that dispersal entails the movement of both an individual propagule (i.e., a pollen grain or a seed) and a distinct set of genes (i.e., the male genotype in case of pollen, or a seed genotype). Thus, dispersal entails simultaneous demographic and genetic effects through recruitment of new individuals in the population and through contributions to gene flow (Harper, 1977). When considered its combined influence on demography and population genetics, the concept of LDD nicely bridges these two paradigms embedded in the biological definition of population (Waples & Gaggiotti, 2006).

Two important components of plant dispersal ecology concern the movement of propagules away from the source population, a type of dispersal relevant to colonization ability and range expansion (Howe & Miriti, 2004), and the movement away from the location of close relatives, i.e., a movement away from the genetic neighborhood (Hardesty *et al.*, 2006; Jones & Muller-Landau, 2008). If we classify dispersal events according to these two spatial frameworks (Table 1) we end up with four distinct types of events depending on whether or not dispersed propagules are disseminated within these reference areas. Setting the limits of a population can be problematic (Waples & Gaggiotti, 2006) yet we can identify with relative

160 ease the geographical limits of plant stands, patches, habitat spots or other types  
161 of habitat or microhabitat discontinuities that determine landmark boundaries of  
162 biological significance (see Kinlan *et al.*, 2005, for further discussion of boundaries  
163 for dispersal). These "frontiers" set biological limits to what a LDD event is in  
164 relation to the geographic limits of the source population. Most plants are dis-  
165 tributed as clumped patches, discrete stands, or relatively isolated populations, so  
166 we may distinguish between short-distance and long-distance dispersal events that  
167 end up with dissemination within or beyond, respectively, the stand or population  
168 geographic boundaries (Table 1,  $SSD_{loc}$  or  $LDD_{loc}$ ) (Figure 2).

169 A second consideration in terms of spatial boundaries, with effects on disper-  
170 sal patterns, is the genetic neighborhood area  $N_e^b$ , i.e., the spatial extent includ-  
171 ing a subset of panmictic individuals within a population (Wright, 1943, 1946).  
172 Thus, the  $N_e^b$  area can be equal to the whole extent of the population whenever  
173 the population is unstructured and there is evidence for random mating events  
174 among all the individuals. However, most populations and stands of long-lived  
175 trees show highly aggregated and clumped distributions (Seidler & Plotkin, 2006),  
176 where relatively long distances may separate groups of individuals within the same  
177 population. In these cases we might expect  $N_e^b$  area to be substantially smaller  
178 than the total population area. Therefore, at least four possible scenarios exist  
179 with distinct implications in terms of consequences for dispersal (Table 1). In the  
180 case of dispersal events not extending beyond the geographic limits of the popu-  
181 lation or reference area, actual LDD events may involve dissemination beyond a  
182 reduced neighborhood area that is smaller than the geographic extent of the pop-

ulation, originating local long-distance ( $LDD_{loc}$ ) dispersal events (Table 1, Fig. 2A). Actual short-distance dispersal would then involve those situations where the propagule is disseminated within *both* the population limits and the genetic neighborhood boundary ( $SDD_{loc}$ ). Along a similar reasoning, dispersal events outside the population limits will not necessarily convey LDD (Table 1, Fig. 2B): this is expected in cases where the genetic neighborhoods are extensive, going beyond the geographic limits of local populations, as in fig trees (Nason *et al.*, 1998), generating LDD events within the genetic neighborhood ( $LDD_{neigh}$ ). Finally, strict-sense LDD events would involve dissemination outside *both* the population limits and the genetic neighborhood boundary ( $LDD_{ss}$ ) (Table 1, Fig. 2A).

While both  $SSD_{loc}$  and  $LDD_{loc}$  can be crucial for assuring the local persistence of populations,  $LDD_{neigh}$  and  $LDD_{ss}$  would be extremely important contributors to the structuring of genetic pools, realized gene flow, and maintaining connectivity in metapopulation scenarios. We argue that both the demographic and the genetic references are relevant for a proper definition of LDD.

## Individual and Population Neighborhoods as Reference

Continuous populations can be modeled with the concepts of isolation by distance and neighborhood size(Wright, 1943, 1946). The former refers to the case that limited gene dispersal in continuous populations produces demes that are panmictic internally, but are isolated to some extent from adjacent demes. Each group of reproducing individuals is the neighborhood, defined as the population of a region

in a continuum, from which the parents of individuals born near the center may be treated as if drawn at random (Wright, 1969). The importance and influence of the dispersal process in determining the size of the neighborhood is given by this equation, which shows how the spatial dispersion (pattern of spatial distribution) of the population influences the effective population size. This influence on the effective size is given by:

$$N_e^b = 4\pi\sigma\delta \quad (1)$$

where  $\delta$  is the density of adults per unit area and  $\sigma$  is the standard deviation of the distance between birth and breeding sites. This formulation is often called the neighborhood size and assumes a normal distribution of distances between parents and offspring (out in a perfect circular shape from the source). Thus, changes in the variance of dispersal distance can affect  $N_e^b$  (highly clumped populations will have reduced  $N_e^b$ ). This is the basic model of "Isolation by Distance" proposed by Wright (1943, 1946). Under this type of model, migration (gene flow) is given by the variance in dispersal, and not by the proportion of the population that is composed of migrants (denoted  $m$ ), as is the case with island models (Slatkin, 1985). With enough distance separating them, two plant individuals have a low probability of mating and can be considered members of distinct genetic populations even if they are not located in geographically distinct populations.

For plants, gene flow may be accomplished by both seeds and pollen, so the variance may be decomposed to account for different patterns of seed and pollen

224 dispersal, and to take into account the mating system (outcrossing rate,  $t$ ). Thus,  
 225 neighborhood size can be defined with the following equation (Crawford 1984) :

$$N_e^b = 4\pi(\sigma_s^2 + \frac{t\sigma_p^2}{2})\delta(1+t) \quad (2)$$

226 where  $\sigma_s$  is the standard deviation of seed dispersal distance,  $\sigma_p$  is the standard  
 227 deviation of pollen dispersal distance, and  $\delta$  is the density of potential parents.

228 Neighborhood size in plants can be estimated by marking pollen and seeds with  
 229 fluorescent dyes, tags, or stable isotope enrichment (Carlo *et al.*, 2009). However,  
 230 these methods do not measure effective pollen or seed movement, but they may  
 231 be combined with genetic analysis to assess genetic identity and relatedness with  
 232 hypervariable DNA markers (Levin, 1988; Nason *et al.*, 1998; Godoy & Jordano,  
 233 2001) to achieve reliable estimates of both effective population size and neighbor-  
 234 hood area.

235 The extent of neighborhood area in plants can be extremely variable, depending  
 236 on life-history attributes such as life-span, spacing patterns, mating system, etc.  
 237 Even a limited sample of available information (Table S1) highlights the fact that  
 238 the size of neighborhood areas can in some cases exceed the geographic limits of  
 239 local populations (Nason *et al.*, 1998). The size of neighborhood areas may en-  
 240 compass at least four orders of magnitude,  $10^{-2} - 10^2$  km in radius, and include  
 241 many individuals. Therefore, reference to this "genetic/evolutionary" paradigm  
 242 and reference to the geographic boundaries (*sensu* Waples & Gaggiotti, 2006) may  
 243 be instrumental to understand the actual role of LDD events in shaping the struc-

244 turing of genetic pools and contributing to gene dispersal.

245 Whenever there is a large discrepancy between population area extent and  $N_e^b$   
246 we might expect the frequency of  $LDL_{loc}$  and  $LDL_{neigh}$  differ enormously. For  
247 example, relatively small  $N_e^b$  may rise the importance of  $LDL_{loc}$  in preserving  
248 scenarios of panmixia within a local population, as most distant dispersal events  
249 will disseminate seeds outside the neighborhood of maternal plants.

## 250 Empirical analysis of contributions to LDD

251 Empirical evaluation of differential contributions to the different forms of LDD  
252 events outlined in Table 1 requires identification of source trees as well as assign-  
253 ment of the dispersed propagules to specific vectors or functional groups of vectors  
254 (Jordano *et al.*, 2007). Recently, DNA-barcoding techniques have been developed  
255 and successfully applied to the identification of frugivore species contributing to  
256 specific seed dispersal events whose source can be identified with genetic, direct  
257 assignment techniques (González-Varo *et al.*, 2014). Otherwise, visual identifica-  
258 tion can reliably assign the genotyped seeds to frugivore species groups based on  
259 specific characteristics of scats and regurgitations (Jordano *et al.*, 2007).

260 We inferred the frugivore groups contributing dispersal events by visually iden-  
261 tifying scats and regurgitations in seed traps and line transects (see Jordano  
262 *et al.*, 2007, and Suppl. Mat. for additional details of methods). These frugivore  
263 functional groups include up to 38 bird and 4 mammal species feeding on *P. ma-*  
264 *haleb* fruits (Jordano & Schupp, 2000). Here, we differentiate four major frugivore

265 groups: large carnivorous mammals (such as foxes, badgers, and stone martens);  
266 two species of medium-sized frugivorous birds, mistle thrushes (*T. viscivorus*),  
267 and carrion crows (*Corvus corone*); and a pool of small-sized frugivorous birds,  
268 including warblers, redstarts, and robins (Jordano *et al.*, 2007).

269 To a large extent, short-distance dispersal events (strict-sense,  $SDD_{loc}$  events)  
270 are contributed by small- and medium-sized (*Turdus*) frugivorous birds (Table 2).  
271 Given the relatively reduced  $N_e^b$  area of *P. mahaleb* (Suppl. Mat. Table S1),  
272  $< 1km^2$ , well below the extent of the local study population (Garcia *et al.*, 2007,  
273 2005), we cannot estimate  $LDD_{neigh}$  events (Table 2), as all LDD events outside  
274 the reference population occur, by definition, outside the  $N_e^b$  area. Larger fru-  
275 givores such as corvids and the pigeon *Columba palumbus* contribute most LDD  
276 events, and most immigrant seeds potentially dispersed from other populations  
277 (Fig. S2). Notably, strict-sense long-distance dispersal ( $LDD_{ss}$ ) appears consis-  
278 tently associated with large-bodied frugivores (Table 2), most likely associated  
279 with a greater frequency of movements outside the local population (Fig. 4).

280 Empirically mapping of dispersal events for either pollen or seed disseminated by  
281 animals may result in a complex pattern of different combinations of dispersal  
282 events (Fig. S1), as animal movements are overlaid onto plant populations occu-  
283 pying complex landscapes, resulting in different types of SDD and LDD events.

## 284 Long-Distance Dispersal: the ecology of extreme events

285 Long-distance dispersal (LDD) is a major component of the population dynamics,  
286 genetic structure, and biogeographic history of plant species. It determines the  
287 colonization ability of new habitats and the possibilities for fragmented populations  
288 to sustain a cohesive metapopulation by immigration-emigration dynamics that  
289 rely on LDD events (Nathan *et al.*, 2008b; Schurr *et al.*, 2009). Yet our current  
290 understanding of the extent, frequency, and consequences of LDD is very limited.  
291 On one hand, theoretical models fail to predict accurately the behavior of the tail  
292 of the dispersal functions, and thus fail to predict very basic properties of LDD. On  
293 the other hand, we still have very limited documentation of actual LDD events in  
294 natural populations and we still see LDD as a sporadic, rarely far-reaching process  
295 still marked with the stamp of natural history curiosity.

296 Combining spatially-explicit references to the geographic population limits and the  
297 genetic neighborhood area extent ( $N_e^b$ ) helps avoiding some imprecision in setting  
298 distance thresholds to characterize LDD events (Jones & Muller-Landau, 2008).  
299 In addition, the framework outlined in Table 1 bridges the combined demographic  
300 and genetic effects of LDD events. When methods available to assign frugivore  
301 taxa to the analyzed dispersal events, as in the study case with *P. mahaleb*, a  
302 classification in the four categories of events is possible.

303 The frugivore assemblage of *P. mahaleb* is composed by a diversified set of animal  
304 species spanning a wide size range, ca. 12-14000 g in body mass. We might expect  
305 that this extreme variation translates in an ample pattern of foraging modes, move-

306 ment distances, and fruit/seed processing (Jordano & Schupp, 2000). If the results  
307 for *P. mahaleb* are generalizable to other disperser assemblages, it seems that the  
308 functional roles of frugivore species in terms of contributions to LDD events are  
309 structured in two distinct groups: small-bodied frugivores, with substantial con-  
310 tributions to SDD events, and large-bodied species with a disproportionate contri-  
311 bution to LDD events. Both components of this sort of diplochorous (vander Wall  
312 & Longland, 2004) dispersal system are very frequent in fleshy-fruited plants with  
313 diversified frugivore assemblages (Galetti *et al.*, 2013). In such cases, small-bodied  
314 frugivores largely contribute the short-distance dispersal key to support *in situ*  
315 recruitment and population persistence. Yet the large-bodied frugivores distinctly  
316 contribute LDD events that sustain the connectivity of metapopulation scenarios  
317 (Urban & Keitt, 2001). As shown in Table 1, SDD and LDD events can be more  
318 complex when we consider the contributions to gene flow via seed and the conse-  
319 quences in terms of structure and spatial distributions of the genetic pools. For  
320 example, local, within-population, dispersal events may vary enormously in terms  
321 of genetic effects and local structuring of the genetic pools depending on whether  
322 they specifically contribute  $SDD_{loc}$  or instead,  $LDD_{loc}$ . Note that only the latter  
323 actually contribute erasing any form of local genetic structure by contributing to  
324 increased genetic neighborhoods.

325 A number of classic studies have demonstrated that the activity of large frugivores  
326 may also significantly contribute to SDD events and inefficient dispersal because  
327 of, i.e., territorial defence, short gut retention times relative to on-tree foraging,  
328 frequent revisit of same trees and perches, etc., resulting in substantial SDD

events (Pratt & Stiles, 1983; Pratt, 1984; Snow & Snow, 1984, 1988; Wheelwright, 1991). Yet these large-bodied frugivores are crucial for both  $LDD_{loc}$  and  $LDD_{ss}$ , given that extensive movement patterns and extremely large foraging ranges may frequently contribute dissemination beyond distance thresholds defined with either spatial landscape or genetic references. Recent analyses of the movement ecology of large frugivores, coupled with results of their seed dispersal services emphasize that LDD are by no means exceptional, either in terms of frequency and extent (e.g., Westcott *et al.*, 2005; Bueno *et al.*, 2013; Morales *et al.*, 2013; Carlo *et al.*, 2013). In addition, medium-sized birds such as thrushes (*Turdus* spp.) can contribute substantial  $LDD_{loc}$  events, i.e., local LDD events contributing to erase local population genetic structuring, effectively increasing the size of genetic neighborhoods. In the case of *P. mahaleb* up to 55.49% of their dispersal events are  $LDD_{loc}$  events. These birds are efficient seed dispersers of *P. mahaleb* and other fleshy-fruited species (Snow & Snow, 1988; Jordano & Schupp, 2000; Carlo *et al.*, 2013), also showing significant contributions of  $LDD_{ss}$  events.

As defined in our framework (Table 1), LDD, and in particular  $LDD_{ss}$  events are a specific case of extreme events (García & Borda-de Água, 2017) consistently associated with large-sized frugivores, yet including also medium-sized and highly efficient frugivorous bird species. Robustly characterizing the expected frequencies and extent of those extreme events would be crucial to properly assess the functional role of frugivores and the full range of influences (demographic, genetic) in plant populations.

### 351 Challenges and future avenues for research

352 Pollen and seed dispersal in plants are essentially spatially-structured processes  
353 for which the outcomes of interactions with dispersal vectors is intimately linked  
354 to landscape features. Given this mechanistic link between the features of the  
355 vector and the environments where its displacement occurs (Nathan *et al.*, 2008a),  
356 consideration of landscape is key to understand the consequences of LDD events.  
357 Yet these consequences hit two central aspects of plant life-histories: the demo-  
358 graphic recruitment process (Harper, 1977), and the genetic signatures of pollen-  
359 and seed-mediated gene flow in complex landscapes (Sork *et al.*, 1999). Recent  
360 evidences point out that the selective extinction of large-bodied frugivores may  
361 significantly impact plant populations dependent on frugivores both in terms of  
362 recruitment (Traveset *et al.*, 2012; Pérez-Méndez *et al.*, 2015) and genetic con-  
363nectivity (Pérez-Méndez *et al.*, 2016). Frugivore downsizing represents a lasting  
364 challenge for the collapse of seed dispersal processes where  $LDD_{ss}$  events are cru-  
365 cial for population persistence and the cohesion of fragmented populations within  
366 metapopulation scenarios.

367 We advocate (also see Jordano & Godoy, 2002; Nathan *et al.*, 2003; Jones &  
368 Muller-Landau, 2008; Hardesty *et al.*, 2011) a combination of approaches includ-  
369 ing large-scale biotelemetry to characterize animal movement, coupled with large-  
370 scale genetic sampling of dispersed propagules, and demogenetic approaches that  
371 combine both demographic and genetic research. A crucial aspect would be to  
372 effectively associate the role of individual frugivore species to specific dispersal

373 outcomes, by identifying the actual disperser contributing a dissemination event  
374 (González-Varo *et al.*, 2014) and simultaneously characterizing the source maternal  
375 plant (Jordano & Godoy, 2002).

376 The actual challenges to properly characterize the typologies of LDD events out-  
377 lined in Table 1 will probably persist. We need more efficient quantitative ap-  
378 proaches to assess these infrequent events, that occur over enormous spatial scales  
379 and that need to be documented with sample sizes sufficient to facilitate modeling  
380 efforts and robust statistical inferences. These are not trivial difficulties given the  
381 urgency to assess how forest loss, defaunation, genetic purging due to logging, etc.,  
382 alter plant populations.

383 *Acknowledgements.* I am indebted to Cristina García, José A. Godoy, Manolo  
384 Carrión, Juan Luis García-Castaño, Jesús Rodríguez and, especially, Juan Miguel  
385 Arroyo for generous help with field and laboratory work and making possible this  
386 study. I appreciate the help and advice of Cristina García and Etienne Klein dur-  
387 ing the final stages of the manuscript. The study was supported by a Junta de  
388 Andalucía Excellence Grant (RNM-5731), as well as a Severo Ochoa Excellence  
389 Award from the Ministerio de Economía y Competitividad (SEV-2012-0262). The  
390 Agencia de Medio Ambiente, Junta de Andalucía, provided generous facilities that  
391 made possible this study in the Andalusian natural parks (Sierra de Cazorla, Al-  
392 cornocales) and authorized my work there.

## References

- Bueno, R.S., Guevara, R., Ribeiro, M.C., Culot, L., Bufalo, F.S. & Galetti, M. (2013) Functional Redundancy and Complementarities of Seed Dispersal by the Last Neotropical Megafrugivores. *PLoS ONE*, **8**(2), e56252, doi:10.1371/journal.pone.0056252.s003.
- Carlo, T.A., García, D., Martínez, D., Gleditsch, J.M. & Morales, J.M. (2013) Where do seeds go when they go far? Distance and directionality of avian seed dispersal in heterogeneous landscapes. *Ecology*, **94**(2), 301–307.
- Carlo, T., Tewksbury, J., Martínez, C. & Mart (2009) A new method to track seed dispersal and recruitment using  $^{15}\text{N}$  isotope enrichment. *Ecology*, **90**, 3516–3525.
- Clark, J.S., Fastie, C., Hurttt, G.C., Jackson, S.T., Johnson, C., King, G., Lewis, M., Lynch, J., Pacala, S., Prentice, C., Schupp, E., Webb III, T. & Wyckoff, P. (1998) Reid's paradox of rapid plant migration. *BioScience*, **48**, 13–24.
- Clark, J. (1998) Why trees migrate so fast: Confronting theory with dispersal biology and the paleorecord. *The American Naturalist*, **152**, 204–224.
- Cousens, R.D., Dytham, C. & Law, R. (2008) *Dispersal in plants. A population perspective*. Oxford University Press, Oxford, UK.
- D'Aloia, C.C., Bogdanowicz, S.M., Majoris, J.E., Harrison, R.G. & Buston, P.M. (2013) Self-recruitment in a Caribbean reef fish: a method for approximating dispersal kernels accounting for seascape. *Molecular Ecology*, n/a–n/a, doi:10.1111/mec.12274.

Galetti, M., Guevara, R., Côrtes, M.C., Fadini, R., Von Matter, S., Leite, A.B., Labecca, F., Ribeiro, T., Carvalho, C.S., Collevatti, R.G., Pires, M.M., Guimaraes, P.R., Brancalion, P.H., Ribeiro, M.C. & Jordano, P. (2013) Functional extinction of birds drives rapid evolutionary changes in seed size. *Science*, **340**(6136), 1086–1090, doi:10.1126/science.1233774.

Garcia, C., Arroyo, J., Godoy, J. & Jordano, P. (2005) Mating patterns, pollen dispersal, and the ecological maternal neighbourhood in a *Prunus mahaleb* L. population. *Molecular Ecology*, **14**, 1821–1830, doi:10.1111/j.1365-294X.2005.02542.x.

Garcia, C., Jordano, P. & Godoy, J.A. (2007) Contemporary pollen and seed dispersal in a *Prunus mahaleb* population: patterns in distance and direction. *Molecular Ecology*, **16**, 1947–1955, doi:10.1111/j.1365-294X.2006.03126.x.

García, C. & Borda-de Água, L. (2017) Extended dispersal kernels: insights from statistics of extremes. *Journal of Ecology*, **00**, 000–000.

Godoy, J.A. & Jordano, P. (2001) Seed dispersal by animals: exact identification of source trees with endocarp DNA microsatellites. *Molecular Ecology*, **10**(9), 2275–2283.

González-Varo, J.P., Arroyo, J.M. & Jordano, P. (2014) Who dispersed the seeds? The use of DNA barcoding in frugivory and seed dispersal studies. *Methods in Ecology and Evolution*, **5**(8), 806–814, doi:10.1111/2041-210X.12212.

Hampe, A. & Jump, A.S. (2011) Climate Relicts: Past, Present, Future. *An-*

nual Review of Ecology Evolution and Systematics, **42**(1), 313–333, doi:10.1146/annurev-ecolsys-102710-145015.

Hardesty, B.D., Hubbell, S.P. & Bermingham, E. (2006) Genetic evidence of frequent long-distance recruitment in a vertebrate-dispersed tree. *Ecology Letters*, **9**, 516–525.

Hardesty, B.D., Metcalfe, S.S. & Westcott, D.A. (2011) Persistence and spread in a new landscape: Dispersal ecology and genetics of Miconia invasions in Australia. *Acta Oecologica*, **37**(6), 657–665, doi:10.1016/j.actao.2011.06.006.

Harper, J. (1977) *Population Biology of Plants*, 1st edn. Academic Press, London.

Howe, H.F. & Miriti, M.N. (2004) When seed dispersal matters. *BioScience*, **54**(7), 651–660.

Jones, F.A. & Muller-Landau, H.C. (2008) Measuring long-distance seed dispersal in complex natural environments: an evaluation and integration of classical and genetic methods. *Journal of Ecology*, **96**(4), 642–652, doi:10.1111/j.1365-2745.2008.01400.x.

Jordano, P., García, C., Godoy, J.A. & García-Castaño, J.L. (2007) Differential contribution of frugivores to complex seed dispersal patterns. *Proceedings Of The National Academy Of Sciences Of The United States Of America*, **104**(9), 3278–3282, doi:10.1073/pnas.0606793104.

Jordano, P. & Godoy, J. (2002) Frugivore-generated seed shadows: a landscape view of demographic and genetic effects. *Seed dispersal and frugivory: ecology,*

*evolution, and conservation* (eds. D. Levey, W. Silva & M. Galetti). Book chapter, Wallingford, UK, pp. 305–321.

Jordano, P. & Schupp, E.W. (2000) Seed disperser effectiveness: the quantity component and patterns of seed rain for *Prunus mahaleb*. *Ecological Monographs*, **70**(4), 591–615, doi:10.2307/2657187?ref=no-x-route:1d70d282eb4ed20cce6eac7274a631bb.

Kinlan, B., Gaines, S. & Lester, S. (2005) Propagule dispersal and the scales of marine community process. *Diversity and Distributions*, **11**, 139–148, doi: 10.1111/j.1366-9516.2005.00158.x.

Kramer, A., Ison, J., Ashley, M. & Howe, H. (2008) The paradox of forest fragmentation genetics. *Conservation Biology*, 8, doi:10.1111/j.1523-1739.2008.00944.x.

Levin, D.A. (1988) The paternity pools of plants. *The American Naturalist*, **132**, 309–317, doi:10.2307/2461984.

Morales, J.M., García, D., Martínez, D., Rodríguez-Pérez, J. & Herrera, J.M. (2013) Frugivore Behavioural Details Matter for Seed Dispersal: A Multi-Species Model for Cantabrian Thrushes and Trees. *PLoS ONE*, **8**(6), e65216, doi:10.1371/journal.pone.0065216.s005.

Nason, J.D., Herre, E. & Hamrick, J.L. (1998) The breeding structure of a tropical keystone plant resource. *Nature*, **391**, 685–687.

Nathan, R., Perry, G., Cronin, J.T., Strand, A.E. & Cain, M.L. (2003) Methods for estimating long-distance dispersal. *Oikos*, **103**, 261–273.

Nathan, R. (2005) Long-distance dispersal research: building a network of yellow brick roads. *Diversity and Distributions*, **11**(2), 125–130, doi:10.1111/j.1366-9516.2005.00159.x.

Nathan, R. (2006) Long-distance dispersal of plants. *Science*, **313**, 786–788.

Nathan, R., Getz, W.M., Revilla, E., Holyoak, M., Kadmon, R., Saltz, D. & Smouse, P.E. (2008a) A movement ecology paradigm for unifying organismal movement research. *Proceedings of the National Academy of Sciences USA*, **105**(49), 19052–19059, doi:10.1073/pnas.0800375105.

Nathan, R., Schurr, F.M., Spiegel, O., Steinitz, O., Trakhtenbrot, A. & Tsoar, A. (2008b) Mechanisms of long-distance seed dispersal. *Trends In Ecology & Evolution*, **23**(11), 638–647, doi:10.1016/j.tree.2008.08.003.

Pérez-Méndez, N., Jordano, P. & Valido, A. (2015) Downsized mutualisms: Consequences of seed dispersers' body-size reduction for early plant recruitment. *Perspectives In Plant Ecology Evolution And Systematics*, **17**(2), 151–159, doi: 10.1016/j.ppees.2014.12.001.

Pérez-Méndez, N., Jordano, P. & Valido, A. (2016) Persisting in defaunated landscapes: reduced plant population connectivity after seed dispersal collapse. *In 2nd. review.*

Pratt, T. (1984) Examples of tropical frugivores defending fruit-bearing plants. *Condor*, **86**, 123–129.

- Pratt, T. & Stiles, E. (1983) How long fruit-eating birds stay in the plants where they feed - implications for seed dispersal. *The American Naturalist*, **122**(6), 797–805.
- Schupp, E. (1995) Seed–seedling conflicts, habitat choice, and patterns of plant recruitment. *American Journal of Botany*, **82**, 399–409, doi:10.2307%2F2445586.
- Schurr, F.M., Spiegel, O., Steinitz, O., Trakhtenbrot, A., Tsoar, A. & Nathan, R. (2009) Long-distance seed dispersal. *Annual Plant Reviews*, **38**, 204–237.
- Seidler, T.G. & Plotkin, J.B. (2006) Seed dispersal and spatial pattern in tropical trees. *PLoS Biology*, **4**, 2132–2137, doi:10.1371/journal.pbio.0040344.
- Slatkin, M. (1985) Gene flow in natural populations. *Annual Review of Ecology and Systematics*, doi:10.2307/2097054.
- Snow, B. & Snow, D. (1984) Long-term defence of fruit by mistle thrushes *Turdus viscivorus*. *Ibis*, **126**, 39–49.
- Snow, B. & Snow, D. (1988) *Birds and berries*. Poyser, Calton, UK.
- Sork, V., Nason, J., Campbell, D. & Fernández, J. (1999) Landscape approaches to historical and contemporary gene flow in plants. *Trends In Ecology & Evolution*, **14**, 224–224.
- Traveset, A., González-Varo, J.P. & Valido, A. (2012) Long-term demographic consequences of a seed dispersal disruption. *Proceedings Of The Royal Society B-Biological Sciences*, doi:10.1098/rspb.2012.0535.

Urban, D. & Keitt, T. (2001) Landscape connectivity: a graph-theoretic perspective. *Ecology*, **82**, 1205–1218.

vander Wall, S. & Longland, W. (2004) Diplochory: are two seed dispersers better than one? *Trends In Ecology & Evolution*, **19**, 155–161, doi:10.1016/j.tree.2003.12.004.

Waples, R. & Gaggiotti, O. (2006) What is a population? An empirical evaluation of some genetic methods for identifying the number of gene pools and their degree of connectivity. *Molecular Ecology*, **15**, 1419–1439, doi:10.1111/j.1365-294X.2006.02890.x.

Westcott, D.A., Bentupperbaumer, J., Bradford, M.G. & McKeown, A. (2005) Incorporating patterns of disperser behaviour into models of seed dispersal and its effects on estimated dispersal curves. *Oecologia*, **146**, 57–67.

Wheelwright, N. (1991) How long do fruit-eating birds stay in the plants where they feed? *Biotropica*, **23**(1), 29–40.

Wright, S. (1943) Isolation by distance. *Genetics*, **28**, 114–138.

Wright, S. (1946) Isolation by distance under diverse systems of mating. *Genetics*, **31**, 39–59.

Wright, S. (1969) *Evolution and the genetics of populations. Vol. 2: The theory of gene frequencies*. University of Chicago Press, Chicago, USA.

**Table 1:** Types of dispersal as a function of population area limits and genetic neighborhood limits. See Fig. 2 for a graphical representation of the four scenarios.

Genetic neighborhood limit	Population geographic limit
Within	Within
Outside	Outside
Within	Local, short-distance dispersal, $SDD_{loc}$
Outside	Local, long-distance dispersal, $LDD_{loc}$
Within	Within neighborhood, long-distance dispersal, $LDD_{neigh}$
Outside	Strict sense long-distance dispersal, $LDD_{ss}$

**Table 2:** Relative frequencies of *Prunus mahaleb* seed dispersal events for different frugivore groups according to population area limits and genetic neighborhood limits. See Fig. 2 for a graphical representation of the four scenarios.  $N = 655$  seeds (see Table 1 in Jordano *et al.*, 2007). Given that the estimated neighborhood size is smaller than population area,  $LDD_{neigh}$  would be zero.

Frugivore group	Within-population, within-neighborhood $SDD_{loc}$	Within-population, long-distance $LDD_{loc}$	Outside-population, within-neighborhood $LDD_{neigh}$	Strict-sense long-distance $LDD_{ss}$	$N$ seeds
Small-birds	0.7842	0.0171	0.00	0.1986	292
<i>Turdus</i>	0.2370	0.5549	0.00	0.2081	173
Large-birds	0.0435	0.3913	0.00	0.5652	23
Mammals	0.0120	0.2455	0.00	0.7425	167

## Figures

393 **Figure 1.** The two approaches used in analyses of dispersal processes in plants.  
394 A, the “forward” approach attempts to track the dispersal events away from the  
395 known sources, e.g., by tracking the movement patterns of frugivores as they leave  
396 fruiting plants after feeding. B, the “backward” approach attempts to reconstruct  
397 the most likely source of a dispersed propagule by inferring the sources given the  
398 propagule delivery pattern, the fecundity of potential sources, and the dispersal  
399 function. The main technical challenge in A is to sample enough dispersal events  
400 away from the source to be able to fully characterize the tail (long-distance dis-  
401 persal, LDD, events) of the dispersal function. In B, the main challenge is to have  
402 a robust sampling scheme with propagule collectors (e.g., seed traps) and a good  
403 characterization of the potential sources to derive robust estimates of the actual  
404 sources with inverse-modeling techniques.

405

406 **Figure 2.** Schematic representation of different types of long-distance dispersal  
407 events in relation to the geographical limits of local populations (dashed lines)  
408 and the genetic neighborhood area  $N_e^b$  (grey area) of specific individual plants  
409 (squares). Dispersal events (arrows) can be classified depending on their actual  
410 incidence on propagule movement outside these spatially-explicit reference areas  
411 (Table 1). Strict-sense long-distance dispersal events ( $LDD_{ss}$ ) just include the  
412 LDD events that disseminate propagules out of *both* the population and genetic  
413 neighborhood boundaries. A, the neighborhood area is included within the geo-

414 graphic limits of the population, with some dispersal events potentially contribut-  
415 ing local LDD; B, the neighborhood area is much larger than the geographic limits  
416 of the population.

417

418 **Figure 3.** Empirical frequency distributions of seed dispersal events as a function  
419 of dispersal distance by animal frugivores consuming *Prunus mahaleb* fruits. In  
420 red, left (inset), frequencies of within-population dispersal events inferred from di-  
421 rect assignment based on seed endocarp genotypes and maternal trees genotypes.  
422 Larger frame, left, contributions of four functional frugivore groups (small birds,  
423 medium- and large-sized birds, and mammals) to seed dissemination and propor-  
424 tional contributions (right bar) to dispersal of the inferred immigrant seeds (i.e.,  
425 those not matching any maternal tree in the study population) (Jordano *et al.*,  
426 2007).

427

428 **Figure 4.** Differential contributions of functional groups of frugivores to the  
429 four combinations of *Prunus mahaleb* seed dispersal events outlined in Table 1.  
430 These result from dissemination within (yellow) or outside (blue) the population  
431 geographic limits; within-population dispersal events can either be short-distance  
432 ( $SDD_{loc}$ ) or local LDD ( $LDD_{loc}$ ) depending on the size of the genetic neigborhood.  
433 Dispersal outside the local population can entail short-distance dispersal, if within  
434 the genetic neighborhood area limits ( $SDD_{neigh}$ ) (yellow) or represent strict-sense  
435 LDD ( $LDD_{ss}$ ) (blue).

436

<sup>437</sup> **Online Support Material and data accessibility**

<sup>438</sup> This review does not use new raw data, but includes some re-analyses of pre-  
<sup>439</sup> viously published material. All the original data supporting the paper, R code,  
<sup>440</sup> supplementary figures, and summaries of analytical protocols is available at the  
<sup>441</sup> author's GitHub repository ([https://github.com/pedroj/MS\\_LDD](https://github.com/pedroj/MS_LDD)), with DOI:  
<sup>442</sup> #/zenodo.#.







