

PLOS Biology

Chasing Ecological Interactions

--Manuscript Draft--

Manuscript Number:	PBIOLOGY-D-16-01135R1
Full Title:	Chasing Ecological Interactions
Article Type:	Research Matters (invitation only)
Corresponding Author:	Pedro Jordano Estacion Biologica de Doñana CSIC Sevilla, Sevilla SPAIN
First Author:	Pedro Jordano
Order of Authors:	Pedro Jordano
Abstract:	Basic research on biodiversity has concentrated on individual species--naming new species, studying distribution patterns, and analyzing their evolutionary relationships. Yet biodiversity is more than a collection of individual species; it is the combination of biological entities and processes that support life on Earth. To understand biodiversity, we must catalog it but we must also assess the ways species interact with other species to provide functional support for the Tree of Life. Ecological interactions may be lost well before the species involved in those interactions go extinct; their ecological functions disappear even though they remain. Here I address the challenges in studying the functional aspects of species interactions and how basic research is helping us address the fast-paced extinction of species due to human activities.
Additional Information:	
Question	Response
<p>Ethics Statement</p> <p>All research involving human participants must have been approved by the authors' institutional review board or equivalent committee(s) and that board must be named by the authors in the manuscript. For research involving human participants, informed consent must have been obtained (or the reason for lack of consent explained, e.g. the data were analyzed anonymously) and all clinical investigation must have been conducted according to the principles expressed in the Declaration of Helsinki. Authors should submit a statement from their ethics committee or institutional review board indicating the approval of the research. We also encourage authors to submit a sample of a patient consent form and may require submission of completed forms on particular occasions.</p> <p>All animal work must have been conducted according to relevant national and international guidelines. In accordance with the recommendations of the Weatherall report, "The use of non-human primates in research" we specifically require authors to include details of animal welfare and steps taken</p>	N/A

<p>to ameliorate suffering in all work involving non-human primates. The relevant guidelines followed and the committee that approved the study should be identified in the ethics statement.</p> <p>Please enter your ethics statement below and place the same text at the beginning of the Methods section of your manuscript (with the subheading Ethics Statement). Enter "N/A" if you do not require an ethics statement.</p>	
<p>Competing Interest</p> <p>For yourself and on behalf of all the authors of this manuscript, please declare below any competing interests as described in the "PLoS Policy on Declaration and Evaluation of Competing Interests."</p> <p>You are responsible for recognizing and disclosing on behalf of all authors any competing interest that could be perceived to bias their work, acknowledging all financial support and any other relevant financial or competing interests.</p> <p>If no competing interests exist, enter: "The authors have declared that no competing interests exist."</p> <p>If you have competing interests to declare, please fill out the text box completing the following statement: "I have read the journal's policy and have the following conflicts"</p> <p>* typeset</p>	<p>The authors have declared that no competing interests exist.</p>

Cover letter

August 6, 2016

Dear editor,

please find enclosed a revised version of my MS entitled ” *Chasing Ecological Interactions*”, which I submit to your consideration for publication in **PLoS Biology**, as an article contribution to the section *Research Matters*. As you can see, I have accepted all the suggestions by the associated editor. I very much appreciate the help and comments, which have improved very much the text.

I hope you consider this version acceptable for publication.

Sincerely,
Pedro Jordano

Response to Reviewers

August 7, 2016

Dear editor,

As you can see, all the suggestions by the associated editor have been incorporated in this revised version. I very much appreciate the help and comments, which have improved very much the text.

I'm including also the figures and maintext manuscript in the format requested by the production.

Please find also in this submission the authorizations received from the authors of the photos used in Fig. 1.

I hope you consider this version acceptable for publication.

Sincerely,
Pedro Jordano

Chasing Ecological Interactions

Pedro Jordano^{1*}

1 Integrative Ecology Group, Estación Biológica de Doñana, CSIC, Pabellón del Perú, Avda. María Luisa, s/n, E-41013 Sevilla, Spain

* jordano@ebd.csic.es

Abstract

Basic research on biodiversity has concentrated on individual species--naming new species, studying distribution patterns, and analyzing their evolutionary relationships. Yet biodiversity is more than a collection of individual species; it is the combination of biological entities and processes that support life on Earth. To understand biodiversity, we must catalog it but we must also assess the ways species interact with other species to provide functional support for the Tree of Life. Ecological interactions may be lost well before the species involved in those interactions go extinct; their ecological functions disappear even though they remain. Here I address the challenges in studying the functional aspects of species interactions and how basic research is helping us address the fast-paced extinction of species due to human activities.

I am tempted to give one more instance showing how plants and animals, most remote in the scale of nature, are bound together by a web of complex relations.
(Darwin Ch. 1860. On the origin of species by means of natural selection. Chapter 3, p. 75).

There is a much more insidious kind of extinction: the extinction of ecological interactions.
(Janzen, D.H. 1974. The deflowering of Central America. *Natural History*, 83: 48-53).

Suppose you want to build the LEGO® *Triceratops Trapper* model. It has 256 pieces, corresponding to 74 distinct parts. It's a relatively simple model, yet impossible to build by assembling these 256 pieces at random. To build the fully functional Triceratops Trapper we need to know more than the inventory of its parts: we also need to know how the different pieces fit together. We'll have a functional Triceratops Trapper only if we assemble the model connecting its component pieces the right way. Like the pieces of our Triceratops Trapper, species in ecosystems are not connected (linked) by random interactions. They fit together to form functional units that share some basic properties, independent on the type of ecosystem and even on the type of ecological interaction. This Web of Life [1] shapes the wireframe that supports biodiversity, and ecological interactions among the species that make up this web provide essential services and functions that support its persistence. Interactions might be lost (go extinct) even well before the species. For instance, the "empty forest syndrome" [2] describes situations where animals and plants may persist in disturbed areas (e.g., a tropical forest fragment) yet in such reduced abundance that their functional ecological role is lost. Interaction extinction may result in the loss of important ecological functions such as pollination and seed dispersal that are crucial for forest regeneration and ecosystem persistence. These losses of interactions cause unprecedented changes in cascade in natural communities (ie, like trophic cascades), implying losses of ecological functions. Foreexample, just imagine the myriad consequences of extinctions of pollinators and frugivore seed dispersers for ecosystems like tropical rainforests, where more than 90% of the woody plant species depend on frugivores to support their life cycles. The disappearance of frugivores could unleash a cascade of effects with unprecedented changes in the structure and function of an ecosystem.

No single species on Earth lives without interacting with other species. These interactions are the backbone of biodiversity and create the architectural foundation of ecosystems. Exploring and inventorying biodiversity represents a fundamental challenge

for basic research in ecology and conservation biology. This basic knowledge is urgently required to properly diagnose the status of biodiversity conservation and develop early warning signals for the potential disappearance of interactions and collapse of communities.

Diversity: of species and their interactions

The Web of Life results from the assembly of species that interact with each other in a variety of ways, forming complex interaction networks (Fig. 1). A myriad of interaction modes exist in nature, reflecting the complexity of natural histories of partner species. Interactions take the form of predation, competition, commensalism, amensalism, mutualism, symbiosis, parasitism and in all cases involve reciprocal effects for the interacting species. Recent basic research on the topology and structure of these networks has revealed universal patterns that ultimately affect their stability and resilience. Yet we are far from fully documenting all the types of interaction modes that exist even in simple ecosystems.

Just as we sample individuals of free-living species to estimate the diversity in a particular area or ecosystem, we can sample interactions. In this way we can assess the full complexity of ecosystem structure. Yet the exercise is not trivial. We are far from fully understanding the minimum set of functional links that are needed to support and restore damaged ecosystems. We don't even have robust estimates of the total number of species living on Earth. Assessing the diversity of their interactions is a far more daunting task.

Life on Earth is supported by zillions of interactions among species. Understanding these complex systems demands that a large fraction of these interactions be experimentally or computationally probed. This is very difficult, as rapid and effective actions or conservation and restoration of human-disturbed ecosystems urgently require the identification of the minimum amount of complexity that has to be restored to facilitate an ecosystem's persistence.

Considering all the distinct ways in which such highly complex, interactive systems can be decomposed into parts cannot be done on a statistical basis (as, for example, with an ideal gas) because each interaction is particular; genomes, proteins, cells, and species interact in specific ways. The number of ways a complex system can be partitioned is known as Bell's number B_n [5]. For three elements, there are $B_3 = 5$ such partitions; three species may not interact at all, or any two of them can interact, or all three, for a total of five possibilities. Understanding the system requires measuring the probability and magnitude of each of these five. This is out of the feasibility limits for any study [5], given that Bell's number scales supra-exponentially with the number of components. In real plant-animal webs the number of actual pairwise interactions among species in local assemblages scales with species richness (Fig. 1). These real ecological systems would be within the range of $n = 10^3 - 10^5$ or even $n = 10^4 - 10^{6.5}$ components, depending on spatial scale when we move from local to regional and up to continental spatial scales. To fully quantify the size of these interactomes we thus need to focus on what we know about the macroscopic properties of complex ecological interaction networks [1].

Basic conservation science in the Anthropocene: challenges

Most discussions about the effects of the biodiversity crisis have largely focused on the loss of species, ignoring the extinction of ecological interactions. In our lifetimes, a myriad of vertebrate species have experienced population declines and eventual extinctions matching human expansion, i.e. the "Anthropocene defaunation" [3]. Large bodied vertebrates have been especially hard hit, and as a result, many disturbed ecosystems currently host only small- to medium-bodied species, or population sizes so reduced that the species are no longer functional in their ecosystem [2]. If the remnant, extant species fail to provide pivotal services formerly assisted by vanishing large vertebrates, human-driven defaunation may trigger negative cascading effects on ecosystem dynamics. This downgrading process is expected to result, for example, in collapsed mutualisms of pollination and seed dispersal for plants depending on them for regeneration.

Early signals of the collapse of mutualistic interactions may trigger severe population declines which lag behind the loss of seed dispersers or pollinators. Consider, for example, that deforestation, logging, fragmentation and climate change are already having significant impacts on tropical carbon stocks. Further reductions in carbon storage could follow the disruption of mutualisms resulting from the loss of animal pollinators and frugivores acting as seed dispersers for diverse forest plants because forest regeneration may collapse. Loss of key ecological interactions may precede the local extinction of partner species that depend on the key ecological services provided. Without basic ecological and biodiversity research, we'd have no hope of forecasting the consequences and rebuilding functional ecosystems.

Acknowledgments

My work was funded by grant CGL2013-47429P and a Severo-Ochoa Excellence Grant (SEV2012-0262) from the Spanish






Ministerio de Economía y Competitividad (MINECO), and RNM-5731 from the Junta de Andalucía. I am very grateful to Guto Balieiro, Denise Lange and Kleber del-Claro for generously providing photos for Figure 1.

References

1. Thompson JN. The coevolving web of life. *The American Naturalist* 2009; 173: 125–140.
doi:10.1086/595752
2. Redford KH. The empty forest. *Bioscience*. 1992; 42: 412–422.
3. Dirzo R, Young HS, Galetti M, Ceballos G, Isaac NJB, Collen B. Defaunation in the Anthropocene. *Science*. 2014; 345: 401–406. doi: 10.1126/science.1251817
4. Lima-Mendez G, Faust K, Henry N, Decelle J, Colin S, Carcillo F, et al. Ocean plankton. Determinants of community structure in the global plankton interactome. *Science*. 2015; 348: 1262073–1262073. doi:10.1126/science.1262073
5. Koch, C. Modular biological complexity. *Science*. 2012; 337: 531–532.
doi:10.1126/science.1218616.

Figure legend

Figure 1. The structure of ecological interactions. Top, examples of ecological interactions between plants and animals. a, Ariel toucan (*Ramphastos ariel*) (Ramphastidae) feeding on palmito juçara fruits *Euterpe edulis* (Arecaceae) in the Brazilian Atlantic rainforest; b, *Ectatomma tuberculatum* over extra-floral nectaries at the base of a leaf of *Qualea multiflora* (Vochysiaceae) in the cerrado vegetation (Brazil); c, a carpenter bee *Xylocopa violacea* (Apidae) visiting an *Allium ampeloprasum* (Liliaceae) (wild garlic) inflorescence in SE Spain. Bottom, different visualizations of the complexity of interaction networks among species (colored spheres) illustrated by their actual links (light green lines): d, food webs typically describe all the interactions occurring in a given ecosystem with multiple trophic levels; e, most plant–animal interactions can be displayed as bipartite graphs describing the pairwise pattern of mutual interdependencies among two distinct sets of animals (orange nodes) and plants (yellow); f, interactions among species with a higher degree of intimacy, such as ant-plants show a distinct pattern of structure, often with multiple distinct groups (modules) of closely intimate associations. Photos: a, José Augusto Balieiro, with permission; b, Denise Lange and Kleber del-Claro, with permission; c, Pedro Jordano. Images d, e, and c produced with FoodWeb3D, written by R.J. Williams and provided by the Pacific Ecoinformatics and Computational Ecology Lab (www.foodwebs.org).







[Click here to access/download](#)

Other

[Authorizations_photos_CCBY.pdf](#)

