

Subtrees with small branching number

Pedro Marun
Carnegie Mellon University

March 23, 2023

Definition

- ▶ A *tree* is a partially ordered set (T, \leq) such that for every $x \in T$, the set $\{y \in T : y < x\}$ is well-ordered.

Definition

- ▶ A *tree* is a partially ordered set (T, \leq) such that for every $x \in T$, the set $\{y \in T : y < x\}$ is well-ordered.
- ▶ The *height* of a point $x \in T$, denoted $\text{ht}_T(x)$, is the order type of $\{y \in T : y < x\}$.

Definition

- ▶ A *tree* is a partially ordered set (T, \leq) such that for every $x \in T$, the set $\{y \in T : y < x\}$ is well-ordered.
- ▶ The *height* of a point $x \in T$, denoted $\text{ht}_T(x)$, is the order type of $\{y \in T : y < x\}$.
- ▶ For an ordinal α , the α *th level* of T is the set $T_\alpha = \{x \in T : \text{ht}(x) = \alpha\}$.

Definition

- ▶ A *tree* is a partially ordered set (T, \leq) such that for every $x \in T$, the set $\{y \in T : y < x\}$ is well-ordered.
- ▶ The *height* of a point $x \in T$, denoted $\text{ht}_T(x)$, is the order type of $\{y \in T : y < x\}$.
- ▶ For an ordinal α , the α *th level* of T is the set $T_\alpha = \{x \in T : \text{ht}(x) = \alpha\}$.
- ▶ The *height of the tree* T is the ordinal $\text{ht}(T) = \min\{\alpha : T_\alpha = \emptyset\}$.

Definition

- ▶ A *tree* is a partially ordered set (T, \leq) such that for every $x \in T$, the set $\{y \in T : y < x\}$ is well-ordered.
- ▶ The *height* of a point $x \in T$, denoted $\text{ht}_T(x)$, is the order type of $\{y \in T : y < x\}$.
- ▶ For an ordinal α , the α *th level* of T is the set $T_\alpha = \{x \in T : \text{ht}(x) = \alpha\}$.
- ▶ The *height of the tree* T is the ordinal $\text{ht}(T) = \min\{\alpha : T_\alpha = \emptyset\}$.
- ▶ For $X \subseteq \text{ht}(T)$, define $T \upharpoonright X = \{x \in T : \text{ht}_T(x) \in X\}$.

Definition

Let T be a tree.

- ▶ Given a regular cardinal κ , T is a κ -tree iff $\text{ht}(T) = \kappa$ and $|T_\alpha| < \kappa$ for all $\alpha < \kappa$.

Definition

Let T be a tree.

- ▶ Given a regular cardinal κ , T is a κ -tree iff $\text{ht}(T) = \kappa$ and $|T_\alpha| < \kappa$ for all $\alpha < \kappa$.
- ▶ A *chain* in T is a linearly ordered subset of T .

Definition

Let T be a tree.

- ▶ Given a regular cardinal κ , T is a κ -tree iff $\text{ht}(T) = \kappa$ and $|T_\alpha| < \kappa$ for all $\alpha < \kappa$.
- ▶ A *chain* in T is a linearly ordered subset of T .
- ▶ A *cofinal branch* in T is a chain that meets every level.

Definition

Let T be a tree.

- ▶ Given a regular cardinal κ , T is a κ -tree iff $\text{ht}(T) = \kappa$ and $|T_\alpha| < \kappa$ for all $\alpha < \kappa$.
- ▶ A *chain* in T is a linearly ordered subset of T .
- ▶ A *cofinal branch* in T is a chain that meets every level.

Lemma (König [3])

Let T be an \aleph_0 -tree. Then T has a cofinal branch.

Definition

Let T be a tree.

- ▶ Given a regular cardinal κ , T is a κ -tree iff $\text{ht}(T) = \kappa$ and $|T_\alpha| < \kappa$ for all $\alpha < \kappa$.
- ▶ A *chain* in T is a linearly ordered subset of T .
- ▶ A *cofinal branch* in T is a chain that meets every level.

Lemma (König [3])

Let T be an \aleph_0 -tree. Then T has a cofinal branch.

Theorem (Aronszajn [6])

There exists an \aleph_1 -tree with no cofinal branches.

Definition

- ▶ Let T be a tree and $x \in T$. The set of *immediate successors* of T is denoted by $I_T(x)$.

Definition

- ▶ Let T be a tree and $x \in T$. The set of *immediate successors* of T is denoted by $I_T(x)$.
- ▶ Let λ be a cardinal. A tree T is λ -*branching* [respectively $<\lambda$ -*branching*] iff for every $x \in T$, $|I_T(x)| = \lambda$ [respectively $|I_T(x)| < \lambda$].

Definition

- ▶ Let T be a tree and $x \in T$. The set of *immediate successors* of T is denoted by $I_T(x)$.
- ▶ Let λ be a cardinal. A tree T is λ -*branching* [respectively $<\lambda$ -*branching*] iff for every $x \in T$, $|I_T(x)| = \lambda$ [respectively $|I_T(x)| < \lambda$].
- ▶ A *subtree* of a tree T is a subset S of T such that for every $s \in S$ and $t \in T$, if $t < s$, then $t \in S$.

Definition

- ▶ Let T be a tree and $x \in T$. The set of *immediate successors* of T is denoted by $I_T(x)$.
- ▶ Let λ be a cardinal. A tree T is λ -*branching* [respectively $<\lambda$ -*branching*] iff for every $x \in T$, $|I_T(x)| = \lambda$ [respectively $|I_T(x)| < \lambda$].
- ▶ A *subtree* of a tree T is a subset S of T such that for every $s \in S$ and $t \in T$, if $t < s$, then $t \in S$.

Lemma

Let T be an \aleph_1 -tree. If T has no uncountable 1-branching subtrees, then T is Aronszajn.

Definition

- ▶ Let T be a tree and $x \in T$. The set of *immediate successors* of T is denoted by $I_T(x)$.
- ▶ Let λ be a cardinal. A tree T is λ -*branching* [respectively $<\lambda$ -*branching*] iff for every $x \in T$, $|I_T(x)| = \lambda$ [respectively $|I_T(x)| < \lambda$].
- ▶ A *subtree* of a tree T is a subset S of T such that for every $s \in S$ and $t \in T$, if $t < s$, then $t \in S$.

Lemma

Let T be an \aleph_1 -tree. If T has no uncountable 1-branching subtrees, then T is Aronszajn.

Proof.

If b is a cofinal branch, then b is an uncountable 1-branching subtree.



Definition

An infinitely branching \aleph_1 -tree T is *Lindelöf* iff every finitely branching subtree of T is countable.

Definition

An infinitely branching \aleph_1 -tree T is *Lindelöf* iff every finitely branching subtree of T is countable.

The previous lemma shows that

$$\{\text{Lindelöf}\} \subseteq \{\text{Aronszajn}\}$$

Definition

An infinitely branching \aleph_1 -tree T is *Lindelöf* iff every finitely branching subtree of T is countable.

The previous lemma shows that

$$\{\text{Lindelöf}\} \subseteq \{\text{Aronszajn}\}$$

We will show that the inclusion is proper.

Definition

An infinitely branching \aleph_1 -tree T is *Lindelöf* iff every finitely branching subtree of T is countable.

The previous lemma shows that

$$\{\text{Lindelöf}\} \subseteq \{\text{Aronszajn}\}$$

We will show that the inclusion is proper.

First, we explain the terminology:

Definition

An infinitely branching \aleph_1 -tree T is *Lindelöf* iff every finitely branching subtree of T is countable.

The previous lemma shows that

$$\{\text{Lindelöf}\} \subseteq \{\text{Aronszajn}\}$$

We will show that the inclusion is proper.

First, we explain the terminology: Trees are Lindelöf if and only if they have the Lindelöf property with respect to a natural topology.

Definition

An infinitely branching \aleph_1 -tree T is *Lindelöf* iff every finitely branching subtree of T is countable.

The previous lemma shows that

$$\{\text{Lindelöf}\} \subseteq \{\text{Aronszajn}\}$$

We will show that the inclusion is proper.

First, we explain the terminology: Trees are Lindelöf if and only if they have the Lindelöf property with respect to a natural topology.

Definition (Nyikos)

Let T be a tree. The *fine wedge topology* on T is generated by all sets of the form $\uparrow x$ and their complements, where

$\uparrow x = \{y \in T : x \leq y\}$ and $x \in T$.

Definition

An infinitely branching \aleph_1 -tree T is *Lindelöf* iff every finitely branching subtree of T is countable.

The previous lemma shows that

$$\{\text{Lindelöf}\} \subseteq \{\text{Aronszajn}\}$$

We will show that the inclusion is proper.

First, we explain the terminology: Trees are Lindelöf if and only if they have the Lindelöf property with respect to a natural topology.

Definition (Nyikos)

Let T be a tree. The *fine wedge topology* on T is generated by all sets of the form $\uparrow x$ and their complements, where

$\uparrow x = \{y \in T : x \leq y\}$ and $x \in T$.

If $X \subseteq T$, write $\uparrow X = \{y \in T : \exists x \in X (x \leq y)\}$.

Lemma

If $x \in T$, the family $\{\uparrow x \setminus \uparrow F : F \in [I(x)]^{<\omega}\}$ is a local basis of open neighbourhoods of x . In particular, the topology is Hausdorff.

Lemma

If $x \in T$, the family $\{\uparrow x \setminus \uparrow F : F \in [I(x)]^{<\omega}\}$ is a local basis of open neighbourhoods of x . In particular, the topology is Hausdorff.

Proof.


Let $x \in W = (\uparrow x_0) \cap \cdots \cap (\uparrow x_{n-1}) \cap (\uparrow y_0)^c \cap \cdots \cap (\uparrow y_{m-1})^c$.

Lemma

If $x \in T$, the family $\{\uparrow x \setminus \uparrow F : F \in [I(x)]^{<\omega}\}$ is a local basis of open neighbourhoods of x . In particular, the topology is Hausdorff.

Proof.

Let $x \in W = (\uparrow x_0) \cap \cdots \cap (\uparrow x_{n-1}) \cap (\uparrow y_0)^c \cap \cdots \cap (\uparrow y_{m-1})^c$.




Lemma

If $x \in T$, the family $\{\uparrow x \setminus \uparrow F : F \in [I(x)]^{<\omega}\}$ is a local basis of open neighbourhoods of x . In particular, the topology is Hausdorff.

Proof.

Let $x \in W = (\uparrow x_0) \cap \cdots \cap (\uparrow x_{n-1}) \cap (\uparrow y_0)^c \cap \cdots \cap (\uparrow y_{m-1})^c$.

Since T is a tree, $\{x_i : i < n\}$ is a chain, say with maximum x_0 .




Lemma

If $x \in T$, the family $\{\uparrow x \setminus \uparrow F : F \in [I(x)]^{<\omega}\}$ is a local basis of open neighbourhoods of x . In particular, the topology is Hausdorff.

Proof.

Let $x \in W = (\uparrow x_0) \cap (\uparrow y_0)^c \cap \cdots \cap (\uparrow y_{m-1})^c$.




Lemma

If $x \in T$, the family $\{\uparrow x \setminus \uparrow F : F \in [I(x)]^{<\omega}\}$ is a local basis of open neighbourhoods of x . In particular, the topology is Hausdorff.

Proof.

Let $x \in W = (\uparrow x_0) \cap (\uparrow y_0)^c \cap \cdots \cap (\uparrow y_{m-1})^c$.




Lemma

If $x \in T$, the family $\{\uparrow x \setminus \uparrow F : F \in [I(x)]^{<\omega}\}$ is a local basis of open neighbourhoods of x . In particular, the topology is Hausdorff.

Proof.

Let $x \in W = (\uparrow x) \cap (\uparrow y_0)^c \cap \cdots \cap (\uparrow y_{m-1})^c$.




Lemma

If $x \in T$, the family $\{\uparrow x \setminus \uparrow F : F \in [I(x)]^{<\omega}\}$ is a local basis of open neighbourhoods of x . In particular, the topology is Hausdorff.

Proof.

Let $x \in W = (\uparrow x) \cap (\uparrow y_0)^c \cap \cdots \cap (\uparrow y_{m-1})^c$.




Lemma

If $x \in T$, the family $\{\uparrow x \setminus \uparrow F : F \in [I(x)]^{<\omega}\}$ is a local basis of open neighbourhoods of x . In particular, the topology is Hausdorff.

Proof.

Let $x \in W = (\uparrow x) \cap (\uparrow y_0)^c \cap \cdots \cap (\uparrow y_{m-1})^c$.




Lemma

If $x \in T$, the family $\{\uparrow x \setminus \uparrow F : F \in [I(x)]^{<\omega}\}$ is a local basis of open neighbourhoods of x . In particular, the topology is Hausdorff.

Proof.

Let $x \in W = (\uparrow x) \cap (\uparrow y_0)^c \cap \cdots \cap (\uparrow y_{m-1})^c$.




Lemma

If $x \in T$, the family $\{\uparrow x \setminus \uparrow F : F \in [I(x)]^{<\omega}\}$ is a local basis of open neighbourhoods of x . In particular, the topology is Hausdorff.

Proof.

Let $x \in W = (\uparrow x) \cap (\uparrow y_0)^c \cap \cdots \cap (\uparrow y_{m-1})^c$.



It therefore suffices to study open covers of the form $\mathcal{U}_f = \{\uparrow x \setminus \uparrow f(x) : x \in T\}$, where $f \in \prod_{x \in T} [I(x)]^{<\omega}$.

It therefore suffices to study open covers of the form $\mathcal{U}_f = \{\uparrow x \setminus \uparrow f(x) : x \in T\}$, where $f \in \prod_{x \in T} [I(x)]^{<\omega}$.

Definition

Given $f \in \prod_{x \in T} [I(x)]^{<\omega}$, we will say that $x \in T$ is *safe* iff for every $y < x$, $x \in \uparrow f(y)$.

It therefore suffices to study open covers of the form $\mathcal{U}_f = \{\uparrow x \setminus \uparrow f(x) : x \in T\}$, where $f \in \prod_{x \in T} [I(x)]^{<\omega}$.

Definition

Given $f \in \prod_{x \in T} [I(x)]^{<\omega}$, we will say that $x \in T$ is *safe* iff for every $y < x$, $x \in \uparrow f(y)$.

Note that the set of safe points is a finitely branching subtree of T .

It therefore suffices to study open covers of the form $\mathcal{U}_f = \{\uparrow x \setminus \uparrow f(x) : x \in T\}$, where $f \in \prod_{x \in T} [I(x)]^{<\omega}$.

Definition

Given $f \in \prod_{x \in T} [I(x)]^{<\omega}$, we will say that $x \in T$ is *safe* iff for every $y < x$, $x \in \uparrow f(y)$.

Note that the set of safe points is a finitely branching subtree of T .

Lemma

Let T be an infinitely branching \aleph_1 -tree and $f \in \prod_{x \in T} [I(x)]^{<\omega}$.
The following are equivalent:

It therefore suffices to study open covers of the form $\mathcal{U}_f = \{\uparrow x \setminus \uparrow f(x) : x \in T\}$, where $f \in \prod_{x \in T} [I(x)]^{<\omega}$.

Definition

Given $f \in \prod_{x \in T} [I(x)]^{<\omega}$, we will say that $x \in T$ is *safe* iff for every $y < x$, $x \in \uparrow f(y)$.

Note that the set of safe points is a finitely branching subtree of T .

Lemma

Let T be an infinitely branching \aleph_1 -tree and $f \in \prod_{x \in T} [I(x)]^{<\omega}$.
The following are equivalent:

1. f has no countable subcover.

It therefore suffices to study open covers of the form $\mathcal{U}_f = \{\uparrow x \setminus \uparrow f(x) : x \in T\}$, where $f \in \prod_{x \in T} [I(x)]^{<\omega}$.

Definition

Given $f \in \prod_{x \in T} [I(x)]^{<\omega}$, we will say that $x \in T$ is *safe* iff for every $y < x$, $x \in \uparrow f(y)$.

Note that the set of safe points is a finitely branching subtree of T .

Lemma

Let T be an infinitely branching \aleph_1 -tree and $f \in \prod_{x \in T} [I(x)]^{<\omega}$.
The following are equivalent:

1. f has no countable subcover.
2. Every level has a safe point.

It therefore suffices to study open covers of the form $\mathcal{U}_f = \{\uparrow x \setminus \uparrow f(x) : x \in T\}$, where $f \in \prod_{x \in T} [I(x)]^{<\omega}$.

Definition

Given $f \in \prod_{x \in T} [I(x)]^{<\omega}$, we will say that $x \in T$ is *safe* iff for every $y < x$, $x \in \uparrow f(y)$.

Note that the set of safe points is a finitely branching subtree of T .

Lemma

Let T be an infinitely branching \aleph_1 -tree and $f \in \prod_{x \in T} [I(x)]^{<\omega}$. The following are equivalent:

1. f has no countable subcover.
2. Every level has a safe point.
3. The set of $\{ht(x) : x \text{ is safe}\}$ is uncountable.

Theorem (M.)

Let T be an infinitely branching \aleph_1 -tree. Then T is a Lindelöf tree if and only if it has the Lindelöf property with respect to the fine wedge topology.

Theorem (M.)

Let T be an infinitely branching \aleph_1 -tree. Then T is a Lindelöf tree if and only if it has the Lindelöf property with respect to the fine wedge topology.

Proof.

\Rightarrow) Let f code a cover with no countable subcover. Let S be the set of safe points. By the previous lemma, S is uncountable.

Theorem (M.)

Let T be an infinitely branching \aleph_1 -tree. Then T is a Lindelöf tree if and only if it has the Lindelöf property with respect to the fine wedge topology.

Proof.

\Rightarrow) Let f code a cover with no countable subcover. Let S be the set of safe points. By the previous lemma, S is uncountable.

\Leftarrow) Let S be an uncountable finitely branching subtree of T .

Theorem (M.)

Let T be an infinitely branching \aleph_1 -tree. Then T is a Lindelöf tree if and only if it has the Lindelöf property with respect to the fine wedge topology.

Proof.

\Rightarrow) Let f code a cover with no countable subcover. Let S be the set of safe points. By the previous lemma, S is uncountable.

\Leftarrow) Let S be an uncountable finitely branching subtree of T . Define $f(x) = \emptyset$ for $x \in T \setminus S$ and $f(x) = I_S(x)$ for $x \in S$.

Theorem (M.)

Let T be an infinitely branching \aleph_1 -tree. Then T is a Lindelöf tree if and only if it has the Lindelöf property with respect to the fine wedge topology.

Proof.

\Rightarrow) Let f code a cover with no countable subcover. Let S be the set of safe points. By the previous lemma, S is uncountable.

\Leftarrow) Let S be an uncountable finitely branching subtree of T .

Define $f(x) = \emptyset$ for $x \in T \setminus S$ and $f(x) = I_S(x)$ for $x \in S$.

If $x \in S$ and $y < x$, then $x \in \uparrow f(y)$, so x is safe for S . So

$S \subseteq \{x \in T : x \text{ is safe}\}$, hence the latter is uncountable. □

Recall that if T is a tree, an *antichain* is a set of pairwise incomparable elements of T . A *Suslin tree* is a tree with no uncountable chains or antichains.

Recall that if T is a tree, an *antichain* is a set of pairwise incomparable elements of T . A *Suslin tree* is a tree with no uncountable chains or antichains.

If T is a tree, \mathbb{P}_T is the dual order.

Recall that if T is a tree, an *antichain* is a set of pairwise incomparable elements of T . A *Suslin tree* is a tree with no uncountable chains or antichains.

If T is a tree, \mathbb{P}_T is the dual order.

Lemma (folklore [2])

Let T be a Suslin tree. The poset \mathbb{P}_T has the ccc and is countably distributive. Moreover, if $D \subseteq \mathbb{P}_T$ is dense and open, then there exists $\alpha < \omega_1$ such that $T \restriction [\alpha, \omega_1) \subseteq D$.

Recall that if T is a tree, an *antichain* is a set of pairwise incomparable elements of T . A *Suslin tree* is a tree with no uncountable chains or antichains.

If T is a tree, \mathbb{P}_T is the dual order.

Lemma (folklore [2])

Let T be a Suslin tree. The poset \mathbb{P}_T has the ccc and is countably distributive. Moreover, if $D \subseteq \mathbb{P}_T$ is dense and open, then there exists $\alpha < \omega_1$ such that $T \restriction [\alpha, \omega_1) \subseteq D$.

Lemma (folklore [2])

Let T be a Suslin tree in the universe V . If W is an outer model and $b \in W$ is a cofinal branch through T , then b is \mathbb{P}_T -generic over V .

Theorem (M.)

Let T be an infinitely branching Suslin tree. Then T is Lindelöf.

Theorem (M.)

Let T be an infinitely branching Suslin tree. Then T is Lindelöf.

Proof.

Suppose not. Let $S \subseteq T$ be a finitely branching uncountable subtree. Then S is also Suslin. Force with \mathbb{P}_S to add a branch b . By the previous lemma, b is \mathbb{P}_T -generic over V . But S is finitely branching and T is infinitely branching, so b is disjoint from S above some node of T , by a density argument. □

Theorem (M.)

Let T be an infinitely branching Suslin tree. Then T is Lindelöf.

Proof.

Suppose not. Let $S \subseteq T$ be a finitely branching uncountable subtree. Then S is also Suslin. Force with \mathbb{P}_S to add a branch b . By the previous lemma, b is \mathbb{P}_T -generic over V . But S is finitely branching and T is infinitely branching, so b is disjoint from S above some node of T , by a density argument. □

$$\{\text{Suslin}\} \subseteq \{\text{Lindelöf}\} \subseteq \{\text{Aronszajn}\}$$

An example of a Lindelöf tree

Let $\vec{e} = \langle e_\alpha : \alpha < \omega_1 \rangle$ be a *coherent sequence of injections*, so $e_\alpha : \alpha \rightarrow \omega$ is injective and $\alpha < \beta$ implies $e_\alpha =^* e_\beta$, that is $\{\xi < \alpha : e_\alpha(\xi) = e_\beta(\xi)\}$ is finite.

An example of a Lindelöf tree

Let $\vec{e} = \langle e_\alpha : \alpha < \omega_1 \rangle$ be a *coherent sequence of injections*, so $e_\alpha : \alpha \rightarrow \omega$ is injective and $\alpha < \beta$ implies $e_\alpha =^* e_\beta$, that is $\{\xi < \alpha : e_\alpha(\xi) = e_\beta(\xi)\}$ is finite.

It is well-known that the set

$$T^{\vec{e}} = \bigcup_{\alpha < \omega_1} \{t \in {}^\alpha \omega : t \text{ is injective} \wedge t =^* e_\alpha\}$$

is an infinitely branching Aronszajn tree.

An example of a Lindelöf tree

Let $\vec{e} = \langle e_\alpha : \alpha < \omega_1 \rangle$ be a *coherent sequence of injections*, so $e_\alpha : \alpha \rightarrow \omega$ is injective and $\alpha < \beta$ implies $e_\alpha =^* e_\beta$, that is $\{\xi < \alpha : e_\alpha(\xi) = e_\beta(\xi)\}$ is finite.

It is well-known that the set

$$T^{\vec{e}} = \bigcup_{\alpha < \omega_1} \{t \in {}^\alpha \omega : t \text{ is injective} \wedge t =^* e_\alpha\}$$

is an infinitely branching Aronszajn tree.

Let \mathbb{C} be Cohen forcing, instantiated as the set of finite partial functions from ω to ω .

An example of a Lindelöf tree

Let $\vec{e} = \langle e_\alpha : \alpha < \omega_1 \rangle$ be a *coherent sequence of injections*, so $e_\alpha : \alpha \rightarrow \omega$ is injective and $\alpha < \beta$ implies $e_\alpha =^* e_\beta$, that is $\{\xi < \alpha : e_\alpha(\xi) = e_\beta(\xi)\}$ is finite.

It is well-known that the set

$$T^{\vec{e}} = \bigcup_{\alpha < \omega_1} \{t \in {}^\alpha \omega : t \text{ is injective} \wedge t =^* e_\alpha\}$$

is an infinitely branching Aronszajn tree.

Let \mathbb{C} be Cohen forcing, instantiated as the set of finite partial functions from ω to ω .

Theorem (Todorćević)

If \vec{e} is a coherent sequence of injections and $r : \omega \rightarrow \omega$ is Cohen generic over V , then

$$\{r \circ t : t \in T^{\vec{e}}\}$$

is an infinitely branching Suslin tree in $V[r]$.

An example of a Lindelöf tree

Theorem (M.)

Let \vec{e} be a coherent sequence of injections. Then $T^{\vec{e}}$ is a Lindelöf tree.

An example of a Lindelöf tree

Theorem (M.)

Let \vec{e} be a coherent sequence of injections. Then $T^{\vec{e}}$ is a Lindelöf tree.

Proof.

It is clear that $T^{\vec{e}}$ is infinitely branching. Suppose now that $S \subseteq T$ is an uncountable finitely branching subtree. Force to add a Cohen real $r : \omega \rightarrow \omega$. In $V[r]$, $T^* = \{r \circ t : t \in T\}$ is infinitely branching by genericity and $S^* = \{r \circ s : s \in S\}$ is a finitely branching uncountable subtree of T^* . But T^* is a Suslin tree in $V[r]$, which is a contradiction. □

An example of a Lindelöf tree

Theorem (M.)

Let \vec{e} be a coherent sequence of injections. Then $T^{\vec{e}}$ is a Lindelöf tree.

Proof.

It is clear that $T^{\vec{e}}$ is infinitely branching. Suppose now that $S \subseteq T$ is an uncountable finitely branching subtree. Force to add a Cohen real $r : \omega \rightarrow \omega$. In $V[r]$, $T^* = \{r \circ t : t \in T\}$ is infinitely branching by genericity and $S^* = \{r \circ s : s \in S\}$ is a finitely branching uncountable subtree of T^* . But T^* is a Suslin tree in $V[r]$, which is a contradiction. □

Since $T^{\vec{e}}$ is never Suslin,

$$\{\text{Suslin}\} \subsetneq \{\text{Lindelöf}\} \subseteq \{\text{Aronszajn}\}$$

Theorem

There exists an Aronszajn non-Lindelöf tree.

Theorem

There exists an Aronszajn non-Lindelöf tree.

Proof sketch.

Fix a coherent sequence of injection \vec{e} and use a bijection $f : \omega_1 \times \omega \rightarrow \omega_1$ to transfer \vec{e} to a coherent sequence $\langle f[e_\alpha] : \alpha < \omega_1 \rangle$, which is then used to build a binary Aronszajn tree T .


Theorem

There exists an Aronszajn non-Lindelöf tree.

Proof sketch.

Fix a coherent sequence of injection \vec{e} and use a bijection $f : \omega_1 \times \omega \rightarrow \omega_1$ to transfer \vec{e} to a coherent sequence $\langle f[e_\alpha] : \alpha < \omega_1 \rangle$, which is then used to build a binary Aronszajn tree T .

Now recursively build, level by level, an infinitely branching tree U which has T as a subtree. \square




Theorem

There exists an Aronszajn non-Lindelöf tree.

Proof sketch.

Fix a coherent sequence of injection \vec{e} and use a bijection $f : \omega_1 \times \omega \rightarrow \omega_1$ to transfer \vec{e} to a coherent sequence $\langle f[e_\alpha] : \alpha < \omega_1 \rangle$, which is then used to build a binary Aronszajn tree T .

Now recursively build, level by level, an infinitely branching tree U which has T as a subtree. \square




Theorem

There exists an Aronszajn non-Lindelöf tree.

Proof sketch.

Fix a coherent sequence of injection \vec{e} and use a bijection $f : \omega_1 \times \omega \rightarrow \omega_1$ to transfer \vec{e} to a coherent sequence $\langle f[e_\alpha] : \alpha < \omega_1 \rangle$, which is then used to build a binary Aronszajn tree T .

Now recursively build, level by level, an infinitely branching tree U which has T as a subtree. \square




Theorem

There exists an Aronszajn non-Lindelöf tree.

Proof sketch.

Fix a coherent sequence of injections \vec{e} and use a bijection $f : \omega_1 \times \omega \rightarrow \omega_1$ to transfer \vec{e} to a coherent sequence $\langle f[e_\alpha] : \alpha < \omega_1 \rangle$, which is then used to build a binary Aronszajn tree T .

Now recursively build, level by level, an infinitely branching tree U which has T as a subtree. \square



$$\{\text{Suslin}\} \subsetneq \{\text{Lindelöf}\} \subsetneq \{\text{Aronszajn}\}$$

Thank you :)

References

- [1] Tomek Bartoszyński and Haim Judah. *Set Theory*. A K Peters, Ltd., Wellesley, MA, 1995. xii+546.
- [2] Keith J. Devlin and Håvard Johnsbråten. *The Souslin Problem*. Lecture Notes in Mathematics 405. Springer, 1974. 132 pp.
- [3] Dénes König. “Über Eine Schlussweise Aus Dem Endlichen Ins Unendliche”. In: *Acta Scientiarum Mathematicarum (Szeged)* 3 (1927).
- [4] Pedro E. Marun. “Square Compactness and Lindelöf Trees”. In: *Submitted for publication* (2023).
- [5] Peter J. Nyikos. “Various Topologies on Trees”. In: *Proceedings of the Tennessee Topology Conference (Nashville, TN, 1996)*. 1997.
- [6] Ernst Specker. “Sur Un Problème de Sikorski”. In: *Colloquium Mathematicae* 2 (1949).