

# Subtrees with small branching number

Pedro Marun

Carnegie Mellon University

March 23, 2023

## Definition

- ▶ Let  $T$  be a tree and  $x \in T$ . The set of *immediate successors* of  $T$  is denoted by  $I_T(x)$ .

## Definition

- ▶ Let  $T$  be a tree and  $x \in T$ . The set of *immediate successors* of  $T$  is denoted by  $I_T(x)$ .
- ▶ Let  $\lambda$  be a cardinal. A tree  $T$  is  $\lambda$ -*branching* [respectively  $<\lambda$ -*branching*] iff for every  $x \in T$ ,  $|I_T(x)| = \lambda$  [respectively  $|I_T(x)| < \lambda$ ].

## Definition

- ▶ Let  $T$  be a tree and  $x \in T$ . The set of *immediate successors* of  $T$  is denoted by  $I_T(x)$ .
- ▶ Let  $\lambda$  be a cardinal. A tree  $T$  is  $\lambda$ -*branching* [respectively  $<\lambda$ -*branching*] iff for every  $x \in T$ ,  $|I_T(x)| = \lambda$  [respectively  $|I_T(x)| < \lambda$ ].
- ▶ A tree  $T$  is *finitely branching* if it is  $<\aleph_0$ -branching. It is *infinitely branching* if  $|I_T(x)| \geq \aleph_0$  for every  $x \in T$ .
- ▶ A *subtree* of a tree  $T$  is a subset  $S$  of  $T$  such that for every  $s \in S$  and  $t \in T$ , if  $t < s$ , then  $t \in S$ .

## Definition

- ▶ Let  $T$  be a tree and  $x \in T$ . The set of *immediate successors* of  $T$  is denoted by  $I_T(x)$ .
- ▶ Let  $\lambda$  be a cardinal. A tree  $T$  is  $\lambda$ -*branching* [respectively  $<\lambda$ -*branching*] iff for every  $x \in T$ ,  $|I_T(x)| = \lambda$  [respectively  $|I_T(x)| < \lambda$ ].
- ▶ A tree  $T$  is *finitely branching* if it is  $<\aleph_0$ -branching. It is *infinitely branching* if  $|I_T(x)| \geq \aleph_0$  for every  $x \in T$ .
- ▶ A *subtree* of a tree  $T$  is a subset  $S$  of  $T$  such that for every  $s \in S$  and  $t \in T$ , if  $t < s$ , then  $t \in S$ .

## Lemma

Let  $T$  be an  $\aleph_1$ -tree. If  $T$  has no uncountable 1-branching subtrees, then  $T$  is Aronszajn.

## Definition

- ▶ Let  $T$  be a tree and  $x \in T$ . The set of *immediate successors* of  $T$  is denoted by  $I_T(x)$ .
- ▶ Let  $\lambda$  be a cardinal. A tree  $T$  is  $\lambda$ -*branching* [respectively  $<\lambda$ -*branching*] iff for every  $x \in T$ ,  $|I_T(x)| = \lambda$  [respectively  $|I_T(x)| < \lambda$ ].
- ▶ A tree  $T$  is *finitely branching* if it is  $<\aleph_0$ -branching. It is *infinitely branching* if  $|I_T(x)| \geq \aleph_0$  for every  $x \in T$ .
- ▶ A *subtree* of a tree  $T$  is a subset  $S$  of  $T$  such that for every  $s \in S$  and  $t \in T$ , if  $t < s$ , then  $t \in S$ .

## Lemma

Let  $T$  be an  $\aleph_1$ -tree. If  $T$  has no uncountable 1-branching subtrees, then  $T$  is Aronszajn.

## Proof.

If  $b$  is a cofinal branch, then  $b$  is an uncountable 1-branching subtree.



## Definition

An infinitely branching  $\aleph_1$ -tree  $T$  is *Lindelöf* iff every finitely branching subtree of  $T$  is countable.

## Definition

An infinitely branching  $\aleph_1$ -tree  $T$  is *Lindelöf* iff every finitely branching subtree of  $T$  is countable.

The previous lemma shows that

$$\{\text{Lindelöf}\} \subseteq \{\text{Aronszajn}\}$$



## Definition

An infinitely branching  $\aleph_1$ -tree  $T$  is *Lindelöf* iff every finitely branching subtree of  $T$  is countable.

The previous lemma shows that

$$\{\text{Lindelöf}\} \subseteq \{\text{Aronszajn}\}$$

We will show that the inclusion is proper.

## Definition

An infinitely branching  $\aleph_1$ -tree  $T$  is *Lindelöf* iff every finitely branching subtree of  $T$  is countable.

The previous lemma shows that

$$\{\text{Lindelöf}\} \subseteq \{\text{Aronszajn}\}$$

We will show that the inclusion is proper.

First, we explain the terminology:

## Definition

An infinitely branching  $\aleph_1$ -tree  $T$  is *Lindelöf* iff every finitely branching subtree of  $T$  is countable.

The previous lemma shows that

$$\{\text{Lindelöf}\} \subseteq \{\text{Aronszajn}\}$$

We will show that the inclusion is proper.

First, we explain the terminology: Trees are Lindelöf if and only if they have they are Lindelöf spaces with respect to a natural topology.

## Definition

An infinitely branching  $\aleph_1$ -tree  $T$  is *Lindelöf* iff every finitely branching subtree of  $T$  is countable.

The previous lemma shows that

$$\{\text{Lindelöf}\} \subseteq \{\text{Aronszajn}\}$$

We will show that the inclusion is proper.

First, we explain the terminology: Trees are Lindelöf if and only if they have the Lindelöf property with respect to a natural topology.

## Definition (Nyikos)

Let  $T$  be a tree. The *fine wedge topology* on  $T$  is generated by all sets of the form  $\uparrow x$  and their complements, where

$\uparrow x = \{y \in T : x \leq y\}$  and  $x \in T$ .

## Definition

An infinitely branching  $\aleph_1$ -tree  $T$  is *Lindelöf* iff every finitely branching subtree of  $T$  is countable.

The previous lemma shows that

$$\{\text{Lindelöf}\} \subseteq \{\text{Aronszajn}\}$$

We will show that the inclusion is proper.

First, we explain the terminology: Trees are Lindelöf if and only if they have the Lindelöf property with respect to a natural topology.

## Definition (Nyikos)

Let  $T$  be a tree. The *fine wedge topology* on  $T$  is generated by all sets of the form  $\uparrow x$  and their complements, where

$\uparrow x = \{y \in T : x \leq y\}$  and  $x \in T$ .

If  $X \subseteq T$ , write  $\uparrow X = \{y \in T : \exists x \in X (x \leq y)\}$ .

## Lemma

*If  $x \in T$ , the family  $\{\uparrow x \setminus \uparrow F : F \in [I(x)]^{<\omega}\}$  is a local basis of open neighbourhoods of  $x$ . In particular, the topology is Hausdorff.*

## Lemma

*If  $x \in T$ , the family  $\{\uparrow x \setminus \uparrow F : F \in [I(x)]^{<\omega}\}$  is a local basis of open neighbourhoods of  $x$ . In particular, the topology is Hausdorff.*

## Proof.


Let  $x \in W = (\uparrow x_0) \cap \cdots \cap (\uparrow x_{n-1}) \cap (\uparrow y_0)^c \cap \cdots \cap (\uparrow y_{m-1})^c$ .

## Lemma

*If  $x \in T$ , the family  $\{\uparrow x \setminus \uparrow F : F \in [I(x)]^{<\omega}\}$  is a local basis of open neighbourhoods of  $x$ . In particular, the topology is Hausdorff.*

## Proof.

Let  $x \in W = (\uparrow x_0) \cap \cdots \cap (\uparrow x_{n-1}) \cap (\uparrow y_0)^c \cap \cdots \cap (\uparrow y_{m-1})^c$ .






## Lemma

If  $x \in T$ , the family  $\{\uparrow x \setminus \uparrow F : F \in [I(x)]^{<\omega}\}$  is a local basis of open neighbourhoods of  $x$ . In particular, the topology is Hausdorff.

## Proof.

Let  $x \in W = (\uparrow x_0) \cap \cdots \cap (\uparrow x_{n-1}) \cap (\uparrow y_0)^c \cap \cdots \cap (\uparrow y_{m-1})^c$ .

Since  $T$  is a tree,  $\{x_i : i < n\}$  is a chain, say with maximum  $x_0$ .




## Lemma

If  $x \in T$ , the family  $\{\uparrow x \setminus \uparrow F : F \in [I(x)]^{<\omega}\}$  is a local basis of open neighbourhoods of  $x$ . In particular, the topology is Hausdorff.

## Proof.

Let  $x \in W = (\uparrow x_0) \cap (\uparrow y_0)^c \cap \cdots \cap (\uparrow y_{m-1})^c$ .




## Lemma

If  $x \in T$ , the family  $\{\uparrow x \setminus \uparrow F : F \in [I(x)]^{<\omega}\}$  is a local basis of open neighbourhoods of  $x$ . In particular, the topology is Hausdorff.

## Proof.

Let  $x \in W = (\uparrow x_0) \cap (\uparrow y_0)^c \cap \cdots \cap (\uparrow y_{m-1})^c$ .




### Lemma

If  $x \in T$ , the family  $\{\uparrow x \setminus \uparrow F : F \in [I(x)]^{<\omega}\}$  is a local basis of open neighbourhoods of  $x$ . In particular, the topology is Hausdorff.

### Proof.

Let  $x \in W = (\uparrow x) \cap (\uparrow y_0)^c \cap \cdots \cap (\uparrow y_{m-1})^c$ .




## Lemma

If  $x \in T$ , the family  $\{\uparrow x \setminus \uparrow F : F \in [I(x)]^{<\omega}\}$  is a local basis of open neighbourhoods of  $x$ . In particular, the topology is Hausdorff.

## Proof.

Let  $x \in W = (\uparrow x) \cap (\uparrow y_0)^c \cap \cdots \cap (\uparrow y_{m-1})^c$ .




## Lemma

If  $x \in T$ , the family  $\{\uparrow x \setminus \uparrow F : F \in [I(x)]^{<\omega}\}$  is a local basis of open neighbourhoods of  $x$ . In particular, the topology is Hausdorff.

## Proof.

Let  $x \in W = (\uparrow x) \cap (\uparrow y_0)^c \cap \cdots \cap (\uparrow y_{m-1})^c$ .




## Lemma

If  $x \in T$ , the family  $\{\uparrow x \setminus \uparrow F : F \in [I(x)]^{<\omega}\}$  is a local basis of open neighbourhoods of  $x$ . In particular, the topology is Hausdorff.

## Proof.

Let  $x \in W = (\uparrow x) \cap (\uparrow y_0)^c \cap \cdots \cap (\uparrow y_{m-1})^c$ .




## Lemma

If  $x \in T$ , the family  $\{\uparrow x \setminus \uparrow F : F \in [I(x)]^{<\omega}\}$  is a local basis of open neighbourhoods of  $x$ . In particular, the topology is Hausdorff.

## Proof.

Let  $x \in W = (\uparrow x) \cap (\uparrow y_0)^c \cap \cdots \cap (\uparrow y_{m-1})^c$ .





It therefore suffices to study open covers of the form  $\mathcal{U}_f = \{\uparrow x \setminus \uparrow f(x) : x \in T\}$ , where  $f \in \prod_{x \in T} [I(x)]^{<\omega}$ .

It therefore suffices to study open covers of the form  $\mathcal{U}_f = \{\uparrow x \setminus \uparrow f(x) : x \in T\}$ , where  $f \in \prod_{x \in T} [I(x)]^{<\omega}$ .

### Definition

Given  $f \in \prod_{x \in T} [I(x)]^{<\omega}$ , we will say that  $x \in T$  is *safe* iff for every  $y < x$ ,  $x \in \uparrow f(y)$ .

It therefore suffices to study open covers of the form  $\mathcal{U}_f = \{\uparrow x \setminus \uparrow f(x) : x \in T\}$ , where  $f \in \prod_{x \in T} [I(x)]^{<\omega}$ .

### Definition

Given  $f \in \prod_{x \in T} [I(x)]^{<\omega}$ , we will say that  $x \in T$  is *safe* iff for every  $y < x$ ,  $x \in \uparrow f(y)$ .

Note that the set of safe points is a finitely branching subtree of  $T$ .

It therefore suffices to study open covers of the form  $\mathcal{U}_f = \{\uparrow x \setminus \uparrow f(x) : x \in T\}$ , where  $f \in \prod_{x \in T} [I(x)]^{<\omega}$ .

### Definition

Given  $f \in \prod_{x \in T} [I(x)]^{<\omega}$ , we will say that  $x \in T$  is *safe* iff for every  $y < x$ ,  $x \in \uparrow f(y)$ .

Note that the set of safe points is a finitely branching subtree of  $T$ .

### Lemma

Let  $T$  be an infinitely branching  $\aleph_1$ -tree and  $f \in \prod_{x \in T} [I(x)]^{<\omega}$ .  
The following are equivalent:

It therefore suffices to study open covers of the form  $\mathcal{U}_f = \{\uparrow x \setminus \uparrow f(x) : x \in T\}$ , where  $f \in \prod_{x \in T} [I(x)]^{<\omega}$ .

### Definition

Given  $f \in \prod_{x \in T} [I(x)]^{<\omega}$ , we will say that  $x \in T$  is *safe* iff for every  $y < x$ ,  $x \in \uparrow f(y)$ .

Note that the set of safe points is a finitely branching subtree of  $T$ .

### Lemma

Let  $T$  be an infinitely branching  $\aleph_1$ -tree and  $f \in \prod_{x \in T} [I(x)]^{<\omega}$ .  
The following are equivalent:

1.  $f$  has no countable subcover.

It therefore suffices to study open covers of the form  $\mathcal{U}_f = \{\uparrow x \setminus \uparrow f(x) : x \in T\}$ , where  $f \in \prod_{x \in T} [I(x)]^{<\omega}$ .

### Definition

Given  $f \in \prod_{x \in T} [I(x)]^{<\omega}$ , we will say that  $x \in T$  is *safe* iff for every  $y < x$ ,  $x \in \uparrow f(y)$ .

Note that the set of safe points is a finitely branching subtree of  $T$ .

### Lemma

Let  $T$  be an infinitely branching  $\aleph_1$ -tree and  $f \in \prod_{x \in T} [I(x)]^{<\omega}$ .  
The following are equivalent:

1.  $f$  has no countable subcover.
2. Every level has a safe point.

It therefore suffices to study open covers of the form  $\mathcal{U}_f = \{\uparrow x \setminus \uparrow f(x) : x \in T\}$ , where  $f \in \prod_{x \in T} [I(x)]^{<\omega}$ .

### Definition

Given  $f \in \prod_{x \in T} [I(x)]^{<\omega}$ , we will say that  $x \in T$  is *safe* iff for every  $y < x$ ,  $x \in \uparrow f(y)$ .

Note that the set of safe points is a finitely branching subtree of  $T$ .

### Lemma

Let  $T$  be an infinitely branching  $\aleph_1$ -tree and  $f \in \prod_{x \in T} [I(x)]^{<\omega}$ . The following are equivalent:

1.  $f$  has no countable subcover.
2. Every level has a safe point.
3. The set of  $\{ht(x) : x \text{ is safe}\}$  is uncountable.

## Theorem (M.)

*Let  $T$  be an infinitely branching  $\aleph_1$ -tree. Then  $T$  is a Lindelöf tree if and only if it has the Lindelöf property with respect to the fine wedge topology.*



## Theorem (M.)

*Let  $T$  be an infinitely branching  $\aleph_1$ -tree. Then  $T$  is a Lindelöf tree if and only if it has the Lindelöf property with respect to the fine wedge topology.*

## Proof.

$\Rightarrow$ ) Let  $f$  code a cover with no countable subcover. Let  $S$  be the set of safe points. By the previous lemma,  $S$  is uncountable.

## Theorem (M.)

*Let  $T$  be an infinitely branching  $\aleph_1$ -tree. Then  $T$  is a Lindelöf tree if and only if it has the Lindelöf property with respect to the fine wedge topology.*

## Proof.

$\Rightarrow$ ) Let  $f$  code a cover with no countable subcover. Let  $S$  be the set of safe points. By the previous lemma,  $S$  is uncountable.

$\Leftarrow$ ) Let  $S$  be an uncountable finitely branching subtree of  $T$ .

## Theorem (M.)

*Let  $T$  be an infinitely branching  $\aleph_1$ -tree. Then  $T$  is a Lindelöf tree if and only if it has the Lindelöf property with respect to the fine wedge topology.*

### Proof.

$\Rightarrow$ ) Let  $f$  code a cover with no countable subcover. Let  $S$  be the set of safe points. By the previous lemma,  $S$  is uncountable.

$\Leftarrow$ ) Let  $S$  be an uncountable finitely branching subtree of  $T$ . Define  $f(x) = \emptyset$  for  $x \in T \setminus S$  and  $f(x) = I_S(x)$  for  $x \in S$ .

## Theorem (M.)

*Let  $T$  be an infinitely branching  $\aleph_1$ -tree. Then  $T$  is a Lindelöf tree if and only if it has the Lindelöf property with respect to the fine wedge topology.*

### Proof.

$\Rightarrow$ ) Let  $f$  code a cover with no countable subcover. Let  $S$  be the set of safe points. By the previous lemma,  $S$  is uncountable.

$\Leftarrow$ ) Let  $S$  be an uncountable finitely branching subtree of  $T$ .

Define  $f(x) = \emptyset$  for  $x \in T \setminus S$  and  $f(x) = I_S(x)$  for  $x \in S$ .

If  $x \in S$  and  $y < x$ , then  $x \in \uparrow f(y)$ , so  $x$  is safe for  $S$ . So

$S \subseteq \{x \in T : x \text{ is safe}\}$ , hence the latter is uncountable. □

Recall that if  $T$  is a tree, an *antichain* is a set of pairwise incomparable elements of  $T$ . A *Suslin tree* is a tree with no uncountable chains or antichains.

Recall that if  $T$  is a tree, an *antichain* is a set of pairwise incomparable elements of  $T$ . A *Suslin tree* is a tree with no uncountable chains or antichains.

If  $T$  is a tree,  $\mathbb{P}_T$  is the dual order.

Recall that if  $T$  is a tree, an *antichain* is a set of pairwise incomparable elements of  $T$ . A *Suslin tree* is a tree with no uncountable chains or antichains.

If  $T$  is a tree,  $\mathbb{P}_T$  is the dual order.

### Lemma (folklore [2])

*Let  $T$  be a Suslin tree. The poset  $\mathbb{P}_T$  has the ccc and is countably distributive. Moreover, if  $D \subseteq \mathbb{P}_T$  is dense and open, then there exists  $\alpha < \omega_1$  such that  $T \restriction [\alpha, \omega_1) \subseteq D$ .*

Recall that if  $T$  is a tree, an *antichain* is a set of pairwise incomparable elements of  $T$ . A *Suslin tree* is a tree with no uncountable chains or antichains.

If  $T$  is a tree,  $\mathbb{P}_T$  is the dual order.

### Lemma (folklore [2])

*Let  $T$  be a Suslin tree. The poset  $\mathbb{P}_T$  has the ccc and is countably distributive. Moreover, if  $D \subseteq \mathbb{P}_T$  is dense and open, then there exists  $\alpha < \omega_1$  such that  $T \restriction [\alpha, \omega_1) \subseteq D$ .*

### Lemma (folklore [2])

*Let  $T$  be a Suslin tree in the universe  $V$ . If  $W$  is an outer model and  $b \in W$  is a cofinal branch through  $T$ , then  $b$  is  $\mathbb{P}_T$ -generic over  $V$ .*



## Theorem (M.)

*Let  $T$  be an infinitely branching Suslin tree. Then  $T$  is Lindelöf.*

## Theorem (M.)

*Let  $T$  be an infinitely branching Suslin tree. Then  $T$  is Lindelöf.*

### Proof.

Suppose not. Let  $S \subseteq T$  be a finitely branching uncountable subtree. Then  $S$  is also Suslin. Force with  $\mathbb{P}_S$  to add a branch  $b$ . By the previous lemma,  $b$  is  $\mathbb{P}_T$ -generic over  $V$ . But  $S$  is finitely branching and  $T$  is infinitely branching, so  $b$  is disjoint from  $S$  above some node of  $T$ , by a density argument. □

## Theorem (M.)

*Let  $T$  be an infinitely branching Suslin tree. Then  $T$  is Lindelöf.*

### Proof.

Suppose not. Let  $S \subseteq T$  be a finitely branching uncountable subtree. Then  $S$  is also Suslin. Force with  $\mathbb{P}_S$  to add a branch  $b$ . By the previous lemma,  $b$  is  $\mathbb{P}_T$ -generic over  $V$ . But  $S$  is finitely branching and  $T$  is infinitely branching, so  $b$  is disjoint from  $S$  above some node of  $T$ , by a density argument. □

$$\{\text{Suslin}\} \subseteq \{\text{Lindelöf}\} \subseteq \{\text{Aronszajn}\}$$

# A non-Lindelöf Aronszajn tree

Let  $\vec{e} = \langle e_\alpha : \alpha < \omega_1 \rangle$  be a coherent sequence of injections, i.e.  $e_\alpha : \alpha \rightarrow \omega$  is injective and  $e_\alpha =^* e_\beta$ .

# A non-Lindelöf Aronszajn tree

Let  $\vec{e} = \langle e_\alpha : \alpha < \omega_1 \rangle$  be a coherent sequence of injections, i.e.  $e_\alpha : \alpha \rightarrow \omega$  is injective and  $e_\alpha =^* e_\beta$ .


Use a bijection  $f : \omega_1 \times \omega \rightarrow \omega_1$  to transfer  $\vec{e}$  to a coherent sequence  $\langle f[e_\alpha] : \alpha < \omega_1 \rangle$ , which is then used to build a binary Aronszajn tree  $T$ .

# A non-Lindelöf Aronszajn tree

Let  $\vec{e} = \langle e_\alpha : \alpha < \omega_1 \rangle$  be a coherent sequence of injections, i.e.  $e_\alpha : \alpha \rightarrow \omega$  is injective and  $e_\alpha =^* e_\beta$ .

Use a bijection  $f : \omega_1 \times \omega \rightarrow \omega_1$  to transfer  $\vec{e}$  to a coherent sequence  $\langle f[e_\alpha] : \alpha < \omega_1 \rangle$ , which is then used to build a binary Aronszajn tree  $T$ .

Now recursively build, level by level, an infinitely branching tree  $U$  which has  $T$  as a subtree.




# A non-Lindelöf Aronszajn tree

Let  $\vec{e} = \langle e_\alpha : \alpha < \omega_1 \rangle$  be a coherent sequence of injections, i.e.  $e_\alpha : \alpha \rightarrow \omega$  is injective and  $e_\alpha =^* e_\beta$ .

Use a bijection  $f : \omega_1 \times \omega \rightarrow \omega_1$  to transfer  $\vec{e}$  to a coherent sequence  $\langle f[e_\alpha] : \alpha < \omega_1 \rangle$ , which is then used to build a binary Aronszajn tree  $T$ .

Now recursively build, level by level, an infinitely branching tree  $U$  which has  $T$  as a subtree.




# A non-Lindelöf Aronszajn tree

Let  $\vec{e} = \langle e_\alpha : \alpha < \omega_1 \rangle$  be a coherent sequence of injections, i.e.  $e_\alpha : \alpha \rightarrow \omega$  is injective and  $e_\alpha =^* e_\beta$ .

Use a bijection  $f : \omega_1 \times \omega \rightarrow \omega_1$  to transfer  $\vec{e}$  to a coherent sequence  $\langle f[e_\alpha] : \alpha < \omega_1 \rangle$ , which is then used to build a binary Aronszajn tree  $T$ .

Now recursively build, level by level, an infinitely branching tree  $U$  which has  $T$  as a subtree.






# A non-Lindelöf Aronszajn tree

Let  $\vec{e} = \langle e_\alpha : \alpha < \omega_1 \rangle$  be a coherent sequence of injections, i.e.  $e_\alpha : \alpha \rightarrow \omega$  is injective and  $e_\alpha =^* e_\beta$ .

Use a bijection  $f : \omega_1 \times \omega \rightarrow \omega_1$  to transfer  $\vec{e}$  to a coherent sequence  $\langle f[e_\alpha] : \alpha < \omega_1 \rangle$ , which is then used to build a binary Aronszajn tree  $T$ .

Now recursively build, level by level, an infinitely branching tree  $U$  which has  $T$  as a subtree.




# A non-Lindelöf Aronszajn tree

Let  $\vec{e} = \langle e_\alpha : \alpha < \omega_1 \rangle$  be a coherent sequence of injections, i.e.  $e_\alpha : \alpha \rightarrow \omega$  is injective and  $e_\alpha =^* e_\beta$ .

Use a bijection  $f : \omega_1 \times \omega \rightarrow \omega_1$  to transfer  $\vec{e}$  to a coherent sequence  $\langle f[e_\alpha] : \alpha < \omega_1 \rangle$ , which is then used to build a binary Aronszajn tree  $T$ .

Now recursively build, level by level, an infinitely branching tree  $U$  which has  $T$  as a subtree.



$$\{\text{Suslin}\} \subseteq \{\text{Lindelöf}\} \subsetneq \{\text{Aronszajn}\}$$

## ( $\diamond$ ) A special Lindelöf tree

Let  $\vec{f} = \langle f_\alpha : \alpha \in \text{lim}(\omega_1) \rangle$  be such  $f_\alpha : \alpha \rightarrow [\alpha]^{<\omega}$  and  $\vec{f}$  guesses any  $f : \omega_1 \rightarrow [\omega_1]^{<\omega}$  stationarily often.

## ( $\diamond$ ) A special Lindelöf tree

Let  $\vec{f} = \langle f_\alpha : \alpha \in \text{lim}(\omega_1) \rangle$  be such  $f_\alpha : \alpha \rightarrow [\alpha]^{<\omega}$  and  $\vec{f}$  guesses any  $f : \omega_1 \rightarrow [\omega_1]^{<\omega}$  stationarily often.

Build an infinitely branching tree  $T$  with underlying set  $\omega_1$  together with a specializing function  $\varphi : T \rightarrow \mathbb{Q}$  by recursion on levels, maintaining that if  $\varphi(x) < q$  then there is some  $y > x$  with  $\varphi(y) = q$ .

## ( $\diamond$ ) A special Lindelöf tree

Let  $\vec{f} = \langle f_\alpha : \alpha \in \lim(\omega_1) \rangle$  be such  $f_\alpha : \alpha \rightarrow [\alpha]^{<\omega}$  and  $\vec{f}$  guesses any  $f : \omega_1 \rightarrow [\omega_1]^{<\omega}$  stationarily often.

Build an infinitely branching tree  $T$  with underlying set  $\omega_1$  together with a specializing function  $\varphi : T \rightarrow \mathbb{Q}$  by recursion on levels, maintaining that if  $\varphi(x) < q$  then there is some  $y > x$  with  $\varphi(y) = q$ .

At stage  $\alpha = \omega \cdot \alpha$ , for each pair  $(x, q) \in (T \restriction \alpha) \times \mathbb{Q}$  with  $\varphi(x) < q$ , choose a cofinal branch  $b$  through  $T \restriction \alpha$  such that  $x \in b$ ,  $\sup(\varphi \restriction b) = q$  and **the unique point of  $b$  immediately above  $x$  is not in  $f_\alpha(x)$** . Then put a new node  $y \in T_\alpha$  above  $b$  and let  $\varphi(y) = q$ .

## ( $\diamond$ ) A special Lindelöf tree

Let  $\vec{f} = \langle f_\alpha : \alpha \in \lim(\omega_1) \rangle$  be such  $f_\alpha : \alpha \rightarrow [\alpha]^{<\omega}$  and  $\vec{f}$  guesses any  $f : \omega_1 \rightarrow [\omega_1]^{<\omega}$  stationarily often.

Build an infinitely branching tree  $T$  with underlying set  $\omega_1$  together with a specializing function  $\varphi : T \rightarrow \mathbb{Q}$  by recursion on levels, maintaining that if  $\varphi(x) < q$  then there is some  $y > x$  with  $\varphi(y) = q$ .

At stage  $\alpha = \omega \cdot \alpha$ , for each pair  $(x, q) \in (T \restriction \alpha) \times \mathbb{Q}$  with  $\varphi(x) < q$ , choose a cofinal branch  $b$  through  $T \restriction \alpha$  such that  $x \in b$ ,  $\sup(\varphi \restriction b) = q$  and **the unique point of  $b$  immediately above  $x$  is not in  $f_\alpha(x)$** . Then put a new node  $y \in T_\alpha$  above  $b$  and let  $\varphi(y) = q$ . Every  $y \in T_\alpha$  is obtained as above from some  $(x, q)$  and  $y \in (\uparrow x) \setminus \uparrow f_\alpha(x)$ , hence  $f_\alpha \restriction (T \restriction \alpha)$  covers  $T$ .

## ( $\diamond$ ) A special Lindelöf tree

Let  $\vec{f} = \langle f_\alpha : \alpha \in \lim(\omega_1) \rangle$  be such  $f_\alpha : \alpha \rightarrow [\alpha]^{<\omega}$  and  $\vec{f}$  guesses any  $f : \omega_1 \rightarrow [\omega_1]^{<\omega}$  stationarily often.

Build an infinitely branching tree  $T$  with underlying set  $\omega_1$  together with a specializing function  $\varphi : T \rightarrow \mathbb{Q}$  by recursion on levels, maintaining that if  $\varphi(x) < q$  then there is some  $y > x$  with  $\varphi(y) = q$ .

At stage  $\alpha = \omega \cdot \alpha$ , for each pair  $(x, q) \in (T \restriction \alpha) \times \mathbb{Q}$  with  $\varphi(x) < q$ , choose a cofinal branch  $b$  through  $T \restriction \alpha$  such that  $x \in b$ ,  $\sup(\varphi \restriction b) = q$  and **the unique point of  $b$  immediately above  $x$  is not in  $f_\alpha(x)$** . Then put a new node  $y \in T_\alpha$  above  $b$  and let  $\varphi(y) = q$ .

Every  $y \in T_\alpha$  is obtained as above from some  $(x, q)$  and  $y \in (\uparrow x) \setminus \uparrow f_\alpha(x)$ , hence  $f_\alpha \restriction (T \restriction \alpha)$  covers  $T$ .

Therefore,  $T$  is Lindelöf and so, under  $\diamond$ ,

$$\{\text{Suslin}\} \subsetneq \{\text{Lindelöf}\} \subsetneq \{\text{Aronszajn}\}$$

## Definition

Let  $T$  be a normal infinitely branching  $\aleph_1$ -tree. Consider the following poset  $\mathbb{P}$ : conditions are functions  $p \in \prod_{x \in F} [I(x)]^{<\omega}$ , where  $F \in [T]^{<\omega}$ , such that:

$$\forall x, y \in \text{dom}(p) (x < y \rightarrow y \restriction (\text{ht}(x) + 1) \in p(x)).$$



## Definition

Let  $T$  be a normal infinitely branching  $\aleph_1$ -tree. Consider the following poset  $\mathbb{P}$ : conditions are functions  $p \in \prod_{x \in F} [I(x)]^{<\omega}$ , where  $F \in [T]^{<\omega}$ , such that:

$$\forall x, y \in \text{dom}(p) (x < y \rightarrow y \restriction (\text{ht}(x) + 1) \in p(x)).$$


The order on  $\mathbb{P}$  is inclusion,  $p \leq q \iff p \supseteq q$ .

## Definition

Let  $T$  be a normal infinitely branching  $\aleph_1$ -tree. Consider the following poset  $\mathbb{P}$ : conditions are functions  $p \in \prod_{x \in F} [I(x)]^{<\omega}$ , where  $F \in [T]^{<\omega}$ , such that:

$$\forall x, y \in \text{dom}(p) (x < y \rightarrow y \restriction (\text{ht}(x) + 1) \in p(x)).$$

The order on  $\mathbb{P}$  is inclusion,  $p \leq q \iff p \supseteq q$ .




## Definition

Let  $T$  be a normal infinitely branching  $\aleph_1$ -tree. Consider the following poset  $\mathbb{P}$ : conditions are functions  $p \in \prod_{x \in F} [I(x)]^{<\omega}$ , where  $F \in [T]^{<\omega}$ , such that:

$$\forall x, y \in \text{dom}(p) (x < y \rightarrow y \restriction (\text{ht}(x) + 1) \in p(x)).$$

The order on  $\mathbb{P}$  is inclusion,  $p \leq q \iff p \supseteq q$ .




## Definition

Let  $T$  be a normal infinitely branching  $\aleph_1$ -tree. Consider the following poset  $\mathbb{P}$ : conditions are functions  $p \in \prod_{x \in F} [I(x)]^{<\omega}$ , where  $F \in [T]^{<\omega}$ , such that:

$$\forall x, y \in \text{dom}(p) (x < y \rightarrow y \restriction (\text{ht}(x) + 1) \in p(x)).$$

The order on  $\mathbb{P}$  is inclusion,  $p \leq q \iff p \supseteq q$ .




The generic subtree will be  $\dot{S} = \bigcup \{ \text{dom}(p) : p \in \dot{G} \}$ .

## Definition

Let  $T$  be a normal infinitely branching  $\aleph_1$ -tree. Consider the following poset  $\mathbb{P}$ : conditions are functions  $p \in \prod_{x \in F} [I(x)]^{<\omega}$ , where  $F \in [T]^{<\omega}$ , such that:

$$\forall x, y \in \text{dom}(p) (x < y \rightarrow y \restriction (\text{ht}(x) + 1) \in p(x)).$$

The order on  $\mathbb{P}$  is inclusion,  $p \leq q \iff p \supseteq q$ .



The generic subtree will be  $\dot{S} = \bigcup \{ \text{dom}(p) : p \in \dot{G} \}$ .

If  $p \in \mathbb{P}$  and  $x \in \text{dom}(p)$ , then  $p$  is a promise that  $p(x) = I_{\dot{S}}(x)$ .

## Theorem (M.)

*If  $T$  is a normal, infinitely branching Aronszajn tree, then  $\mathbb{P}$  has the ccc and  $\Vdash_{\mathbb{P}} \dot{S}$  is a finitely branching normal subtree of  $T$ .*

## Theorem (M.)

*If  $T$  is a normal, infinitely branching Aronszajn tree, then  $\mathbb{P}$  has the ccc and  $\Vdash_{\mathbb{P}} \dot{S}$  is a finitely branching normal subtree of  $T$ .*

Is  $T$  still Aronszajn in  $V^{\mathbb{P}}$ ?

## Theorem (M.)

*If  $T$  is a normal, infinitely branching Aronszajn tree, then  $\mathbb{P}$  has the ccc and  $\Vdash_{\mathbb{P}} \dot{S}$  is a finitely branching normal subtree of  $T$ .*

Is  $T$  still Aronszajn in  $V^{\mathbb{P}}$ ?

## Corollary

*Let  $\mathbb{S}$  be Baumgartner's poset for specializing  $T$  with finite conditions. Then  $\mathbb{S} \times \mathbb{P}$  is a ccc poset which forces that  $T$  is a special non-Lindelöf Aronszajn tree.*



## Theorem (M.)

*If  $T$  is a normal, infinitely branching Aronszajn tree, then  $\mathbb{P}$  has the ccc and  $\Vdash_{\mathbb{P}} \dot{S}$  is a finitely branching normal subtree of  $T$ .*

Is  $T$  still Aronszajn in  $V^{\mathbb{P}}$ ?

## Corollary

*Let  $\mathbb{S}$  be Baumgartner's poset for specializing  $T$  with finite conditions. Then  $\mathbb{S} \times \mathbb{P}$  is a ccc poset which forces that  $T$  is a special non-Lindelöf Aronszajn tree.*

## Corollary

*If  $\text{MA}_{\aleph_1}$  holds, then there are no Lindelöf trees.*

# Other ways of adding subtrees?

## Theorem (M.)

*Let  $T$  be an infinitely branching  $\aleph_1$  tree. Suppose  $\mathbb{P}$  is a poset,  $G$  is  $\mathbb{P}$ -generic over  $V$  and  $S \in V[G]$  is a finitely branching subtree of  $T$ .*

## Other ways of adding subtrees?

### Theorem (M.)

*Let  $T$  be an infinitely branching  $\aleph_1$  tree. Suppose  $\mathbb{P}$  is a poset,  $G$  is  $\mathbb{P}$ -generic over  $V$  and  $S \in V[G]$  is a finitely branching subtree of  $T$ . Suppose that  $\mathbb{P}$  is either*

- ▶ countably closed*
- ▶ strongly proper for a stationary set of countable elementary substructures of some (large)  $H_\lambda$ .*

*Then  $S \in V$ .*

Thank you :)

# References

- [1] Tomek Bartoszyński and Haim Judah. *Set Theory*. A K Peters, Ltd., Wellesley, MA, 1995. xii+546.
- [2] Keith J. Devlin and Håvard Johnsbråten. *The Souslin Problem*. Lecture Notes in Mathematics 405. Springer, 1974. 132 pp.
- [3] Pedro E. Marun. “Square Compactness and Lindelöf Trees”. In: *In preparation* (2023).
- [4] Peter J. Nyikos. “Various Topologies on Trees”. In: *Proceedings of the Tennessee Topology Conference (Nashville, TN, 1996)*. 1997.