

Models of the STLC

P. Rocha

February 20, 2017

1 Comments

If we were asked if the λ terms

$$\lambda f : o \rightarrow o. \lambda g : o \rightarrow o. \lambda x : o. f(g(x)) \text{ and } \lambda f : o \rightarrow o. \lambda g : o \rightarrow o. \lambda x : o. g(f(x))$$

were equal, we would promptly answer *no* since in general function composition is not commutative. If we were asked to prove it by syntactical means only we would be with a lot of trouble.

One of the advantages of having denotational models of the lambda calculus lies precisely in the fact that we can easily answer questions like that once we have a model that matches our intuition on what a lambda term means. Another is the connection established between the denotational models and the operational meaning that somehow guarantees that the operation rules that we devised were not randomly chosen, that they make sense.

When we think of an interpretation of the simply typed λ -calculus we naturally interpret types as sets and terms of a given type as elements of the corresponding set. The idea is formalized through the notion of a *type frame*.

Definition 1. A *pre-frame* is a pair of functions (\mathcal{A}, A) where \mathcal{A} acts on types and A on pair of types for which the following holds

- For every type t , $\mathcal{A}[[t]] \neq \emptyset$
- $A^{s,t} : \mathcal{A}[[s \rightarrow t]] \times \mathcal{A}[[s]] \rightarrow \mathcal{A}[[t]]$ (intuitively gives an interpretation of function application)
- If $f, g \in \mathcal{A}[[s \rightarrow t]]$ and $A^{s,t}(f, x) = A^{s,t}(g, x)$, for every $x \in \mathcal{A}[[s]]$, then $f = g$ (extensionality property)

□

We can then extend the function \mathcal{A} to interpret terms. But we have to be careful since a term might not be closed and therefore its interpretation will depend on the type of its free variables. Thus, we will not interpret terms but typing derivations. For that, the notion of H -environment will play a key role.

Definition 2. Let H be a type assignment and (\mathcal{A}, A) a type frame. The function ρ that acts on variables is said to be a H -environment if for every $x : t \in H$ we have

$$\rho(x) \in \mathcal{A}[[t]]$$

□

Definition 3. A *type frame* (or *frame*) is a pre-frame (\mathcal{A}, A) in which we can extend \mathcal{A} to act on triples $H \triangleright M : t$, where $H \vdash M : t$ and the following holds

$$(F1) \quad \mathcal{A}[[H \triangleright x : t]]\rho = \rho(x)$$

$$(F2) \quad \mathcal{A}[[H \triangleright M(N)]]\rho = A^{s,t}(\mathcal{A}[[H \triangleright M : s \rightarrow t]]\rho, \mathcal{A}[[H \triangleright N : s]]\rho), \text{ where } s \text{ is the unique type such that } H \vdash M : s \rightarrow t$$

$$(F3) \quad A^{s,t}(\mathcal{A}[[H \triangleright \lambda x : s. M : s \rightarrow t]]\rho, d) = \mathcal{A}[[H, x : s \triangleright M : t]]\rho[x \mapsto d]$$

□

As it was proved in Lemma 2.14 [Gun], if a pre-frame has an extension to a frame, then that extension must be unique. However, not every pre-frame can be extended to a frame. For example, if we let

$$\mathcal{A}[[t]] = \mathbb{N}$$

for every type t and define $A^{s,t}$ as being the natural numbers multiplication, that is

$$A^{s,t}(m, n) = m * n$$

then we may observe that (\mathcal{A}, A) fulfils every condition to be considered a pre-frame. In particular, the extensionality property holds since if

$$A^{s,t}(m, x) = A^{s,t}(m, x)$$

for every $x \in \mathbb{N}$, then $m * 1 = n * 1$ and consequently, $m = n$.

If now we assume that (\mathcal{A}, A) is also a frame, then we get from condition (F3):

$$\begin{aligned} (\mathcal{A}[[y : o \triangleright \lambda x : o \rightarrow o. y]]\rho) * n &= \mathcal{A}[[y : o, x : o \rightarrow o \triangleright y]]\rho[x \mapsto n] \\ &= \rho(y) \end{aligned}$$

which is a contradiction since we may vary n so as to get $\rho(y) = 2 * \rho(y)$, for example.

Definition 4. Given a frame \mathcal{A} and an equation $(H \triangleright M = N : t)$ (where $H \models M, N : t$) we define the satisfiability relation \models by

$$\mathcal{A} \models (H \triangleright M = N : t) \Leftrightarrow \mathcal{A} \llbracket H \triangleright M : t \rrbracket = \mathcal{A} \llbracket H \triangleright N : t \rrbracket$$

We extend \models to sets of equations by

$$\mathcal{A} \models T$$

if $\mathcal{A} \models x$ for every $x \in T$. □

The first correspondence between the denotational models of the simply typed λ -calculus and the operational meaning is established by the soundness Theorem 2.15 [Gun]. We present it here and give it a proof. But before we must prove some auxiliary results.

Lemma 1. Suppose $H \vdash M : t$, $x \notin H$ and $d \in \llbracket s \rrbracket$. Then,

$$\mathcal{A} \llbracket H, x : s \triangleright M : t \rrbracket \rho[x \mapsto d] = \mathcal{A} \llbracket H \triangleright M : t \rrbracket \rho$$

for every frame \mathcal{A} and every H -environment ρ .

Proof. The proof is done by induction on the height of the derivation tree of $H \vdash M : t$.

If $H \vdash y : t$, then $y \neq x$ and

$$\begin{aligned} \mathcal{A} \llbracket H, x : s \triangleright y : t \rrbracket \rho[x \mapsto d] &= \rho(y) \\ &= \mathcal{A} \llbracket H \triangleright y : t \rrbracket \rho \end{aligned}$$

If $H \vdash M_1 M_2 : t$, then $H \vdash M_1 : r \rightarrow t$ and $H \vdash M_2 : r$ for some type r . Thus,

$$\begin{aligned} &\mathcal{A} \llbracket H, x : s \triangleright M_1 M_2 : t \rrbracket \rho[x \mapsto d] \\ &= A^{r,t}(\mathcal{A} \llbracket H, x : s \triangleright M_1 : r \rightarrow t \rrbracket \rho[x \mapsto d], \mathcal{A} \llbracket H, x : s \triangleright M_2 : r \rrbracket \rho[x \mapsto d]) \\ &= A^{r,t}(\mathcal{A} \llbracket H \triangleright M_1 : r \rightarrow t \rrbracket \rho, \mathcal{A} \llbracket H \triangleright M_2 : r \rrbracket \rho) \quad (\text{induction}) \\ &= \mathcal{A} \llbracket H \triangleright M_1 M_2 : t \rrbracket \rho \end{aligned}$$

If $H \vdash \lambda y : r. M : t$, we may assume that $y \neq x$. Then, for every $d_1 \in \llbracket r \rrbracket$, we have

$$\begin{aligned} &A^{r,t}(\mathcal{A} \llbracket H \triangleright \lambda y : r. M : t \rrbracket \rho, d_1) \\ &= \mathcal{A} \llbracket H, y : r \triangleright M : t \rrbracket \rho[y \mapsto d_1] \\ &= \mathcal{A} \llbracket H, y : r, x : s \triangleright M : t \rrbracket \rho[y \mapsto d_1][x \mapsto d] \quad (\text{induction}) \\ &= \mathcal{A} \llbracket H, x : s, y : r \triangleright M : t \rrbracket \rho[x \mapsto d][y \mapsto d_1] \\ &= A^{r,t}(\mathcal{A} \llbracket H, x : s \triangleright \lambda y : r. M : r \rightarrow t \rrbracket \rho[x \mapsto d], d_1) \end{aligned}$$

As the extensionality property holds we obtain the desired result. □

Lemma 2 (Substitution). *Suppose $H, x : s \vdash M : t$ and $H \vdash N : s$. Then, for every H -environment ρ ,*

$$\mathcal{A}[[H, x : s \triangleright M : t]]\rho[x \mapsto \mathcal{A}[[H \triangleright N : t]]\rho] = \mathcal{A}[[H \triangleright [N/x]M : t]]\rho$$

Proof. The proof is done by induction on the structure of M . For every case, let

$$\rho^* = \rho[x \mapsto \mathcal{A}[[H \triangleright N : t]]\rho]$$

- If $M = y$ and $y = x$, then

$$\begin{aligned} & \mathcal{A}[[H, x : s \triangleright x : s]]\rho^* \\ &= \rho[x \mapsto \mathcal{A}[[H \triangleright N : t]]\rho](x) \\ &= \mathcal{A}[[H \triangleright N : t]]\rho \\ &= \mathcal{A}[[H \triangleright [N/x]M : s]]\rho \end{aligned}$$

On the other hand, if $y \neq x$, then by applying Lemma (1) we obtain

$$\begin{aligned} & \mathcal{A}[[H, x : s \triangleright y : t]]\rho^* \\ &= \mathcal{A}[[H \triangleright y : t]]\rho \\ &= \mathcal{A}[[H \triangleright [N/x]y : t]]\rho \end{aligned}$$

- Suppose $M = M_1 M_2$, $H, x : s \vdash M_1 : r \rightarrow t$ and $H, x : s \vdash M_2 : r$. Then,

$$\begin{aligned} & \mathcal{A}[[H, x : s \triangleright M_1 M_2 : t]]\rho^* \\ &= A^{r,t}(\mathcal{A}[[H, x : s \triangleright M_1 : r \rightarrow t]]\rho^*, \mathcal{A}[[H, x : s \triangleright M_2 : r]]\rho^*) \\ &= A^{r,t}(\mathcal{A}[[H \triangleright [N/x]M_1 : r \rightarrow t]]\rho, \mathcal{A}[[H \triangleright [N/x]M_2 : r]]\rho) \\ & \quad \text{(induction)} \\ &= \mathcal{A}[[H \triangleright ([N/x]M_1)([N/x]M_2) : t]]\rho \\ &= \mathcal{A}[[H \triangleright [N/x](M_1 M_2) : t]]\rho \end{aligned}$$

- If $M = \lambda y : r. M_1 : r \rightarrow t$ we may assume that $y \neq x$ (if it were not the case, we could choose an α -equivalent term such that the inequality would hold). Then, for every $d \in \llbracket r \rrbracket$ we have

$$\begin{aligned} & A^{r,t}(\mathcal{A}[[H, x : s \triangleright \lambda y. M_1 : r \rightarrow t]]\rho^*, d) \\ &= \mathcal{A}[[H, x : s, y : r \triangleright M_1 : t]]\rho^*[y \mapsto d] \\ &= \mathcal{A}[[H, y : r \triangleright [N/x]M_1 : t]\rho[y \mapsto d] \quad \text{(induction)} \\ &= A^{r,t}(\mathcal{A}[[H \triangleright [N/x]\lambda y. M_1 : r \rightarrow t]]\rho, d) \end{aligned}$$

From the extensionality property it follows

$$\mathcal{A}[\![H, x : s \triangleright \lambda y. M_1 : r \rightarrow t]\!] \rho^* = \mathcal{A}[\![H \triangleright [N/x] \lambda y. M_1 : r \rightarrow t]\!] \rho$$

□

Lemma 3 (Soundness). *For any theory T and frame \mathcal{A} , if $\mathcal{A} \models T$ and $T \vdash (H \triangleright M = N : t)$, then $\mathcal{A} \models (H \triangleright M = N : t)$.*

Proof. The proof is done by induction on the height of the derivation $T \vdash (H \triangleright M = N : t)$. We prove that \models is closed under the rules of Table 2.2 [Gun]. It's almost a direct consequence of the properties of \models for the rules $\{Axiom\}$, $\{Permute\}$, $\{Refl\}$, $\{Sym\}$ and $\{Trans\}$. Thus we only prove closure for the remaining rules.

- $\{Add\}$. Suppose $\mathcal{A} \models (H \triangleright M = N : t)$ and $x \notin H$. Since every $H, x : s$ -environment is also a H -environment we conclude that $\mathcal{A} \models (H, x : s \triangleright M = N : t)$.
- $\{Drop\}$. Suppose $\mathcal{A} \models (H, x : s \triangleright M = N : t)$ and $x \notin FV(M) \cup FV(N)$. Let $d \in \llbracket s \rrbracket$. For every H -environment ρ

$$\begin{aligned} \mathcal{A}[\![H \triangleright M : t]\!] \rho &= \llbracket H, x : s \triangleright M : t \rrbracket \rho[x \mapsto d] && (\text{Lemma (1)}) \\ &= \llbracket H, x : s \triangleright N : t \rrbracket \rho[x \mapsto d] \\ &= \llbracket H \triangleright M : t \rrbracket \rho && (\text{Lemma (1)}) \end{aligned}$$

- $\{Cong\}$. Suppose $\mathcal{A} \models (H \triangleright M_1 = M_2 : s \rightarrow t)$ and $\mathcal{A} \models (H \triangleright N_1 = N_2 : s)$. Then, for every H -environment ρ

$$\begin{aligned} &\mathcal{A}[\![H \triangleright M_1(N_1) : t]\!] \rho \\ &= A^{s,t}(\mathcal{A}[\![H \triangleright M_1 : s \rightarrow t]\!] \rho, \mathcal{A}[\![H \triangleright N_1 : s]\!] \rho) \\ &= A^{s,t}(\mathcal{A}[\![H \triangleright M_2 : s \rightarrow t]\!] \rho, \mathcal{A}[\![H \triangleright N_2 : s]\!] \rho) \\ &= \mathcal{A}[\![H \triangleright M_2(N_2) : t]\!] \rho \end{aligned}$$

- $\{\xi\}$. Suppose $\mathcal{A} \models (H, x : s \triangleright M = N : t)$. Then, for every H -environment ρ and $d \in \llbracket s \rrbracket$

$$\begin{aligned} &A^{s,t}(\mathcal{A}[\![H \triangleright \lambda x : s. M : s \rightarrow t]\!] \rho, d) \\ &= \mathcal{A}[\![H, x : s \triangleright M : t]\!] \rho[x \mapsto d] \\ &= \mathcal{A}[\![H, x : s \triangleright N : t]\!] \rho[x \mapsto d] \\ &= A^{s,t}(\mathcal{A}[\![H \triangleright \lambda x : s. M : s \rightarrow t]\!] \rho, d) \end{aligned}$$

Hence, by the extensionality property follows the desired result.

- $\{\beta\}$. Suppose $H, x : s \vdash M : t$ and $H \vdash N : s$. Then,

$$\begin{aligned}
& \mathcal{A}[\![H \triangleright (\lambda x : s. M)N : t]\!]\rho \\
&= A^{s,t}(\mathcal{A}[\![H \triangleright \lambda x : s. M : s \rightarrow t]\!]\rho, \mathcal{A}[\![H \triangleright N : s]\!]\rho) \\
&= \mathcal{A}[\![H, x : s \triangleright M : t]\!]\rho[x \mapsto \mathcal{A}[\![H \triangleright N : s]\!]\rho] \\
&= \mathcal{A}[\![H \triangleright [N/x]M : t]\!]\rho \quad (\text{Lemma (2)})
\end{aligned}$$

- $\{\eta\}$. Suppose $H \vdash M : s \rightarrow t$ and $x \notin FV(M)$. Then, for every H -environment ρ and $d \in \llbracket s \rrbracket$

$$\begin{aligned}
& A^{s,t}(\mathcal{A}[\![H \triangleright \lambda x : s. M(x)]\!]\rho, d) \\
&= \mathcal{A}[\![H, x : s \triangleright M(x) : t]\!]\rho[x \mapsto d] \\
&= A^{s,t}(\mathcal{A}[\![H, x : s \triangleright M : s \rightarrow t]\!]\rho[x \mapsto d], \mathcal{A}[\![H, x : s \triangleright x : t]\!]\rho[x \mapsto d]) \\
&= A^{s,t}(\mathcal{A}[\![H \triangleright M : s \rightarrow t]\!]\rho, d) \quad (\text{Lemma (1)})
\end{aligned}$$

By the extensionality property we get the desired result.

□

2 Exercises

Definition 5. Let \mathcal{A} and \mathcal{B} be frames. A family of relations R^s indexed over types s is said to be a logical relation between \mathcal{A} and \mathcal{B} if

- $R^s \subseteq \mathcal{A}[[s]] \times \mathcal{B}[[s]]$
- $fR^{s \rightarrow t}$ iff $\forall x, y \ xR^s y \implies A^{s,t}(f, x)R^t B^{s,t}(g, y)$

□

Exercise 2.16. The notion of a partial homomorphism is a special instance of a logical relation. Explain why this is the case and conjecture a version of Lemma 2.22 about partial homomorphisms that would apply to logical relations. Prove your version of the lemma.

Solution.

Definition 6. We say that a binary relation $R \subseteq A \times B$ is *surjective* if

$$\forall b \in B \ \exists a \in A \ aRb$$

□

Definition 7. We say that a binary relation $R \subseteq A \times B$ is *function-like* if

$$\forall b \in B \ \forall a_1, a_2 \in A \ a_1 R b \text{ and } a_2 R b \implies a_1 = a_2$$

□

Observe that every partial homomorphism $\Phi : \mathcal{A} \rightarrow \mathcal{B}$ induces a function-like surjective logical relation R in the following way. For every Φ^s define R^s by

$$xR^s y \Leftrightarrow \Phi^s(x) = y$$

□

Exercise 2.15. Show that Theorem 2.25 fails when X is not infinite

Solution. For example, if X is a singleton set, then for every H -environment ρ , where $H = x : o, y : o$, we have

$$\begin{aligned} \mathcal{F}_X[[H \triangleright x : o]]\rho &= \rho(x) \\ &= \rho(y) \\ &= \mathcal{F}_X[[H \triangleright y : o]]\rho \end{aligned}$$

Thus,

$$\mathcal{F}_X \models (H \triangleright x = y : o)$$

But as we saw in Theorem 2.13, the simply-type λ -calculus is non-trivial, therefore it is not the case that

$$\vdash (H \triangleright x = y : o)$$

For the most general case in which X is finite but not necessarily a singleton set, consider the following Church encoding of the natural numbers

$$\begin{aligned} c_0 &= \lambda f : o \rightarrow o. \lambda x : o. x \\ c_n &= \lambda f : o \rightarrow o. \lambda x : o. f(c_{n-1}fx), \quad n \geq 1 \end{aligned}$$

Observe that if X_1 and X_2 are finite, then so it is $X_2^{X_1}$, the set of functions from X_1 to X_2 . Thus, for every type t , $\mathcal{F}_X \llbracket t \rrbracket$ must be finite.

Observe also that since each c_n is a closed term, the denotation on \mathcal{F}_X will be independent of the environment, that is, for any two environments ρ_1 and ρ_2

$$\mathcal{F}_X \llbracket \triangleright c_n : t \rrbracket \rho_1 = \mathcal{F}_X \llbracket \triangleright c_n : t \rrbracket \rho_2$$

for each $n \in \mathbb{N}$, where $t = ((o \rightarrow o) \rightarrow o) \rightarrow o$.

Since we can choose finitely many elements out of $\mathcal{F}_X \llbracket t \rrbracket$, then it must be the case that for some $n, m \in \mathbb{N}$ with $n < m$ we have

$$\mathcal{F}_X \models (\triangleright c_n = c_m : t)$$

But clearly that fails to hold in every frame. For example, in $\mathcal{F}_{\mathbb{N}}$, if we let $s : \mathbb{N} \rightarrow \mathbb{N}$ be the successor function, then

$$\begin{aligned} (\mathcal{F}_{\mathbb{N}} \llbracket \triangleright c_n : t \rrbracket \rho s) 0 &= n \\ &\neq m \\ &= (\mathcal{F}_{\mathbb{N}} \llbracket \triangleright c_m : t \rrbracket \rho s) 0 \end{aligned}$$

Thus,

$$\mathcal{F}_{\mathbb{N}} \not\models (\triangleright c_n = c_m : t)$$

and, consequently, the soundness result implies that

$$\not\vdash (\triangleright c_n = c_m : t)$$

Hence, Theorem 2.25 fails. □