

# Spectral relaxations of persistent rank invariants


With a focus on *parameterized* settings

Matt Piekenbrock<sup>†</sup> & Jose Perea<sup>‡</sup>

## Vectorizing diagrams

There are many mappings from dgm's to function spaces (e.g. Hilbert spaces)

- Persistence Landscapes (Bubenik 2020)




$$\lambda(k, t) = \sup\{h \geq 0 \mid \text{rank}(H_p^{i-h} \rightarrow H_p^{i+h}) \geq k\}$$

## Vectorizing diagrams

There are many mappings from dgm's to function spaces (e.g. Hilbert spaces)

- Persistence Landscapes (Bubenik 2020) + Learning applications (Kim et al. 2020)

$$\lambda(k, t) = \sup\{h \geq 0 \mid \text{rank}(H_p^{i-h} \rightarrow H_p^{i+h}) \geq k\}$$




## Vectorizing diagrams

There are many mappings from dgm's to function spaces (e.g. Hilbert spaces)

Persistence Landscapes (Bubenik 2020) + Learning applications (Kim et al. 2020)

Persistence Images (Adams et al. 2017)



$$\rho(f, \phi) = \sum_{(i,j) \in \text{dgm}} f(i, j) \phi(|j - i|)$$


## Vectorizing diagrams

There are many mappings from dgm's to function spaces (e.g. Hilbert spaces)

Persistence Landscapes (Bubenik 2020) + Learning applications (Kim et al. 2020)

Persistence Images (Adams et al. 2017) + Learning applications (Som et al. 2020)

$$\rho(f, \phi) = \sum_{(i,j) \in \text{dgm}} f(i, j) \phi(|j - i|)$$




## Vectorizing diagrams

There are many mappings from dgm's to function spaces (e.g. Hilbert spaces)

Persistence Landscapes (Bubenik 2020) + Learning applications (Kim et al. 2020)

Persistence Images (Adams et al. 2017) + Learning applications (Som et al. 2020)

A few others...<sup>1</sup>



See (Bubenik 2020) for an overview.

## Many goals in common...

- Vectorize persistence information
- Optimize persistence invariants
- Leverage the stability of persistence
- Connect to other areas of mathematics