

# Solutions

FREEMAN CHENG

7 August 2024

## Problem (4-3).

For notation, take  $\langle \cdot, \cdot \rangle$  to be  $T(\cdot, \cdot)$ . Let  $e_1, \dots, e_n$  be an orthonormal basis for  $V$ . By definition,  $|\omega(e_1, \dots, e_n)| = 1$ , and applying Theorem 4-6 gives

$$|\omega(e_1, \dots, e_n)| = \left| \det \begin{bmatrix} \ddots & & \\ & \langle w_i, e_j \rangle & \\ & & \ddots \end{bmatrix} \right|.$$

Using the common identity (see any quantum mechanics textbook)

$$\begin{bmatrix} \ddots & & \\ & \langle w_i, e_j \rangle & \\ & & \ddots \end{bmatrix} \begin{bmatrix} \ddots & & \\ & \langle e_i, w_j \rangle & \\ & & \ddots \end{bmatrix} = \begin{bmatrix} \ddots & & \\ & \langle w_i, w_j \rangle & \\ & & \ddots \end{bmatrix},$$

we arrive at the desired result.

## Problem (4-4).

$$\begin{aligned} f^*\omega(v_1, \dots, v_n) &= \omega(f(v_1), \dots, f(v_n)) \\ &= \det \begin{bmatrix} \ddots & & \\ & T(f(v_i), f(e_j)) & \\ & & \ddots \end{bmatrix} \omega(f(e_1), \dots, f(e_n)) \quad (\text{Theorem 4-3}) \\ &= \det \begin{bmatrix} \ddots & & \\ & T(f(v_i), f(e_j)) & \\ & & \ddots \end{bmatrix} \\ &= \det \begin{bmatrix} \ddots & & \\ & \langle v_i, e_j \rangle & \\ & & \ddots \end{bmatrix}, \end{aligned}$$

where the last two equalities come from the fact that  $\omega$  is a volume element and  $f^*T(\cdot, \cdot) = \langle \cdot, \cdot \rangle$ .

## Problem (4-5).

Because  $\det$  is continuous, the image of  $\det \circ \gamma$  on the path must be of the same sign.

## Problem (4-6).