

Fast high-order solvers on simplices for the de Rham complex


Patrick E. Farrell^{1,2} Pablo Brubeck¹ Charles Parker¹ Rob Kirby³



¹University of Oxford


²Charles University

³Baylor University



High-order finite element discretisations have important advantages:

- ✓ rapid convergence,
- ✓ high arithmetic intensity.




High-order finite element discretisations have important advantages:

- ✓ rapid convergence,
- ✓ high arithmetic intensity.

But there are practical challenges to using them:

- ✗ require high-order meshes,
- ✗ **require fast, matrix-free solvers.**



High-order finite element discretisations have important advantages:

- ✓ rapid convergence,
- ✓ high arithmetic intensity.

But there are practical challenges to using them:

- ✗ require high-order meshes,
- ✗ **require fast, matrix-free solvers.**

This talk ...

... is about fast solvers for high-order discretisations of some canonical PDEs on simplices.

The PDEs we consider derive from the de Rham complex

$$H(\text{grad}) \xrightarrow{\text{grad}} H(\text{curl}) \xrightarrow{\text{curl}} H(\text{div}) \xrightarrow{\text{div}} L^2.$$

The PDEs we consider derive from the de Rham complex

$$H(\text{grad}) \xrightarrow{\text{grad}} H(\text{curl}) \xrightarrow{\text{curl}} H(\text{div}) \xrightarrow{\text{div}} L^2.$$

They are the *Riesz maps*

$$\text{find } u \in H(\text{grad}) : (\beta u, v) + (\alpha \text{grad } u, \text{grad } v) = (f, v) \quad \forall v \in H(\text{grad}),$$

$$\text{find } \mathbf{u} \in H(\text{curl}) : (\beta \mathbf{u}, \mathbf{v}) + (\alpha \text{curl } \mathbf{u}, \text{curl } \mathbf{v}) = (\mathbf{f}, \mathbf{v}) \quad \forall \mathbf{v} \in H(\text{curl}),$$

$$\text{find } \mathbf{u} \in H(\text{div}) : (\beta \mathbf{u}, \mathbf{v}) + (\alpha \text{div } \mathbf{u}, \text{div } \mathbf{v}) = (\mathbf{f}, \mathbf{v}) \quad \forall \mathbf{v} \in H(\text{div}).$$

The PDEs we consider derive from the de Rham complex

$$H(\text{grad}) \xrightarrow{\text{grad}} H(\text{curl}) \xrightarrow{\text{curl}} H(\text{div}) \xrightarrow{\text{div}} L^2.$$

They are the *Riesz maps*

$$\text{find } u \in H(\text{grad}) : (\beta u, v) + (\alpha \text{grad } u, \text{grad } v) = (f, v) \quad \forall v \in H(\text{grad}),$$

$$\text{find } \mathbf{u} \in H(\text{curl}) : (\beta \mathbf{u}, \mathbf{v}) + (\alpha \text{curl } \mathbf{u}, \text{curl } \mathbf{v}) = (\mathbf{f}, \mathbf{v}) \quad \forall \mathbf{v} \in H(\text{curl}),$$

$$\text{find } \mathbf{u} \in H(\text{div}) : (\beta \mathbf{u}, \mathbf{v}) + (\alpha \text{div } \mathbf{u}, \text{div } \mathbf{v}) = (\mathbf{f}, \mathbf{v}) \quad \forall \mathbf{v} \in H(\text{div}).$$

These Riesz maps are key building blocks in developing block preconditioners:

The PDEs we consider derive from the de Rham complex

$$H(\text{grad}) \xrightarrow{\text{grad}} H(\text{curl}) \xrightarrow{\text{curl}} H(\text{div}) \xrightarrow{\text{div}} L^2.$$

They are the *Riesz maps*

$$\text{find } u \in H(\text{grad}) : (\beta u, v) + (\alpha \text{grad } u, \text{grad } v) = (f, v) \quad \forall v \in H(\text{grad}),$$

$$\text{find } \mathbf{u} \in H(\text{curl}) : (\beta \mathbf{u}, \mathbf{v}) + (\alpha \text{curl } \mathbf{u}, \text{curl } \mathbf{v}) = (\mathbf{f}, \mathbf{v}) \quad \forall \mathbf{v} \in H(\text{curl}),$$

$$\text{find } \mathbf{u} \in H(\text{div}) : (\beta \mathbf{u}, \mathbf{v}) + (\alpha \text{div } \mathbf{u}, \text{div } \mathbf{v}) = (\mathbf{f}, \mathbf{v}) \quad \forall \mathbf{v} \in H(\text{div}).$$

These Riesz maps are key building blocks in developing block preconditioners:



-  R. Hiptmair. "Operator Preconditioning". In: *Computers & Mathematics with Applications* 52.5 (2006), pp. 699–706.
-  K.-A. Mardal and R. Winther. "Preconditioning discretizations of systems of partial differential equations". In: *Numerical Linear Algebra with Applications* 18.1 (2011), pp. 1–40.

Headline result

Co-designed basis + solver with $\mathcal{O}(p^6)$ flops and $\mathcal{O}(p^4)$ storage (vs $\mathcal{O}(p^9)$ and $\mathcal{O}(p^6)$).



Headline result

Co-designed basis + solver with $\mathcal{O}(p^6)$ flops and $\mathcal{O}(p^4)$ storage (vs $\mathcal{O}(p^9)$ and $\mathcal{O}(p^6)$).

Flop count, $H(\text{grad})$ Peak memory, $H(\text{grad})$



Headline result

Co-designed basis + solver with $\mathcal{O}(p^6)$ flops and $\mathcal{O}(p^4)$ storage (vs $\mathcal{O}(p^9)$ and $\mathcal{O}(p^6)$).

Flop count, $H(\text{curl})$ Peak memory, $H(\text{curl})$

Headline result

Co-designed basis + solver with $\mathcal{O}(p^6)$ flops and $\mathcal{O}(p^4)$ storage (vs $\mathcal{O}(p^9)$ and $\mathcal{O}(p^6)$).

Flop count, $H(\text{div})$ Peak memory, $H(\text{div})$

Full disclosure:

Full disclosure:

Offline setup costs

involve solving $\mathcal{O}(1)$ small eigenproblems, with complexities $\mathcal{O}(p^9)$ time and $\mathcal{O}(p^6)$ storage.

Full disclosure:

Offline setup costs

involve solving $\mathcal{O}(1)$ small eigenproblems, with complexities $\mathcal{O}(p^9)$ time and $\mathcal{O}(p^6)$ storage.

Matrix-vector products

We don't know how to sum-factorise the basis, so matvecs are $\mathcal{O}(p^6)$ time and $\mathcal{O}(p^3)$ storage.

Full disclosure:

Offline setup costs

involve solving $\mathcal{O}(1)$ small eigenproblems, with complexities $\mathcal{O}(p^9)$ time and $\mathcal{O}(p^6)$ storage.

Matrix-vector products

We don't know how to sum-factorise the basis, so matvecs are $\mathcal{O}(p^6)$ time and $\mathcal{O}(p^3)$ storage.

Open question

Can we achieve $\mathcal{O}(p^4)$ flops and $\mathcal{O}(p^3)$ storage for solver + matvecs on simplices?

Full disclosure:

Offline setup costs

involve solving $\mathcal{O}(1)$ small eigenproblems, with complexities $\mathcal{O}(p^9)$ time and $\mathcal{O}(p^6)$ storage.

Matrix-vector products

We don't know how to sum-factorise the basis, so matvecs are $\mathcal{O}(p^6)$ time and $\mathcal{O}(p^3)$ storage.

Open question

Can we achieve $\mathcal{O}(p^4)$ flops and $\mathcal{O}(p^3)$ storage for solver + matvecs on simplices? (Hopefully.)

Section 2

The key idea

All multigrid/domain decomposition solvers must do some relaxation on fine grids.

All multigrid/domain decomposition solvers must do some relaxation on fine grids.

Star patches give effective and p -robust relaxation methods.

All multigrid/domain decomposition solvers must do some relaxation on fine grids.

Star patches give effective and p -robust relaxation methods.

Definition (star)


The $\text{star } *e$ of an entity e of dimension k is the union of all entities of dimension $k' \geq k$ containing e .

All multigrid/domain decomposition solvers must do some relaxation on fine grids.

Star patches give effective and p -robust relaxation methods.

Definition (star)

The $\star e$ of an entity e of dimension k is the union of all entities of dimension $k' \geq k$ containing e .



\star vertex


$k = 0$

All multigrid/domain decomposition solvers must do some relaxation on fine grids.

Star patches give effective and p -robust relaxation methods.

Definition (star)

The $\star e$ of an entity e of dimension k is the union of all entities of dimension $k' \geq k$ containing e .




All multigrid/domain decomposition solvers must do some relaxation on fine grids.


Star patches give effective and p -robust relaxation methods.

Definition (star)


The $\star e$ of an entity e of dimension k is the union of all entities of dimension $k' \geq k$ containing e .



\star vertex



\star edge



\star cell

$k = 0$

$k = 1$


$k = 2$

All multigrid/domain decomposition solvers must do some relaxation on fine grids.

Star patches give effective and p -robust relaxation methods.


Definition (star)

The $\star e$ of an entity e of dimension k is the union of all entities of dimension $k' \geq k$ containing e .



\star vertex

$$k = 0$$



\star edge

$$k = 1$$



\star cell

$$k = 2$$

Let's see some examples of how star patches are useful in solvers.

For the Poisson problem in $H(\text{grad})$, the space decomposition

$$X_p^0 = X_1^0 + \sum_{v \in \Delta_0(\mathcal{T}_h)} X_p^0|_{\star v}$$



is p -robust: only $\mathcal{O}(1)$ conjugate gradient iterations.

Here $\Delta_k(\mathcal{T}_h)$ is the set of k -entities in the mesh \mathcal{T}_h .

Joachim Schöberl



For the Poisson problem in $H(\text{grad})$, the space decomposition

$$X_p^0 = X_1^0 + \sum_{v \in \Delta_0(\mathcal{T}_h)} X_p^0|_{\star v}$$



Joachim Schöberl

is p -robust: only $\mathcal{O}(1)$ conjugate gradient iterations.

Here $\Delta_k(\mathcal{T}_h)$ is the set of k -entities in the mesh \mathcal{T}_h .

For the $H(\text{curl})$ ($k = 1$) and $H(\text{div})$ ($k = 2$) Riesz maps, the space decomposition

$$X_p^k = X_1^k + \sum_{J \in \Delta_{k-1}(\mathcal{T}_h)} d^{k-1} X_p^{k-1}|_{\star J} + \sum_{L \in \Delta_k(\mathcal{T}_h)} X_p^k|_{\star L}$$



is p -robust (but the proof is an open problem).

If the stiffness matrix on a cell is dense, star patch problems have $\mathcal{O}(p^d)$ unknowns.

If the stiffness matrix on a cell is dense, star patch problems have $\mathcal{O}(p^d)$ unknowns.

Bad news

Doing dense linear algebra with $\mathcal{O}(p^d)$ unknowns costs $\mathcal{O}(p^{3d})$ flops and $\mathcal{O}(p^{2d})$ storage.

If the stiffness matrix on a cell is dense, star patch problems have $\mathcal{O}(p^d)$ unknowns.

Bad news

Doing dense linear algebra with $\mathcal{O}(p^d)$ unknowns costs $\mathcal{O}(p^{3d})$ flops and $\mathcal{O}(p^{2d})$ storage.

A single vertex star can typically have around 24 tetrahedra. Memory, setup, and solution costs become prohibitive as p increases.

If the stiffness matrix on a cell is dense, star patch problems have $\mathcal{O}(p^d)$ unknowns.

Bad news

Doing dense linear algebra with $\mathcal{O}(p^d)$ unknowns costs $\mathcal{O}(p^{3d})$ flops and $\mathcal{O}(p^{2d})$ storage.

A single vertex star can typically have around 24 tetrahedra. Memory, setup, and solution costs become prohibitive as p increases.

Our response

Solve lower-dimensional patch problems: $\mathcal{O}(p^{3(d-1)})$ flops and $\mathcal{O}(p^{2(d-1)})$ storage.

Complexities for some p -robust high-order solvers in three dimensions

		Year	Cell	Problem	Time	Storage
►	Pavarino & Widlund	1993		$H(\text{grad})$	p^9	p^6
	Arnold, Falk & Winther	2000		all	p^9	p^6
	Beuchler & Pillwein	2007		$H(\text{grad})$	-	-
	Schöberl et al.	2008		$H(\text{grad})$	p^9	p^6
	Farrell & Brubeck	2024		all	p^4	p^3
	This work	2025		all	p^6	p^4

Pavarino & Widlund

proved that two-level p -multigrid with *star patches* on quads and hexes is p -robust.

Complexities for some p -robust high-order solvers in three dimensions

		Year	Cell	Problem	Time	Storage
	Pavarino & Widlund	1993		$H(\text{grad})$	p^9	p^6
►	Arnold, Falk & Winther	2000		all	p^9	p^6
	Beuchler & Pillwein	2007		$H(\text{grad})$	-	-
	Schöberl et al.	2008		$H(\text{grad})$	p^9	p^6
	Farrell & Brubeck	2024		all	p^4	p^3
	This work	2025		all	p^6	p^4

Arnold, Falk & Winther

proved that *star patches* are good h -multigrid relaxation for $H(\text{curl})$ and $H(\text{div})$.

Complexities for some p -robust high-order solvers in three dimensions

		Year	Cell	Problem	Time	Storage
Pavarino & Widlund		1993		$H(\text{grad})$	p^9	p^6
Arnold, Falk & Winther		2000		all	p^9	p^6
► Beuchler & Pillwein		2007		$H(\text{grad})$	-	-
Schöberl et al.		2008		$H(\text{grad})$	p^9	p^6
Farrell & Brubeck		2024		all	p^4	p^3
This work		2025		all	p^6	p^4

Beuchler & Pillwein

proposed a basis in which cell interior problems can be solved in $\mathcal{O}(p^6)$ time.

Complexities for some p -robust high-order solvers in three dimensions

		Year	Cell	Problem	Time	Storage
Pavarino & Widlund		1993		$H(\text{grad})$	p^9	p^6
Arnold, Falk & Winther		2000		all	p^9	p^6
Beuchler & Pillwein		2007		$H(\text{grad})$	-	-
► Schöberl et al.		2008		$H(\text{grad})$	p^9	p^6
Farrell & Brubeck		2024		all	p^4	p^3
This work		2025		all	p^6	p^4

Schöberl, Melenk, Pechstein, & Zaglmayr

proved the Pavarino–Widlund result (star patches are p -robust) on simplices.

Complexities for some p -robust high-order solvers in three dimensions

		Year	Cell	Problem	Time	Storage
Pavarino & Widlund		1993		$H(\text{grad})$	p^9	p^6
Arnold, Falk & Winther		2000		all	p^9	p^6
Beuchler & Pillwein		2007		$H(\text{grad})$	-	-
Schöberl et al.		2008		$H(\text{grad})$	p^9	p^6
► Farrell & Brubeck		2024		all	p^4	p^3
This work		2025		all	p^6	p^4

Farrell & Brubeck

propose tensor product bases with the same complexities as matrix-vector products.

Complexities for some p -robust high-order solvers in three dimensions

		Year	Cell	Problem	Time	Storage
Pavarino & Widlund	1993		$H(\text{grad})$	p^9	p^6	
Arnold, Falk & Winther	2000		all	p^9	p^6	
Beuchler & Pillwein	2007		$H(\text{grad})$	-	-	
Schöberl et al.	2008		$H(\text{grad})$	p^9	p^6	
Farrell & Brubeck	2024		all	p^4	p^3	
► This work	2025		all	p^6	p^4	

This work

much harder than hexes: suboptimal complexity, but much better than a naïve approach!

Split our patch problem into interior (\mathcal{I}) and interface (Γ) unknowns.

Split our patch problem into interior (\mathcal{I}) and interface (Γ) unknowns.

We will design new bases for the standard finite element spaces with the property that

Split our patch problem into interior (\mathcal{I}) and interface (Γ) unknowns.

We will design new bases for the standard finite element spaces with the property that

Interior decoupling on a cell

$$\text{cond} \left(\begin{bmatrix} A_{\Gamma\Gamma} & 0 \\ 0 & \text{diag}(A_{\mathcal{I}\mathcal{I}}) \end{bmatrix}^{-1} \begin{bmatrix} A_{\Gamma\Gamma} & A_{\Gamma\mathcal{I}} \\ A_{\mathcal{I}\Gamma} & A_{\mathcal{I}\mathcal{I}} \end{bmatrix} \right) \leq C \max \left\{ 1, \frac{\beta}{\alpha} h^2 \right\},$$

where C is p -robust.

Split our patch problem into interior (\mathcal{I}) and interface (Γ) unknowns.

We will design new bases for the standard finite element spaces with the property that

Interior decoupling on a cell

$$\text{cond} \left(\begin{bmatrix} A_{\Gamma\Gamma} & 0 \\ 0 & \text{diag}(A_{\mathcal{I}\mathcal{I}}) \end{bmatrix}^{-1} \begin{bmatrix} A_{\Gamma\Gamma} & A_{\Gamma\mathcal{I}} \\ A_{\mathcal{I}\Gamma} & A_{\mathcal{I}\mathcal{I}} \end{bmatrix} \right) \leq C \max \left\{ 1, \frac{\beta}{\alpha} h^2 \right\},$$

where C is p -robust.

In other words,

- ▶ the interior block is weakly coupled (the diagonal is good enough);

Split our patch problem into interior (\mathcal{I}) and interface (Γ) unknowns.

We will design new bases for the standard finite element spaces with the property that

Interior decoupling on a cell

$$\text{cond} \left(\begin{bmatrix} A_{\Gamma\Gamma} & 0 \\ 0 & \text{diag}(A_{\mathcal{I}\mathcal{I}}) \end{bmatrix}^{-1} \begin{bmatrix} A_{\Gamma\Gamma} & A_{\Gamma\mathcal{I}} \\ A_{\mathcal{I}\Gamma} & A_{\mathcal{I}\mathcal{I}} \end{bmatrix} \right) \leq C \max \left\{ 1, \frac{\beta}{\alpha} h^2 \right\},$$

where C is p -robust.

In other words,

- ▶ the interior block is weakly coupled (the diagonal is good enough);
- ▶ the interface block is equivalent to the Schur complement (static condensation).

Split our patch problem into interior (\mathcal{I}) and interface (Γ) unknowns.

We will design new bases for the standard finite element spaces with the property that

Interior decoupling on a cell

$$\text{cond} \left(\begin{bmatrix} A_{\Gamma\Gamma} & 0 \\ 0 & \text{diag}(A_{\mathcal{I}\mathcal{I}}) \end{bmatrix}^{-1} \begin{bmatrix} A_{\Gamma\Gamma} & A_{\Gamma\mathcal{I}} \\ A_{\mathcal{I}\Gamma} & A_{\mathcal{I}\mathcal{I}} \end{bmatrix} \right) \leq C \max \left\{ 1, \frac{\beta}{\alpha} h^2 \right\},$$

where C is p -robust.


In other words,

- ▶ the interior block is weakly coupled (the diagonal is good enough);
- ▶ the interface block is equivalent to the Schur complement (static condensation).


So we pay the dense linear algebra costs for a problem *one dimension lower*, on Γ .

Sparsity patterns for $(du, dv)_K$ on reference cell, $p = 10$


Lagrange, $d = \text{grad}$



Nédélec, $d = \text{curl}$




Raviart–Thomas, $d = \text{div}$



Section 3

First numerical example

A p -robust solver for the Riesz maps:



CG iteration counts (rel. tol. = 10^{-8}).

d	p	$H(\text{grad})$	$H(\text{curl})$	$H(\text{div})$
2	4	20		24
	7	20		23
	10	20		23
	14	20		23
3	4	23	39	19
	7	23	42	19
	10	23	42	19


Additive space decompositions on the interface Schur complement

- ▶ Vertex-star + Lowest-order (à la Schöberl et al.)
- ▶ Edge-star + Vertex-star on grad $H(\text{grad})$ + Lowest-order (à la Hiptmair)
- ▶ Edge-star + Lowest-order (à la Arnold–Falk–Winther)

Section 4


Construction of the basis

We're going to build *different bases* for the *usual spaces*



and their first-kind brothers.

We're going to build *different bases* for the *usual spaces*



and their first-kind brothers.

We do this by *choosing different degrees of freedom*.

Definition (Finite element)

A finite element is a triple $(K, \mathcal{V}, \mathcal{L})$ where

- The cell K is a bounded, closed subset of \mathbb{R}^d with nonempty connected interior and piecewise smooth boundary;



Philippe Ciarlet

Definition (Finite element)

A finite element is a triple $(K, \mathcal{V}, \mathcal{L})$ where

- ▶ The cell K is a bounded, closed subset of \mathbb{R}^d with nonempty connected interior and piecewise smooth boundary;
- ▶ The space $\mathcal{V} = \mathcal{V}(K)$ is a finite dimensional function space on K of dimension m ;



Philippe Ciarlet

Definition (Finite element)

A finite element is a triple $(K, \mathcal{V}, \mathcal{L})$ where

- ▶ The cell K is a bounded, closed subset of \mathbb{R}^d with nonempty connected interior and piecewise smooth boundary;
- ▶ The space $\mathcal{V} = \mathcal{V}(K)$ is a finite dimensional function space on K of dimension m ;
- ▶ The set of degrees of freedom $\mathcal{L} = \{\ell_1, \dots, \ell_m\}$ is a basis for \mathcal{V}^* , the dual space of \mathcal{V} .



Philippe Ciarlet

Definition (Finite element)

A finite element is a triple $(K, \mathcal{V}, \mathcal{L})$ where

- ▶ The cell K is a bounded, closed subset of \mathbb{R}^d with nonempty connected interior and piecewise smooth boundary;
- ▶ The space $\mathcal{V} = \mathcal{V}(K)$ is a finite dimensional function space on K of dimension m ;
- ▶ The set of degrees of freedom $\mathcal{L} = \{\ell_1, \dots, \ell_m\}$ is a basis for \mathcal{V}^* , the dual space of \mathcal{V} .



Philippe Ciarlet

Intuition

The degrees of freedom are what you store to uniquely specify a function on the cell.

Definition (Finite element)

A finite element is a triple $(K, \mathcal{V}, \mathcal{L})$ where


- ▶ The cell K is a bounded, closed subset of \mathbb{R}^d with nonempty connected interior and piecewise smooth boundary;
- ▶ The space $\mathcal{V} = \mathcal{V}(K)$ is a finite dimensional function space on K of dimension m ;
- ▶ The set of degrees of freedom $\mathcal{L} = \{\ell_1, \dots, \ell_m\}$ is a basis for \mathcal{V}^* , the dual space of \mathcal{V} .




Philippe Ciarlet

Basis functions are biorthogonal

The basis functions $\{\phi_j\}$ are constructed to satisfy $\ell_i(\phi_j) = \delta_{ij}$.



The quadratic Lagrange finite element CG_2 in two dimensions.



The quadratic Lagrange finite element CG_2 in two dimensions.

Example (CG_2 in 2D)

$$K = \Delta, \mathcal{V} = \text{span}(1, x, y, x^2, y^2, xy), \mathcal{L} = \{\ell_1, \dots, \ell_6\}.$$

Each ℓ_i evaluates the function at a vertex or edge midpoint.

Subsection 1

Degrees of freedom for $H(\text{grad})$

Usually CG_p just has point evaluations as degrees of freedom.



Leszek Demkowicz



Peter Monk 20 / 38

Usually CG_p just has point evaluations as degrees of freedom.

Demkowicz et al. proposed different degrees of freedom for the whole de Rham complex that induce discrete commuting de Rham complexes on each entity. Each entity can have a different polynomial degree.



Leszek Demkowicz



Usually CG_p just has point evaluations as degrees of freedom.



Demkowicz et al. proposed different degrees of freedom for the whole de Rham complex that induce discrete commuting de Rham complexes on each entity. Each entity can have a different polynomial degree.

The degrees of freedom are pointwise evaluation at vertices

$$\ell_V : v \mapsto v(V) \quad \forall V \in \Delta_0(\triangle),$$

Leszek Demkowicz



Usually CG_p just has point evaluations as degrees of freedom.



Demkowicz et al. proposed different degrees of freedom for the whole de Rham complex that induce discrete commuting de Rham complexes on each entity. Each entity can have a different polynomial degree.

The degrees of freedom are pointwise evaluation at vertices

$$\ell_V : v \mapsto v(V) \quad \forall V \in \Delta_0(\triangle),$$

Leszek Demkowicz

and moments on each higher-dimensional entity

$$\ell_S : v \mapsto (\text{grad}_S q, \text{grad}_S v)_S \quad \forall S \in \bigcup_{k=1}^d \Delta_k(\triangle) \quad \forall q \in \mathbb{P}_{p,0}(S),$$



where $\mathbb{P}_{p,0}(S)$ is the *bubble space* on S

$$\mathbb{P}_{p,0}(S) := \{v \in \mathbb{P}_p(S) : v = 0 \text{ on } \partial S\}.$$

This construction works for any choice of basis for the bubble space.

This construction works for any choice of basis for the bubble space.

Our idea

Choose the basis for the bubble space by solving small eigenproblems.

This construction works for any choice of basis for the bubble space.

Our idea

Choose the basis for the bubble space by solving small eigenproblems.

We choose the basis for $\mathbb{P}_{p,0}(S)$ such that

$$(\text{grad}_S \phi_j^S, \text{grad}_S \phi_i^S)_S = \delta_{ij}, \quad (\phi_j^S, \phi_i^S)_S = \lambda_j \delta_{ij}.$$

This construction works for any choice of basis for the bubble space.

Our idea

Choose the basis for the bubble space by solving small eigenproblems.

We choose the basis for $\mathbb{P}_{p,0}(S)$ such that

$$(\text{grad}_S \phi_j^S, \text{grad}_S \phi_i^S)_S = \delta_{ij}, \quad (\phi_j^S, \phi_i^S)_S = \lambda_j \delta_{ij}.$$

In other words, the basis for the bubble space should be orthogonal in both inner products.

This construction works for any choice of basis for the bubble space.

Our idea

Choose the basis for the bubble space by solving small eigenproblems.

We choose the basis for $\mathbb{P}_{p,0}(S)$ such that


$$(\text{grad}_S \phi_j^S, \text{grad}_S \phi_i^S)_S = \delta_{ij}, \quad (\phi_j^S, \phi_i^S)_S = \lambda_j \delta_{ij}.$$

In other words, the basis for the bubble space should be orthogonal in both inner products.


This requires one small dense offline eigensolve for each type of entity S ($\mathcal{O}(p^9)$ offline cost).

our basis



stiffness



mass




hierarchical




$H(\text{grad}), d = 2, p = 10$



our basis



mass



hierarchical



$H(\text{grad}), d = 3, p = 10$

Subsection 2

Degrees of freedom for $H(\text{curl})$ and $H(\text{div})$

For $H(\text{curl})$, our eigenidea with Demkowicz et al. is to employ tangential moments on edges

$$\ell_E : v \mapsto (q, v \cdot t)_E \quad \forall E \in \Delta_1(\triangle) \quad \forall q \in \mathbb{P}_p(E),$$

For $H(\text{curl})$, our eigenidea with Demkowicz et al. is to employ tangential moments on edges

$$\ell_E : v \mapsto (q, v \cdot t)_E \quad \forall E \in \Delta_1(\mathbb{D}) \quad \forall q \in \mathbb{P}_p(E),$$

and two kinds of moments on entities of higher dimension

$$\ell_{S,1} : v \mapsto (\text{grad}_S \phi_j^S, v)_S \quad \forall S \in \bigcup_{k=2}^d \Delta_k(\mathbb{D}) \quad \forall \phi_j^S,$$

where $\{\phi_j^S\}$ are the eigenfunctions from $H(\text{grad})$, and

For $H(\text{curl})$, our eigenidea with Demkowicz et al. is to employ tangential moments on edges

$$\ell_E : v \mapsto (q, v \cdot t)_E \quad \forall E \in \Delta_1(\mathbb{D}) \quad \forall q \in \mathbb{P}_p(E),$$

and two kinds of moments on entities of higher dimension

$$\ell_{S,1} : v \mapsto (\text{grad}_S \phi_j^S, v)_S \quad \forall S \in \bigcup_{k=2}^d \Delta_k(\mathbb{D}) \quad \forall \phi_j^S,$$

where $\{\phi_j^S\}$ are the eigenfunctions from $H(\text{grad})$, and

$$\ell_{S,2} : v \mapsto (\text{curl}_S \Phi_j^S, \text{curl } v)_S \quad \forall S \in \bigcup_{k=2}^d \Delta_k(\mathbb{D}) \quad \forall \Phi_j^S,$$

where $\{\text{curl}_S \Phi_j^S\}$ is a basis for $\text{curl}_S \mathbb{X}$, $\mathbb{X} = [\mathbb{P}_p(S)]^d \cap H_0(\text{curl}, S)$, such that

$$(\text{curl}_S \Phi_j^S, \text{curl}_S \Phi_i^S)_S = \delta_{ij}, \quad (\Phi_j^S, \Phi_i^S)_S = \lambda_j \delta_{ij}, \quad \Phi_j^S \times \mathbf{n} = 0 \text{ on } \partial S.$$

For $H(\text{div})$, our eigenidea with Demkowicz et al. is to employ normal moments on faces

$$\ell_F : v \mapsto (q, v \cdot n)_F \quad \forall F \in \Delta_2(\triangle) \quad \forall q \in \mathbb{P}_p(F),$$

For $H(\text{div})$, our eigenidea with Demkowicz et al. is to employ normal moments on faces

$$\ell_F : v \mapsto (q, v \cdot n)_F \quad \forall F \in \Delta_2(\triangle) \quad \forall q \in \mathbb{P}_p(F),$$

and two kinds of moments on the only entity of higher dimension, the interior K

$$\ell_{K,1} : v \mapsto (\text{curl } \Phi_j^K, v) \quad \forall \Phi_j^K$$

where $\{\Phi_j^K\}$ are the eigenfunctions from $H(\text{curl})$, and

For $H(\text{div})$, our eigenidea with Demkowicz et al. is to employ normal moments on faces

$$\ell_F : v \mapsto (q, v \cdot n)_F \quad \forall F \in \Delta_2(\triangle) \quad \forall q \in \mathbb{P}_p(F),$$

and two kinds of moments on the only entity of higher dimension, the interior K





$$\ell_{K,1} : v \mapsto (\text{curl } \Phi_j^K, v) \quad \forall \Phi_j^K$$

where $\{\Phi_j^K\}$ are the eigenfunctions from $H(\text{curl})$, and

$$\ell_{K,2} : v \mapsto (\text{div } \Psi_j^K, \text{div } v) \quad \forall \Psi_j^K,$$

where $\{\text{div } \Psi_j^K\}$ is a basis for $\text{div } \mathbb{Y}$, $\mathbb{Y} = [\mathbb{P}_p(K)]^d \cap H_0(\text{div}, K)$, such that

$$(\text{div } \Psi_j^K, \text{div } \Psi_i^K)_K = \delta_{ij}, \quad (\Psi_j^K, \Psi_i^K)_K = \lambda_j \delta_{ij}, \quad \Psi_j^K \cdot \mathbf{n} = 0 \text{ on } \partial K.$$

space	stiffness	mass
$H(\text{curl})$	 A sparse matrix plot for the $H(\text{curl})$ stiffness matrix. It has a total size of 780x780, with non-zero elements concentrated in several horizontal bands. The first band (rows 1-420) contains a 4x4 block diagonal pattern. The second band (rows 421-780) is mostly zero except for a few diagonal entries. The matrix is symmetric about the main diagonal. nz = 31366	 A sparse matrix plot for the $H(\text{curl})$ mass matrix. It has a total size of 780x780, with non-zero elements forming a banded structure. The main diagonal is solid blue. There are several off-diagonal blocks of varying sizes, primarily along the second band. The matrix is symmetric. nz = 334940
$H(\text{div})$	 A sparse matrix plot for the $H(\text{div})$ stiffness matrix. It has a total size of 715x715, showing a very sparse structure with a single dominant diagonal band. A few small off-diagonal blocks are visible near the top. nz = 235	 A sparse matrix plot for the $H(\text{div})$ mass matrix. It has a total size of 715x715, featuring a large solid blue block on the main diagonal. A few smaller off-diagonal blocks are located near the top and bottom of the matrix. nz = 132599

$$d = 3, p = 10$$

Section 5

Analysis

Lemma (Unisolvence)

The degrees of freedom are unisolvant.

Lemma (Unisolvence)

The degrees of freedom are unisolvent.

Theorem (Commuting diagram (Demkowicz et al., 2000))

The spaces form a commuting subcomplex of the de Rham complex.

Lemma (Unisolvence)

The degrees of freedom are unisolvent.

Theorem (Commuting diagram (Demkowicz et al., 2000))

The spaces form a commuting subcomplex of the de Rham complex.

Theorem (Trace similarity)

The degrees of freedom are trace similar: e.g. the degrees of freedom built for $d = 2$ are present on each face of the reference tetrahedron in $d = 3$.

Lemma (Unisolvence)

The degrees of freedom are unisolvent.

Theorem (Commuting diagram (Demkowicz et al., 2000))

The spaces form a commuting subcomplex of the de Rham complex.

Theorem (Trace similarity)

The degrees of freedom are trace similar: e.g. the degrees of freedom built for $d = 2$ are present on each face of the reference tetrahedron in $d = 3$.

Theorem (Kernel and range)

The kernel and range of \mathbf{d} have a low-order part from the Whitney forms, and a high-order part we characterise precisely (as certain basis functions).

Interior decoupling on a physical cell

$$\text{cond} \left(\begin{bmatrix} \text{diag}(A_{\mathcal{I}\mathcal{I}}) & 0 \\ 0 & A_{\Gamma\Gamma} \end{bmatrix}^{-1} \begin{bmatrix} A_{\mathcal{I}\mathcal{I}} & A_{\mathcal{I}\Gamma} \\ A_{\Gamma\mathcal{I}} & A_{\Gamma\Gamma} \end{bmatrix} \right) \leq C \max \left\{ 1, \frac{\beta}{\alpha} h^2 \right\},$$

where C depends on shape regularity, d , k , and $\text{diam}(\Omega)$.

Interior decoupling on a physical cell

$$\text{cond} \left(\begin{bmatrix} \text{diag}(A_{\mathcal{I}\mathcal{I}}) & 0 \\ 0 & A_{\Gamma\Gamma} \end{bmatrix}^{-1} \begin{bmatrix} A_{\mathcal{I}\mathcal{I}} & A_{\mathcal{I}\Gamma} \\ A_{\Gamma\mathcal{I}} & A_{\Gamma\Gamma} \end{bmatrix} \right) \leq C \max \left\{ 1, \frac{\beta}{\alpha} h^2 \right\},$$

where C depends on shape regularity, d , k , and $\text{diam}(\Omega)$.

How do we use this to efficiently implement patch solves?

Theorem (interior-interface splitting)

Given an energy-stable space decomposition

$$X^k = \sum_{m=1}^M X_m^k$$

the *interior-interface split decomposition*

$$X^k = \sum_{m=1}^M X_m^k \cap X_\Gamma^k + X_\mathcal{I}^k$$

is also energy-stable provided $\beta h^2/\alpha \leq 1$.

The ratio of stability constants is independent of h , p , α , and β .

Theorem (interior-interface splitting)

Given an energy-stable space decomposition

$$X^k = \sum_{m=1}^M X_m^k$$

the *interior-interface split decomposition*

$$X^k = \sum_{m=1}^M X_m^k \cap X_\Gamma^k + X_\mathcal{I}^k$$

is also energy-stable provided $\beta h^2/\alpha \leq 1$.

The ratio of stability constants is independent of h , p , α , and β .

We can take a robust space decomposition and robustly split it for fast patch solves!

Section 6

Application: the Hodge Laplacian

We employ our Riesz map solvers in a new application of operator preconditioning.

We employ our Riesz map solvers in a new application of operator preconditioning.

Operator preconditioning

Let \mathbb{Z} be a Hilbert space. Consider the problem: find $z \in \mathbb{Z}$ such that

$$a(z, w) = L(w) \quad \forall w \in \mathbb{Z},$$

where a is a symmetric bilinear form and $L \in \mathbb{Z}^*$.

We employ our Riesz map solvers in a new application of operator preconditioning.

Operator preconditioning

Let \mathbb{Z} be a Hilbert space. Consider the problem: find $z \in \mathbb{Z}$ such that

$$a(z, w) = L(w) \quad \forall w \in \mathbb{Z},$$

where a is a symmetric bilinear form and $L \in \mathbb{Z}^*$.

We assume this is well-posed, with continuity and inf-sup constants C, β , measured with $\|\cdot\|_{\mathbb{Z}}$.

We employ our Riesz map solvers in a new application of operator preconditioning.

Operator preconditioning

Let \mathbb{Z} be a Hilbert space. Consider the problem: find $z \in \mathbb{Z}$ such that

$$a(z, w) = L(w) \quad \forall w \in \mathbb{Z},$$

where a is a symmetric bilinear form and $L \in \mathbb{Z}^*$.

We assume this is well-posed, with continuity and inf-sup constants C, β , measured with $\|\cdot\|_{\mathbb{Z}}$.

Using Z , the Riesz map for \mathbb{Z} , as a preconditioner for the operator A associated with a yields

$$\text{cond}(Z^{-1}A) \leq C\beta^{-1}.$$

We consider the *Hodge Laplacian*, a fundamental PDE associated with a complex:
find $(\sigma, u) \in X^{k-1} \times X^k$ such that

$$\begin{aligned} -(\sigma, \tau) + (u, d^{k-1} \tau) &= 0 & \forall \tau \in X^{k-1}, \\ (d^{k-1} \sigma, v) + (d^k u, d^k v) &= F(v) & \forall v \in X^k. \end{aligned}$$

We consider the *Hodge Laplacian*, a fundamental PDE associated with a complex:
find $(\sigma, u) \in X^{k-1} \times X^k$ such that

$$\begin{aligned} -(\sigma, \tau) + (u, d^{k-1} \tau) &= 0 & \forall \tau \in X^{k-1}, \\ (d^{k-1} \sigma, v) + (d^k u, d^k v) &= F(v) & \forall v \in X^k. \end{aligned}$$

This is a mixed formulation for the problems

$$-\operatorname{div} \operatorname{grad} u = f, \quad \operatorname{curl} \operatorname{curl} \mathbf{u} - \operatorname{grad} \operatorname{div} \mathbf{u} = \mathbf{f}.$$

We consider the *Hodge Laplacian*, a fundamental PDE associated with a complex:
 find $(\sigma, u) \in X^{k-1} \times X^k$ such that

$$\begin{aligned} -(\sigma, \tau) + (u, d^{k-1} \tau) &= 0 & \forall \tau \in X^{k-1}, \\ (d^{k-1} \sigma, v) + (d^k u, d^k v) &= F(v) & \forall v \in X^k. \end{aligned}$$

This is a mixed formulation for the problems

$$-\operatorname{div} \operatorname{grad} u = f, \quad \operatorname{curl} \operatorname{curl} \mathbf{u} - \operatorname{grad} \operatorname{div} \mathbf{u} = \mathbf{f}.$$

These are well-posed with

$$\mathbb{Z} = \begin{cases} H(\operatorname{grad}) \times H(\operatorname{curl}) & k = 1, \\ H(\operatorname{curl}) \times H(\operatorname{div}) & k = 2, \\ H(\operatorname{div}) \times L^2 & k = 3. \end{cases}$$

A standard application of the operator preconditioning framework yields that the Riesz map

$$\langle (\sigma, u), (\tau, v) \rangle_{\mathbb{Z}} = (\sigma, \tau) + (\mathrm{d}^{k-1} \sigma, \mathrm{d}^{k-1} \tau) + (u, v) + (\mathrm{d}^k u, \mathrm{d}^k v)$$

is a good preconditioner with

$$\mathrm{cond}(Z^{-1} A) \leq C$$

for some constant C (provably independent of h , observed independent of p).

A standard application of the operator preconditioning framework yields that the Riesz map

$$\langle (\sigma, u), (\tau, v) \rangle_{\mathbb{Z}} = (\sigma, \tau) + (\mathrm{d}^{k-1} \sigma, \mathrm{d}^{k-1} \tau) + (u, v) + (\mathrm{d}^k u, \mathrm{d}^k v)$$

is a good preconditioner with

$$\mathrm{cond}(Z^{-1} A) \leq C$$

for some constant C (provably independent of h , observed independent of p).

But we can do a bit better, for free!

A standard application of the operator preconditioning framework yields that the Riesz map

$$\langle(\sigma, u), (\tau, v)\rangle_{\mathbb{Z}} = (\sigma, \tau) + (\mathrm{d}^{k-1} \sigma, \mathrm{d}^{k-1} \tau) + (u, v) + (\mathrm{d}^k u, \mathrm{d}^k v)$$

is a good preconditioner with

$$\text{cond}(Z^{-1} A) \leq C$$

for some constant C (provably independent of h , observed independent of p).

But we can do a bit better, for free!

Theorem (weighted inner product)

For $\gamma > 0$, let \mathbb{Z}_γ be the Hilbert space with weighted inner product

$$\langle(\sigma, u), (\tau, v)\rangle_{\mathbb{Z}_\gamma} = (\sigma, \tau) + \gamma(\mathrm{d}^{k-1} \sigma, \mathrm{d}^{k-1} \tau) + \gamma^{-1}(u, v) + (\mathrm{d}^k u, \mathrm{d}^k v).$$

A standard application of the operator preconditioning framework yields that the Riesz map

$$\langle(\sigma, u), (\tau, v)\rangle_{\mathbb{Z}} = (\sigma, \tau) + (\mathrm{d}^{k-1} \sigma, \mathrm{d}^{k-1} \tau) + (u, v) + (\mathrm{d}^k u, \mathrm{d}^k v)$$

is a good preconditioner with

$$\text{cond}(Z^{-1} A) \leq C$$

for some constant C (provably independent of h , observed independent of p).

But we can do a bit better, for free!

Theorem (weighted inner product)

For $\gamma > 0$, let \mathbb{Z}_γ be the Hilbert space with weighted inner product

$$\langle(\sigma, u), (\tau, v)\rangle_{\mathbb{Z}_\gamma} = (\sigma, \tau) + \gamma(\mathrm{d}^{k-1} \sigma, \mathrm{d}^{k-1} \tau) + \gamma^{-1}(u, v) + (\mathrm{d}^k u, \mathrm{d}^k v).$$

Then all eigenvalues of $Z_\gamma^{-1} A$ converge to ± 1 as $\gamma \rightarrow \infty$ like $\mathcal{O}(\gamma^{-1})$.

A standard application of the operator preconditioning framework yields that the Riesz map

$$\langle(\sigma, u), (\tau, v)\rangle_{\mathbb{Z}} = (\sigma, \tau) + (\mathrm{d}^{k-1} \sigma, \mathrm{d}^{k-1} \tau) + (u, v) + (\mathrm{d}^k u, \mathrm{d}^k v)$$

is a good preconditioner with

$$\text{cond}(Z^{-1} A) \leq C$$

for some constant C (provably independent of h , observed independent of p).

But we can do a bit better, for free!

Theorem (weighted inner product)


For $\gamma > 0$, let \mathbb{Z}_γ be the Hilbert space with weighted inner product

$$\langle(\sigma, u), (\tau, v)\rangle_{\mathbb{Z}_\gamma} = (\sigma, \tau) + \gamma(\mathrm{d}^{k-1} \sigma, \mathrm{d}^{k-1} \tau) + \gamma^{-1}(u, v) + (\mathrm{d}^k u, \mathrm{d}^k v).$$

Then all eigenvalues of $Z_\gamma^{-1} A$ converge to ± 1 as $\gamma \rightarrow \infty$ like $\mathcal{O}(\gamma^{-1})$.

For both Riesz maps, $\gamma \gg 1$ takes us in to the regime our solvers work best.

We consider the Fichera cube with right hand side $f = 1$ or $\mathbf{f} = (1, 1, 1)^\top$.



The reentrant corner causes singularities and low regularity in the solution.

MINRES iteration counts: interior-interface split (not split)

p	$\text{CG}_p \times \text{Ned}_p^1$		$\text{RT}_p \times \text{DG}_{p-1}$	
	$\gamma = 1$	$\gamma = 10^3$	$\gamma = 1$	$\gamma = 10^3$
1	6 (6)	3 (3)	5 (5)	3 (3)
2	41 (41)	37 (37)	3 (25)	14 (16)
3	46 (48)	40 (41)	26 (28)	17 (17)
4	52 (50)	44 (44)	29 (28)	19 (18)
5	59 (51)	51 (44)	31 (29)	19 (19)
6	68 (52)	57 (44)	31 (29)	20 (19)

Observations:

- ▶ $\gamma = 10^3$ gives perfect control of the Schur complement for $p = 1$.

MINRES iteration counts: interior-interface split (not split)

p	$\text{CG}_p \times \text{Ned}_p^1$		$\text{RT}_p \times \text{DG}_{p-1}$	
	$\gamma = 1$	$\gamma = 10^3$	$\gamma = 1$	$\gamma = 10^3$
1	6 (6)	3 (3)	5 (5)	3 (3)
2	41 (41)	37 (37)	3 (25)	14 (16)
3	46 (48)	40 (41)	26 (28)	17 (17)
4	52 (50)	44 (44)	29 (28)	19 (18)
5	59 (51)	51 (44)	31 (29)	19 (19)
6	68 (52)	57 (44)	31 (29)	20 (19)

Observations:

- ▶ $\gamma = 10^3$ gives perfect control of the Schur complement for $p = 1$.
- ▶ Moving from $\gamma = 1$ to $\gamma = 10^3$ saves ~ 10 iterations, for free.

MINRES iteration counts: interior-interface split (not split)

p	$\text{CG}_p \times \text{Ned}_p^1$		$\text{RT}_p \times \text{DG}_{p-1}$	
	$\gamma = 1$	$\gamma = 10^3$	$\gamma = 1$	$\gamma = 10^3$
1	6 (6)	3 (3)	5 (5)	3 (3)
2	41 (41)	37 (37)	3 (25)	14 (16)
3	46 (48)	40 (41)	26 (28)	17 (17)
4	52 (50)	44 (44)	29 (28)	19 (18)
5	59 (51)	51 (44)	31 (29)	19 (19)
6	68 (52)	57 (44)	31 (29)	20 (19)

Observations:

- ▶ $\gamma = 10^3$ gives perfect control of the Schur complement for $p = 1$.
- ▶ Moving from $\gamma = 1$ to $\gamma = 10^3$ saves ~ 10 iterations, for free.
- ▶ The interior-interface split is *much* cheaper to compute than the unsplit solver.

Section 7

Conclusion

Riesz maps

We have a solver with $\mathcal{O}(p^6)$ flops and $\mathcal{O}(p^4)$ storage for the Riesz maps on tetrahedra.

Riesz maps

We have a solver with $\mathcal{O}(p^6)$ flops and $\mathcal{O}(p^4)$ storage for the Riesz maps on tetrahedra.

Operator preconditioning

These can be used anywhere operator preconditioning calls for Riesz map solvers.

Riesz maps

We have a solver with $\mathcal{O}(p^6)$ flops and $\mathcal{O}(p^4)$ storage for the Riesz maps on tetrahedra.

Operator preconditioning

These can be used anywhere operator preconditioning calls for Riesz map solvers.

Take-home message

For efficiency, design formulation + basis + solver together!