

PTCB

Study Guide

Simulation Test

© Copyright 2026 by MED PREP – All rights reserved.

This publication is designed to support individuals preparing for the **Pharmacy Technician Certification Exam (PTCE)**. The content is intended solely for educational and informational purposes and should be used as a supplementary study resource alongside official materials, accredited training programs, and institutional instruction.

Every effort has been made to ensure that the information presented is accurate, relevant, and aligned with commonly tested concepts. However, MED PREP does not guarantee that all content reflects the most current exam blueprint or that it fully covers all topics included in the PTCE. Certification requirements and exam content are subject to change, and candidates are responsible for verifying current guidelines with the **Pharmacy Technician Certification Board (PTCB)**.

All practice questions, scenarios, and explanatory materials contained in this book have been independently developed for study purposes. They do not represent actual PTCE exam questions and are not derived from or affiliated with any official examination content. This publication is not endorsed by, affiliated with, or sponsored by the **Pharmacy Technician Certification Board (PTCB)** or any related organization.

The information provided in this book is not intended to serve as medical advice, clinical instruction, or a substitute for professional training. Readers are responsible for applying the material appropriately within their scope of practice and in accordance with applicable laws, regulations, and workplace policies.

MED PREP and the author assume no responsibility or liability for any outcomes related to exam performance, certification results, employment decisions, or any actions taken based on the use of this material.

No part of this publication may be reproduced, stored, or transmitted in any form or by any means without prior written permission from the publisher, except as permitted by applicable copyright laws.

All trademarks, certification marks, and brand names referenced in this publication are the property of their respective owners. Their inclusion is for identification and educational purposes only and does not imply endorsement or affiliation.

All rights reserved.

ONLINE PRACTICE & PRINTABLE SIMULATION TESTS

This PDF includes **34 PTCB simulation tests in printable PDF format** (which are added to the 2 tests present in the book), designed to help you **build skills, reinforce concepts, and gain confidence step by step**.

In addition, **4 carefully selected simulation tests are also available online** in a structured, timed format.

This is intentional.

Online simulations are most effective **only when you are already prepared**, not while you're still learning. They are designed to reflect **PTCB-style questions and timing**, but they are **not an official exam interface**.

That's why the online tests are reserved for the **final stage of preparation**, when you need to:

- Practice under time pressure
- Test your readiness in one sitting
- Experience exam-like focus without interruptions
- Confirm that you're ready for test day

The remaining simulation tests are provided **exclusively in PDF format** so you can:

- Pause and reflect
- Review explanations carefully
- Learn from mistakes without pressure

In short:

You train with the PDFs.

You validate readiness with the online simulations.

This approach ensures that online practice is **meaningful, realistic, and confidence-building** — not overwhelming or premature.



NEED HELP OR HAVE A QUESTION?

If you experience any issues accessing the online practice tests, have a question about the material, or need assistance, you can contact us directly:

 **Email:** dunya84@gmail.com

We're happy to help and typically respond within 24–48 hours.

PTCB Sim Test 01

1) A patient requests the “white water pill” they always get, but today’s label prints for furosemide 20 mg and the previous fill was hydrochlorothiazide 25 mg. Which is the best technician action before filling?

- a) Fill furosemide because both are diuretics
- b) Change to hydrochlorothiazide because it matches the prior fill
- c) Stop and compare the new prescription details to the profile; ensure the prescriber, drug, and directions match the new order before proceeding
- d) Ask the patient which one “worked better” and dispense that

2) A prescription reads: “amoxicillin 400 mg/5 mL: take 6 mL PO BID x 10 days.” How many mL should be dispensed?

- a) 100 mL
- b) 120 mL
- c) 150 mL
- d) 200 mL

3) A pharmacy receives an e-prescription for “Xarelto 20 mg daily.” Which generic should be selected during entry?

- a) Dabigatran
- b) Rivaroxaban
- c) Apixaban
- d) Edoxaban

4) During checkout, a customer wants to buy pseudoephedrine. The register prompts for ID. What is required at the point of sale under the Combat Methamphetamine Epidemic Act?

- a) Record the sale with product/quantity, verify ID, and obtain the purchaser’s signature (electronic or written)
- b) Require a prescription and store the product in the C-II safe
- c) Allow sale only if the patient is already in the pharmacy database
- d) Limit to a 3-day supply for all purchasers regardless of product strength

5) While processing a new prescription, the technician sees “allergy: penicillin— anaphylaxis.” The e-prescription is for cephalexin. What should the technician do?

- a) Enter and fill because cephalexin is not penicillin
- b) Delete the allergy because it is non-specific
- c) Add an auxiliary label “take with food” and proceed
- d) Hold the order for pharmacist review due to allergy severity and cross-reactivity concern

6) A prescription reads: “levothyroxine 0.125 mg PO daily.” Which strength should be selected?

- a) 125 mcg
- b) 12.5 mcg
- c) 1.25 mg
- d) 0.0125 mg

7) A technician is billing insulin glargine pens: "Inject 24 units SC nightly. Dispense 1 box." A box contains 5 pens of 3 mL each (100 units/mL). What days' supply should be submitted?

- a) 30
- b) 45
- c) 50
- d) 62

8) A prescriber's office calls and asks to "phone in hydrocodone/acetaminophen 5/325, #20." What is the appropriate technician response?

- a) Accept the verbal order and enter it for the pharmacist to verify later
- b) Transfer the call to the pharmacist because verbal controlled-substance prescriptions must be taken by a pharmacist
- c) Ask for the prescriber's DEA number and proceed if provided
- d) Tell the caller to leave a voicemail and the pharmacy will process it

9) A patient asks for "Lipitor." The profile shows atorvastatin is covered but brand is not. What is the technician's best response?

- a) Explain that atorvastatin is the generic of Lipitor and ask if they prefer generic today
- b) Dispense brand Lipitor and bill the generic NDC to get coverage
- c) Tell the patient generics are weaker and should be avoided
- d) Change the prescription to "DAW" so insurance will pay for brand

10) A technician is entering: "Take 1 tab PO qid prn pain." What does "qid" mean?

- a) Every other day
- b) At bedtime
- c) Four times daily
- d) Three times daily

11) The pharmacy receives a written prescription for alprazolam with the patient name, drug, strength, directions, quantity, and prescriber signature, but missing the prescriber address. What is the best next step?

- a) Process it; address is not required if DEA is on the pad
- b) Add the clinic address from memory and proceed
- c) Treat it as invalid and discard it
- d) Hold for pharmacist review to determine if the missing prescriber information can be obtained/verified before dispensing

12) A label prints for "hydromorphone 2 mg" but the stock bottle in hand is "hydroxyzine 25 mg." Which step best prevents a LASA dispensing error?

- a) Compare only the first 4 letters of the drug names to confirm similarity
- b) Verify the NDC on the bottle against the NDC selected on the label/screen before counting
- c) Choose based on tablet color described in the monograph
- d) Use whichever is stored in the controlled-substance section

13) A prescription reads: "prednisone 10 mg: take 4 tabs daily x 3 days, then 3 tabs daily x 3 days, then 2 tabs daily x 3 days, then 1 tab daily x 3 days." How many tablets should be

dispensed?

- a) 24
- b) 30
- c) 40
- d) 42

14) A pharmacy is transferring out a patient's noncontrolled prescription to another pharmacy. Which information is typically necessary for the receiving pharmacy to create the new record?

- a) Patient's insurance BIN/PCN
- b) Prescriber's DEA number
- c) Original Rx date, last fill date, quantity/remaining refills, and directions
- d) Patient's diagnosis code

15) A patient wants to return an opened bottle of tablets for a refund. What is the most accurate statement for typical community pharmacy practice?

- a) It can be returned to stock if the cap is child-resistant
- b) It can be returned to stock if fewer than 10 tablets were used
- c) It must be destroyed immediately without documentation
- d) It generally cannot be restocked once dispensed; handle according to the pharmacy's refund/disposal process

16) A technician receives a claim rejection: "Days supply exceeds plan limit." Rx: albuterol HFA "2 puffs q4h prn." Inhaler contains 200 puffs. For billing when PRN frequency is present, days supply is commonly calculated using the maximum frequency. What days' supply is most appropriate?

- a) 8
- b) 12
- c) 16
- d) 30

17) A prescription is for "metoprolol succinate ER 25 mg daily." The technician selects metoprolol tartrate 25 mg. What should be done?

- a) Keep tartrate and change sig to BID
- b) Keep tartrate and label "ER" to match the order
- c) Dispense both tartrate and succinate to ensure coverage
- d) Correct the selection to metoprolol succinate ER and continue normal verification steps

18) A prescriber sends "Eliquis 5 mg BID." Which generic is correct?

- a) Apixaban
- b) Rivaroxaban
- c) Edoxaban
- d) Dabigatran

19) In a hospital, a technician notices an ADC pocket labeled "morphine 2 mg/mL" contains hydromorphone syringes. What is the best technician action?

- a) Remove the incorrect items, secure them per procedure, correct the pocket, and report the discrepancy through the medication safety process

- b) Leave the pocket as-is and tell the nurse to double-check before removal
- c) Swap the pocket label to match the contents to avoid delays
- d) Move the syringes to a nearby pocket that has space

20) A prescription reads: "Apply 1 patch transdermal q72h. Dispense #10 patches." What is the days' supply?

- a) 10
- b) 20
- c) 30
- d) 72

21) A technician is preparing to compound a nonsterile oral solution and cannot locate the Master Formulation Record. What should the technician do?

- a) Use a commonly available online recipe and document it later
- b) Ask another technician what was used last time and proceed
- c) Stop and obtain the required formulation documentation before compounding
- d) Make a half-size batch and adjust as needed

22) A patient presents a Schedule II prescription and requests it be transferred to a different pharmacy. What is the correct technician response?

- a) Transfers are allowed if the other pharmacy is in the same chain
- b) Transfers are allowed once if the prescription has not been filled
- c) Transfers are allowed if the patient signs a transfer consent form
- d) Schedule II prescriptions generally cannot be transferred; route to the pharmacist for appropriate handling options

23) A new prescription comes in for "Januvia 100 mg daily." Which generic should be selected?

- a) Saxagliptin
- b) Sitagliptin
- c) Linagliptin
- d) Alogliptin

24) The refrigerator temperature log shows 12°C overnight for vaccines labeled "store at 2°C to 8°C." What is the best immediate action?

- a) Continue using them because they were not frozen
- b) Place them in the freezer to rapidly correct temperature
- c) Quarantine the vaccines and escalate for cold-chain assessment before use
- d) Discard all vaccine inventory immediately

25) A technician is entering "0.5 mg" tablets. The prescription requires a total dose of 15 mg. How many tablets are needed?

- a) 15
- b) 20
- c) 25
- d) 30

26) A prescription is written: "nitroglycerin SL: dissolve 1 tab under tongue at onset of chest

pain; may repeat q5 min x 3." What does "SL" indicate?

- a) Sustained release
- b) Sublingual
- c) Subcutaneous
- d) Sterile liquid

27) The pharmacy receives a wholesaler invoice for Schedule II items. How long must federal law generally require controlled substance records be retained?

- a) 6 months
- b) 1 year
- c) 2 years
- d) 5 years

28) A technician is processing "Plavix 75 mg daily." Which generic is correct?

- a) Ticagrelor
- b) Prasugrel
- c) Cilostazol
- d) Clopidogrel

29) A patient's spouse arrives to pick up the patient's medications. What best supports HIPAA compliance at pickup?

- a) Refuse to release any prescriptions unless the patient is present
- b) Release after verifying appropriate pickup authorization per workflow while limiting information disclosed
- c) Provide the patient's full medication profile so the spouse can confirm each item
- d) Require the spouse to sign a HIPAA form at the register for each pickup

30) A prescription reads: "Instill 1 gtt OU qhs." What does this mean?

- a) Instill 1 drop in both eyes at bedtime
- b) Instill 1 drop in the right eye at bedtime
- c) Instill 1 drop in both ears at bedtime
- d) Instill 1 drop in the left eye every morning

31) A third-party claim rejects: "Non-matched NDC." What is the most likely cause?

- a) Quantity exceeds plan limit
- b) The prescriber wrote brand but generic was selected
- c) The NDC billed does not match the NDC of the product selected/dispensed
- d) The prescriber's NPI is missing from the e-prescription

32) A new prescription is received for "sertraline 50 mg daily," but the profile shows "paroxetine 20 mg daily" filled 5 days ago. What is the best technician action?

- a) Fill both because SSRIs are commonly combined
- b) Flag for pharmacist review due to possible therapeutic duplication or switch
- c) Inactivate paroxetine and fill sertraline without review
- d) Dispense sertraline and add "do not take with paroxetine" auxiliary label

33) A prescription reads: "cephalexin 250 mg/5 mL: take 5 mL PO QID x 7 days." How many mL should be dispensed?

- a) 120 mL
- b) 140 mL
- c) 150 mL
- d) 180 mL

34) Which medication is a REMS program medication where the dispensing process can include specific documentation and timing requirements?

- a) Isotretinoin
- b) Amoxicillin
- c) Metformin
- d) Lisinopril

35) A controlled substance inventory is due. Which schedules are included in the federal biennial inventory requirement?

- a) Schedule II only
- b) Schedule I–V
- c) Schedule III–V only
- d) Schedule II–III only

36) A patient asks whether they can crush bupropion XL tablets. What is the best technician response?

- a) Yes, crushing improves absorption
- b) Yes, but only if mixed with water
- c) No, but splitting the tablet in half is fine
- d) Refer the patient to the pharmacist for counseling

37) The pharmacy receives a recall notice: “Quarantine all lots 7K12 of atorvastatin 20 mg.” The technician finds an open bottle with lot 7K12. What should be done?

- a) Use it first so none is wasted
- b) Move it to the back of the shelf and avoid selecting it
- c) Remove and quarantine it per recall procedure for return/credit and documentation
- d) Repackage into amber vials so the lot number is not visible

38) A prescription is for warfarin. The technician notices an active new prescription for trimethoprim-sulfamethoxazole from an outside provider entered in the profile. What should the technician do?

- a) Proceed because interaction checks are only for pharmacists
- b) Add a note/flag for pharmacist review due to increased bleeding risk
- c) Suggest the patient separate doses by 2 hours
- d) Inactivate warfarin until the antibiotic is finished

39) A patient’s label indicates “metformin ER.” During filling, the technician pulls metformin IR. What is the best technician action?

- a) Continue because both are metformin
- b) Adjust directions to “take BID” to make IR equivalent
- c) Replace with metformin ER and continue routine verification steps
- d) Dispense IR and add “do not crush” label

40) A prescription reads: "Take 1.5 tablets PO daily. Dispense #45." What is the days' supply?

- a) 15
- b) 20
- c) 30
- d) 45

41) A technician is asked which DEA form is used (paper) to order Schedule II controlled substances from a wholesaler.

- a) DEA Form 106
- b) DEA Form 41
- c) DEA Form 222
- d) DEA Form 363

42) A patient presents a prescription for oxycodone that is missing the date written. What is the appropriate technician action?

- a) Enter today's date and process
- b) Ask the patient for the date and enter it
- c) Hold and contact the prescriber to obtain the missing required element before dispensing
- d) Dispense a partial supply and fix the date later

43) A prescription is for "Prozac 20 mg daily." Which generic is correct?

- a) Sertraline
- b) Paroxetine
- c) Citalopram
- d) Fluoxetine

44) A patient drops off a prescription for "Adderall XR 20 mg DAW." Only generic mixed amphetamine salts ER is in stock. What should the technician do?

- a) Dispense the generic and keep DAW on the claim
- b) Change DAW code to allow substitution and process
- c) Contact the prescriber directly to change DAW
- d) Route to the pharmacist to determine substitution options and next steps

45) A technician is calculating an IV rate: 1,000 mL over 8 hours. What is the rate in mL/hr?

- a) 125 mL/hr
- b) 100 mL/hr
- c) 150 mL/hr
- d) 80 mL/hr

46) A patient asks to buy a sharps container and dispose of used fentanyl patches at the pharmacy. What is the correct approach?

- a) Accept and place them in regular trash if wrapped
- b) Accept and return to stock if sealed
- c) Use the pharmacy's authorized medication disposal process if available and document according to procedure
- d) Tell the patient to flush patches immediately and do not document

47) While pulling will-call, a technician finds a refrigerated biologic left at room temperature in will-call for 5 days. What should be done?

- a) Return it to the refrigerator and dispense as normal
- b) Quarantine it and escalate for stability assessment before dispensing/restocking
- c) Dispense if the patient arrives within the next hour
- d) Apply a new beyond-use date label and restock

48) A prescription reads: "valacyclovir 1 g: take 2 tablets PO BID for 1 day." What quantity should be dispensed?

- a) 2
- b) 4
- c) 6
- d) 8

49) During data entry, the technician sees two patient profiles with the same name but different DOBs; the e-prescription DOB doesn't match the selected profile. What is the best action?

- a) Use the address to guess which one is correct and continue
- b) Continue and correct after dispensing
- c) Create a third profile to avoid mixing records
- d) Stop and resolve the identity mismatch before processing

50) A prescription is for "Toprol-XL 50 mg daily." Which is the correct generic formulation?

- a) Metoprolol tartrate
- b) Metoprolol succinate ER
- c) Atenolol
- d) Propranolol ER

51) A prescription is for nitroglycerin SL tablets. The manufacturer bottle is designed to protect from moisture/light. How should it be dispensed when required for stability?

- a) Count into an amber vial to improve child-resistance
- b) Dispense in original manufacturer container when required
- c) Combine partial bottles if appearance matches
- d) Place a desiccant packet into a standard vial and dispense

52) A technician receives a fax requesting a patient's refill history from a prescriber for continuity of care. What is the best privacy-compliant action?

- a) Fax the full patient profile including payment methods and immunizations
- b) Email the information because it is faster and more secure
- c) Fax the minimum necessary information to a verified fax number associated with the requester
- d) Refuse unless the patient signs a release in person

53) A prescription reads: "hydrochlorothiazide 25 mg daily." During selection, the technician accidentally clicks hydroxychloroquine 200 mg. What is the best prevention step?

- a) Select the first "hydro-" option to reduce time
- b) Confirm the full drug name, strength, and indication clues (when available) before saving
- c) Rely on the pharmacist final check to correct entry errors

d) Use tablet color to confirm selection

54) A prescription reads: "Take 2.5 mL PO daily." The concentration is 100 mg/5 mL. How many mg per dose?

- a) 25 mg
- b) 50 mg
- c) 75 mg
- d) 100 mg

55) Which medication is a calcium channel blocker commonly used for hypertension?

- a) Lisinopril
- b) Hydrochlorothiazide
- c) Metoprolol
- d) Amlodipine

56) A prescription for gabapentin reads "300 mg: take 1 capsule TID. Dispense #270." What days' supply is this?

- a) 30
- b) 60
- c) 90
- d) 270

57) A patient's prescription is billed for #90 tablets, but the plan limit is #30 per fill. What is the best technician action?

- a) Dispense #90 and bill #30 without telling the patient
- b) Change quantity to #30 for this fill and process; ensure the remaining therapy is addressed via refills/new Rx per workflow
- c) Keep resubmitting the claim until it pays for #90
- d) Reverse the claim and dispense cash only

58) A patient asks: "Is Macrobid the same as nitrofurantoin?" What is the best technician response?

- a) Yes, Macrobid is a nitrofurantoin product
- b) No, Macrobid is a cephalosporin antibiotic
- c) Only if taken with milk
- d) Only for kidney infections, not bladder infections

59) A technician sees a bottle labeled "morphine ER 30 mg" placed in the bin for "morphine IR 30 mg." What is the best action?

- a) Leave it; the pharmacist final check will catch it
- b) Move both products into the same bin to reduce confusion
- c) Remove and restock correctly, and report/record the near-miss per quality process
- d) Put a handwritten note on the shelf and continue filling

60) A prescription reads: "insulin lispro U-100 vial: inject 8 units before meals TID. Dispense 10 mL." What days' supply should be billed?

- a) 28
- b) 35

- c) 41
- d) 50

61) Which medication is an SSRI antidepressant?

- a) Fluoxetine
- b) Amitriptyline
- c) Bupropion
- d) Duloxetine

62) Which action is required when a Medication Guide is flagged as required at dispensing?

- a) Provide it only on the first fill
- b) Provide it only when the patient requests it
- c) Provide it with each dispensing when required for that medication
- d) Provide it only for brand-name products

63) A new prescription reads: "omeprazole 20 mg daily," and the patient is also receiving clopidogrel. What is the best technician action?

- a) Process both and add "separate by 2 hours" label
- b) Substitute pantoprazole automatically during entry
- c) Refuse to fill omeprazole because it is OTC
- d) Flag for pharmacist review due to potential interaction/therapy concern

64) A written prescription for methylphenidate is presented without a prescriber signature. What should the technician do?

- a) Fill it since the prescriber name is printed on the pad
- b) Ask the patient to sign where the prescriber would sign
- c) Accept verbal confirmation from the office and dispense
- d) Hold it as invalid until a properly signed prescription is obtained

65) A prescription reads: "Take 5 mL PO TID x 7 days." How many total mL should be dispensed (no extra)?

- a) 70 mL
- b) 90 mL
- c) 105 mL
- d) 120 mL

66) A patient requests an early refill for clonazepam stating the medication was stolen. The last fill was 12 days ago for a 30-day supply. What should the technician do next?

- a) Process the refill now and note "stolen"
- b) Offer a small emergency supply as cash without review
- c) Tell the patient the pharmacy is required to replace it once
- d) Route the request for pharmacist evaluation before any controlled-substance early refill is processed

67) A prescription reads: "KCl 10 mEq PO daily." Only potassium chloride ER 20 mEq tablets are in stock. What should the technician do?

- a) Dispense 20 mEq and instruct "take every other day"
- b) Dispense 20 mEq and instruct "split tablet"

- c) Enter the available product and route to the pharmacist for decision on equivalence/therapy plan
- d) Change the dose to 20 mEq and reduce quantity by half

68) A prescription for “Zithromax Z-Pak” is entered. Which directions match standard Z-Pak dosing?

- a) Take 1 tablet twice daily for 7 days
- b) Take 2 tablets on day 1, then 1 tablet daily on days 2–5
- c) Take 2 tablets daily for 3 days
- d) Take 1 tablet daily for 10 days

69) A patient asks what atorvastatin is used for. What is the best technician response?

- a) Explain it lowers cholesterol and prevents heart attacks
- b) Tell the patient it treats high blood pressure
- c) Refer the patient to the pharmacist for counseling
- d) Tell the patient to stop taking it if muscle pain occurs

70) A pharmacy receives a shipment of refrigerated insulin, and the temperature indicator shows out-of-range exposure in transit. What should the technician do?

- a) Refrigerate and dispense first to minimize waste
- b) Freeze it for 30 minutes to restore stability
- c) Accept it if the boxes are unopened and cold to touch
- d) Quarantine the shipment and escalate for guidance before use/stocking

71) A technician is asked which drug is a high-alert anticoagulant commonly associated with serious harm if mis-dosed.

- a) Warfarin
- b) Loratadine
- c) Docusate
- d) Amoxicillin

72) A technician is entering: “ondansetron ODT 4 mg: dissolve 1 tab on tongue q8h prn nausea.” Which dosage form is correct?

- a) Oral capsule
- b) Immediate-release tablet
- c) Orally disintegrating tablet
- d) Buccal patch

73) A pharmacist asks the technician to gather transfer details for a Schedule IV prescription being transferred. Which item must be communicated in the transfer record?

- a) Patient’s diagnosis
- b) Prescriber’s state license expiration date
- c) Patient’s insurance group number
- d) Original prescription number and original date of issuance

74) A prescription is for “Advair Diskus.” Which dosage form is it?

- a) Metered-dose inhaler (MDI)
- b) Dry powder inhaler (DPI)

- c) Nebulizer solution
- d) Nasal spray

75) A technician notices a label for "Lasix 40 mg" was placed on a bottle of Lexapro 10 mg, caught before bagging. What should be done immediately?

- a) Remove the incorrect label, correct the labeling/product selection, and document as a near miss per QA process
- b) Leave it for the pharmacist to catch at final verification
- c) Bag it and instruct the cashier to confirm at pickup
- d) Discard the medication bottle to prevent confusion

76) A prescription reads: "Apply 2 g topically BID." The tube contains 60 g. Approximately how many days' supply is one tube?

- a) 7 days
- b) 10 days
- c) 15 days
- d) 30 days

77) A patient asks the technician to "run my friend's insurance just this once." What is the best response?

- a) Do it if the friend verbally agrees by phone
- b) Run it and then rebill correctly later
- c) Decline because it can constitute fraud; escalate according to workflow
- d) Use any discount card automatically without telling the patient

78) A prescription is entered as "azithromycin 500 mg daily x 3 days." The pharmacy has 250 mg tablets. What is the best technician action?

- a) Change directions to "take 2 tablets daily" and dispense 6 tablets
- b) Dispense 3 tablets and tell patient to take two on day 1 only
- c) Dispense 6 tablets with directions unchanged
- d) Route to the pharmacist to confirm appropriate substitution and labeling

79) A technician is asked about partial filling a Schedule II prescription due to insufficient stock. Which statement best reflects federal rules?

- a) Partial fills of Schedule II are never allowed
- b) Partial fills may be allowed under specific conditions with required documentation and time limits
- c) Partial fills are allowed only when the patient pays cash
- d) A partial fill converts the prescription to Schedule III

80) A prescription reads: "dorzolamide/timolol: instill 1 gtt OS BID." What does "OS" mean?

- a) Right eye
- b) Left ear
- c) Left eye
- d) Both eyes

81) A patient picks up two prescriptions and the technician realizes afterward the wrong leaflet (patient education) was stapled to one bag, but the medication label and patient name

on the vial are correct. What is the best quality action?

- a) Ignore it because the medication was correct
- b) Document and correct the paperwork error per QA process because it can confuse patients and cause misuse
- c) Call the patient and tell them to read the bottle label only
- d) Reprint the leaflet for the next fill and do nothing now

82) A new prescription is for "amlodipine 10 mg daily." Which class is amlodipine?

- a) ACE inhibitor
- b) Calcium channel blocker
- c) Beta blocker
- d) Thiazide diuretic

83) A technician is asked which action best supports DSCSA compliance when receiving product missing the required 2D data matrix on the package.

- a) Create a new barcode from the NDC and apply it
- b) Put it on the shelf but mark "do not use"
- c) Dispense it only if the patient signs acknowledging the missing code
- d) Quarantine the product and escalate for proper investigation/handling

84) A prescription reads: "metformin ER 500 mg: take 2 tablets by mouth every evening. Dispense #60." What days' supply should be entered?

- a) 15
- b) 30
- c) 45
- d) 60

85) A patient with the same first and last name as another patient arrives to pick up oxycodone. What is the best first step before retrieving any bag?

- a) Ask for two identifiers such as date of birth and address
- b) Ask which strength they take and select based on that
- c) Pull both patients' bags and ask which looks correct
- d) Ask only for a phone number

86) A prescription reads: "insulin aspart: inject 12 units SC before meals TID." How many total units are used per day?

- a) 12
- b) 24
- c) 36
- d) 48

87) A technician is asked where to file executed DEA Form 222 and Schedule II invoices so they are compliant and retrievable. Which is correct?

- a) File with general purchasing records and discard after one year
- b) Store in a manner that is readily retrievable and retained for at least 2 years
- c) Mail to the wholesaler after receiving the order
- d) Keep only the electronic scan; destroy the paper immediately

88) A prescription for "Synthroid 125 mcg DAW" is received. What should the technician do during product selection and billing?

- a) Select generic levothyroxine because it is equivalent
- b) Change DAW to allow generic substitution
- c) Select the brand Synthroid product and ensure DAW is processed correctly
- d) Dispense generic but write "brand requested" on the label

89) A prescription reads: "Take 1 tab PO qod." How should "qod" be interpreted?

- a) Four times daily
- b) Every other day
- c) Every day
- d) Once weekly

90) A prescription reads: "amoxicillin/clavulanate 600-42.9 mg/5 mL: take 7.5 mL PO BID x 10 days." How many mL should be dispensed (no extra)?

- a) 120 mL
- b) 150 mL
- c) 160 mL
- d) 180 mL

Answer Key

1) c	31) c	61) a
2) b	32) b	62) c
3) b	33) b	63) d
4) a	34) a	64) d
5) d	35) b	65) c
6) a	36) d	66) d
7) c	37) c	67) c
8) b	38) b	68) b
9) a	39) c	69) c
10) c	40) c	70) d
11) d	41) c	71) a
12) b	42) c	72) c
13) b	43) d	73) d
14) c	44) d	74) b
15) d	45) a	75) a
16) c	46) c	76) c
17) d	47) b	77) c
18) a	48) b	78) d
19) a	49) d	79) b
20) c	50) b	80) c
21) c	51) b	81) b
22) d	52) c	82) b
23) b	53) b	83) d
24) c	54) b	84) b
25) d	55) d	85) a
26) b	56) c	86) c
27) c	57) b	87) b
28) d	58) a	88) c
29) b	59) c	89) b
30) a	60) c	90) b

PTCB Sim Test 02

1) While filling, the barcode scan pulls up “metformin ER 500 mg,” but the label is for “metformin 500 mg (IR).” What should the technician do next?

- a) Continue if the strength matches
- b) Change the directions to “once daily” to match ER use
- c) Stop and select the correct IR product/NDC before counting
- d) Dispense ER and mark “substitution allowed”

2) A patient wants to purchase pseudoephedrine, but their government-issued ID is expired. The requested quantity is within daily limits. What is the correct action?

- a) Complete the sale and record the ID number anyway
- b) Decline the sale because valid ID is required to log the transaction
- c) Accept a credit card as verification
- d) Sell it if the patient signs a log statement

3) An e-prescription reads: “Crestor 20 mg PO daily.” Which product should be selected?

- a) Atorvastatin 20 mg
- b) Rosuvastatin 20 mg
- c) Simvastatin 20 mg
- d) Pravastatin 20 mg

4) A caller says they are the patient’s spouse and asks, “Is my husband’s refill ready and how much will it cost?” The spouse is not listed as authorized. What can the technician say?

- a) The refill is ready, but not the medication name
- b) The medication name, but not the price
- c) Nothing about the patient’s prescription status without authorization
- d) The prescriber name only

5) A prescriber’s office phones in a new prescription for amoxicillin. The pharmacist is on the phone with another prescriber. What is the appropriate handling?

- a) The technician may take it because it is not controlled
- b) The technician may take it and the pharmacist can review later
- c) Ask the caller to fax it; phone orders are not allowed
- d) Have the caller hold or call back so the pharmacist can receive the verbal order

6) A patient profile lists “anaphylaxis to sulfa.” A new order arrives for sulfamethoxazole/trimethoprim DS. What should the technician do?

- a) Enter and fill; the prescriber decided benefits outweigh risks
- b) Enter and override the allergy alert to prevent delay
- c) Hold the prescription for pharmacist review before dispensing
- d) Change the antibiotic to nitrofurantoin and proceed

7) A refrigerator temperature log shows 11°C for about 2 hours. Vaccines were stored inside. What should the technician do first?

- a) Move vaccines to the freezer to stabilize them
- b) Keep using vaccines; only freezing is a concern

- c) Quarantine the affected vaccines and alert the pharmacist promptly
- d) Discard the vaccines immediately

8) An insurance claim rejects "PA required." The patient hands the technician a manufacturer copay card and asks to "just run it instead." What is the best next step within technician scope?

- a) Run the copay card as primary to bypass the PA
- b) Change the days' supply until the claim pays
- c) Offer the cash price and route the situation for pharmacist follow-up on PA options
- d) Enter a different diagnosis code to force approval

9) A prescription is for "Toprol-XL 50 mg daily." During entry, the technician selected metoprolol tartrate 50 mg. What is the correct action?

- a) Keep tartrate and change sig to BID
- b) Keep tartrate; they are equivalent
- c) Add an "ER" auxiliary label and dispense tartrate
- d) Correct the selection to metoprolol succinate ER before continuing

10) At pickup, two will-call bags have similar names (e.g., "Maria P." and "Marian P."). What is the best verification method?

- a) Verify last name only
- b) Ask what medication they're expecting
- c) Verify name and date of birth before handing over any bag
- d) Verify prescriber name only

11) A written Schedule II prescription is missing the patient address. What must occur before dispensing?

- a) The pharmacy may add the address from the patient profile and proceed
- b) The patient may write in the address at pickup
- c) The prescriber must provide a valid prescription containing required information before dispensing
- d) Dispense a partial fill and obtain the address later

12) Prescription: "Amoxicillin 400 mg/5 mL: Take 7.5 mL PO BID x 10 days." How many mL should be dispensed?

- a) 150 mL
- b) 75 mL
- c) 120 mL
- d) 200 mL

13) A prescription is written: "Azithromycin Z-Pak" with no directions and no quantity. What is the appropriate technician action?

- a) Enter standard Z-Pak directions and #6 automatically
- b) Enter #5 because it is a 5-day course
- c) Substitute azithromycin 500 mg daily x 3 days
- d) Set aside for pharmacist because required directions/quantity are missing

14) A patient profile shows warfarin active. A new prescription arrives for metronidazole 500

mg BID x 7 days. What should the technician do?

- a) Process normally because it is short duration
- b) Route for pharmacist review due to significant interaction risk
- c) Tell the patient to skip warfarin while on metronidazole
- d) Change to clindamycin without contacting the prescriber

15) During a controlled substance count, the perpetual inventory shows 250 tablets, but the physical count is 246. What should the technician do?

- a) Adjust the perpetual count to 246 and continue
- b) Recount and promptly report the discrepancy for pharmacist review
- c) Ignore it; small discrepancies are expected
- d) Dispense 4 tablets to "fix" the count

16) Prescription: "Ventolin HFA: Inhale 2 puffs QID." Inhaler contains 200 actuations. What is the days' supply?

- a) 12 days
- b) 20 days
- c) 25 days
- d) 50 days

17) A trading partner requests DSCSA transaction documentation for a recalled lot your pharmacy received last month. What should the technician retrieve?

- a) Only the purchase price and payment record
- b) The transaction information/transaction statement maintained by the pharmacy
- c) The patient copays for dispensed prescriptions from that lot
- d) Only the controlled-substance invoice, if any

18) A customer asks for an OTC product for new chest tightness with sweating and nausea. What is the best technician action?

- a) Suggest antacids and rest
- b) Suggest ibuprofen and fluids
- c) Suggest a cough/cold product for symptom relief
- d) Get the pharmacist immediately and advise urgent medical evaluation

19) A prescription is for "prednisone 20 mg tablets," but the computer selection is prednisolone oral solution 15 mg/5 mL. What should the technician do?

- a) Convert the dose to mL and dispense the solution
- b) Continue; both are corticosteroids
- c) Ask the patient which form they prefer and dispense that
- d) Correct the drug/form selection to match the prescription before processing

20) Prescription: "Take 2 tablets PO q6h x 5 days." How many tablets are needed?

- a) 20
- b) 30
- c) 40
- d) 60

21) A patient requests an early refill on a Schedule IV medication because they are leaving

town tomorrow. What is the appropriate technician action?

- a) Process the early refill automatically with a note "travel"
- b) Refuse; early refills are never allowed for controls
- c) Gather travel details and route for pharmacist decision/override per workflow
- d) Split the fill into two prescriptions to make it legal

22) A prescription reads: "Eliquis 5 mg." Which generic name is correct?

- a) Apixaban
- b) Rivaroxaban
- c) Dabigatran
- d) Edoxaban

23) In a unit-dose cart, a blister pack is missing a readable lot number. What should the technician do?

- a) Use it if the drug name and strength are readable
- b) Write the lot number based on a similar blister
- c) Set it aside for proper identification/traceability before use
- d) Use it only for PRN doses

24) A patient requests a transfer of a noncontrolled prescription to a different pharmacy. Which information is essential to communicate?

- a) Patient diagnosis
- b) Original prescription number and remaining refills
- c) Prescriber's DEA number
- d) Patient insurance BIN/PCN

25) Insulin glargine U-100 pens: "Inject 24 units SC daily. Dispense 1 box." One box = 5 pens × 3 mL each. What days' supply should be entered?

- a) 25 days
- b) 31 days
- c) 62 days
- d) 75 days

26) A patient drops off a prescription for "omeprazole 20 mg daily." Which class best matches this medication?

- a) H2 blocker
- b) Proton pump inhibitor
- c) Antispasmodic
- d) Prostaglandin analog

27) A faxed prescription for a controlled medication has a signature that appears copied. What is the best technician action?

- a) Fill it if the prescriber is in the system
- b) Call the patient to confirm it is legitimate
- c) Proceed if the insurance claim pays
- d) Set it aside for pharmacist review as potentially suspicious

28) Prescription: "Augmentin 875/125 mg: 1 tab q12h x 10 days," but quantity is written as

#14. What should the technician do?

- a) Fill #14 because quantity is explicit
- b) Fill #20 because duration is explicit
- c) Change the directions to "x 7 days" to match #14
- d) Route to the pharmacist because the quantity conflicts with directions

29) A prescription for isotretinoin triggers "REMS requirements not met." What should the technician do?

- a) Bypass the alert; eRx means requirements were met
- b) Fill as cash to avoid REMS restrictions
- c) Hold processing and route for pharmacist to address REMS steps before dispensing
- d) Substitute topical tretinoin

30) A prescription is for Norvasc 10 mg daily. Which generic is correct?

- a) Amlodipine
- b) Losartan
- c) Lisinopril
- d) Diltiazem

31) Fentanyl patches: "Apply 1 patch every 72 hours. Dispense #10." What days' supply should be entered?

- a) 10 days
- b) 20 days
- c) 30 days
- d) 72 days

32) The label prints "Take 1 tablet BID," but the eRx directions are "Take 1 tablet daily." What should the technician do?

- a) Dispense and let the patient choose
- b) Correct the directions to match the prescription before it moves forward
- c) Change the quantity to match the printed label
- d) Keep the printed label; changes are not allowed after printing

33) Which pair represents a common LASA (look-alike/sound-alike) risk?

- a) Hydroxyzine and hydralazine
- b) Sertraline and senna
- c) Amoxicillin and acetaminophen
- d) Ibuprofen and isosorbide

34) A patient has "ACE inhibitor—angioedema" listed as a severe allergy. A new prescription arrives for lisinopril. What should the technician do?

- a) Process; it is likely intentional
- b) Delete the allergy since it causes claim problems
- c) Change to losartan automatically
- d) Route for pharmacist review before filling due to high-severity allergy history

35) An IV order states: "0.9% NaCl 1,000 mL infuse over 8 hours." What rate is correct?

- a) 100 mL/hr

- b) 120 mL/hr
- c) 125 mL/hr
- d) 150 mL/hr

36) A patient returns an unopened controlled substance requesting a refund and for it to be returned to stock. What should the technician do?

- a) Return it to inventory if the seal is intact
- b) Accept it and place it back into will-call
- c) Inform them it cannot be returned to stock; route to pharmacist for disposal options
- d) Take it back only if they bring the receipt

37) Which medication is a benzodiazepine?

- a) Buspirone
- b) Zolpidem
- c) Lorazepam
- d) Hydroxyzine

38) A stock bottle arrives with a broken seal and tablets visible. What is the correct technician response?

- a) Use it if tablets appear clean
- b) Repackage into a new bottle and label as pharmacy stock
- c) Use it only for noncontrolled prescriptions
- d) Quarantine as suspect product and bring to pharmacist/vendor return process

39) A prescription is for "ketorolac 0.5% ophthalmic solution." The technician selected ketorolac 10 mg tablets. What should be done?

- a) Change directions and dispense tablets
- b) Correct the dosage form and product selection before proceeding
- c) Dispense tablets; same drug name is sufficient
- d) Ask the patient which form they want

40) Prescription: "Test blood glucose TID. Dispense #100." What days' supply should be entered?

- a) 25 days
- b) 30 days
- c) 33 days
- d) 90 days

41) A patient says, "My doctor told me twice a day, but the label says once a day." What is the best first step for the technician?

- a) Tell the patient to follow what the doctor said
- b) Tell the patient to follow the label and ignore the doctor
- c) Collect details and bring the discrepancy to the pharmacist for assessment
- d) Reprint the label with BID directions

42) A recall notice lists a specific lot of metformin ER 500 mg. What should the technician do?

- a) Pull only the affected lot from inventory and quarantine it

- b) Pull all metformin products regardless of lot
- c) Continue dispensing until stock is gone
- d) Put a warning sticker on the shelf and keep filling

43) A Schedule II e-prescription arrives with directions but no quantity. What is required before dispensing?

- a) Default to #30
- b) Dispense a 7-day supply and complete later
- c) Obtain prescriber clarification for the missing required element
- d) Ask the patient what quantity they want

44) A patient requests counseling on a new anticoagulant at pickup. What should the technician do?

- a) Provide counseling on bleeding precautions
- b) Offer counseling only if the patient asks twice
- c) Refer the patient to the pharmacist for counseling
- d) Hand them a medication guide and complete the sale

45) Which medication is an SSRI?

- a) Amitriptyline
- b) Sertraline
- c) Duloxetine
- d) Bupropion

46) An opioid-naïve 15-year-old has a new prescription for morphine ER 30 mg. What should the technician do?

- a) Enter it and expedite filling to avoid delaying therapy
- b) Reduce the dose to 15 mg ER and process
- c) Convert to morphine IR and process
- d) Route for pharmacist review due to high-risk therapy concerns

47) A profile lists "anaphylaxis to penicillin." A prescription arrives for cephalexin. What should the technician do?

- a) Fill it; cephalexin is not penicillin
- b) Override the allergy because cross-reactivity is uncommon
- c) Hold and route for pharmacist review due to severe allergy history
- d) Call the prescriber and request azithromycin

48) Cyanocobalamin 1,000 mcg/mL: "Inject 1 mL monthly." How many mg is each dose?

- a) 0.1 mg
- b) 0.5 mg
- c) 1 mg
- d) 10 mg

49) Heparin infusion: 25,000 units in 250 mL D5W. Ordered rate: 1,500 units/hr. What pump rate (mL/hr) is correct?

- a) 10 mL/hr
- b) 12 mL/hr

- c) 15 mL/hr
- d) 20 mL/hr

50) A prescription is for clonidine 0.1 mg/24 hr patch weekly #4. The technician selected clonidine 0.1 mg tablets. What should be done?

- a) Convert to tablets because clonidine is clonidine
- b) Change quantity to #28 and proceed
- c) Keep tablets and add "do not crush" label
- d) Correct the dosage form selection before proceeding

51) A patient asks if they can move nitroglycerin SL tablets into a weekly pill organizer. What is the most appropriate technician response?

- a) Yes; this prevents missed doses
- b) No; they should store it in the bathroom medicine cabinet
- c) Refer to the pharmacist; these are generally best kept in the original container
- d) Yes, as long as the organizer is plastic

52) Which medication is considered high-alert in most settings?

- a) Fluticasone nasal spray
- b) Docusate
- c) Insulin
- d) Loratadine

53) Lactulose 10 g/15 mL: "Take 30 mL daily." How many grams per dose?

- a) 10 g
- b) 15 g
- c) 20 g
- d) 30 g

54) A prescriber's office requests the patient's complete medication profile by fax for "chart updating," but there is no patient authorization on file. What should the technician do?

- a) Fax it because it is for treatment
- b) Fax only controlled substances
- c) Fax it if the office confirms DOB and address
- d) Route through the pharmacist/privacy workflow to verify permissible disclosure before sending

55) Diltiazem infusion: 125 mg in 100 mL. Ordered 5 mg/hr. What is the infusion rate?

- a) 2 mL/hr
- b) 4 mL/hr
- c) 6 mL/hr
- d) 8 mL/hr

56) Testosterone cypionate 200 mg/mL: "Inject 0.75 mL IM weekly." How many mg per dose?

- a) 75 mg
- b) 100 mg
- c) 150 mg

d) 200 mg

57) Which medication is an ACE inhibitor?

- a) Losartan
- b) Amlodipine
- c) Propranolol
- d) Lisinopril

58) A patient requests transfer of a Schedule III prescription to another pharmacy. Which statement is correct?

- a) It can be transferred only if zero refills remain
- b) A technician may transfer it independently if refills remain
- c) It may be transferred once between pharmacies (if allowed by state law) through the pharmacist transfer process
- d) Schedule III prescriptions can never be transferred

59) An eye drop bottle is 10 mL and delivers about 20 drops per mL. Approximately how many drops are in the bottle?

- a) 100 drops
- b) 150 drops
- c) 200 drops
- d) 250 drops

60) A patient profile shows sertraline active. A new prescription from a different prescriber is for fluoxetine. What should the technician do?

- a) Fill both; they treat different symptoms
- b) Inactivate sertraline automatically and fill fluoxetine
- c) Dispense fluoxetine and tell the patient to stop sertraline
- d) Route for pharmacist review for possible duplicate SSRI therapy/transition

61) Directions: "Take 1.5 tablets daily." Quantity dispensed: 45 tablets. What days' supply is correct?

- a) 15 days
- b) 30 days
- c) 45 days
- d) 60 days

62) Which generic is Lasix?

- a) Hydrochlorothiazide
- b) Spironolactone
- c) Bumetanide
- d) Furosemide

63) Prescription: "omeprazole 20 mg daily #90." The technician selected esomeprazole 20 mg and the patient says "either is fine." What should the technician do?

- a) Dispense esomeprazole because the patient agreed
- b) Correct to omeprazole as prescribed; any interchange requires proper authorization/workflow

- c) Change the drug and document "patient preference"
- d) Dispense esomeprazole and annotate "therapeutic interchange"

64) A patient scheduled for vaccination has a severe latex allergy on profile. What should the technician do?

- a) Ignore it; latex is irrelevant to vaccines
- b) Remove the allergy because it is not drug-related
- c) Ask the patient to bring non-latex gloves
- d) Flag it so product/packaging can be checked before administration

65) Cream directions: "Apply 2 g topically BID." Tube contains 60 g. What days' supply should be entered?

- a) 10 days
- b) 12 days
- c) 15 days
- d) 30 days

66) Third-party requires days' supply for an albuterol inhaler. Prescription: "Inhale 2 puffs q4h PRN." Inhaler contains 120 puffs. If billed using a maximum of 6 doses/day, what days' supply is most appropriate?

- a) 5 days
- b) 10 days
- c) 15 days
- d) 30 days

67) During entry, the system auto-populates "cephalexin 500 mg TID," but the prescription says "500 mg QID x 10 days #40." What should the technician do?

- a) Keep TID because it is common dosing
- b) Change quantity to #30 to match TID x 10 days
- c) Correct the directions to QID to match the prescription
- d) Leave it; the pharmacist will fix it later if needed

68) A parent asks for an OTC product for a child's fever but does not know the child's weight. What should the technician do?

- a) Recommend aspirin for faster fever reduction
- b) Recommend splitting an adult acetaminophen tablet
- c) Recommend a multi-symptom cold/flu product
- d) Refer to the pharmacist for appropriate product and dosing guidance

69) Which brand matches rivaroxaban?

- a) Xarelto
- b) Eliquis
- c) Pradaxa
- d) Savaysa

70) Metformin ER 500 mg daily is prescribed, but the technician pulled metformin IR 500 mg from an adjacent shelf. What should the technician do?

- a) Dispense IR and add "do not crush" label

- b) Change the directions to BID and dispense IR
- c) Correct to ER and verify with barcode/NDC scan before filling
- d) Dispense IR and note "equivalent generic"

71) Prescription: "Take 1 tablet q6h PRN pain. Dispense #24." What is the maximum days' supply at maximum use?

- a) 2 days
- b) 4 days
- c) 6 days
- d) 8 days

72) The hard copy reads "hydralazine 25 mg TID," but the label prints "hydroxyzine 25 mg." What should the technician do?

- a) Dispense; pharmacist will catch it at final check
- b) Correct the drug selection and reprint labels as needed before it proceeds
- c) Change the sig to match hydroxyzine common dosing
- d) Ask the patient at pickup which one they take

73) An e-prescription for a controlled medication arrives, and the patient profile lacks date of birth. What should the technician do before completing processing?

- a) Proceed; DOB is optional
- b) Use an estimated DOB so the claim will adjudicate
- c) Confirm DOB from a reliable source to prevent misidentification and processing errors
- d) Wait until pickup and update after dispensing

74) A prescription is for Lantus SoloStar. Which generic name should be selected?

- a) Insulin lispro
- b) Insulin glargine
- c) Insulin aspart
- d) Insulin NPH

75) Eye drops: "Instill 1 drop OU BID." Bottle size 5 mL. Drop factor 20 drops/mL. What days' supply should be entered (use 1 drop = 1 gtt)?

- a) 12 days
- b) 25 days
- c) 50 days
- d) 100 days

76) A partial fill of a Schedule II prescription was dispensed due to insufficient stock 18 days ago. The patient requests the remainder today. What should the technician do?

- a) Dispense the remainder now; partial fills can be completed anytime
- b) Create a new prescription for the remainder
- c) Transfer the remainder to another pharmacy
- d) Route for pharmacist review to confirm whether completion is allowed within applicable time limits

77) Which medication is a beta blocker commonly used for hypertension and rate control?

- a) Losartan

- b) Metoprolol
- c) Amlodipine
- d) Hydrochlorothiazide

78) The prescription is entered as bupropion SR 150 mg, but the prescriber's note says "XL preferred." What should the technician do?

- a) Change SR to XL; they are both extended-release
- b) Fill SR because it is close enough
- c) Route for pharmacist clarification because SR vs XL is a formulation difference
- d) Substitute IR and change directions to TID

79) Two different manufacturers' bottles of the same generic/strength are on the shelf. The claim and label are linked to a specific NDC. What is best practice?

- a) Combine the bottles to simplify inventory
- b) Use either manufacturer; generics are interchangeable for billing
- c) Disable scanning to prevent delays
- d) Scan the stock bottle used to ensure the dispensed NDC matches what is billed/printed

80) Which medication is an inhaled corticosteroid used for asthma maintenance?

- a) Albuterol
- b) Ipratropium
- c) Fluticasone
- d) Montelukast

81) Which auxiliary label is most appropriate for doxycycline?

- a) Avoid excessive sunlight/use sunscreen
- b) Take only at bedtime
- c) Avoid grapefruit juice
- d) May cause drowsiness

82) A patient asks the technician to "change the label to a lower dose" because the medication makes them sleepy. What should the technician do?

- a) Change the directions in the computer to a lower dose
- b) Suggest taking half the tablet until the next appointment
- c) Tell the patient to stop immediately
- d) Refer the request to the pharmacist for assessment and counseling

83) A new prescription is for rosuvastatin 20 mg daily. The profile shows atorvastatin 40 mg filled last week and still active. What should the technician do?

- a) Inactivate atorvastatin and fill rosuvastatin
- b) Fill both; two statins may be used together routinely
- c) Change rosuvastatin to atorvastatin automatically
- d) Route for pharmacist review for possible therapy change/duplicate therapy

84) Which medication is a thiazide diuretic?

- a) Furosemide
- b) Bumetanide
- c) Hydrochlorothiazide

d) Spironolactone

85) A Schedule II eRx for Adderall XR 20 mg arrives. The address on the eRx does not match the address in the patient profile. What should the technician do?

- a) Ignore the mismatch if the DOB matches
- b) Update the profile to match the eRx without verification
- c) Continue and confirm the address at pickup after dispensing
- d) Pause processing and verify correct patient/profile information before proceeding

86) A patient asks for an OTC sleep aid that is safe to take with a new opioid prescription. What should the technician do?

- a) Recommend diphenhydramine because it is OTC
- b) Recommend doxylamine because it is stronger
- c) Recommend melatonin because it is "natural"
- d) Refer the patient to the pharmacist for interaction/safety screening and recommendation

87) While restocking, a technician sees potassium chloride injection stored next to saline flush syringes. What is the best action to reduce error risk?

- a) Keep them together for faster restocking
- b) Separate and store potassium chloride injection away from flushes
- c) Store both in the refrigerator
- d) Remove the flushes and keep potassium chloride in the bin

88) Which medication is an opioid antagonist used for overdose reversal?

- a) Buprenorphine
- b) Naloxone
- c) Methadone
- d) Naltrexone

89) A caller asks if the pharmacy can confirm whether their spouse picked up a prescription earlier today. The caller is not authorized on the profile. What should the technician do?

- a) Confirm pickup status only
- b) Confirm if the caller knows the date of birth
- c) Confirm the total price only
- d) Decline to disclose and advise the patient contact the pharmacy directly

90) A technician is typing a controlled-substance prescription and notices the prescriber's DEA number is missing one digit on the hard copy. What should the technician do?

- a) Add the missing digit based on the prescriber's NPI pattern
- b) Use the pharmacy DEA number to submit the claim
- c) Enter the incomplete DEA; it will still validate later
- d) Verify the prescriber identifier through appropriate records before completing controlled-substance processing requirements

Answer Key

1) c	31) c	61) b
2) b	32) b	62) d
3) b	33) a	63) b
4) c	34) d	64) d
5) d	35) c	65) b
6) c	36) c	66) b
7) c	37) c	67) c
8) c	38) d	68) d
9) d	39) b	69) a
10) c	40) c	70) c
11) c	41) c	71) b
12) a	42) a	72) b
13) d	43) c	73) c
14) b	44) c	74) b
15) b	45) b	75) c
16) c	46) d	76) d
17) b	47) c	77) b
18) d	48) c	78) c
19) d	49) c	79) d
20) c	50) d	80) c
21) c	51) c	81) a
22) a	52) c	82) d
23) c	53) b	83) d
24) b	54) d	84) c
25) c	55) b	85) d
26) b	56) c	86) d
27) d	57) d	87) b
28) d	58) c	88) b
29) c	59) c	89) d
30) a	60) d	90) d

PTCB Sim Test 03

1) A new eRx reads: "Synthroid 75 mcg: take 1 tablet daily." The product pulled is Euthyrox 75 mcg. The patient profile shows "DAW=1 (brand medically necessary)" from the prescriber. What should be dispensed?

- a) Euthyrox 75 mcg
- b) Any levothyroxine 75 mcg generic
- c) Synthroid 75 mcg
- d) Liothyronine 75 mcg

2) A patient asks to purchase pseudoephedrine 30 mg, quantity 96 tablets. Your state uses an electronic NPLEx log. What must occur before completing the sale?

- a) Confirm the patient's diagnosis and document it
- b) Record the sale in the system and verify government-issued photo ID
- c) Obtain a written prescription from the prescriber
- d) Process the claim through the patient's insurance plan first

3) While entering "metoprolol succinate ER 50 mg daily," you notice the system selected "metoprolol tartrate 50 mg." The printed label has not been generated yet. What is the most appropriate technician action?

- a) Change the product selection to metoprolol succinate ER 50 mg
- b) Keep tartrate but change directions to twice daily
- c) Keep tartrate because the milligram strength matches
- d) Delete the prescription and ask the prescriber to resend

4) A pharmacist asks you to file controlled substance invoices. Which schedule requires invoices to be readily retrievable and maintained separately from noncontrolled records (or clearly identifiable)?

- a) Schedule II
- b) Schedule IV
- c) Schedule V
- d) All schedules have identical filing requirements under federal law

5) Rx: "Amoxicillin/clavulanate 875/125 mg: take 1 tablet PO BID x 10 days." What quantity should be dispensed?

- a) 14
- b) 20
- c) 28
- d) 30

6) During pickup, the patient asks, "Is it okay to take this new antibiotic with my warfarin?" What should the technician do?

- a) Tell the patient to separate doses by 2 hours
- b) Tell the patient to stop warfarin until the antibiotic is finished
- c) Provide a printed interaction list and let the patient decide
- d) Offer to connect the patient with the pharmacist for counseling

7) A hospital floor sends a request for “insulin lispro U-100” but the technician accidentally selects “insulin glargine U-100.” What type of error is this most directly?

- a) Wrong route
- b) Wrong drug selection
- c) Wrong patient
- d) Wrong prescriber

8) A technician receives a sealed tote from the wholesaler. The DSCSA system flags one bottle as “product identifier could not be verified.” What should the technician do with the bottle?

- a) Place it in active stock if the lot and expiration look normal
- b) Return it to the shelf but add a note to scan again later
- c) Quarantine it in the designated area for investigation
- d) Dispense it only for cash transactions to avoid insurance audits

9) Rx: “Albuterol HFA: inhale 2 puffs q4h PRN.” The inhaler contains 200 actuations. If used at the maximum frequency, what is the days’ supply?

- a) 16 days
- b) 20 days
- c) 25 days
- d) 30 days

10) A patient’s profile shows anaphylaxis to penicillin. A new prescription arrives for cephalexin. What should the technician do before it is filled?

- a) Change cephalexin to azithromycin and continue
- b) Add an auxiliary label “watch for rash” and fill
- c) Hold the order for pharmacist review of the allergy risk
- d) Fill it since cephalexin is not a penicillin

11) In a long-term care facility, a faxed Schedule II prescription is received for a resident. Under federal rules, when can a fax serve as the original for dispensing?

- a) Never; a fax is not acceptable for Schedule II
- b) Only if the patient is in a long-term care facility or for hospice/emergency exceptions as applicable
- c) Only if the prescriber also calls to confirm it
- d) Only if the pharmacy keeps the fax for 1 year

12) A prescription label prints “predniSONE 10 mg,” but the bottle pulled is “prednisoLONE 10 mg.” What is the best technician action?

- a) Dispense prednisolone; they are the same drug
- b) Change the directions to match prednisolone and proceed
- c) Continue filling and let the patient decide at pickup
- d) Stop and correct the product selection to match the prescription

13) A patient asks for the generic name of Crestor. Which is correct?

- a) Rosuvastatin
- b) Atorvastatin
- c) Simvastatin

d) Pravastatin

14) You are filling “diltiazem ER 240 mg” and the prescriber’s eRx does not specify the ER type. The patient previously received “diltiazem CD 240 mg.” What is the best technician action?

- a) Select any ER diltiazem product because they are interchangeable
- b) Choose the least expensive NDC to reduce copay
- c) Match the last-filled product (CD) without checking
- d) Route to the pharmacist to confirm the intended ER formulation

15) A pharmacy has a partial fill of a Schedule II due to insufficient stock. Which statement is correct federally for a typical community setting?

- a) The remainder must be dispensed within 72 hours or the balance is void
- b) The remainder can be filled anytime within 6 months
- c) The remainder becomes a refill and can be transferred
- d) Partial fills are never permitted for Schedule II prescriptions

16) Rx: “Azithromycin 200 mg/5 mL: take 12.5 mL day 1, then 6.25 mL daily days 2–5.” How many mL should be dispensed?

- a) 25 mL
- b) 30 mL
- c) 37.5 mL
- d) 45 mL

17) A prescription reads: “Instill 1 gtt OS BID.” What does OS mean?

- a) Left eye
- b) Right eye
- c) Both eyes
- d) Left ear

18) A technician is asked which of these medications is a proton pump inhibitor.

- a) Famotidine
- b) Omeprazole
- c) Dicyclomine
- d) Furosemide

19) A patient requests a copy of their adult daughter’s medication profile. The daughter is not present and no authorization is on file. What should the technician do?

- a) Provide the list if the caller can confirm address and DOB
- b) Provide only noncontrolled medications
- c) Decline and request authorization from the patient or have the patient contact the pharmacy
- d) Provide the last 3 fills only since that is not “full PHI”

20) Rx: “Warfarin 5 mg: take as directed. Dispense #30.” The directions are not specific and the patient is new to the pharmacy. What is the best next step?

- a) Process as written because warfarin is commonly “as directed”
- b) Enter “take 1 tablet daily” to create a calculable days’ supply

- c) Route for pharmacist clarification so the exact dosing schedule is documented
- d) Dispense #30 and mark days' supply as 30

21) A technician is restocking and notices hydrALAZINE 25 mg and hydrOXYzine 25 mg stored adjacent with similar shelf labels. What action best reduces selection errors?

- a) Store them together so the staff can compare carefully
- b) Separate them and apply LASA/Tall Man safeguards per workflow
- c) Remove one medication from inventory
- d) Re-label both as "H-25" to simplify

22) Which medication is a direct oral anticoagulant (DOAC)?

- a) Apixaban
- b) Enoxaparin
- c) Warfarin
- d) Heparin

23) A prescriber's office leaves a voicemail with a new prescription for tramadol. What is the correct workflow?

- a) A technician may enter and process it as a new verbal prescription
- b) A pharmacist must take the verbal order (or directly receive it) before it can be processed
- c) Any staff member can take controlled substance voicemails as long as they document it
- d) The patient can sign a form to validate the voicemail

24) Rx: "Cephalexin 250 mg/5 mL: take 10 mL PO TID x 7 days." How many mL should be dispensed?

- a) 180 mL
- b) 200 mL
- c) 210 mL
- d) 240 mL

25) A technician enters "digoxin 0.25 mg" as "2.5 mg." Which best prevents this error?

- a) Use trailing zeros (0.250 mg)
- b) Avoid leading zeros (.25 mg)
- c) Use leading zeros and avoid trailing zeros (0.25 mg)
- d) Round doses to the nearest whole number

26) The pharmacist asks you to identify which medication is a statin.

- a) Losartan
- b) Rosuvastatin
- c) Metformin
- d) Levothyroxine

27) A prescription is presented for alprazolam with 6 refills authorized. What is the correct next step?

- a) Fill with 6 refills because the prescriber authorized it
- b) Reduce to 5 refills and proceed without documentation
- c) Dispense only a 30-day supply and keep 6 refills on file
- d) Route for correction because Schedule IV is limited to 5 refills in 6 months federally

28) A patient's label prints with an NDC that does not match the bottle scanned at filling, though the drug and strength appear identical. What should the technician do?

- a) Ignore it if the imprint matches
- b) Reprint/adjust to the correct product/NDC before completion so the record matches the dispensed item
- c) Change the NDC after dispensing to save time
- d) Cover the NDC on the bottle to prevent confusion

29) Which auxiliary statement is most appropriate for nitroglycerin sublingual tablets for chest pain?

- a) "Swallow whole; do not crush"
- b) "Take with food to prevent nausea"
- c) "Store in the bathroom medicine cabinet for convenience"
- d) "Call 911 if chest pain continues after the first dose"

30) A technician is asked which product should be refrigerated before dispensing (unopened).

- a) Latanoprost ophthalmic solution
- b) Albuterol HFA inhaler
- c) Omeprazole DR capsules
- d) Amlodipine tablets

31) A patient says their bottle was stolen and requests an early refill of a noncontrolled maintenance medication. Insurance rejects "refill too soon." What is the technician's best next step?

- a) Select a lost-medication override code and rebill without documentation
- b) Gather details and route to the pharmacist to determine appropriate handling/override documentation
- c) Tell the patient it is impossible to refill early
- d) Change the days' supply to 1 so insurance pays

32) A prescription is received for "Plavix 75 mg daily." Which generic is correct?

- a) Clopidogrel
- b) Clonidine
- c) Cilostazol
- d) Carvedilol

33) A controlled substance prescription hardcopy is missing the prescriber's signature. What should be done?

- a) Fill it and obtain the signature at pickup
- b) Enter it and leave it in will-call pending signature
- c) Hold it for correction; it cannot be dispensed without a valid signature
- d) Ask the patient to sign for the prescriber

34) While counting tablets, a technician drops one onto the floor. What is the most appropriate action?

- a) Return it to the tray if it looks intact

- b) Place it back after wiping with alcohol
- c) Set it aside and continue counting; include it in the total
- d) Discard it according to procedure and replace from stock

35) A new eRx reads "methotrexate 2.5 mg: take 1 tablet daily." The indication field says rheumatoid arthritis. What should the technician do?

- a) Process as written; daily dosing is standard for RA
- b) Change directions to weekly and fill
- c) Route to the pharmacist due to high-risk dosing frequency concern
- d) Dispense a 7-day supply only

36) Rx: "Fluticasone nasal spray 50 mcg: 2 sprays each nostril daily." The bottle provides 120 sprays. What is the days' supply?

- a) 15 days
- b) 30 days
- c) 45 days
- d) 60 days

37) A technician is asked to identify a beta blocker for the drug class field. Which is a beta blocker?

- a) Amlodipine
- b) Lisinopril
- c) Metoprolol
- d) Hydrochlorothiazide

38) A coworker bypasses barcode scanning because it "slows the line down." What is the most appropriate technician response?

- a) Ignore it if the pharmacist will check later
- b) Speak up and escalate through the appropriate supervisor/pharmacist because it increases error risk
- c) Only scan controlled substances and skip the rest
- d) Scan the label, not the stock bottle, to save time

39) A patient requests transfer of a Schedule III prescription to another pharmacy. Under federal law (if state law allows), how many times can it be transferred?

- a) Zero; Schedule III can never be transferred
- b) Once
- c) Twice
- d) Unlimited

40) A patient profile has two patients with the same last name and first initial. The address differs by one digit and the DOB differs by one year. What is the best step to prevent wrong-patient entry?

- a) Select the profile that last filled a prescription
- b) Confirm two patient identifiers before selecting the profile for processing
- c) Merge the profiles to prevent duplicates
- d) Use only the address to choose

41) A prescription reads: "Take 2 tsp PO q6h PRN cough." How many mL is 2 tsp?

- a) 5 mL
- b) 10 mL
- c) 15 mL
- d) 30 mL

42) Which medication is an SSRI?

- a) Sertraline
- b) Bupropion
- c) Amitriptyline
- d) Diazepam

43) The pharmacy receives a manufacturer recall for a specific lot of losartan. What is the most appropriate technician action?

- a) Pull all losartan regardless of lot number and return it
- b) Quarantine only the affected lots and assist with identifying impacted dispensing records as directed
- c) Continue dispensing until replacements arrive
- d) Dispense recalled product first to reduce waste

44) A technician is asked which DEA form is used to order Schedule II medications in paper form.

- a) DEA Form 41
- b) DEA Form 106
- c) DEA Form 222
- d) DEA Form 363

45) Rx: "Lantus SoloStar: inject 24 units subQ nightly. Dispense 1 box." A box contains 5 pens of 3 mL each (100 units/mL). What is the days' supply?

- a) 31 days
- b) 45 days
- c) 62 days
- d) 75 days

46) A patient wants the "water pill" listed as Lasix on an old bottle. Which generic should you select?

- a) Furosemide
- b) Hydrochlorothiazide
- c) Spironolactone
- d) Torsemide

47) A technician is preparing a compounded suspension. The formula says "add purified water q.s. to 120 mL." What does this mean?

- a) Measure 120 mL water first, then add ingredients
- b) Add water as needed until the final volume is 120 mL
- c) Add exactly 120 mL total liquid including flavoring
- d) Add water only if the mixture becomes too thick

48) Which is a high-alert medication commonly associated with serious harm if mis-dosed?

- a) Insulin
- b) Loratadine
- c) Docusate
- d) Fluticasone nasal spray

49) A patient asks what brand name they might see for apixaban. Which is correct?

- a) Pradaxa
- b) Eliquis
- c) Xarelto
- d) Plavix

50) A technician receives a call from a prescriber's office asking for a patient's full medication list. The caller can only provide the patient's name, no DOB or address. What should the technician do?

- a) Provide the list since it's for treatment
- b) Provide only the last medication filled
- c) Request additional verification/appropriate authorization before disclosing PHI
- d) Provide only controlled substance names to be safe

51) While checking a refrigerator, you find a multidose vaccine vial with no puncture date/time documented. The vial has been opened. What should be done?

- a) Use it if the manufacturer expiration date is valid
- b) Quarantine it for evaluation because BUD cannot be verified
- c) Freeze it to preserve potency
- d) Move it to will-call so it gets used first

52) A prescription comes in for "Zoloft 50 mg daily." Which generic is correct?

- a) Fluoxetine
- b) Citalopram
- c) Paroxetine
- d) Sertraline

53) In a retail pharmacy, which controlled substances must be included in a DEA-required initial inventory?

- a) Only Schedule II
- b) Only Schedule III–V
- c) Schedule II–V on hand
- d) Only controlled substances received in the past 6 months

54) Rx: "Diclofenac 1% gel: apply 2 g to affected area BID." Tube size is 100 g. Approximately how many days will one tube last?

- a) 12.5 days
- b) 25 days
- c) 50 days
- d) 100 days

55) A technician is processing "bupropion XL 150 mg daily" but selects "bupropion SR 150

mg.” What is the best next step?

- a) Continue since both are extended-release products
- b) Change directions to twice daily and proceed
- c) Correct to XL before filling because SR and XL are not interchangeable
- d) Dispense SR but add an auxiliary label “do not take at bedtime”

56) A patient presents a Schedule V codeine cough syrup prescription dated 14 months ago with refills remaining. What is correct federally?

- a) It may be refilled if refills remain
- b) It is no longer valid; controlled substance prescriptions cannot be refilled after 1 year
- c) It can be refilled twice after 1 year
- d) It is valid for 2 years because it is Schedule V

57) Which medication is an ACE inhibitor?

- a) Losartan
- b) Amlodipine
- c) Lisinopril
- d) Metoprolol

58) A technician is about to dispense a medication labeled “for external use only,” but the sig on the label says “instill 2 drops in each eye BID.” What should be done?

- a) Dispense; eye products are external
- b) Change the sig to “apply to eyelids”
- c) Add “do not swallow” and dispense
- d) Stop and verify product selection because the dosage form/use likely does not match

59) A patient on an active warfarin profile has a new prescription for rivaroxaban with a note “switch therapy.” What is the most appropriate technician action?

- a) Inactivate warfarin and fill rivaroxaban
- b) Fill both and let the patient decide which to take
- c) Route to the pharmacist to confirm the switch and avoid duplicate anticoagulation
- d) Put rivaroxaban on hold until warfarin runs out

60) Rx: “Ventolin HFA: inhale 2 puffs q6h PRN.” Inhaler contains 200 actuations. At maximum labeled use, what is the days’ supply?

- a) 12 days
- b) 17 days
- c) 25 days
- d) 50 days

61) The pharmacist asks you to identify which of these is a benzodiazepine.

- a) Alprazolam
- b) Buspirone
- c) Zolpidem
- d) Hydroxyzine

62) A technician is receiving a new C-II shipment. Which action is required as part of standard controlled substance receiving practices?

- a) Store the C-II product in the OTC section to reduce theft
 - b) Leave the shipment sealed until the end of the week
 - c) Document receipt per pharmacy procedure and file the invoice in the designated controlled substance records
 - d) Remove all outer labels from the bottle before stocking
- 63) Order: "Heparin 25,000 units in 250 mL D5W." What is the concentration?
- a) 50 units/mL
 - b) 80 units/mL
 - c) 100 units/mL
 - d) 125 units/mL
- 64) A patient profile shows "easy-open cap requested." The medication arrives in manufacturer packaging with a child-resistant cap. What should the technician do?
- a) Dispense as is; manufacturer cap must remain
 - b) Repackage into an easy-open container if permitted and documented, maintaining proper labeling
 - c) Remove the child-resistant mechanism from the manufacturer bottle
 - d) Tell the patient to bring a tool to open it
- 65) Which medication is a loop diuretic?
- a) Furosemide
 - b) Hydrochlorothiazide
 - c) Spironolactone
 - d) Amiloride
- 66) A technician is verifying a prescriber's DEA number format on a controlled substance prescription. Which is correct format?
- a) 9 digits only
 - b) Two letters followed by seven digits
 - c) Seven letters followed by two digits
 - d) Two letters followed by five digits
- 67) A patient asks where to dispose of unused fentanyl patches. What is an appropriate technician response?
- a) Recommend flushing them immediately
 - b) Recommend cutting patches in half and throwing in the trash
 - c) Direct them to a DEA-authorized take-back option and offer pharmacist assistance for local resources
 - d) Tell them to keep them for future pain
- 68) A prescription for "omeprazole 20 mg daily" is being transferred out at the patient's request. What is the proper process?
- a) A technician may complete the transfer independently
 - b) A pharmacist must complete the transfer communication per requirements
 - c) Transfers are only allowed for controlled substances
 - d) The original prescription must be voided before transfer

69) Rx: "Dorzolamide-timolol drops: instill 1 gtt OU BID." Bottle size 10 mL. Assume 20 gtt = 1 mL. Approximate days' supply?

- a) 25 days
- b) 50 days
- c) 75 days
- d) 100 days

70) Which medication is used for hypothyroidism?

- a) Levothyroxine
- b) Metformin
- c) Allopurinol
- d) Acyclovir

71) A pharmacist asks which schedule is hydrocodone/acetaminophen (most products).

- a) Schedule II
- b) Schedule III
- c) Schedule IV
- d) Schedule V

72) A technician sees two strengths of potassium chloride ER tablets stored next to potassium chloride oral solution. What is the most effective way to reduce wrong-form selection?

- a) Store all potassium products in one bin
- b) Use separate storage locations with clear labels and barcode scanning at selection
- c) Remove auxiliary labels to reduce clutter
- d) Repackage all potassium products into identical containers

73) Rx: "Take 1.5 tablets daily." Quantity dispensed is #45 tablets. How many days' supply?

- a) 15 days
- b) 20 days
- c) 30 days
- d) 45 days

74) A patient is picking up isotretinoin. Which dispensing requirement is commonly associated with this medication?

- a) DEA Form 222 must be completed
- b) Medication Guide must be dispensed with the prescription
- c) Pseudoephedrine logbook entry is required
- d) It must be stored frozen until pickup

75) A prescription is written "Levothyroxine 0.075 mg daily." Which product strength matches?

- a) 7.5 mcg
- b) 75 mcg
- c) 0.75 mg
- d) 750 mcg

76) A technician finds a stock bottle with a torn label and unreadable NDC. Tablets cannot

be confidently identified. What should be done?

- a) Identify by color/shape and relabel
- b) Mix into an existing bottle that looks similar
- c) Quarantine and remove from use until identified/handled properly
- d) Place in back stock and use last

77) A patient presents for a pseudoephedrine purchase and has already hit the daily limit in the log. What should the technician do?

- a) Complete the sale as cash since limits apply only to insurance
- b) Reduce the quantity to a non-log amount and complete the sale
- c) Tell the patient to return tomorrow and sell it then
- d) Decline the sale because the system indicates the legal limit has been reached

78) A prescription reads: "Apply 1 patch transdermal q72h." Quantity dispensed is 10 patches. What is the days' supply?

- a) 10 days
- b) 20 days
- c) 30 days
- d) 72 days

79) Which medication is a calcium channel blocker?

- a) Atenolol
- b) Amlodipine
- c) Lisinopril
- d) Atorvastatin

80) In a community pharmacy, which items must be kept in a tamper-evident manner for OTC products (per federal tamper-evident packaging requirements)?

- a) All OTC products
- b) Only dietary supplements
- c) OTC products sold at retail to consumers (with certain exemptions) must have tamper-evident packaging
- d) Only prescription medications

81) A technician is asked to convert 0.5 g to mg. What is the correct value?

- a) 50 mg
- b) 500 mg
- c) 5,000 mg
- d) 0.5 mg

82) A prescription is entered for "clonazepam 0.5 mg: take 1 tab BID. Dispense #60." How many days' supply?

- a) 15 days
- b) 20 days
- c) 30 days
- d) 60 days

83) Which medication is an antiplatelet agent?

- a) Clopidogrel
- b) Apixaban
- c) Atorvastatin
- d) Metoprolol

84) A technician receives a written prescription for a controlled substance that appears altered (different ink and mismatched handwriting in quantity). What is the best technician action?

- a) Fill it quickly to avoid confrontation
- b) Refuse and return it to the patient without documentation
- c) Continue processing and let insurance reject if needed
- d) Set it aside for pharmacist review before any dispensing steps continue

85) A technician is entering a prescription and sees "Sig: i tab po bid prn." What does "i" mean?

- a) 1
- b) 2
- c) 5
- d) 10

86) A patient is prescribed "atorvastatin 40 mg daily." The patient asks what condition it treats. What should the technician do?

- a) Provide full counseling on risks and benefits
- b) Tell the patient to read the package insert
- c) Provide a brief general use statement and offer pharmacist counseling for detailed questions
- d) Tell the patient it is for blood pressure

87) Which is a correct statement about Schedule II prescription transfers under federal law?

- a) They may be transferred once for the first fill
- b) They may be transferred only between chain pharmacies
- c) They may be transferred only if there are refills remaining
- d) They cannot be transferred

88) A prescription for "gabapentin 300 mg: take 1 capsule TID. Dispense #270." How many days' supply?

- a) 60 days
- b) 75 days
- c) 90 days
- d) 120 days

89) A technician is asked which medication is commonly used for gout.

- a) Allopurinol
- b) Acyclovir
- c) Alendronate
- d) Albuterol

90) A prescription is written for "tramadol" and dated 7 months ago with refills remaining.

Under federal rules, which is true for Schedule IV refills?

- a) Refills are permitted up to 5 times within 6 months; after 6 months it cannot be refilled
- b) Refills are permitted for 1 year if refills remain
- c) Refills are permitted indefinitely for Schedule IV
- d) Refills are never permitted for Schedule IV

Answer Key

1) c	31) b	61) a
2) b	32) a	62) c
3) a	33) c	63) c
4) a	34) d	64) b
5) b	35) c	65) a
6) d	36) d	66) b
7) b	37) c	67) c
8) c	38) b	68) b
9) b	39) b	69) b
10) c	40) b	70) a
11) b	41) b	71) a
12) d	42) a	72) b
13) a	43) b	73) c
14) d	44) c	74) b
15) a	45) c	75) b
16) c	46) a	76) c
17) a	47) b	77) d
18) b	48) a	78) c
19) c	49) b	79) b
20) c	50) c	80) c
21) b	51) b	81) b
22) a	52) d	82) c
23) b	53) c	83) a
24) c	54) b	84) d
25) c	55) c	85) a
26) b	56) b	86) c
27) d	57) c	87) d
28) b	58) d	88) c
29) d	59) c	89) a
30) a	60) b	90) a

PTCB Sim Test 04

1) During data entry, an eRx reads: "Levoxyl 112 mcg take 1 tablet by mouth daily. DAW: 0." Which product should be selected?

- a) Levoxyl 112 mcg with DAW 1
- b) levothyroxine 112 mcg tablet with substitution permitted
- c) liothyronine 25 mcg tablet
- d) levothyroxine 100 mcg tablet

2) A patient's eRx for Humalog KwikPen says: "Inject 8 units SQ before meals TID." How many mL are used per day?

- a) 0.08 mL
- b) 0.16 mL
- c) 0.24 mL
- d) 0.80 mL

3) A patient asks to buy pseudoephedrine 30 mg tablets, 96-count. The log shows they already purchased 2.88 g earlier today. What should the technician do?

- a) Process the sale because each transaction is under 3.6 g
- b) Process the sale and document "travel exception"
- c) Decline the sale because it would exceed the daily limit
- d) Split the sale into two transactions on different registers

4) While filling, the barcode scan shows the NDC pulled does not match the NDC on the label, but the drug name and strength appear the same. What is the best action?

- a) Reprint the label so it matches the scanned NDC
- b) Override the scan because the name/strength match
- c) Continue if the tablets look identical
- d) Stop and pull the exact product/NDC selected for the label

5) A prescription reads: "amoxicillin/clavulanate 600-42.9 mg/5 mL; give 7.5 mL PO BID x 10 days." How many mL should be dispensed?

- a) 75 mL
- b) 100 mL
- c) 150 mL
- d) 200 mL

6) A patient's profile shows "anaphylaxis to sulfa." A new prescription is received for sulfamethoxazole/trimethoprim. What should the technician do?

- a) Enter and fill because TMP-SMX is not a sulfonamide antibiotic
- b) Remove the allergy because it is nonspecific
- c) Route the allergy concern for pharmacist review before dispensing
- d) Change to nitrofurantoin because it is safer

7) A technician receives an eRx for "metoprolol succinate ER 25 mg daily." Which selection best matches?

- a) metoprolol tartrate 25 mg tablet

- b) metoprolol succinate ER 25 mg tablet
- c) carvedilol 3.125 mg tablet
- d) metoprolol succinate ER 50 mg tablet

8) A patient requests a refill of lorazepam that was originally written 8 months ago with refills remaining. Under federal law, what is correct?

- a) Refill is allowed if refills remain
- b) Refill is allowed only if dispensed as a 7-day supply
- c) Refill is not allowed because it is beyond 6 months from issue date
- d) Refill is allowed if the patient pays cash

9) A label reads: "Instill 1 gtt OD qhs." Which eye is OD?

- a) Left eye
- b) Right eye
- c) Both eyes
- d) Left ear

10) A prescription reads: "prednisone 20 mg: take 2 tablets daily x 5 days." Quantity needed is:

- a) 5 tablets
- b) 10 tablets
- c) 14 tablets
- d) 20 tablets

11) A patient presents a paper prescription for oxycodone IR with no prescriber signature. The patient says the office will "sign it later." What should happen?

- a) Fill it and store the hard copy until it's signed
- b) Dispense a partial and request signature for the remainder
- c) Call the prescriber, document verbal approval, and fill
- d) Do not dispense without a valid signed prescription

12) At pickup, two patients share the same last name and similar first names. What is the best verification step before handing over the bag?

- a) Ask for the address only
- b) Ask which doctor wrote it
- c) Confirm full name and date of birth against the bag label
- d) Ask the patient to identify the medication by appearance

13) A technician is entering "levothyroxine 0.025 mg daily." Which strength should be selected?

- a) 25 mcg
- b) 2.5 mcg
- c) 0.25 mcg
- d) 250 mcg

14) A refrigerated vaccine storage log shows 11°C (51.8°F) this morning. What should be done first?

- a) Use the vaccines today and adjust the thermostat

- b) Quarantine the vaccines and document the excursion for guidance
- c) Move vaccines to the freezer to cool them faster
- d) Discard all vaccines immediately

15) An eRx arrives: "Zofran ODT 8 mg take 1 tablet q8h prn nausea." The technician selects ondansetron 8 mg regular tablets (not ODT). What is the issue?

- a) Regular tablets are Schedule II
- b) ODT and regular tablets are different dosage forms that may affect use
- c) Regular tablets are a higher dose than ODT
- d) ODT is not allowed for generic substitution

16) A prescription reads: "latanoprost 0.005% instill 1 gtt OU qhs." Bottle size is 2.5 mL. Assume 20 drops/mL. Days' supply is:

- a) 10 days
- b) 25 days
- c) 50 days
- d) 100 days

17) A patient asks which generic matches Plavix.

- a) clopidogrel
- b) clonidine
- c) clozapine
- d) clomipramine

18) A technician receives a request from a prescriber's office to fax a patient's full medication profile. No patient authorization is on file. What is the best action?

- a) Fax it because it is for treatment
- b) Fax only the last 12 months
- c) Send only noncontrolled meds
- d) Route for pharmacist review to determine permitted disclosure/minimum necessary

19) A prescription says: "Apply 1 patch daily." The prescriber wrote "Dispense: 1 box" for lidocaine 5% patches (30 patches/box). What quantity should be entered?

- a) 1
- b) 10
- c) 12
- d) 30

20) A claim rejects "Refill too soon" for an inhaler the patient says was lost. What is the technician's best next step within workflow?

- a) Reverse the previous claim and rebill
- b) Submit a prior authorization request directly
- c) Check plan options (lost medication override/vacation supply) and gather required details
- d) Change the days' supply to 1 day to force approval

21) A prescription reads: "Ventolin HFA inhale 2 puffs q4h prn." Inhaler contains 200 actuations. If used at maximum frequency, days' supply is:

- a) 8 days

- b) 10 days
- c) 12 days
- d) 25 days

22) A Schedule II eRx is received. The patient asks to transfer it to a different pharmacy across town. Under federal rules, what is correct?

- a) Transfer is allowed one time if both pharmacies are in the same chain
- b) Transfer is allowed if it has not been filled
- c) Transfer is allowed with prescriber verbal approval
- d) It cannot be transferred; a new prescription is required at the receiving pharmacy

23) A prescription reads: "Take ii tablets PO TID." Quantity is 180 tablets. Days' supply is:

- a) 20 days
- b) 30 days
- c) 45 days
- d) 60 days

24) A patient's profile shows warfarin. A new prescription for TMP-SMX is entered today. An interaction alert appears. What should the technician do?

- a) Override the alert because the pharmacist will counsel
- b) Hold warfarin and dispense TMP-SMX only
- c) Route the interaction alert for pharmacist review before completion
- d) Delete TMP-SMX from the queue to prevent harm

25) A shipment arrives and the DSCSA transaction statement is missing for several totes. What is the best action?

- a) Accept the shipment; the invoice is sufficient
- b) Quarantine affected product and request missing documentation
- c) Create a transaction statement from the invoice
- d) Return only controlled substances because DSCSA is for controls

26) A prescription reads: "clindamycin 150 mg caps; take 2 caps QID x 7 days." Quantity needed is:

- a) 28
- b) 42
- c) 56
- d) 84

27) A technician finds a stock bottle containing mixed tablets (different imprints) with the same label. What should be done?

- a) Use an imprint guide and sort them back into stock
- b) Quarantine/remove from dispensing inventory for investigation/disposition
- c) Keep using the tablets that match the label and discard the rest
- d) Continue dispensing and rely on final verification to catch issues

28) A patient asks which medication is a proton pump inhibitor.

- a) famotidine
- b) omeprazole

- c) sucralfate
- d) dicyclomine

29) A prescription reads: "Take 2 tablets PO q6h prn pain; max 8 tablets/day." Quantity is 64 tablets. Maximum days' supply is:

- a) 6 days
- b) 7 days
- c) 8 days
- d) 10 days

30) A technician pulls glyburide 2.5 mg into the tray, but the label is for glimepiride 2 mg.

Best action:

- a) Dispense because both are sulfonylureas
- b) Change the label to glyburide
- c) Stop and retrieve the correct drug/strength matching the label
- d) Dispense and add "take with food" auxiliary label

31) A prescription: "valacyclovir 1 g PO BID x 10 days." Quantity needed is:

- a) 10
- b) 14
- c) 20
- d) 28

32) A controlled substance inventory count shows a C-II bottle with a broken seal and no record of opening. What should the technician do?

- a) Adjust the count to match the expected inventory
- b) Set aside and report to pharmacist/manager for investigation/documentation
- c) Dispense from that bottle first to minimize discrepancy
- d) Transfer contents to a new bottle and relabel

33) A patient asks which medication is an ACE inhibitor.

- a) losartan
- b) amlodipine
- c) lisinopril
- d) metoprolol

34) A prescription reads: "Take 10 mL PO q6h prn cough." Quantity is 180 mL. At maximum frequency, days' supply is:

- a) 3 days
- b) 4 days
- c) 4.5 days
- d) 7 days

35) A technician is processing "diltiazem ER 240 mg daily" but selected diltiazem IR 60 mg.

Best correction:

- a) Change directions to QID and continue
- b) Reselect the ER formulation as written before proceeding
- c) Keep IR and add "ER" to the label

d) Dispense IR and add a note for counseling

36) A written prescription for hydrocodone/APAP has patient name, drug, strength, directions, quantity, and prescriber signature but no date. What is required?

- a) Enter using today's date and fill
- b) Ask the patient for the date and write it on the Rx
- c) Fill and request a corrected copy later
- d) Obtain a properly dated prescription before processing

37) A patient asks which medication is a statin.

- a) rosuvastatin
- b) rivastigmine
- c) ropinirole
- d) rosiglitazone

38) A recall notice is received for a specific lot of atorvastatin. The technician finds that lot on the shelf. What should be done?

- a) Continue dispensing until replacement arrives
- b) Remove and quarantine the affected lot per recall instructions
- c) Repackage into amber vials and relabel with a new lot
- d) Return it to stock after checking expiration date

39) A prescription reads: "warfarin 2.5 mg daily." The technician selected warfarin 5 mg. Best action:

- a) Keep 5 mg and label "take 1/2 tablet daily"
- b) Change the sig to 1/2 tablet without contacting anyone
- c) Reselect the correct 2.5 mg strength as written
- d) Dispense 5 mg and ask the patient to split at home

40) Which law established the federal counseling/offer-to-counseling requirements tied to Medicaid?

- a) Kefauver-Harris Amendment
- b) OBRA '90
- c) FD&C Act of 1938
- d) Controlled Substances Act of 1970

41) A prescription reads: "Use 2 gtt OU QID." How many drops per day?

- a) 8
- b) 12
- c) 16
- d) 24

42) A technician receives a faxed Schedule II opioid prescription as a routine outpatient order (not an allowed fax-as-original exception). What is correct handling?

- a) Treat the fax as the original and dispense
- b) Convert it to a verbal order and document
- c) Use fax as reference only; obtain a valid C-II prescription before dispensing
- d) Dispense a 3-day emergency supply and request the original later

43) A patient asks which medication is a benzodiazepine.

- a) buspirone
- b) clonazepam
- c) bupropion
- d) carbamazepine

44) A prescription: "Augmentin 875/125 mg take 1 tab BID x 7 days." Quantity needed is:

- a) 7
- b) 10
- c) 14
- d) 21

45) An eRx includes "insulin regular U-500." The technician selected insulin regular U-100. Best technician action:

- a) Keep U-100 because U-500 is usually an error
- b) Change directions to multiply dose by 5
- c) Dispense U-100 and add a warning label
- d) Stop and ensure the correct concentration/product is selected before proceeding

46) A patient asks which medication is an SSRI.

- a) amitriptyline
- b) duloxetine
- c) bupropion
- d) sertraline

47) A printed patient leaflet with PHI is left on the counter visible to others. Best immediate action:

- a) Put it into will-call and say nothing
- b) Shred it immediately and document/report per incident process
- c) Fold it and place under the keyboard for later
- d) Hand it to the next customer to avoid waste

48) A prescription reads: "Take 1 tablet PO q8h." Quantity is 90 tablets. Days' supply is:

- a) 20 days
- b) 30 days
- c) 45 days
- d) 90 days

49) A patient asks which generic corresponds to Eliquis.

- a) rivaroxaban
- b) dabigatran
- c) apixaban
- d) edoxaban

50) A compounded oral suspension is requested, but the master formulation record is not available. What should the technician do?

- a) Compound using the previous label as the recipe

- b) Use an online recipe and add documentation later
- c) Substitute an OTC liquid with the same directions
- d) Pause compounding until required documentation is available

51) A prescription reads: "methotrexate 2.5 mg tablets; take 6 tablets daily" for rheumatoid arthritis. Best technician action:

- a) Enter as written because indication is listed
- b) Route for pharmacist review due to high-alert weekly dosing concern
- c) Change "daily" to "weekly" and process
- d) Reduce to 1 tablet daily to be safer

52) Which medication is a high-alert anticoagulant with significant risk if mis-dosed?

- a) loratadine
- b) omeprazole
- c) heparin
- d) fluticasone nasal spray

53) A prescription reads: "Lamictal 100 mg. DAW 1." What should the technician select?

- a) lamotrigine generic 100 mg
- b) Lamictal brand 100 mg with DAW required
- c) lamotrigine ER 100 mg
- d) Lamictal 25 mg and adjust quantity

54) A technician is restocking and notices hydrALAZINE and hydrOXYzine stored next to each other. Best risk-reduction step:

- a) Keep them adjacent to encourage careful reading
- b) Remove tall-man lettering stickers to reduce clutter
- c) Separate storage and rely on barcode scanning during fills
- d) Repackage both into identical amber vials for consistency

55) A prescription reads: "metformin ER 500 mg daily." The technician selected metformin IR 500 mg because it's cheaper. Best action:

- a) Substitute IR and change directions to BID
- b) Dispense IR with ER directions
- c) Reselect ER as prescribed; formulation changes require authorization
- d) Dispense both and let the patient choose

56) A patient asks which medication is a beta blocker.

- a) prazosin
- b) propranolol
- c) prednisone
- d) pregabalin

57) A prescription is for "Bactrim DS." Which components are correct?

- a) sulfamethoxazole/trimethoprim
- b) amoxicillin/clavulanate
- c) doxycycline/minocycline
- d) cephalexin/probenecid

58) A topical cream prescription reads: "Apply 1 gram BID." Quantity is 60 g. Days' supply is:

- a) 15 days
- b) 20 days
- c) 30 days
- d) 60 days

59) During order entry, directions say "take 1 tablet daily," quantity is #180, and a prescriber note says "30-day supply." What should the technician do?

- a) Change quantity to #30 to match the note
- b) Dispense #180 because quantity is the legal order
- c) Dispense #90 as a compromise
- d) Hold and seek clarification due to conflicting information

60) A patient's profile lists sildenafil. A new prescription arrives for nitroglycerin SL. Best technician step:

- a) Fill both because they are for different conditions
- b) Add "avoid alcohol" label and proceed
- c) Ensure the interaction is flagged for pharmacist review before dispensing
- d) Tell the patient to stop sildenafil immediately

61) A prescription reads: "Apply 2 patches every 72 hours." Carton contains 12 patches. Days' supply is:

- a) 12 days
- b) 18 days
- c) 24 days
- d) 36 days

62) A noncontrolled maintenance medication refill request comes in with zero refills remaining. What is the best next step?

- a) Add one emergency refill automatically
- b) Send a refill request to the prescriber per workflow
- c) Transfer the prescription to another pharmacy to generate refills
- d) Tell the patient therapy must stop until the office responds

63) A handwritten list from a patient shows "Celebrex 200 mg" but the technician reads it as "Cerebyx." Best action at intake to reduce LASA risk:

- a) Enter the closest match and let verification catch it
- b) Use strength, dosage form, and indication/patient history to confirm intended drug before entry
- c) Assume Celebrex because it is more common
- d) Enter both medications and let the patient decide at pickup

64) A prescription reads: "Take 0.5 mL IM every month." Concentration is 150 mg/mL. How many mg per dose?

- a) 50 mg
- b) 75 mg
- c) 100 mg

d) 150 mg

65) A will-call bag contains a vial label with a different patient name than the bag label. What should the technician do?

- a) Cross out the name and write the correct one
- b) Verify DOB matches and dispense if it does
- c) Separate items and send through the correction/verification workflow
- d) Leave it until the patient arrives to confirm

66) Which generic matches Klonopin?

- a) clonidine
- b) clonazepam
- c) clopidogrel
- d) clozapine

67) A patient asks which medication is an antiplatelet.

- a) clotrimazole
- b) colchicine
- c) clonidine
- d) clopidogrel

68) A prescription reads: "Take 2 tablets PO once daily." Quantity is #60. Days' supply is:

- a) 15 days
- b) 20 days
- c) 30 days
- d) 60 days

69) A prescription arrives for "Jardiance 10 mg daily #30." The technician selected Januvia 100 mg. Best technician action:

- a) Continue because both treat diabetes
- b) Reduce Januvia dose to be safer
- c) Stop and select empagliflozin 10 mg as written
- d) Dispense and label as a therapeutic interchange

70) A Schedule III–V prescription is being processed and the prescriber's DEA number is missing from the pharmacy's prescriber profile, but it is on the original prescription hard copy. Best action:

- a) Enter the DEA from the original prescription into the system record
- b) Dispense without DEA; it is only required for C-II
- c) Replace DEA with NPI
- d) Request a new prescription containing the DEA

71) A patient asks which medication is an ARB.

- a) lisinopril
- b) hydrochlorothiazide
- c) metoprolol
- d) losartan

72) A prescription: "benzonatate 100 mg; take 1 capsule TID prn cough." Quantity #30.

Days' supply (as directed) is:

- a) 5 days
- b) 7 days
- c) 10 days
- d) 15 days

73) A family member asks for a patient's pickup history, stating they "pay for the meds." No authorization is documented. Best response:

- a) Provide information because payment implies permission
- b) Provide only the total dollar amount
- c) Provide information if they know the DOB
- d) Do not disclose without proper authorization/verified permitted access

74) An eRx for gabapentin 300 mg TID and pregabalin 75 mg BID is received the same day for the same patient. Best technician step:

- a) Fill both because they work differently
- b) Delete one to avoid duplication
- c) Route for pharmacist review for potential duplicate therapy
- d) Change pregabalin to gabapentin for cost

75) A prescription reads: "insulin glargine (Lantus SoloStar) inject 20 units nightly." The technician selects Toujeo SoloStar. Best action:

- a) Dispense Toujeo because both are glargine
- b) Select Lantus as written because Toujeo is not interchangeable unit-for-unit
- c) Dispense Toujeo and adjust dose on the label
- d) Dispense Toujeo and add counseling note about conversion

76) A patient asks which product is an ICS/LABA maintenance inhaler.

- a) albuterol HFA
- b) ipratropium nasal spray
- c) budesonide/formoterol
- d) tiotropium HandiHaler

77) A prescription: "insulin lispro U-100 inject 10 units before meals TID." One pen contains 3 mL. How many days will 1 pen last?

- a) 7 days
- b) 8 days
- c) 9 days
- d) 10 days

78) A technician finds an unreconstituted antibiotic powder bottle that expired last month. What should be done?

- a) Reconstitute today because BUD resets after mixing
- b) Remove from usable inventory for proper disposition
- c) Use only for short courses
- d) Mix with a new bottle to restore potency

79) A prescription reads: "Take 1.5 tablets daily." Each tablet is 8 mg. How many mg per day?

- a) 10 mg
- b) 12 mg
- c) 16 mg
- d) 24 mg

80) A prescription label prints "nitroglycerin SL 0.4 mg." The technician has a bottle that was previously opened and used for counting earlier in the week. What should be dispensed?

- a) The previously opened bottle because stability is unchanged
- b) Tablets poured into an amber vial with a new beyond-use date
- c) A manufacturer's unopened container for the patient
- d) Any bottle as long as the expiration date is valid

81) A Schedule II prescription is partially filled because the pharmacy is out of stock. Under federal rules for a partial due to lack of supply, the remainder must generally be dispensed within:

- a) 72 hours
- b) 7 days
- c) 30 days
- d) 6 months

82) A Schedule III prescription is being transferred to another pharmacy (assume state law allows). Under federal rules, how many transfers are permitted for the prescription?

- a) None
- b) One transfer
- c) Two transfers
- d) Unlimited transfers until refills are used

83) A technician answers a phone call from a prescriber's office wanting to phone in lorazepam. What is the correct handling?

- a) Technician may take it if they read back the order
- b) Technician may take it only if another technician witnesses
- c) Transfer to the pharmacist to take the verbal controlled prescription
- d) Request it be emailed because phone orders are not allowed

84) A prescription reads: "Take 2 tablets PO BID." Quantity is #120. Days' supply is:

- a) 15 days
- b) 20 days
- c) 30 days
- d) 60 days

85) A patient asks which generic matches Xarelto.

- a) dabigatran
- b) apixaban
- c) edoxaban
- d) rivaroxaban

86) An eRx for “sertraline 50 mg daily” is active. A new eRx arrives for “fluoxetine 20 mg daily” with note “start today.” Best technician step:

- a) Fill both because they are antidepressants
- b) Discontinue sertraline in the profile and fill fluoxetine
- c) Route for pharmacist review for possible switch/duplication
- d) Change fluoxetine to escitalopram to avoid interactions

87) A prescription label reads: “Instill 1 gtt OS BID.” What does OS mean?

- a) Right eye
- b) Left eye
- c) Both eyes
- d) Left ear

88) A prescription reads: “morphine sulfate ER 30 mg.” The technician accidentally selects morphine oral solution 20 mg/mL. What is the primary concern?

- a) Both are equivalent because they contain morphine
- b) Different dosage form/route creates high risk of serious dosing error
- c) Oral solution is less potent, so it is acceptable
- d) ER tablets are not allowed for chronic pain

89) A patient requests to return an unopened refrigerated medication they kept at home so the pharmacy can put it back into stock. What is the correct action?

- a) Return to stock if the box is sealed
- b) Return to stock if the patient states it stayed refrigerated
- c) Do not return to stock; handle as a return per proper disposition process
- d) Return to stock after checking the expiration date

90) A prescription reads: “Take 2 mL PO daily.” Concentration is 10 mg/5 mL. How many mg per dose?

- a) 2 mg
- b) 4 mg
- c) 5 mg
- d) 10 mg

Answer Key

1) b	31) c	61) b
2) c	32) b	62) b
3) c	33) c	63) b
4) d	34) b	64) b
5) c	35) b	65) c
6) c	36) d	66) b
7) b	37) a	67) d
8) c	38) b	68) c
9) b	39) c	69) c
10) b	40) b	70) a
11) d	41) d	71) d
12) c	42) c	72) c
13) a	43) b	73) d
14) b	44) c	74) c
15) b	45) d	75) b
16) c	46) d	76) c
17) a	47) b	77) d
18) d	48) b	78) b
19) d	49) c	79) b
20) c	50) d	80) c
21) b	51) b	81) a
22) d	52) c	82) b
23) b	53) b	83) c
24) c	54) c	84) c
25) b	55) c	85) d
26) d	56) b	86) c
27) b	57) a	87) b
28) b	58) c	88) b
29) c	59) d	89) c
30) c	60) c	90) a

PTCB Sim Test 05

- 1) While typing an eRx for “metoprolol succinate ER 25 mg tablet,” the technician accidentally selects metoprolol tartrate 25 mg. What is the best next step before continuing?
- a) Keep the selection and change the frequency to BID
 - b) Correct the product selection to succinate ER and continue entry
 - c) Add “ER” in the patient directions to clarify
 - d) Leave it for pharmacist check at final verification
- 2) A patient’s friend calls asking, “Is Maria’s antibiotic ready, and what is it for?” The friend is not listed as an authorized representative. What can the technician disclose?
- a) The drug name only
 - b) The indication only
 - c) Only that information cannot be provided without authorization
 - d) The price and prescriber name only
- 3) A refrigerated shipment arrives with an insulin carton that feels warm and has no cold pack in the tote. The invoice shows it left the wholesaler yesterday. What should the technician do first?
- a) Place it in the refrigerator and dispense it before other stock
 - b) Quarantine the item and notify the appropriate supervisor/pharmacist for guidance/credit
 - c) Freeze it for 30 minutes, then refrigerate
 - d) Return it to shelf stock if the seal is intact
- 4) Rx: amoxicillin/clavulanate 600 mg/42.9 mg per 5 mL. Sig: 7.5 mL PO BID x 10 days. How many mL should be dispensed?
- a) 150 mL
 - b) 120 mL
 - c) 180 mL
 - d) 75 mL
- 5) A patient profile lists “ACE inhibitors—angioedema.” A new eRx arrives for lisinopril 10 mg daily. What should the technician do?
- a) Enter and bypass the alert because it’s prescriber ordered
 - b) Hold and route for pharmacist review due to recorded severe allergy
 - c) Change to losartan because it’s commonly substituted
 - d) Delete the allergy because it’s not a single drug name
- 6) A prescriber voicemail for alprazolam 0.5 mg includes patient name, directions, quantity, and prescriber DEA, but the patient address is missing. What is the appropriate next step?
- a) Process now and obtain address at pickup
 - b) Call the prescriber’s office to obtain the missing required information before dispensing
 - c) Enter using the pharmacy address temporarily
 - d) Convert it to a noncontrolled prescription so it can be filled
- 7) A claim rejects “Other coverage required.” The patient says they recently got new insurance but doesn’t have the card and also has a manufacturer savings card. What should

the technician do first?

- a) Run the savings card as primary
- b) Cash out and bypass insurance
- c) Update/confirm primary insurance and coordination of benefits information
- d) Change NDCs until it pays

8) During filling, the technician notices hydrOXYzine and hydrALAZINE are stored close together and look similar. Which control best reduces this selection error at the point of fill?

- a) Store both in the same bin so labels can be compared
- b) Use barcode scanning to verify the stock bottle matches the label
- c) Remove auxiliary labels so the name stands out
- d) Always pick the bottle with the earliest expiration date

9) A prescription is written for Eliquis. Which generic should be selected?

- a) rivaroxaban
- b) dabigatran
- c) apixaban
- d) edoxaban

10) A patient requests an early refill on oxycodone/APAP stating it was stolen. Which action is within technician scope?

- a) Refill if refills are available since it was stolen
- b) Decline immediately because early refills are never allowed
- c) Document the request details and route to the pharmacist for decision-making
- d) Call the prescriber and request a replacement controlled prescription

11) The refrigerator log shows 52°F this morning, and no one documented when the excursion started. Vaccines are stored in the unit. What should the technician do first?

- a) Move vaccines to the freezer to rapidly correct temperature
- b) Quarantine affected products and escalate for stability guidance before use
- c) Continue using them because 52°F is close to range
- d) Discard all vaccines immediately

12) Rx: prednisone 10 mg tabs. Sig: 4 tabs daily x 3 days, then 3 tabs daily x 3 days, then 2 tabs daily x 3 days, then 1 tab daily x 3 days. Quantity?

- a) 30
- b) 42
- c) 24
- d) 36

13) Under federal rules (if state law allows), a Schedule III prescription may be transferred:

- a) once between two pharmacies for the purpose of refill dispensing
- b) unlimited times until refills are used
- c) never; controlled prescriptions cannot be transferred
- d) only if the original pharmacy closes

14) A patient profile shows sertraline 100 mg daily active. A new eRx arrives for paroxetine 20 mg daily with no note indicating a switch. What should the technician do?

- a) Process both since they are in the same class
- b) Discontinue sertraline in the system and fill paroxetine
- c) Route for review due to potential therapeutic duplication
- d) Tell the patient SSRIs can't be changed and place on hold

15) Which product is typically stored refrigerated before dispensing (unopened)?

- a) clopidogrel tablets
- b) insulin glargine pens
- c) levothyroxine tablets
- d) omeprazole capsules

16) An eRx is for metformin ER 500 mg daily. The patient requests the "regular" version due to swallowing difficulty. What should the technician do?

- a) Change to metformin IR 500 mg daily and continue
- b) Enter as written and advise the patient to split/crush the tablet
- c) Route the request for pharmacist assessment and possible prescriber contact
- d) Reduce the dose to 250 mg so the tablet is smaller

17) A customer requests pseudoephedrine. Which is required federally under CMEA?

- a) Require a written prescription for all pseudoephedrine products
- b) Record the sale, require valid photo ID, and enforce daily/monthly limits
- c) Store pseudoephedrine only in the controlled substance safe
- d) Report each sale to the DEA within 24 hours

18) A new order arrives for methotrexate 2.5 mg with directions "Take 1 tablet daily," and the profile indicates rheumatoid arthritis. What should the technician do?

- a) Process as written because RA dosing is daily
- b) Change the sig to weekly because that is typical
- c) Route for review because daily dosing may represent a serious error
- d) Dispense and add "take with food" to improve tolerance

19) Order: vancomycin 1 g IV q12h for 5 days. Total grams needed?

- a) 12 g
- b) 20 g
- c) 5 g
- d) 10 g

20) Medication Guides must be provided:

- a) with each dispensing when required for that medication
- b) only when the patient requests one
- c) only for new prescriptions, not refills
- d) only for controlled substances

21) During entry, the technician selects valganciclovir instead of valacyclovir because the names look similar. What is the main risk?

- a) They are therapeutically equivalent so the risk is minimal
- b) Different uses/toxicity; wrong selection could cause significant harm
- c) Insurance will always reject so the patient is protected

d) Dosing is the same so it's unlikely to matter

22) An eRx is for ciprofloxacin 0.3% ophthalmic solution. The patient says it's "for an ear infection." What should the technician do?

- a) Change to the otic product and fill to match the complaint
- b) Fill as written and tell the patient it can be used in the ear
- c) Route for clarification because route/formulation may be incorrect
- d) Cancel the prescription because eye drops can never be used in the ear

23) A perpetual inventory count for a Schedule II medication is short by 6 tablets during a cycle count. What should the technician do?

- a) Adjust the on-hand count to match the computer to balance it
- b) Document the discrepancy and escalate immediately per diversion procedure
- c) Wait until the end-of-month inventory to see if it resolves
- d) Recount later and ignore if it's close

24) Rx: azithromycin 200 mg/5 mL. Sig: 12.5 mL PO daily x 3 days. How many mL should be dispensed?

- a) 60 mL
- b) 30 mL
- c) 75 mL
- d) 37.5 mL

25) Two patient profiles appear duplicated (same name and DOB; different addresses; different allergies listed). What is the best action before processing refills?

- a) Merge immediately to prevent billing issues
- b) Delete the older profile to reduce clutter
- c) Verify identifiers and route for proper merge/cleanup
- d) Ignore unless an insurance rejection occurs

26) A new Rx arrives for clopidogrel 75 mg daily. The profile shows omeprazole 20 mg daily. What is the best technician action?

- a) Fill both; interactions are not addressed during dispensing
- b) Flag for pharmacist review due to potential interaction reducing clopidogrel effect
- c) Tell the patient to stop omeprazole immediately
- d) Switch omeprazole to pantoprazole automatically

27) Lorazepam is classified federally as:

- a) Schedule IV
- b) Schedule II
- c) Schedule III
- d) Schedule V

28) Eye drops days' supply: Sig: 1 drop OU BID. Bottle size: 5 mL. Use 20 drops/mL. What days' supply should be billed?

- a) 25 days
- b) 50 days
- c) 10 days

d) 100 days

29) A voicemail for hydrocodone/APAP cuts off before the quantity is stated. What should the technician do?

- a) Use the last quantity filled and process
- b) Enter a typical quantity for the directions
- c) Obtain the missing quantity from the prescriber before dispensing
- d) Dispense a partial fill and complete when the office calls back

30) During unit-dose cart fill, morphine ER 30 mg and morphine IR 30 mg are stocked in adjacent bins. Which practice best reduces selection errors?

- a) Pull both and let the nurse decide what was ordered
- b) Use barcode scanning and visually verify ER vs IR before packaging
- c) Choose the product with the earliest expiration date
- d) Store them together so they can be compared side-by-side at pick

31) A patient requests a refill of nitroglycerin SL tablets last filled 14 months ago. The label states "discard 6 months after opening." What should the technician do?

- a) Process the refill and ensure pharmacist addresses replacement timing/storage
- b) Recommend using old tablets until finished
- c) Substitute nitroglycerin ointment to reduce replacement frequency
- d) Refuse because nitroglycerin cannot be refilled

32) DEA Form 222 is used for ordering:

- a) Schedule II controlled substances
- b) Schedule IV controlled substances
- c) Schedule III controlled substances
- d) Schedule V controlled substances

33) Rx is for Lamictal. Which indication best matches lamotrigine?

- a) hypothyroidism
- b) epilepsy/bipolar disorder
- c) benign prostatic hyperplasia
- d) bacterial pneumonia

34) A compound formula calls for 30% w/v. This means:

- a) 30 g in 100 mL of solution
- b) 30 mL in 100 mL of solution
- c) 30 g in 100 g of solution
- d) 30 mEq in 100 mL of solution

35) A patient picking up isotretinoin says they "missed their iPLEDGE window." What should the technician do?

- a) Sell it because the prescription is already verified
- b) Dispense but have the patient sign a waiver
- c) Route immediately because dispensing requires current REMS authorization/timing
- d) Tell the patient to return in 30 days and try again

36) Ventolin HFA: "Inhale 2 puffs q4h PRN." Inhaler contains 200 actuations. If used at the maximum frequency, what is the days' supply?

- a) 12 days
- b) 25 days
- c) 16 days
- d) 8 days

37) A patient returns tablets that smell strongly of vinegar and look discolored. What should the technician do?

- a) Restock if the NDC matches the current shelf stock
- b) Quarantine and escalate; do not return to stock
- c) Tell the patient it's normal for older tablets
- d) Replace from an open stock bottle and discard the returned tablets

38) A caregiver listed as an authorized representative calls and requests the medication name and last fill date for the patient's antidepressant. What should the technician do?

- a) Provide the information after confirming identity per procedure
- b) Provide only the medication name, not the date
- c) Refuse all disclosures over the phone
- d) Provide the prescriber's name instead

39) The technician enters levothyroxine but accidentally selects liothyronine due to a search result mix-up. Which step best prevents this error during data entry?

- a) Confirm drug, strength, and dosage form against the prescription image/hardcopy
- b) Turn off DUR alerts to reduce distractions
- c) Ask the patient at pickup to verify what they usually take
- d) Use brand names only during entry

40) For a written controlled-substance prescription, which element is required?

- a) patient diagnosis code
- b) prescriber signature
- c) patient weight
- d) pharmacist NPI

41) Augmentin corresponds to:

- a) cefdinir
- b) amoxicillin
- c) azithromycin
- d) amoxicillin/clavulanate

42) The label prints "Take 1 tablet twice daily," but the hardcopy shows "Take 1 tablet once daily." What should the technician do?

- a) Correct it and continue because it's clearly a typo
- b) Bag it and let counseling address it
- c) Stop and route the discrepancy for resolution before dispensing
- d) Print a label with both directions listed

43) A patient requests a printed medication profile at the register while other customers are

nearby. What is the best technician response?

- a) Print immediately and hand it over to keep the line moving
- b) Verify identity and provide it in a privacy-protective manner
- c) Refuse because profiles can never be printed
- d) Read the profile aloud to confirm it before printing

44) Rx: cephalexin 250 mg/5 mL. Sig: 10 mL PO TID x 7 days. Quantity to dispense?

- a) 280 mL
- b) 200 mL
- c) 210 mL
- d) 140 mL

45) Which is considered a high-alert medication that commonly requires extra safeguards?

- a) docusate
- b) famotidine
- c) loratadine
- d) heparin

46) A patient presents a paper prescription for Adderall dated 45 days ago. Federally, which statement is correct?

- a) Schedule II prescriptions expire after 30 days
- b) Schedule II prescriptions expire after 60 days
- c) There is no federal expiration date; state law/policy may restrict filling
- d) Schedule II prescriptions may be refilled within 6 months

47) A prescriber wrote atorvastatin 30 tablets with 2 refills. Insurance requests a 90-day supply today. What should the technician do?

- a) Change quantity to 90 and refills to 0 without contacting anyone
- b) Submit three 30-day claims on the same date
- c) Dispense 90 and write "insurance request" on the hardcopy
- d) Route for authorization/clarification before changing quantity/refills

48) An eRx for valsartan includes "switch from losartan." An active losartan Rx remains on the profile. What is the best technician action?

- a) Fill both so the patient can choose
- b) Inactivate losartan on the profile per the prescriber note and proceed with valsartan
- c) Replace valsartan with losartan since it's already on file
- d) Transfer losartan to another pharmacy

49) Ritalin LA corresponds to which generic?

- a) dexmethylphenidate
- b) methylphenidate
- c) lisdexamfetamine
- d) amphetamine/dextroamphetamine

50) After counseling, a patient asks the technician if alcohol is safe with metronidazole. What should the technician do?

- a) Advise avoiding alcohol during therapy and for 72 hours after the last dose

- b) Advise alcohol is safe if spaced 2 hours from the dose
- c) Refer the question to the pharmacist
- d) Recommend taking metronidazole with milk before drinking

51) A DSCSA alert flags a potentially suspect lot number currently on the shelf. What should the technician do first?

- a) Continue dispensing until a recall is issued
- b) Quarantine the affected lot to prevent dispensing and initiate the internal process
- c) Remove the lot number sticker from the shelf label
- d) Send it back to the wholesaler immediately without documentation

52) A new warfarin prescription is written "Take as directed" for a new patient, with no dosing schedule included. What should the technician do?

- a) Enter "UD" and fill; counseling will cover dosing
- b) Route because directions are incomplete for safe dispensing
- c) Convert to "Take 1 tablet daily" to avoid delay
- d) Dispense with extra warning labels and proceed

53) In the sterile compounding area, the hood's certification date is past due. What should the technician do?

- a) Proceed; certification is optional if the hood is cleaned
- b) Compound only antibiotics in the hood and avoid high-risk CSPs
- c) Stop sterile compounding and escalate before the hood is used
- d) Work quickly to reduce contamination risk

54) Suboxone contains:

- a) buprenorphine/naloxone
- b) buprenorphine/methadone
- c) naltrexone/naloxone
- d) morphine/naloxone

55) Convert 0.5 g to mg.

- a) 50 mg
- b) 500 mg
- c) 5,000 mg
- d) 500,000 mg

56) In a state where gabapentin is treated as a controlled substance, how should refills be processed?

- a) Apply federal Schedule V rules nationwide
- b) Apply the state-specific controlled requirements in addition to federal rules
- c) Ignore the state rule because it isn't federally scheduled
- d) Treat it as Schedule II for maximum safety

57) An order is for potassium chloride ER tablets. The shelf has KCl ER 20 mEq tablets and KCl oral solution 20 mEq/15 mL. What should the technician do?

- a) Select the solution since mEq is equivalent
- b) Select the tablet and ensure dosage form matches the order

- c) Dispense whichever is in stock to avoid delays
- d) Ask the patient which form they prefer and dispense that

58) Metformin refill rejects "refill too soon." The patient states the prescriber increased the dose yesterday. What is the best next step?

- a) Use an override code to force payment
- b) Obtain a new prescription reflecting the updated directions and process accordingly
- c) Tell the patient to pay cash until refill date
- d) Split the claim into two smaller fills on the same day

59) An unlabelled bottle of tablets is found on the counter with unknown identity. What should be done?

- a) Restock based on imprint recognition by the technician
- b) Place it in return-to-stock to be handled later
- c) Quarantine it for pharmacist evaluation; do not restock
- d) Use it as a temporary loaner bottle

60) A prescription for furosemide 40 mg is entered as torsemide 40 mg. What type of error is this?

- a) wrong route
- b) wrong drug
- c) wrong patient
- d) wrong prescriber

61) A patient asks to transfer a Schedule II prescription from another pharmacy to yours. Federally, which is correct?

- a) It can be transferred once if refills remain
- b) It can be transferred if both pharmacies share a database
- c) It can be transferred only if the prescriber calls the receiving pharmacy
- d) It cannot be transferred; a new prescription is required

62) Insulin lispro U-100 vial 10 mL. Sig: Inject 8 units subcutaneously TID with meals. What is the days' supply for 1 vial?

- a) 41 days
- b) 12 days
- c) 25 days
- d) 125 days

63) A prescription requires REMS documentation before dispensing. Which task is most appropriate for a technician?

- a) Determine whether the patient clinically qualifies for therapy
- b) Complete required administrative steps and route to the pharmacist as required
- c) Adjust dosing to meet REMS requirements
- d) Approve dispensing once forms are signed

64) A patient profile documents OTC ibuprofen use. A new Rx is for meloxicam 15 mg daily. What should the technician do?

- a) Process normally because OTC use doesn't matter

- b) Suggest taking both for better pain control
- c) Remove ibuprofen from the profile to prevent alerts
- d) Flag for review due to duplicate NSAID risk

65) Which is a combination inhaler used for maintenance therapy?

- a) albuterol HFA
- b) ipratropium nasal spray
- c) fluticasone/salmeterol
- d) fluticasone nasal spray

66) The pharmacy receives a Class I recall for a specific lot of a blood pressure medication. What should the technician do first?

- a) Quarantine the affected lot and stop dispensing it
- b) Keep dispensing until replacement arrives
- c) Remove lot number information and continue dispensing
- d) Return only opened bottles; unopened bottles are safe

67) Sig: "Take ii capsules PO BID." How many capsules per day?

- a) 2
- b) 4
- c) 6
- d) 8

68) A patient says their amoxicillin looks different this month. The NDC is different from last fill. What should the technician do?

- a) Tell the patient not to take it because it's not the same medication
- b) Explain manufacturer variation and offer pharmacist verification/counseling
- c) Tell the patient color changes mean the dose is stronger
- d) Assure the patient no verification is needed

69) An eRx is for U-500 insulin, but the supplies line lists U-100 syringes. What should the technician do?

- a) Fill as written because syringes are interchangeable
- b) Substitute U-100 insulin to match the syringes
- c) Remove syringes and dispense only the insulin
- d) Route for review due to high-alert insulin concentration/device mismatch

70) Rx: doxycycline 100 mg. Sig: 1 tablet PO BID x 14 days. Quantity?

- a) 30
- b) 21
- c) 28
- d) 14

71) Law enforcement requests a patient's prescription records without a warrant, subpoena, or patient authorization. What should the technician do?

- a) Provide the records to avoid delaying an investigation
- b) Provide only controlled-substance fills
- c) Provide de-identified records with names removed

d) Decline and route the request for proper handling/verification

72) Which filing method is acceptable under federal rules?

- a) All controlled prescriptions must be filed separately from noncontrols
- b) Schedule II must be separate; Schedule III–V may be filed with noncontrols if readily retrievable
- c) Schedule III–V must be separate; Schedule II may be filed with noncontrols
- d) All prescriptions must be filed together with no special markings

73) In a hospital, a nurse calls the pharmacy with a new medication order for a patient that is not in the prescriber order system. What should the technician do?

- a) Enter it because nurses can issue medication orders
- b) Enter it as a verbal order and send immediately
- c) Ask another technician to confirm and then process
- d) Route it because orders must come from an authorized prescriber per facility process

74) Tamsulosin is most commonly used for:

- a) hyperlipidemia
- b) GERD
- c) BPH urinary symptoms
- d) migraine prevention

75) You need 240 mL of a diluted solution using a 1:3 ratio (1 part concentrate to 3 parts diluent). How many mL of concentrate are needed?

- a) 80 mL
- b) 60 mL
- c) 120 mL
- d) 180 mL

76) An eRx is for bupropion SR 150 mg, but the technician selects bupropion XL 150 mg because the patient previously took XL. What should the technician do?

- a) Continue; SR and XL are interchangeable
- b) Change the sig to once daily to match XL and fill
- c) Correct to the prescribed formulation before processing
- d) Split XL tablets to approximate SR release

77) While stocking, nitrofurantoin monohydrate/macrocrystals is found in the nitroglycerin section. What should the technician do?

- a) Leave it since both begin with “nitro-”
- b) Move it to the correct location to reduce selection error risk
- c) Put a warning sticker on it and keep it there
- d) Dispense it first to clear shelf space

78) A new Rx for lithium is received. The profile shows hydrochlorothiazide. What should the technician do?

- a) Fill normally; no meaningful interaction exists
- b) Discontinue hydrochlorothiazide in the system and fill lithium
- c) Flag for review due to increased lithium level/toxicity risk

d) Suggest the patient drink more water and proceed

79) A prescriber writes "1 tsp" for an oral liquid dose. How many mL is 1 tsp?

- a) 10 mL
- b) 2 mL
- c) 15 mL
- d) 5 mL

80) A patient requests a refill on a medication marked "discontinued—therapy changed, do not refill." What should the technician do?

- a) Refill if refills remain because it was previously authorized
- b) Create a new prescription entry to bypass discontinued status
- c) Tell the patient the prescriber is wrong and refill
- d) Route for review to confirm whether therapy should remain discontinued

81) A patient asks the technician whether buprenorphine/naloxone will show on a drug test. What should the technician do?

- a) Answer that it never shows up on any test
- b) Answer that it always shows as heroin
- c) Refer the question to the pharmacist
- d) Suggest skipping doses before the test

82) The pharmacy receives a faxed prescription for tramadol that lacks a prescriber signature. What should the technician do?

- a) Fill it because tramadol is noncontrolled
- b) Add the prescriber name from the directory and proceed
- c) Obtain proper authorization (valid signature/requirements) before dispensing
- d) Treat it as valid since the fax number is known

83) A prescription is for oral vancomycin capsules, but the technician selects vancomycin injection during product selection. What should the technician do?

- a) Dispense injection and label "for oral use"
- b) Stop and correct the dosage form before proceeding
- c) Continue; vancomycin is the same regardless of dosage form
- d) Substitute metronidazole and fill

84) A profile shows simvastatin 40 mg daily. A new prescription arrives for clarithromycin. What should the technician do?

- a) Process normally; antibiotic interactions are rare
- b) Hold the simvastatin refill and fill the antibiotic only
- c) Change clarithromycin to azithromycin automatically
- d) Flag for review due to significant interaction risk

85) The pharmacy has expired controlled substances used for spill recovery. What is the correct next step?

- a) Dispose in regular trash after removing patient information
- b) Pour into the sink with water running
- c) Return to the wholesaler in the next order tote

d) Follow controlled-substance destruction procedure with required documentation/witnessing

86) A prescription is for a 30 g cream. Directions: apply 2 g daily. What is the days' supply?

- a) 10 days
- b) 20 days
- c) 15 days
- d) 30 days

87) A new eRx for digoxin is 0.25 mg daily; the patient history shows 0.125 mg daily for years. What should the technician do?

- a) Change to 0.125 mg because it matches history
- b) Fill 0.25 mg because the prescriber likely increased it
- c) Flag for review because digoxin is narrow therapeutic index/high risk
- d) Instruct the patient to take half a tablet

88) A neighbor calls asking whether another patient has picked up an antibiotic. What should the technician do?

- a) Confirm pickup status to be helpful
- b) Confirm only whether it was billed to insurance
- c) Ask the neighbor to bring the patient to the phone and then disclose
- d) Decline; do not disclose any protected information

89) A new eRx arrives for Januvia 100 mg daily. The profile shows Janumet 50/1000 mg BID active. What should the technician do?

- a) Process both; they work differently for diabetes
- b) Discontinue Janumet and fill Januvia only
- c) Change Januvia to Jardiance and fill
- d) Flag for review due to duplicate sitagliptin therapy risk

90) Rx: clindamycin 150 mg caps. Sig: Take 2 caps PO QID x 7 days. Quantity?

- a) 42
- b) 56
- c) 84
- d) 28

Answer Key

1) b	31) a	61) d
2) c	32) a	62) a
3) b	33) b	63) b
4) a	34) a	64) d
5) b	35) c	65) c
6) b	36) c	66) a
7) c	37) b	67) b
8) b	38) a	68) b
9) c	39) a	69) d
10) c	40) b	70) c
11) b	41) d	71) d
12) d	42) c	72) b
13) a	43) b	73) d
14) c	44) c	74) c
15) b	45) d	75) b
16) c	46) c	76) c
17) b	47) d	77) b
18) c	48) b	78) c
19) d	49) b	79) d
20) a	50) c	80) d
21) b	51) b	81) c
22) c	52) b	82) c
23) b	53) c	83) b
24) d	54) a	84) d
25) c	55) b	85) d
26) b	56) b	86) c
27) a	57) b	87) c
28) b	58) b	88) d
29) c	59) c	89) d
30) b	60) b	90) b

PTCB Sim Test 06

1) An eRx arrives: "metoprolol succinate ER 50 mg: take 1 tablet by mouth twice daily. #60." What is the best technician action?

- a) Enter as written and send to filling
- b) Change to metoprolol tartrate 50 mg to match twice-daily dosing
- c) Route for pharmacist review because ER succinate is typically dosed once daily and may need clarification
- d) Change directions to "take 2 tablets once daily" and proceed

2) A prescriber writes "Keflex 500 mg: 1 cap po qid x 7 days." What is the correct generic product to select?

- a) Ceftriaxone
- b) Cephalexin
- c) Cefdinir
- d) Cefuroxime

3) A patient asks to purchase pseudoephedrine. The amount is within daily and 30-day limits, and the patient presents a valid passport as ID. What should the technician do?

- a) Complete the sale and record the required purchaser and product information in the log/electronic system
- b) Refuse because only a state driver's license is acceptable
- c) Complete the sale but do not log because the amount is under the limit
- d) Refuse because passports are not allowed for OTC purchases

4) A prescription reads: "amoxicillin/clavulanate 600-42.9 mg/5 mL: take 7 mL po bid x 10 days." How many mL should be dispensed (round up to the nearest 5 mL)?

- a) 140 mL
- b) 150 mL
- c) 160 mL
- d) 175 mL

5) A hardcopy for hydrocodone/acetaminophen is presented. The date and patient address are present, but the prescriber's signature is missing. What should the technician do?

- a) Enter and fill; signature is only required for Schedule II
- b) Ask the patient to have the prescriber sign later and dispense today
- c) Hold the prescription and route for pharmacist handling because a signature is required for validity
- d) Accept if the prescriber name is printed and the DEA number is present

6) A patient's profile lists anaphylaxis to penicillin. An eRx arrives for amoxicillin 500 mg. What should the technician do?

- a) Process normally because allergies are patient-reported and may be inaccurate
- b) Change to azithromycin and continue
- c) Delete the allergy since prescriber ordered a penicillin
- d) Stop processing and route to the pharmacist for clinical assessment

7) A label direction reads: "Instill 1 gtt OS qid." Which is the correct interpretation of OS?

- a) Both eyes
- b) Left eye
- c) Right eye
- d) Both ears

8) A prescription is written: "prednisone 10 mg tabs: 4 tabs daily x 3 days, then 3 daily x 3 days, then 2 daily x 3 days, then 1 daily x 3 days." What quantity should be dispensed?

- a) 24 tablets
- b) 28 tablets
- c) 30 tablets
- d) 36 tablets

9) During receiving, a DSCSA scan returns "Product identifier not recognized." What is the best technician action?

- a) Stock it; scan failures are common and not meaningful
- b) Remove the 2D barcode label and scan only the NDC
- c) Quarantine the item as suspect and route to the appropriate person/workflow for investigation
- d) Dispense it first to avoid shortages, then investigate

10) An eRx reads: "Lantus SoloStar: inject 30 units subcutaneously at bedtime." No DAW is indicated. Which selection is most appropriate?

- a) Insulin glargine-yfgn (Semglee) pens
- b) Insulin aspart pens
- c) Insulin lispro pens
- d) Insulin glargine U-300 (Toujeo) pens

11) A patient arrives for pickup and asks what "take 1 tab po bid prn" means. What is the technician's best response?

- a) Take 1 tablet by mouth twice daily as needed
- b) Take 1 tablet by mouth daily at bedtime
- c) Take 2 tablets by mouth twice daily
- d) Take 1 tablet by mouth every 2 hours as needed

12) A prescriber sends an eRx for isotretinoin. The patient is new to therapy. Before dispensing, what additional process is required?

- a) No extra steps; it is a standard oral antibiotic workflow
- b) REMS/iPLEDGE requirements must be completed/confirmed prior to dispensing
- c) Only a HIPAA signature is needed
- d) The technician can dispense a 7-day emergency supply first

13) A patient requests an early refill for alprazolam. The last fill was 10 days ago for a 30-day supply. What should the technician do?

- a) Refill as cash because insurance timing is the only limitation
- b) Tell the patient it is always illegal to refill early
- c) Process as a partial fill without any review
- d) Route the request through the controlled-substance assessment/documentation workflow

for pharmacist review

14) Which brand/generic pair is correct?

- a) Zoloft / sertraline
- b) Celexa / fluoxetine
- c) Norvasc / losartan
- d) Lipitor / rosuvastatin

15) A prescription reads: "fentanyl 25 mcg/hr patch: apply 1 patch every 72 hours. Qty 10." What is the days' supply?

- a) 10 days
- b) 20 days
- c) 30 days
- d) 35 days

16) A new prescription: "methotrexate 2.5 mg: take 6 tablets by mouth once weekly." The patient's old directions in the profile show "6 tablets daily." What should the technician do?

- a) Keep the old directions for consistency
- b) Enter as once weekly and route for pharmacist verification due to high-alert frequency risk
- c) Replace directions with "take as directed" and continue
- d) Change to "take 3 tablets twice weekly" to reduce toxicity

17) A prescriber writes "metformin XR 500 mg: take 2 tablets daily with evening meal. #180." What is the days' supply?

- a) 45
- b) 60
- c) 90
- d) 180

18) A prescription is "azithromycin 250 mg: take 2 tablets once, then 1 tablet daily for 4 days." What quantity is required?

- a) 5
- b) 6
- c) 7
- d) 10

19) While restocking, a technician finds two different manufacturers' tablets mixed in one stock bottle labeled amlodipine 10 mg. What is the best action?

- a) Continue using it because the drug and strength are the same
- b) Separate by appearance and keep using both in the same bottle
- c) Pour into a new bottle and relabel with "mixed manufacturers"
- d) Remove from use and route for resolution per inventory quality workflow

20) A provider writes: "warfarin 5 mg: take as directed." The patient asks what dose to take tonight. What should the technician do?

- a) Tell them to take 1 tablet nightly until their next INR
- b) Ask for last INR and recommend a dose
- c) Refer the patient to the pharmacist for counseling and dosing clarification

d) Advise them to skip tonight's dose to be safe

21) A prescription reads: "Ventolin HFA: inhale 2 puffs q4h prn. 200 inhalations." If used at the maximum frequency, what is the days' supply?

- a) 8
- b) 12
- c) 16
- d) 25

22) A patient's spouse requests a printed medication list. The spouse is not listed as authorized, and the patient is not present. What should the technician do?

- a) Provide it if the spouse knows the date of birth
- b) Provide only non-controlled medications
- c) Decline to release without patient authorization
- d) Provide it but omit directions and quantities

23) Which DEA form is used to order Schedule II controlled substances from a wholesaler?

- a) DEA Form 41
- b) DEA Form 106
- c) DEA Form 222
- d) DEA Form 224

24) A pharmacy receives a call from another pharmacy requesting transfer of remaining refills for clonazepam. What is the appropriate response?

- a) Transfer immediately; any staff member may transfer Schedule IV refills
- b) Route to a pharmacist (or authorized individual per state law) to complete the controlled-substance transfer process
- c) Refuse; controlled substances can never be transferred
- d) Provide the prescription image by fax and consider it transferred

25) A patient returns unused oxycodone tablets and asks for a refund. What is the correct technician response?

- a) Accept the return and place back into stock if tablets are intact
- b) Accept the return only if the bottle is unopened, then restock
- c) Decline return for resale and provide information on authorized disposal options
- d) Accept and quarantine to re-dispense after visual inspection

26) A prescription reads: "Take 1 tablet po qod." How should qod be interpreted for safer labeling?

- a) Every day
- b) Four times daily
- c) Every other day
- d) Once weekly

27) A prescription is written for "diltiazem ER (CD) 240 mg daily." Which selection best matches?

- a) Diltiazem immediate-release tablet
- b) Diltiazem ER capsule (CD)

- c) Verapamil ER tablet
- d) Diltiazem topical gel

28) A multi-dose vaccine vial was first punctured 30 days ago. The manufacturer states "discard 28 days after first puncture." The vial has been refrigerated. What should the technician do?

- a) Use it because it was refrigerated
- b) Use it if it looks clear with no particles
- c) Move it to the front and use it today only
- d) Discard it and document per vaccine inventory procedure

29) An eRx arrives for "ondansetron 8 mg tablet by mouth q8h prn N/V." The system defaults to "ondansetron ODT 8 mg." What should the technician do?

- a) Accept the default because the strength is the same
- b) Change the route to SL to match ODT
- c) Select the dosage form that matches the prescription before processing
- d) Substitute promethazine due to cost

30) Which medication is a DOAC?

- a) Apixaban
- b) Warfarin
- c) Clopidogrel
- d) Aspirin

31) A technician notices "hydroxyzine 25 mg tid" entered, but the eRx is "hydralazine 25 mg tid." What is the best technician action?

- a) Leave it; pharmacist verification will catch it later
- b) Correct the selection before it proceeds, using LASA verification steps
- c) Dispense the entered drug and add a note for the patient
- d) Change to "take as directed" to minimize risk

32) A prescription reads: "Take 5 mL po q6h prn fever." The bottle is 120 mL. If used around-the-clock, what is the maximum days' supply?

- a) 4
- b) 5
- c) 6
- d) 8

33) A patient has an active profile medication: OTC acetaminophen 500 mg "take 2 tablets three times daily." A new Rx is for hydrocodone/acetaminophen 5/325 mg q6h prn. What should the technician do?

- a) Process normally because OTC meds are irrelevant
- b) Delete the OTC acetaminophen automatically
- c) Route to the pharmacist due to cumulative acetaminophen risk
- d) Change to oxycodone without acetaminophen and continue

34) A prescription is for "ciprofloxacin 500 mg: 1 tab po bid x 7 days." Only 250 mg tablets are in stock. What is the best next step for a technician?

- a) Dispense 250 mg tablets and change directions to "2 tabs bid" without review
- b) Dispense 250 mg tablets and adjust quantity only
- c) Cancel and tell the patient to go elsewhere
- d) Route for pharmacist review because changing strength/directions requires authorization

35) A patient wants to pick up a prescription for "my husband," but cannot provide address or date of birth. What should the technician do?

- a) Release it if it's not a controlled substance
- b) Ask only for last name and proceed
- c) Request adequate identifiers/authorization before releasing or discussing details
- d) Provide the medication name but not the quantity

36) A prescription reads: "latanoprost 0.005%: instill 1 gtt OU qhs." Bottle size is 2.5 mL. Assume 20 gtt = 1 mL. What is the days' supply?

- a) 12
- b) 25
- c) 50
- d) 100

37) A manufacturer recall notice states "Lot XJ004 only" for losartan 50 mg. What should the technician do?

- a) Remove all losartan from shelves
- b) Quarantine only the affected lot and route to the recall workflow
- c) Continue dispensing until replacement arrives
- d) Return the product to the manufacturer without documentation

38) A prescription reads: "QVAR ReditHaler 80 mcg: inhale 2 puffs bid. 120 actuations." What is the days' supply?

- a) 15
- b) 20
- c) 30
- d) 60

39) A Schedule III prescription has refills remaining but was written 8 months ago. Under federal law, what is the correct action?

- a) Refill because Schedule III prescriptions are valid for 1 year
- b) Refill once as a courtesy and document
- c) Transfer it to restart the 6-month window
- d) Do not refill; it is beyond the 6-month refill window

40) A patient asks which OTC product to take for severe chest pressure with shortness of breath that began 20 minutes ago. What should the technician do?

- a) Recommend omeprazole and avoid acidic foods
- b) Recommend ibuprofen and rest
- c) Direct the patient to seek emergency care immediately and alert the pharmacist
- d) Recommend aspirin 81 mg daily and schedule a clinic visit

41) A prescription reads: "spironolactone 25 mg daily." The technician accidentally selected

spironolactone/HCTZ 25/25 mg. What should be done?

- a) Keep the combo; it may help blood pressure
- b) Correct the product selection to spironolactone-only before proceeding
- c) Reduce the dose to every other day and continue
- d) Dispense and add "drink extra water" auxiliary label

42) A patient's insulin pens were found left at room temperature overnight, about 12 hours, before being returned to the refrigerator. What should the technician do?

- a) Return to stock immediately; refrigeration fixes it
- b) Mark them "room temp ok" and dispense first
- c) Quarantine and route for a stability/disposition decision
- d) Discard all refrigerated medications from that shelf

43) Which medication is a proton pump inhibitor?

- a) Famotidine
- b) Omeprazole
- c) Sucralfate
- d) Aluminum hydroxide

44) A technician is asked to accept a new verbal prescription over the phone from a nurse for an outpatient antibiotic. What should the technician do?

- a) Take the order and enter it if it is not a controlled substance
- b) Take the order only if the nurse provides the prescriber's NPI
- c) Decline and route to the pharmacist/authorized personnel per practice setting rules
- d) Take the order and send it directly to filling to avoid delay

45) A prescription is written: "Nitrostat 0.4 mg SL." Which package size is most typical?

- a) Bottle of 25 tablets
- b) Bottle of 30 tablets
- c) Bottle of 60 tablets
- d) Bottle of 100 tablets

46) A patient is prescribed "Crestor 10 mg daily" with no DAW. What should be selected?

- a) Brand Crestor only
- b) Atorvastatin 10 mg because it is a similar statin
- c) Rosuvastatin 10 mg
- d) Hold until prescriber confirms generic substitution

47) A Schedule II prescription is presented as a photocopy. The patient says the original is "in the car." What should the technician do?

- a) Process from the photocopy if the signature is visible
- b) Call the prescriber and accept verbal confirmation in place of the original
- c) Fill a partial now and complete later with the original
- d) Decline to process from the copy and route for pharmacist handling

48) A patient is on simvastatin. A new eRx is received for fluconazole 150 mg "take once today." What should the technician do?

- a) Ignore because it is only one dose

- b) Route for pharmacist review due to interaction risk
- c) Remove simvastatin from the profile and fill fluconazole
- d) Reduce the fluconazole dose and continue

49) A prescription reads: "gabapentin 300 mg: 1 cap qhs x 3 days, then 1 cap bid x 3 days, then 1 cap tid thereafter." How many capsules are needed for the first 6 days?

- a) 6
- b) 9
- c) 12
- d) 18

50) A technician drops a tablet on the counter while counting. What should be done?

- a) Put it back if the counter was recently disinfected
- b) Return it to stock if it's not a hazardous drug
- c) Place it in a separate vial for later use
- d) Discard it per procedure and replace with a new tablet

51) A patient's eRx reads: "Eliquis 5 mg: take 1 tablet by mouth twice daily. #60." The profile shows active warfarin therapy with recent INR documentation. What should the technician do?

- a) Process both medications as active therapy and fill
- b) Enter and route for pharmacist review due to duplicate anticoagulation risk
- c) Discontinue warfarin on the profile and fill Eliquis
- d) Hold in data entry until the patient chooses one anticoagulant

52) A prescription reads: "Apply 1 patch every 72 hours." Quantity dispensed is 5 patches. What is the days' supply?

- a) 12
- b) 15
- c) 20
- d) 10

53) A patient asks whether they can cut a bupropion SR tablet in half to "make it easier to swallow." What should the technician do?

- a) Advise cutting is fine if swallowed quickly
- b) Advise crushing and mixing with applesauce
- c) Advise taking every other day instead
- d) Refer to the pharmacist for counseling; SR tablets generally should not be split/crushed unless product-specific guidance allows

54) A prescription for "lamotrigine 25 mg" is received without DAW. The system is set to dispense brand Lamictal by default. What is the best technician action?

- a) Keep brand because lamotrigine is "sensitive"
- b) Select generic lamotrigine per substitution rules unless DAW requires otherwise
- c) Dispense brand and add DAW to match
- d) Call the prescriber to confirm generic substitution is allowed

55) During data entry, an insulin prescription is written "inject 10 U subcutaneously nightly."

How should the technician enter the dose to reduce error?

- a) "10 U"
- b) "10.0 units"
- c) "10 units"
- d) "0.1 mL"

56) A patient asks to transfer a non-controlled maintenance medication with refills to another pharmacy. What should the technician do?

- a) Transfer it independently without pharmacist involvement
- b) Give the patient the Rx number and have the other pharmacy "pull it" without any transfer process
- c) Refuse because only controlled substances can be transferred
- d) Route the transfer request through the authorized transfer workflow for your pharmacy

57) A pharmacy shipment arrives with torn cartons and one bottle with a broken seal. What should the technician do?

- a) Stock all intact bottles and discard the broken one without documentation
- b) Re-seal the bottle and stock it
- c) Quarantine affected items and route for return/integrity review
- d) Stock everything after removing damaged cartons

58) Which medication is an ACE inhibitor?

- a) Losartan
- b) Lisinopril
- c) Amlodipine
- d) Metoprolol

59) A prescription reads: "clopidogrel 75 mg daily." The profile shows aspirin 81 mg daily active. What should the technician do?

- a) Delete aspirin because clopidogrel replaces it
- b) Process both without any flags because the combination is always appropriate
- c) Route for pharmacist review because dual antiplatelet therapy may be intentional but needs assessment
- d) Tell the patient to stop aspirin at pickup

60) A controlled substance is discovered missing during inventory reconciliation and is suspected theft. Which DEA form is used to report theft or significant loss?

- a) DEA Form 41
- b) DEA Form 106
- c) DEA Form 222
- d) DEA Form 224

61) A prescription reads: "valacyclovir 1 g: take 1 tab po bid x 7 days." What quantity should be dispensed?

- a) 7
- b) 10
- c) 14
- d) 21

62) A technician finds an unlabeled bottle of tablets with unknown identity on the shelf. What should be done?

- a) Use an online pill identifier and relabel it for stock
- b) Compare to pictures and use if it "looks right"
- c) Put it in the back until someone recognizes it
- d) Quarantine and route; do not use unidentified medication

63) A patient is picking up a controlled prescription and store policy requires photo ID and documentation. The person cannot provide ID but knows the patient's address. What should the technician do?

- a) Release it because address matches
- b) Release it if they can state the patient's date of birth
- c) Obtain required ID and documentation before release
- d) Refuse all third-party pickups for controlled substances in all cases

64) A prescription reads: "heparin 1,000 units/mL vials." The bin contains vials labeled "10,000 units/mL." What should the technician do?

- a) Stock them; nurses will adjust doses
- b) Place them behind the 1,000 units/mL to use later
- c) Move to the correct bin and continue without reporting
- d) Treat as a critical mismatch, segregate, and route immediately for correction

65) A prescription is "levothyroxine 75 mcg daily." Which patient counseling point should be directed to the pharmacist (not handled independently by the technician)?

- a) What the medication is for and how to take it with respect to food/other meds
- b) The price and expected pickup time
- c) Whether the bottle is child-resistant
- d) How to sign up for text alerts

66) A patient's prescription is "lithium 300 mg capsules," but the patient previously received lithium 300 mg tablets. What should the technician do?

- a) Substitute tablets because it's the same strength
- b) Substitute capsules because it's the same strength
- c) Open capsules and pour into a bottle to mimic tablets
- d) Route for pharmacist review because dosage form changes can affect product selection and counseling

67) A prescription is "Augmentin 875/125 mg: 1 tab po bid x 10 days." The patient requests the cheapest option. What should the technician do?

- a) Automatically change to amoxicillin 875 mg because it's cheaper
- b) Select generic amoxicillin/clavulanate if available and no DAW; otherwise follow the plan's formulary process
- c) Change to cephalexin 500 mg qid because it's cheaper
- d) Dispense brand only; generics are not equivalent for this drug

68) A technician is reconstituting an antibiotic suspension. The label says "Add 71 mL of water to yield 100 mL." The technician accidentally adds water to the final volume line first

and then adds powder. What should be done?

- a) Shake well and dispense; concentration will be close enough
- b) Add extra powder to correct it
- c) Add another 71 mL water to dilute properly
- d) Discard and restart using correct reconstitution steps

69) A patient brings a bottle back saying the tablets look different than last month. What is the first technician step?

- a) Reassure them that generics always change and it doesn't matter
- b) Exchange the medication for a different manufacturer immediately
- c) Compare the NDC/description dispensed in the system to the tablets and packaging, then escalate if inconsistent
- d) Tell the patient to stop taking it until they see the prescriber

70) A prescription reads: "Take 1.5 tablets by mouth daily." The sig field does not allow decimals. What should the technician do?

- a) Enter "take 1 tablet daily" and dispense extra tablets
- b) Enter "take 1 tablet alternating with 2 tablets daily" without review
- c) Change automatically to a liquid formulation
- d) Route for pharmacist input on the safest accurate directions for the label

71) Which medication is a benzodiazepine?

- a) Buspirone
- b) Clonazepam
- c) Hydroxyzine
- d) Quetiapine

72) A technician is asked to generate a medication history report for a patient who wants it emailed to their employer. What is the best action?

- a) Email it if the employer name is provided
- b) Fax it without consent since it's for the patient's benefit
- c) Provide it only after verifying and documenting the patient's authorization and using an approved secure method
- d) Refuse all requests for records because they are always protected

73) An eRx arrives for "insulin lispro U-200 KwikPen." The patient's prior fills were U-100 pens. What should the technician do?

- a) Substitute U-100 because the patient has used it before
- b) Substitute U-200 because it's the same insulin
- c) Adjust the dose to half and proceed
- d) Route for pharmacist review because concentration changes are high-risk

74) A patient is buying a dextromethorphan-containing cough product and appears younger than the store's age minimum for sale. What should the technician do?

- a) Complete the sale because it is OTC
- b) Ask for acceptable proof of age before completing the sale
- c) Sell only one bottle and proceed
- d) Suggest cough drops and refuse the sale

75) Which brand name matches atorvastatin?

- a) Crestor
- b) Zocor
- c) Lipitor
- d) Pravachol

76) A prescriber wrote "sertraline 50 mg daily." The patient profile shows escitalopram active and sertraline discontinued last month. What should the technician do?

- a) Discontinue escitalopram and fill sertraline
- b) Fill both; dual SSRI therapy is common
- c) Ask the patient which they prefer and fill that one
- d) Route for pharmacist review for duplicate therapy/transition plan

77) A prescription reads: "Codeine/guaifenesin syrup" and the hardcopy is missing the patient address. What should the technician do?

- a) Fill; patient address is optional
- b) Fill if the patient provides their address verbally and the technician writes it in
- c) Route for pharmacist handling because required elements for controlled prescriptions must be complete/validated
- d) Substitute an OTC cough product and proceed

78) A prescription reads: "Take 2 cap po tid." Which is the correct label instruction?

- a) Take 2 capsules by mouth two times daily
- b) Take 2 capsules by mouth three times daily
- c) Take 2 capsules under the tongue three times daily
- d) Take 2 capsules by mouth at bedtime

79) A technician is packaging a hazardous drug for unit-dose use. Which practice is most appropriate?

- a) Count on the main tray to save time
- b) Use bare hands for better control, then sanitize
- c) Use appropriate PPE and dedicated equipment/area as required
- d) Skip PPE if the stock bottle is sealed

80) A patient's prescription is "fluticasone/salmeterol 250/50: inhale 1 puff twice daily." Which product is appropriate to select?

- a) Advair Diskus
- b) Spiriva HandiHaler
- c) Symbicort
- d) Ventolin HFA

81) A prescription reads: "Apply 1 thin layer to affected area bid." The patient asks how many grams to apply each time. What should the technician do?

- a) Estimate a gram amount based on the tube size
- b) Tell the patient to use as much as needed until it feels covered
- c) Suggest using a kitchen measuring spoon
- d) Refer the patient to the pharmacist for counseling on appropriate amount/use

82) A patient asks for the “generic for Plavix.” What should the technician select?

- a) Clopidogrel
- b) Clonidine
- c) Clotrimazole
- d) Clopiramide

83) A prescription reads: “sertraline 100 mg #30: take 1 tablet daily.” The prescriber wrote DAW 1. What should be dispensed?

- a) Generic sertraline
- b) Brand Zoloft
- c) Any SSRI equivalent
- d) Split dose using two 50 mg generics

84) A prescription is “tamsulosin 0.4 mg: take 1 capsule daily after the same meal. #90.” What is the days’ supply?

- a) 30
- b) 45
- c) 60
- d) 90

85) A technician sees two patient profiles with similar names and same birth year. One profile has a severe allergy to sulfa. What is the best step to prevent wrong-patient processing?

- a) Proceed and rely on the patient to correct it at pickup
- b) Use at least two identifiers (e.g., full DOB and address/phone) before entering/filling
- c) Choose the profile with the most recent activity
- d) Merge the profiles to avoid confusion

86) Which medication is a sulfonylurea?

- a) Metformin
- b) Sitagliptin
- c) Glipizide
- d) Acarbose

87) A patient reports a serious adverse reaction and asks how to report it to the FDA. What should the technician do?

- a) Tell them to contact the manufacturer only
- b) Submit a VAERS report for any medication reaction
- c) Tell them to post it online so others are warned
- d) Provide MedWatch reporting information and route to the pharmacist as appropriate

88) A prescription is “albuterol 2.5 mg/3 mL neb: use 1 vial via nebulizer q4-6h prn. #1 box.” The product is 25 vials per box. If used every 4 hours around-the-clock, what is the days’ supply?

- a) 4
- b) 5
- c) 8

d) 10

89) A long-term care facility sends a request for an emergency supply of a Schedule II medication because the delivery is delayed. Under federal law, which statement is correct?

- a) A technician may authorize an emergency supply for any Schedule II
- b) A pharmacist may dispense an emergency quantity only with a properly authorized emergency oral Schedule II prescription, followed by a written prescription within the required timeframe
- c) A Schedule II can be faxed as a refill authorization without any follow-up
- d) Emergency supplies are allowed only for Schedule III–V

90) A prescription reads: "warfarin 2 mg: take 1 tablet daily. #45." The patient says they take "1.5 tablets" some days per clinic. The technician sees the current sig doesn't match the patient statement. What should the technician do?

- a) Change the sig to the patient's statement and proceed
- b) Leave as written; patient will follow the clinic anyway
- c) Update directions to "use as directed" and fill
- d) Route for pharmacist review to clarify directions and ensure accurate labeling

Answer Key

- | | | |
|-------|-------|-------|
| 1) c | 32) b | 63) c |
| 2) b | 33) c | 64) d |
| 3) a | 34) d | 65) a |
| 4) b | 35) c | 66) d |
| 5) c | 36) c | 67) b |
| 6) d | 37) b | 68) d |
| 7) b | 38) c | 69) c |
| 8) c | 39) d | 70) d |
| 9) c | 40) c | 71) b |
| 10) a | 41) b | 72) c |
| 11) a | 42) c | 73) d |
| 12) b | 43) b | 74) b |
| 13) d | 44) c | 75) c |
| 14) a | 45) a | 76) d |
| 15) c | 46) c | 77) c |
| 16) b | 47) d | 78) b |
| 17) c | 48) b | 79) c |
| 18) b | 49) b | 80) a |
| 19) d | 50) d | 81) d |
| 20) c | 51) b | 82) a |
| 21) a | 52) a | 83) b |
| 22) c | 53) d | 84) d |
| 23) c | 54) b | 85) b |
| 24) b | 55) c | 86) c |
| 25) c | 56) d | 87) d |
| 26) c | 57) c | 88) b |
| 27) b | 58) b | 89) b |
| 28) d | 59) c | 90) d |
| 29) c | 60) b | |
| 30) a | 61) c | |
| 31) b | 62) d | |

PTCB Sim Test 07

1. During final preparation, you notice the label reads “metoprolol succinate ER 50 mg” but the stock bottle on the counting tray is “metoprolol tartrate 50 mg.” What is the best technician action?

- a) Change the label to remove “ER” and continue
- b) Continue because both are metoprolol and the strength matches
- c) Recount using the tartrate stock bottle and note the substitution
- d) Stop the fill and send it for pharmacist check due to formulation mismatch

2. A patient asks to buy pseudoephedrine 30 mg tablets, quantity 96. Which step is required before completing the sale?

- a) Verify a government-issued photo ID and record the sale in the logbook/system
- b) Require a prescription for any pseudoephedrine purchase
- c) Refuse the sale because tablets may only be sold in 24-count packages
- d) Approve only if the patient is at least 21 years old

3. An e-prescription arrives: “Bactrim DS: take 1 tablet PO BID x 10 days.” Which product should be selected?

- a) Sulfamethoxazole/trimethoprim 400 mg/80 mg tablet
- b) Sulfamethoxazole/trimethoprim 800 mg/160 mg tablet
- c) Trimethoprim 100 mg tablet
- d) Sulfasalazine 500 mg tablet

4. While updating allergies, a patient says: “Codeine makes me vomit.” What is the best way to document this in the profile?

- a) Codeine — nausea/vomiting
- b) Codeine — anaphylaxis
- c) Opioids — rash
- d) Acetaminophen — nausea/vomiting

5. A prescription reads: “Augmentin 875/125 mg: take 1 tablet PO BID x 7 days.” How many tablets should be dispensed?

- a) 7
- b) 10
- c) 14
- d) 28

6. A patient is prescribed insulin lispro U-100: “Inject 6 units SC before meals TID.” The prescription is for one 10 mL vial. What days’ supply should be entered?

- a) 18 days
- b) 28 days
- c) 55 days
- d) 83 days

7. A new patient’s profile has no address. A controlled prescription is waiting to be processed and the prescriber’s message also lacks a patient address. What should the

technician do next?

- a) Use the pharmacy's address temporarily to allow processing
- b) Proceed and collect the address at pickup
- c) Update the patient profile with an address from a prior insurance claim
- d) Obtain and enter the patient's address before the prescription can proceed

8. A faxed prescription for clonazepam includes the prescriber's name, address, and signature, but no DEA number is listed. What is the best technician action?

- a) Enter and fill; a DEA number is never required for Schedule IV
- b) Pull the DEA number from the last controlled prescription and enter it
- c) Continue processing and leave the DEA field blank
- d) Send for pharmacist review to determine if prescriber identifiers are sufficient

9. The label prints "Take 1 tablet daily," but the scanned hard copy shows "Take 1 tablet twice daily." What should the technician do?

- a) Correct the sig in the system and proceed without reprinting
- b) Set aside for pharmacist verification because the entry does not match the original
- c) Dispense as the label printed so the patient follows the label
- d) Ask the patient at pickup which directions they intended to follow

10. A spouse requests a complete medication list for the patient "to organize pills." There is no authorization on file and the patient is not present. What should the technician do?

- a) Provide the list because the request is from a spouse
- b) Provide medication names only, without directions
- c) Decline and request the patient provide authorization or be present
- d) Provide the list if the spouse confirms the patient's date of birth

11. A prescription is written for "levothyroxine 0.075 mg." Which strength should be selected?

- a) 7.5 mcg
- b) 75 mcg
- c) 0.75 mg
- d) 750 mcg

12. A patient reports their oxycodone/APAP was stolen and requests a replacement. The prescription was filled 9 days ago for a 30-day supply. What should the technician do?

- a) Process a 3-day supply as an emergency fill
- b) Tell the patient to return with a police report before discussing anything
- c) Direct the patient to contact the prescriber for a new prescription and cancel the old one
- d) Refer the request to the pharmacist for assessment and next steps

13. Directions: "Use 2 sprays in each nostril once daily." The bottle contains 120 sprays. What is the days' supply?

- a) 15
- b) 20
- c) 30
- d) 60

14. While selecting “hydroxyzine 25 mg,” you notice “hydralazine 25 mg” nearby on the shelf. Which step most effectively prevents a wrong-drug selection error?

- a) Compare tablet color/shape to the last fill image only
- b) Scan the stock bottle barcode to confirm the correct NDC/drug/strength
- c) Count twice and initial the vial
- d) Pull both bottles to the counting area and choose later

15. A patient asks, “Is Jardiance the same as empagliflozin?” Which response is correct?

- a) No, Jardiance is canagliflozin
- b) Yes
- c) No, Jardiance is dapagliflozin
- d) Yes, but only the 25 mg strength is empagliflozin

16. Two patients in the queue have similar names (Jon Lee and John Lee) and the same last name. What is the best way to reduce a fill mix-up?

- a) Fill both prescriptions together to save time
- b) Use last name only since both are in the system
- c) Separate the work and confirm two identifiers before pulling and labeling
- d) Place both completed bags in the same will-call bin divider

17. A prescription reads: “Azithromycin 250 mg: take 2 tablets today, then 1 tablet daily on days 2–5.” Quantity to dispense is:

- a) 5
- b) 6
- c) 7
- d) 10

18. A new prescription for apixaban is received. The patient profile shows warfarin filled last week and still active. What should the technician do?

- a) Process both because overlap is common for anticoagulants
- b) Inactivate warfarin automatically and fill apixaban
- c) Hold apixaban until the patient confirms they stopped warfarin
- d) Route for pharmacist review due to potential duplicate anticoagulation

19. A prescription is written for tamsulosin 0.4 mg capsules. Which brand name matches?

- a) Uroxatral
- b) Flomax
- c) Rapaflo
- d) Proscar

20. A topical prescription reads: “Apply 1 gram topically BID. Dispense 60 grams.” What days’ supply should be entered?

- a) 15
- b) 20
- c) 30
- d) 60

21. A prescription for methotrexate 2.5 mg is received with directions “Take 6 tablets daily.”

The indication field shows rheumatoid arthritis. What should the technician do?

- a) Enter as written; RA doses often use multiple tablets
- b) Change "daily" to "weekly" because that is standard for RA
- c) Submit for pharmacist review due to high-risk frequency concern
- d) Ask the patient whether they prefer daily or weekly dosing

22. A manufacturer recall notice specifies: "Losartan 50 mg, lot AB1234 only." What is the most appropriate technician action?

- a) Remove all losartan from the shelf and return it
- b) Quarantine only the affected lot and notify the pharmacist
- c) Keep dispensing until the wholesaler sends a replacement
- d) Remove the lot sticker and place the bottles back on the shelf

23. Prescription: "Symbicort 160/4.5: inhale 2 puffs BID." The inhaler contains 120 actuations. What days' supply should be billed?

- a) 15
- b) 20
- c) 30
- d) 60

24. A new prescription for sertraline 50 mg daily arrives. The profile shows paroxetine 20 mg daily filled 5 days ago. What should the technician do?

- a) Fill sertraline and automatically discontinue paroxetine
- b) Fill both; two SSRIs can be used together for anxiety
- c) Ask the patient which one they want and fill that one
- d) Send to the pharmacist due to possible duplicate therapy/transition plan

25. A patient asks, "Is Crestor the same as rosuvastatin?" Choose the correct answer.

- a) No, Crestor is simvastatin
- b) Yes
- c) No, Crestor is atorvastatin
- d) Yes, but only the 10 mg strength is rosuvastatin

26. Amoxicillin 250 mg/5 mL: "Take 10 mL PO TID x 10 days." How many mL should be dispensed?

- a) 150 mL
- b) 200 mL
- c) 250 mL
- d) 300 mL

27. A prescriber's office calls in a verbal prescription for alprazolam. What should the technician do?

- a) Accept the verbal order and document it under the pharmacist's name
- b) Transfer the call to the pharmacist
- c) Accept only if the patient has filled it before
- d) Ask the prescriber to fax it and then enter it as a verbal

28. Warfarin 5 mg daily is received. The patient filled warfarin 2.5 mg "take as directed" 3

days ago. What should the technician do?

- a) Process the 5 mg and place a note for the patient at pickup
- b) Combine both strengths into one vial to avoid confusion
- c) Send for pharmacist review due to narrow therapeutic index and duplication risk
- d) Delete the 2.5 mg from the patient profile before filling

29. Which medication is commonly considered a high-alert medication requiring extra safeguards?

- a) Docusate
- b) Loratadine
- c) Heparin
- d) Amoxicillin

30. You receive a refrigerated vaccine shipment. The temperature indicator shows an out-of-range excursion during transit. What should the technician do?

- a) Store it in the refrigerator and use it first
- b) Discard immediately without documenting to prevent use
- c) Quarantine the shipment and alert the pharmacist for evaluation
- d) Accept the shipment if the vials are unexpired and appear intact

31. A patient asks if they can take sildenafil while using nitroglycerin SL PRN. What should the technician do?

- a) Advise separating doses by at least 2 hours
- b) Recommend using half a sildenafil tablet
- c) Tell the patient it is safe if nitroglycerin is only used PRN
- d) Refer the question to the pharmacist

32. An FDA-required Medication Guide is associated with a drug being dispensed. When must it be provided?

- a) On first fill only
- b) With each dispensing when required for that medication
- c) Only when the patient requests it
- d) Only for patients over age 65

33. A Schedule II e-prescription arrives with "Refills: 1." What should be done?

- a) Enter 1 refill since it was prescribed
- b) Change to 0 refills and send for pharmacist review of validity
- c) Ignore the refill field and continue without flagging
- d) Call the prescriber for verbal confirmation of the refill

34. Lantus (insulin glargine) U-100: "Inject 20 units SC at bedtime." Dispense 1 vial (10 mL). What is the days' supply?

- a) 25 days
- b) 40 days
- c) 50 days
- d) 30 days

35. A prescription is for prednisone 10 mg tablets. On the shelf you see predniSONE 10 mg

tablets and prednisolone oral solution. What is the best action at product selection?

- a) Select prednisolone since it is the active metabolite
- b) Select prednisone tablets and confirm NDC/strength match the label
- c) Select prednisolone solution and adjust the directions to match dose
- d) Choose whichever is in stock and note the substitution on the label

36. A prescription is for clopidogrel 75 mg daily. Which brand name matches?

- a) Brilinta
- b) Effient
- c) Plavix
- d) Coumadin

37. Which federal requirement applies to Schedule II inventory records?

- a) Must be performed monthly
- b) May be kept with other records only if filed alphabetically
- c) May be recorded in pencil only
- d) Must be kept separate from other records or be readily retrievable

38. An insurance claim rejects "refill too soon." The patient says the prescriber increased omeprazole from once daily to twice daily, but the prescription on file still shows once daily. What is the best technician action?

- a) Change the directions to BID and resubmit without documenting
- b) Submit a lost medication override code
- c) Refer to the pharmacist to assess and obtain updated prescription/override
- d) Tell the patient to purchase OTC and stop using insurance

39. Timolol ophthalmic: "Instill 1 gtt OU BID." What does OU mean?

- a) Left eye
- b) Right eye
- c) Both eyes
- d) Both ears

40. A prescription is for morphine sulfate ER 30 mg. The shelf has morphine IR 30 mg and morphine ER 30 mg. What is the best technician check before counting?

- a) Use whichever bottle is closer since the strength matches
- b) Verify ER dosage form and confirm the NDC/description match the label
- c) Choose IR and add an auxiliary label indicating extended release
- d) Select the bottle stored behind the other since it's likely correct

41. A hard copy prescription is marked "DAW 1." How should it be interpreted?

- a) Dispense generic substitution required
- b) Dispense brand as written; no substitution
- c) Patient requested generic substitution
- d) Use the lowest-cost equivalent

42. Metformin 500 mg: "Take 1 tablet BID with meals." The patient requests a 90-day supply. How many tablets are needed?

- a) 90

- b) 120
- c) 180
- d) 360

43. An e-prescription reads "Keflex 500 mg." Which generic is correct?

- a) Cefdinir
- b) Cephalexin
- c) Cefuroxime
- d) Ceftriaxone

44. A community pharmacy needs to dispose of expired Schedule II stock. Which option is appropriate under federal requirements?

- a) Return to the wholesaler on the next routine return invoice
- b) Mix with an undesirable substance and discard in regular trash
- c) Flush if not on the FDA flush list to avoid diversion
- d) Use a compliant reverse distributor process with required documentation

45. Two patient profiles appear identical for name and DOB but have different phone numbers and addresses. A new prescription is waiting. What is the safest technician action?

- a) Select the profile with the most recent fills
- b) Merge the profiles immediately to prevent future confusion
- c) Confirm additional identifiers before choosing the correct profile
- d) Process under one profile and correct after pickup

46. Albuterol HFA: "Inhale 2 puffs q4h PRN." The inhaler contains 200 actuations. If used at maximum frequency, what is the days' supply?

- a) 8 days
- b) 12 days
- c) 16 days
- d) 25 days

47. A patient asks you to teach them how to use a newly dispensed dry-powder inhaler. What is the correct technician response?

- a) Provide full counseling if you've seen the pharmacist do it before
- b) Demonstrate using a dose from the patient's inhaler
- c) Offer to have the pharmacist counsel the patient
- d) Tell the patient to call the prescriber for instructions

48. Valacyclovir 1 g: "Take 1 tablet PO BID x 10 days." What quantity should be dispensed?

- a) 10
- b) 14
- c) 20
- d) 28

49. A new prescription arrives for benazepril 20 mg daily. The profile shows lisinopril 20 mg daily as active. What should the technician do?

- a) Process both since they are in the same class
- b) Automatically discontinue lisinopril and fill benazepril

- c) Fill the newer prescription and delete the older one from the profile
- d) Route for pharmacist review for possible therapeutic duplication

50. A patient returns an opened bottle of non-controlled tablets and asks for credit. What is the appropriate technician action?

- a) Accept and return to stock if the tablets appear clean
- b) Accept only if the purchase was within 7 days
- c) Decline to restock and route for proper disposition per workflow
- d) Accept if the patient signs a waiver

51. Directions: "Take 1/2 tablet at bedtime." Quantity dispensed is #30 tablets. What is the days' supply?

- a) 15
- b) 30
- c) 45
- d) 60

52. Ondansetron ODT 4 mg: "Dissolve 1 tablet q8h PRN nausea." Quantity #24. What is the maximum days' supply?

- a) 6
- b) 8
- c) 12
- d) 24

53. A sealed injectable arrives from a secondary distributor. DSCSA paperwork is present, but the outer seal appears tampered. What should be done?

- a) Accept it if the transaction history is complete
- b) Place into inventory because tampering is only a manufacturer issue
- c) Quarantine as suspect product and alert the pharmacist for investigation
- d) Dispense it quickly to minimize loss if it expires

54. A patient profile lists "penicillin—anaphylaxis." A prescription arrives for amoxicillin/clavulanate. What should the technician do?

- a) Fill because clavulanate reduces allergy risk
- b) Bypass the alert if the prescriber ordered it
- c) Change to cephalexin and proceed
- d) Route for pharmacist review due to severe documented allergy

55. A syringe is labeled 100 mg/mL. The directions say "Inject 0.6 mL daily." How many mg is each dose?

- a) 6 mg
- b) 60 mg
- c) 100 mg
- d) 160 mg

56. A prescription is for Xarelto 20 mg daily. Which generic name matches?

- a) Rivaroxaban
- b) Apixaban

- c) Dabigatran
- d) Edoxaban

57. An isotretinoin prescription is received and the patient wants it filled immediately. What key dispensing requirement is associated with isotretinoin?

- a) Must be dispensed in non-child-resistant packaging
- b) Requires REMS authorization steps prior to dispensing
- c) Must be dispensed as a 90-day supply only
- d) Must be stored refrigerated after opening

58. A filled prescription bag is found on the counter without a patient label. The contents cannot be reliably matched to a specific patient. What should be done?

- a) Put it back in will-call; the patient will identify it
- b) Open and relabel it to the last patient served
- c) Return the medication to stock immediately
- d) Segregate it for pharmacist review and reprocess per procedure

59. A prescription is written for Vyvanse 40 mg #30 with "Refills: 3." What is the best next step?

- a) Enter 3 refills because the prescriber authorized it
- b) Enter as written but block early refills
- c) Send for pharmacist review because Schedule II prescriptions cannot be refilled
- d) Convert it to a non-controlled stimulant and proceed

60. A reconstituted oral antibiotic label must state "Refrigerate. Discard 14 days after reconstitution." What date should be placed as the discard date?

- a) The manufacturer expiration date on the unreconstituted bottle
- b) A standard 30-day discard date
- c) The date 14 days from the reconstitution date
- d) No discard date is needed if "refrigerate" is listed

61. Latanoprost 0.005%: "Instill 1 drop OU qhs." Dispense 2.5 mL. If 20 drops = 1 mL, what days' supply should be billed?

- a) 25
- b) 40
- c) 50
- d) 100

62. A patient requests a transfer of a Schedule III prescription to another pharmacy. Which statement is correct under federal requirements?

- a) Schedule III prescriptions cannot be transferred
- b) Transfers must be communicated directly between two licensed pharmacists
- c) A technician may complete the transfer if documented in the profile
- d) Transfer is allowed only by fax with the original prescription attached

63. A patient asks, "Is Zoloft the same as sertraline?" Choose the correct answer.

- a) Yes
- b) No, Zoloft is fluoxetine

- c) No, Zoloft is paroxetine
- d) Yes, but only the 25 mg tablet is sertraline

64. During entry, an interaction alert appears for nitroglycerin and tadalafil and both are active in the profile. What should the technician do?

- a) Override because one is PRN
- b) Remove nitroglycerin since it is PRN and continue
- c) Change tadalafil to sildenafil to avoid the interaction
- d) Route the alert to the pharmacist for evaluation

65. Doxycycline 100 mg: "Take 1 capsule PO BID x 7 days." What quantity should be dispensed?

- a) 7
- b) 10
- c) 14
- d) 21

66. Used prescription vials are being discarded. What is the best method to protect patient information?

- a) Remove labels and place vials in regular trash
- b) Black out only the patient name and discard
- c) Place vials in a bag and discard; labels are small and unreadable
- d) Destroy/remove PHI using a secure method (e.g., shredding labels or secured disposal)

67. A prescription is for diltiazem ER 240 mg daily. During entry, diltiazem IR 120 mg was selected. What is the best technician action?

- a) Change directions to twice daily and proceed
- b) Continue because total daily mg can be equivalent
- c) Change the product to ER 240 mg without notifying anyone
- d) Stop and send for pharmacist review due to formulation/strength mismatch

68. Clindamycin 75 mg/5 mL: "Take 10 mL PO TID x 10 days." How many mL should be dispensed?

- a) 150 mL
- b) 200 mL
- c) 250 mL
- d) 300 mL

69. A customer already purchased 2.6 grams of pseudoephedrine today. They now request 96 tablets of 30 mg. What should the technician do?

- a) Approve because the daily limit is 4 grams
- b) Approve because tablet sales don't count toward grams
- c) Deny the sale because it would exceed the 3.6 grams/day limit
- d) Deny only if the customer cannot show ID

70. A hard copy controlled-substance prescription appears altered (quantity overwritten). What should the technician do?

- a) Fill it because the prescriber signature is present

- b) Ask the patient to initial the change and proceed
- c) Reduce the quantity to a safe amount and fill
- d) Refer to the pharmacist for evaluation and prescriber verification

71. A prescription is written for ProAir HFA. Which generic name matches?

- a) Levalbuterol
- b) Albuterol
- c) Budesonide
- d) Ipratropium

72. While restocking an automated dispensing cabinet (ADC), which practice best reduces look-alike/sound-alike selection errors?

- a) Stock all strengths together so nurses can find them quickly
- b) Stock by manufacturer to keep packaging consistent
- c) Use barcode scanning and verify drug/strength/lot during stocking
- d) Place high-alert medications in the most accessible bins

73. A patient asks to transfer gabapentin 300 mg (TID, 5 refills) to another pharmacy. What should the technician do?

- a) Transfer it immediately because it is not Schedule II
- b) Tell the patient transfers are not allowed for nervous system drugs
- c) Print a copy for the patient to bring to the other pharmacy
- d) Route to the pharmacist to complete the transfer per requirements

74. Directions: "Take 2.5 mL PO daily." Concentration is 20 mg/5 mL. How many mg is each dose?

- a) 5 mg
- b) 10 mg
- c) 20 mg
- d) 40 mg

75. A medication being dispensed is flagged "Medication Guide required." How should it be handled?

- a) Provide the Medication Guide with the dispensed prescription
- b) Provide it only on the first fill
- c) Provide it only for patients with Medicaid/Medicare
- d) Provide it only if the patient requests written information

76. A technician enters hydrochlorothiazide 25 mg daily, but the prescriber wrote hydralazine 25 mg TID. Which checkpoint is designed to catch this before the patient receives it?

- a) Insurance adjudication
- b) Wholesaler invoice review
- c) Pharmacist verification against the original prescription and selected product
- d) The patient recognizing the tablet at pickup

77. A prescription reads: "Take 1 tablet PO BID. Dispense #180." What is the days' supply?

- a) 45
- b) 60

- c) 90
- d) 180

78. A pregnant patient asks if they should start a newly dispensed antibiotic. What is within technician scope?

- a) Advise them to stop until after delivery
- b) Recommend taking it only if fever occurs
- c) Refer the patient to the pharmacist for counseling
- d) Suggest splitting doses to reduce fetal exposure

79. A serious adverse event occurs with a non-vaccine prescription medication. Which FDA program is used for reporting?

- a) VAERS
- b) MedWatch
- c) DSCSA
- d) REMS

80. Prednisone 20 mg: "Take 1 tablet daily x 5 days, then stop." What quantity should be dispensed?

- a) 5
- b) 10
- c) 20
- d) 30

81. Which is an acceptable federal filing method for Schedule III–V prescription records?

- a) Must be filed separately from non-controlled prescriptions
- b) May be filed with non-controlled prescriptions if readily retrievable
- c) Must be destroyed after 30 days
- d) Must be stored in a locked safe separate from all other documents

82. A prescription for Lamictal includes "brand medically necessary." What is the best technician action?

- a) Select generic lamotrigine and proceed; notes don't affect billing
- b) Select any generic manufacturer and mark DAW 5
- c) Select brand and proceed without notifying anyone
- d) Send for pharmacist review because brand requirement affects selection and billing

83. An order reads: "Inject 1 mL IM once." The vial concentration is 50 mg/mL. How many mg is the dose?

- a) 25 mg
- b) 50 mg
- c) 100 mg
- d) 150 mg

84. A Schedule II order is received: the invoice lists 2 bottles, but only 1 arrived. What is the best action?

- a) Accept the order and adjust inventory later
- b) Record receipt as 2 because the invoice is the legal record

- c) Report and document the discrepancy immediately per controlled receiving procedures
- d) Ignore it if the missing bottle is low cost

85. A prescription is written for Cipro 500 mg. Which generic is correct?

- a) Ciprofloxacin
- b) Cephalexin
- c) Clarithromycin
- d) Clindamycin

86. A partial fill is needed for a Schedule II prescription due to insufficient stock. Which statement is correct federally?

- a) Partial fills are never permitted for Schedule II prescriptions
- b) Partial fills are permitted; the remainder must be dispensed within 72 hours when partial is due to lack of stock
- c) Partial fills are permitted; remainder may be dispensed anytime within 6 months
- d) Partial fills are permitted only if the patient pays cash

87. Tablets in a stock bottle are discolored and crumbling, but the expiration date is still valid. What should the technician do?

- a) Use tablets from the bottom of the bottle only
- b) Mix with tablets from a new bottle to improve appearance
- c) Dispense if the patient has used it before without problems
- d) Quarantine the bottle and alert the pharmacist for evaluation

88. A technician free-texted a drug name during entry, resulting in a misspelled selection that bypassed the drug file. What best practice prevents this?

- a) Continue using free-text because it is faster
- b) Use standardized drug file/NDC-linked selection and avoid free-text for drug names
- c) Enter the SIG first and choose the product later
- d) Rely on the patient at pickup to verify the medication name

89. Directions: "Inject 8 units SC before breakfast and 8 units before dinner." Dispense one 3 mL insulin pen (U-100). What is the days' supply for one pen?

- a) 9 days
- b) 18 days
- c) 30 days
- d) 37 days

90. An e-prescription for a Schedule II stimulant arrives for a new patient. The eRx is missing the prescriber's address. What should the technician do?

- a) Look up the prescriber online and enter an address to complete processing
- b) Convert it to a fax to make it valid
- c) Fill it and collect the address at pickup
- d) Send to the pharmacist to determine if required elements are met before dispensing

Answer Key

- | | | |
|-------|-------|-------|
| 1) d | 32) b | 63) a |
| 2) a | 33) b | 64) d |
| 3) b | 34) b | 65) c |
| 4) a | 35) b | 66) d |
| 5) c | 36) c | 67) d |
| 6) c | 37) d | 68) d |
| 7) d | 38) c | 69) c |
| 8) d | 39) c | 70) d |
| 9) b | 40) b | 71) b |
| 10) c | 41) b | 72) c |
| 11) b | 42) c | 73) d |
| 12) d | 43) b | 74) b |
| 13) c | 44) d | 75) a |
| 14) b | 45) c | 76) c |
| 15) b | 46) a | 77) c |
| 16) c | 47) c | 78) c |
| 17) b | 48) c | 79) b |
| 18) d | 49) d | 80) a |
| 19) b | 50) c | 81) b |
| 20) c | 51) d | 82) d |
| 21) c | 52) a | 83) b |
| 22) b | 53) c | 84) c |
| 23) c | 54) d | 85) a |
| 24) d | 55) b | 86) b |
| 25) b | 56) a | 87) d |
| 26) d | 57) b | 88) b |
| 27) b | 58) d | 89) b |
| 28) c | 59) c | 90) d |
| 29) c | 60) c | |
| 30) c | 61) c | |
| 31) d | 62) b | |

PTCB Sim Test 08

1. A new e-prescription arrives for losartan 50 mg once daily. The patient profile shows losartan 25 mg once daily filled 5 days ago with directions "take 1 tablet daily." The prescriber note states "increase dose." What should the technician do during processing?

- a) Inactivate the 25 mg prescription and fill the 50 mg as the new therapy
- b) Process the 50 mg prescription and route it for pharmacist review of dose change/duplicate therapy
- c) Combine both prescriptions into one label: "take 25 mg + 50 mg daily"
- d) Change the 50 mg to 25 mg so it matches the active prescription

2. A prescription is for amoxicillin/clavulanate 875/125 mg: take 1 tablet twice daily for 7 days. How many tablets should be dispensed?

- a) 7
- b) 10
- c) 14
- d) 21

3. A patient calls and asks, "Can you tell me what my spouse is picking up today?" The caller is not listed as authorized and cannot provide the patient's DOB. What can the technician say?

- a) The drug name and strength only
- b) Only that a prescription is or isn't ready for pickup, without medication details
- c) The prescriber name and directions, but not the drug name
- d) Full details if they can verify the address

4. While typing a new prescription, the sig shows "take 1 tablet daily" but the quantity and day supply requested by the prescriber match "take 1 tablet twice daily." What is the best technician action?

- a) Enter twice daily because the quantity suggests it
- b) Enter daily because the sig is the legal direction
- c) Enter "use as directed" to avoid conflict
- d) Enter as written in the sig and route for pharmacist review of the inconsistency

5. A prescription is written for "metformin ER 500 mg: take 2 tablets twice daily with meals." Quantity dispensed is #240. What is the days' supply?

- a) 30
- b) 45
- c) 60
- d) 120

6. A patient is prescribed insulin glargine U-100 pen: inject 28 units once daily. One pen contains 3 mL. One box contains 5 pens. What days' supply should be billed for one box?

- a) 16
- b) 19
- c) 26
- d) 53

7. During filling, the barcode scan alerts “wrong NDC selected.” The label is for levothyroxine 88 mcg, but the bottle pulled is levothyroxine 100 mcg. What should the technician do?

- a) Override the scan because levothyroxine strengths are close
- b) Change the directions to equal the intended daily dose
- c) Return the incorrect bottle and pull the correct strength, then rescan
- d) Continue filling and let the pharmacist catch it at final check

8. A new prescription arrives for lisinopril. The patient profile lists “pregnancy—2nd trimester.” What should the technician do?

- a) Process normally because pregnancy is not an allergy
- b) Route the prescription for pharmacist review before it is filled
- c) Remove the pregnancy note because it is not relevant
- d) Change lisinopril to labetalol and proceed

9. A prescription is for albuterol HFA: inhale 2 puffs every 4 hours as needed. The inhaler contains 200 actuations. Using the maximum frequency, what days’ supply should be billed?

- a) 12
- b) 16
- c) 20
- d) 25

10. At pickup, a patient asks, “Is Zoloft the same as sertraline?” What is an appropriate technician response?

- a) Yes, Zoloft is the brand name for sertraline
- b) No, sertraline is the brand name for Zoloft
- c) No, Zoloft is fluoxetine
- d) Yes, but only if taken at night

11. A prescription is for prednisone 10 mg: take 4 tablets daily for 3 days, then 3 tablets daily for 3 days, then 2 tablets daily for 3 days, then 1 tablet daily for 3 days. How many tablets should be dispensed?

- a) 24
- b) 30
- c) 36
- d) 40

12. A patient presents a hardcopy prescription for a Schedule II medication with the patient’s address missing. What is the correct technician action?

- a) Add the address from the patient profile and process
- b) Process it because the drug and quantity are present
- c) Hold and route for pharmacist action to obtain required information before dispensing
- d) Return it to the patient and tell them to write the address in

13. The pharmacy receives a request to transfer a patient’s remaining refills for alprazolam to another pharmacy. What is the appropriate handling?

- a) A technician may transfer because it is Schedule IV
- b) Transfer is prohibited for all controlled substances

- c) A technician may fax the original prescription to complete the transfer
- d) Route to the pharmacist because controlled-substance transfers require pharmacist involvement per law/policy

14. A patient asks which OTC product is best for new chest tightness and shortness of breath that began 30 minutes ago. What is the best technician response?

- a) Recommend an antacid first
- b) Recommend ibuprofen and rest
- c) Encourage immediate medical evaluation and involve the pharmacist promptly
- d) Recommend a decongestant to open the airways

15. An e-prescription for a Schedule II stimulant arrives with a written date missing the year. What should the technician do?

- a) Assume the current year and process
- b) Use the e-prescription received date as the written date
- c) Hold and route for pharmacist/prescriber clarification before dispensing
- d) Change it to a 30-day noncontrolled supply and proceed

16. A technician is preparing pseudoephedrine for sale. The system indicates the patient's requested amount would exceed the legal monthly limit. What should the technician do?

- a) Complete the sale because the patient is paying cash
- b) Reduce the quantity and skip the log entry
- c) Decline the sale for the amount that exceeds the limit
- d) Sell the product but document it later

17. A refrigerator temperature log shows 12°C for 5 hours overnight with vaccines stored inside. What should the technician do first?

- a) Continue using the vaccines because the temperature returned to normal
- b) Immediately return the vaccines to the wholesaler
- c) Separate and label the vaccines "Do Not Use" pending stability guidance
- d) Place the vaccines in the freezer to restore potency

18. A patient requests a non-child-resistant cap for their maintenance medication. What is required?

- a) The prescriber must write "non-safety cap" on the prescription
- b) The patient may request it; document the request per workflow and dispense accordingly
- c) Non-child-resistant caps are allowed only for patients over age 65
- d) Non-child-resistant caps may be used only for OTC products

19. A prescription is for warfarin 5 mg daily. The profile shows a new prescription for trimethoprim/sulfamethoxazole started today. What should the technician do?

- a) Fill both and place them in will-call
- b) Remove the antibiotic because it interacts with warfarin
- c) Add a note and route for pharmacist review of the interaction risk
- d) Tell the patient to skip warfarin while on the antibiotic

20. A prescription reads "azithromycin 200 mg/5 mL: take 12.5 mL on day 1, then 6.25 mL daily on days 2–5." What total volume should be dispensed?

- a) 37.5 mL
- b) 45 mL
- c) 50 mL
- d) 60 mL

21. While filing, a technician finds a controlled-substance invoice. Which information must be readily retrievable on receiving records?

- a) Patient copay and register ID
- b) Wholesaler name, drug name/strength, quantity received, and date received
- c) Prescriber NPI and diagnosis codes
- d) Patient address and signature log

22. A pharmacy receives a manufacturer recall notice for a specific lot of atorvastatin on the shelf. What should the technician do with the affected stock?

- a) Keep dispensing until the lot is depleted
- b) Remove from active stock and quarantine per recall procedure
- c) Relabel the bottles with a new beyond-use date
- d) Move the bottles to the back of the shelf for "use last"

23. A prescription is for clopidogrel 75 mg daily. The patient asks, "Is Plavix the same thing?" What can the technician answer?

- a) Plavix is the brand name for clopidogrel
- b) Clopidogrel is the brand name for Plavix
- c) Plavix is aspirin/dipyridamole
- d) Plavix is prasugrel

24. A prescription is for "diltiazem ER 240 mg once daily," but the product list contains multiple ER versions (CD, LA, XR) and the prescription does not specify. What is the best technician step?

- a) Choose any ER version; they are interchangeable
- b) Select the least expensive ER NDC and process
- c) Hold and route for pharmacist clarification of the intended formulation
- d) Substitute immediate-release divided doses to match the daily dose

25. A label is printed for "metoprolol succinate ER 50 mg once daily." The technician accidentally pulled metoprolol tartrate 50 mg. What should the technician do?

- a) Dispense tartrate but change the directions to twice daily
- b) Dispense tartrate and keep the directions the same
- c) Place the product in the bag and flag for pharmacist to fix
- d) Return the incorrect product and pull/scan the correct ER formulation

26. An order entry technician sees "U" written for units on a heparin order. What is the best action?

- a) Enter exactly as written
- b) Replace "U" with "units" in the entered directions
- c) Change the dose to mL to prevent confusion
- d) Delete the order because abbreviations are not allowed

27. A prescription reads "Take ii capsules PO tid." What does this mean?

- a) Take 1 capsule three times daily
- b) Take 2 capsules twice daily
- c) Take 2 capsules three times daily
- d) Take 3 capsules twice daily

28. A patient's profile shows "anaphylaxis to penicillin." A new prescription arrives for amoxicillin. What should the technician do?

- a) Fill as written and add an allergy auxiliary label
- b) Hold and route for pharmacist review before dispensing
- c) Change to cephalexin and proceed
- d) Remove the allergy because it may be outdated

29. A technician is restocking an automated dispensing cabinet and notices hydromorphone was placed in a pocket labeled morphine. What should the technician do?

- a) Leave it since both are opioids
- b) Remove and correct the stock placement, then report per medication safety process
- c) Change the label on the pocket to hydromorphone
- d) Ask nursing to verify before administration

30. A prescription is for ondansetron ODT 4 mg. On the shelf: ondansetron 4 mg tablets (non-ODT) and ondansetron ODT 8 mg. What should the technician do?

- a) Select ondansetron 4 mg tablets since strength matches
- b) Select ondansetron ODT 8 mg and instruct "take half"
- c) Route for pharmacist decision if ODT 4 mg is unavailable rather than substituting dosage form/strength
- d) Substitute promethazine tablets because it is anti-nausea

31. A prescription is "diclofenac 1% gel: apply 4 g to knee four times daily." Dispense 100 g. What days' supply should be billed?

- a) 6
- b) 8
- c) 10
- d) 12

32. A patient asks, "Is Norvasc the same as amlodipine?" What is appropriate for the technician to say?

- a) Norvasc is the brand name for amlodipine
- b) Amlodipine is the brand name for Norvasc
- c) Norvasc is lisinopril
- d) Norvasc is hydrochlorothiazide

33. A will-call bag is found with a label reading "take 1 tablet daily," but the prescription image states "take 1 tablet twice daily." What should the technician do first?

- a) Reprint the label and swap it without notifying anyone
- b) Pull the bag from will-call and route for correction before dispensing
- c) Wait to address it when the patient arrives
- d) Add an auxiliary label "may take twice daily"

34. A long-term care tote returns with open, mixed, unidentified tablets in a bag. What should the technician do?

- a) Identify by color/shape and return to stock
- b) Use an imprint guide and return to stock
- c) Repackage into a vial labeled "mixed meds"
- d) Dispose per procedure; do not return unidentified medications to inventory

35. A prescription is for "insulin aspart U-100 vial: inject 12 units before meals three times daily." One vial is 10 mL. What is the days' supply for one vial?

- a) 18
- b) 22
- c) 27
- d) 30

36. A patient states, "My levothyroxine tablets look different this month." What should the technician do first?

- a) Tell them generics vary and it's fine
- b) Verify the NDC/manufacturer dispensed versus prior fills and offer pharmacist counseling if needed
- c) Replace it with brand Synthroid automatically
- d) Tell the patient to stop taking it until the next refill

37. A paper prescription for amoxicillin is presented without a prescriber signature. What should the technician do?

- a) Fill it because it is not controlled
- b) Have the patient sign and initial it
- c) Hold and route to obtain a valid prescription before dispensing
- d) Use the signature on file from a previous prescription

38. A patient asks, "Is Xarelto the same as rivaroxaban?" What can the technician answer?

- a) Rivaroxaban is the brand name for Xarelto
- b) Xarelto is the brand name for rivaroxaban
- c) Xarelto is apixaban
- d) Xarelto is warfarin

39. A patient requests an early refill of clonazepam 12 days early, stating their medication was lost. What should the technician do?

- a) Process immediately as cash to avoid insurance restrictions
- b) Refuse because early refills are never allowed for controls
- c) Document "lost meds" and proceed without review
- d) Collect details and route to the pharmacist for assessment and next steps

40. A prescription is for nicotine 21 mg patches: apply 1 patch daily. Quantity dispensed is 28 patches. What is the days' supply?

- a) 14
- b) 21
- c) 28

d) 30

41. A patient profile printout is requested at the counter while other customers are nearby. What is the best approach?

- a) Print and hand it over immediately
- b) Offer a more private method (step aside/sealed envelope) and verify identity before providing
- c) Refuse because profiles cannot be printed
- d) Read the list aloud to confirm accuracy before printing

42. A technician is asked to take a new verbal prescription over the phone. What is the correct response?

- a) Accept it for noncontrolled prescriptions only
- b) Accept it if the prescriber's office is busy
- c) Decline and transfer the call to the pharmacist
- d) Accept it and have the pharmacist sign later

43. A prescription is "amoxicillin 250 mg/5 mL: take 10 mL three times daily for 10 days." What total volume should be dispensed?

- a) 200 mL
- b) 250 mL
- c) 300 mL
- d) 350 mL

44. A technician notices two profiles with the same name and similar birthdates. One has a documented NSAID allergy. The prescription being entered is for naproxen. What should the technician do?

- a) Choose the profile with the most recent fill date
- b) Confirm the correct patient using at least two identifiers before continuing
- c) Merge the profiles immediately and proceed
- d) Pick either profile and add a note about possible duplicate profiles

45. A pharmacy receives a DSCSA request from a trading partner for transaction documentation on a returned prescription drug product. What should the technician do?

- a) Ignore because DSCSA applies only to controlled substances
- b) Provide the patient receipt as the transaction record
- c) Provide/route the request for the required transaction information per DSCSA process
- d) Destroy the returned product to close out the request

46. A technician discovers a near miss: the wrong strength was selected but caught before labeling. What best supports quality assurance?

- a) Put the product back and do not document it
- b) Document the near miss in the pharmacy's QA system per procedure
- c) Delete and re-enter the prescription to remove evidence of the error
- d) Wait until the pharmacist requests documentation

47. A prescription is "methotrexate 2.5 mg: take 6 tablets weekly." During entry, it was mistakenly typed as "daily," but the technician catches it before printing. What should the

technician do?

- a) Leave daily because weekly dosing is uncommon
- b) Correct to weekly and ensure it is flagged for pharmacist review due to high-alert weekly dosing
- c) Print both directions so the pharmacist can choose
- d) Change the quantity to 6 tablets and keep daily directions

48. A prescription is "hydrocortisone 2.5% cream: apply thin layer twice daily." Dispense 30

g. If the patient uses 1 g per application, what days' supply should be billed?

- a) 10
- b) 12
- c) 15
- d) 30

49. A prescriber sends an e-prescription with "DAW 1," but the message text also says "generic ok." What should the technician do?

- a) Dispense brand because DAW 1 indicates brand required
- b) Dispense generic because the message says it's ok
- c) Split-fill brand and generic to satisfy both
- d) Route for pharmacist clarification before changing DAW selection

50. A prescription is for "Advair Diskus 250/50: inhale 1 puff twice daily." The device contains 60 inhalations. What days' supply should be billed?

- a) 15
- b) 20
- c) 30
- d) 60

51. A patient asks, "Is Prilosec the same as omeprazole?" What can the technician answer?

- a) Yes, Prilosec is the brand name for omeprazole
- b) No, Prilosec is pantoprazole
- c) No, omeprazole is the brand name for Prilosec
- d) Yes, but only if taken with food

52. A prescription reads: "Take 2 tsp by mouth three times daily." How many mL is one dose?

- a) 5 mL
- b) 7.5 mL
- c) 10 mL
- d) 15 mL

53. A patient prescription label prints with "take 1 tablet daily," but the prescription image clearly states "take 1 tablet twice daily." The medication is still in the filling area. What should the technician do?

- a) Correct the entry and reprint the label before it goes to will-call
- b) Bag it and fix it at pickup
- c) Add an auxiliary label "take twice daily"
- d) Change the quantity to match the incorrect directions

54. A pharmacy receives a request from law enforcement for a patient's prescription records, but no subpoena/court order or written authorization is provided. What should the technician do?

- a) Provide records immediately to avoid obstruction
- b) Provide only controlled-substance records
- c) Provide records if the officer shows a badge
- d) Route appropriately and release records only with proper legal authorization

55. A prescription is for levetiracetam 500 mg. The patient asks, "Is Keppra the same as levetiracetam?" What can the technician answer?

- a) Keppra is the generic name for levetiracetam
- b) Levetiracetam is the brand name for Keppra
- c) Keppra is the brand name for levetiracetam
- d) Keppra is carbamazepine

56. A prescription is for "Klor-Con M20." Which product should the technician select?

- a) Potassium chloride ER 20 mEq tablet
- b) Potassium chloride 20 mEq/15 mL oral solution
- c) Potassium citrate ER 20 mEq tablet
- d) Sodium chloride 1 g tablet

57. A unit-dose blister pack is torn and tablets are loose in the delivery tote. What should the technician do?

- a) Use the tablets if they appear intact
- b) Repackage into a new unit-dose blister and label it
- c) Place tablets into a vial for later use
- d) Remove from use and replace per procedure; do not dispense compromised unit-dose packaging

58. A patient brings an isotretinoin prescription, but the REMS status shows the patient is not within the allowable dispensing window. What should the technician do?

- a) Dispense with a Medication Guide only
- b) Dispense a partial supply until requirements are complete
- c) Hold the prescription pending completion/verification of REMS requirements before dispensing
- d) Bill as cash to bypass restrictions

59. A prescription is for epinephrine auto-injector 0.3 mg. Two boxes are available: one expires in 2 months and one in 14 months. What should be dispensed?

- a) The one expiring in 2 months to reduce waste
- b) The one expiring in 14 months, rotating stock appropriately
- c) Either is acceptable; expiration is not critical
- d) Both boxes and bill for one

60. Eye drops are prescribed: "instill 1 drop in each eye twice daily." Bottle size is 10 mL. Assume 20 drops/mL. What days' supply is one bottle?

- a) 25

- b) 50
- c) 75
- d) 100

61. A prescription reads “methylprednisolone 4 mg dose pack—use as directed.” What days’ supply should be entered for the standard dose pack?

- a) 5
- b) 6
- c) 7
- d) 10

62. A prescription is for valacyclovir 1 g: take 1 tablet three times daily for 7 days. What quantity should be dispensed?

- a) 14
- b) 21
- c) 28
- d) 42

63. A patient’s diagnosis field says “hypertension,” but the new prescription is for hydroxyzine 25 mg. The patient has a history of hydralazine fills. What should the technician do?

- a) Change hydroxyzine to hydralazine and proceed
- b) Enter hydroxyzine and route for pharmacist review due to possible sound-alike selection risk
- c) Delete the diagnosis field because it conflicts
- d) Ask the patient which drug they want

64. A prescription is for “doxycycline 100 mg capsules,” but only doxycycline 100 mg tablets are in stock. What should the technician do?

- a) Dispense tablets and change the label to “tablets”
- b) Dispense tablets without changing anything since strength matches
- c) Route for pharmacist determination of appropriate substitution before dispensing
- d) Cancel the prescription and tell the patient to call the prescriber

65. A prescription is for “alendronate 70 mg: take 1 tablet weekly.” Quantity is #12. What days’ supply should be billed?

- a) 28
- b) 56
- c) 84
- d) 90

66. A Schedule II prescription is presented with obvious overwriting in the quantity field. What should the technician do?

- a) Process it as written because it has a signature
- b) Fill a smaller quantity to reduce risk
- c) Refuse and return it to the patient immediately
- d) Set it aside and route for pharmacist evaluation/verification before processing

67. The pharmacy receives a faxed request for refills of a Schedule II medication. What is the correct technician action?

- a) Process it if refills are remaining
- b) Convert it to a new prescription and process
- c) Route for pharmacist handling because Schedule II refills are not permitted
- d) Transfer it to another pharmacy for completion

68. A prescription is for "tamsulosin 0.4 mg daily." The patient asks what it is used for. What should the technician do?

- a) Provide the indication and counseling points
- b) Tell the patient it is a blood pressure medication
- c) Offer to have the pharmacist counsel on what it's for
- d) Advise the patient to take it only as needed

69. A prescription is "test strips: test blood glucose four times daily." Quantity is 200. What days' supply should be submitted?

- a) 30
- b) 45
- c) 50
- d) 90

70. A prescription is "furosemide 20 mg: take 1 tablet by mouth daily as needed for swelling." Quantity is #30. The plan requires using the maximum implied daily use. What days' supply should be billed?

- a) 15
- b) 30
- c) 45
- d) 60

71. A patient arrives to pick up a prescription, but the person at the counter cannot confirm DOB or address and is not listed as authorized. What should the technician do?

- a) Release it if they share the last name
- b) Provide the medication name so they can confirm
- c) Release it but require a signature afterward
- d) Do not release or discuss details without proper identification/authorization

72. A sterile compounding technician notices an IV base solution bag is leaking at the port before any additives are injected. What should be done?

- a) Wipe the port with alcohol and proceed
- b) Tape the leak and proceed
- c) Discard the compromised bag and replace it
- d) Add the medication and ask nursing to transfer later

73. A prescription is "oxymetazoline nasal spray: 2 sprays each nostril twice daily for 3 days." What is the best technician action at entry?

- a) Remove the duration because it's OTC
- b) Enter as written and select the correct product/size
- c) Change to fluticasone for long-term therapy

d) Increase duration to 7 days to match common package size

74. A prescription is for "sertraline 100 mg #30" with 0 refills. The patient requests a one-year supply. What can the technician do?

- a) Add refills to allow a year supply
- b) Change quantity to #365 and dispense once
- c) Dispense as written and route the request for additional quantity/refills appropriately
- d) Refuse to fill unless a year supply is authorized

75. The pharmacist steps away and asks the technician to perform the final product verification for a waiting patient. What is the correct action?

- a) Perform the final check if the technician is certified
- b) Perform the final check for noncontrolled prescriptions only
- c) Decline unless working under an authorized tech-check-tech program for that setting
- d) Perform the final check only if the pharmacist will sign later

76. A hospital order is heparin 5,000 units subcutaneously every 8 hours. Multiple heparin concentrations are stored, including 10,000 units/mL vials. What is the best selection safety step for the technician?

- a) Choose the highest concentration to reduce volume
- b) Use barcode/NDC verification and compare concentration to the order before dispensing
- c) Stock all heparin together to speed selection
- d) Select any heparin; nursing will adjust the dose

77. A prescription is "clopidogrel 75 mg daily." The patient profile shows prasugrel 10 mg daily filled 3 days ago. What should the technician do?

- a) Fill clopidogrel and add a bleeding auxiliary label
- b) Fill both because they can be combined
- c) Discontinue prasugrel in the profile and fill clopidogrel
- d) Route for pharmacist review of possible therapeutic duplication

78. A technician is counting tablets and keeps two open stock bottles on the tray to "save time." Which practice best reduces selection error?

- a) Rely on tablet color and shape differences
- b) Keep both open bottles nearby for quick comparison
- c) Use barcode scanning and keep only one open bottle on the counting area at a time
- d) Remove NDC labels to reduce clutter

79. A patient asks, "Is Claritin the same as loratadine?" What can the technician answer?

- a) Claritin is the brand name for loratadine
- b) Loratadine is the brand name for Claritin
- c) Claritin is cetirizine
- d) Claritin is fexofenadine

80. A prescription is "benzonatate 100 mg: take 1 capsule three times daily as needed for cough." Quantity is #30. Using the maximum daily use, what days' supply should be billed?

- a) 7
- b) 10

- c) 14
- d) 30

81. During entry, a technician sees "MS" listed as the drug on a medication order with no further clarification. What should be done?

- a) Enter morphine sulfate
- b) Enter magnesium sulfate
- c) Route for clarification because "MS" is error-prone/ambiguous
- d) Enter both and let the pharmacist decide at verification

82. A prescription is "clindamycin 300 mg every 6 hours for 10 days." What quantity should be dispensed?

- a) 30
- b) 40
- c) 60
- d) 80

83. A patient with a latex allergy is scheduled for vaccination. What should the technician do?

- a) Proceed; latex is not relevant to vaccines
- b) Verify whether vaccine components/supplies contain latex and alert the immunizer/pharmacist
- c) Ask the patient to sign a waiver and proceed
- d) Use latex gloves to prevent contamination

84. A prescription is "mometasone nasal spray: 2 sprays in each nostril once daily." The bottle delivers 120 sprays. What days' supply should be billed?

- a) 15
- b) 20
- c) 30
- d) 60

85. A controlled substance is discovered in the wrong patient's will-call bag before pickup. What is the best technician action?

- a) Switch the bags quietly and continue workflow
- b) Open both bags and compare labels to decide what belongs where
- c) Pull both from will-call and route for pharmacist review while correcting the will-call placement error
- d) Leave them until the patient arrives to confirm

86. A prescription is "gabapentin 300 mg: take 1 capsule at bedtime for 3 days, then 1 capsule twice daily for 3 days, then 1 capsule three times daily thereafter." The prescriber writes to dispense #90. What is the best technician entry approach?

- a) Enter "take as directed" and dispense #90
- b) Enter the full titration directions and dispense #90
- c) Enter "take 1 capsule three times daily" because that is the maintenance dose
- d) Enter once-daily directions to match the starting dose

87. A prescription is "oseltamivir 75 mg: take 1 capsule twice daily for 5 days." What quantity should be dispensed?

- a) 5
- b) 10
- c) 14
- d) 20

88. A patient wants to transfer a Schedule II prescription to a different pharmacy across town. What is the correct handling?

- a) Transfer is allowed if both pharmacies share the same computer system
- b) A technician may transfer it by phone if the patient consents
- c) Schedule II prescriptions cannot be transferred; route to the pharmacist for patient guidance on next steps
- d) Fax the prescription image to the new pharmacy to complete the transfer

89. A prescription is for "Ventolin HFA." Which generic name matches?

- a) Fluticasone/salmeterol
- b) Levalbuterol
- c) Albuterol
- d) Ipratropium

90. A patient asks to pick up a friend's controlled medication and asks you to confirm the medication name. The person is not listed as authorized. What should the technician do?

- a) Confirm the name but not the directions
- b) Do not disclose medication details; follow identity/authorization requirements before release
- c) Confirm everything if they know the patient's phone number
- d) Provide the directions but not the name

Answer Key

1) b	31) a	61) b
2) c	32) a	62) b
3) b	33) b	63) b
4) d	34) d	64) c
5) c	35) c	65) c
6) c	36) b	66) d
7) c	37) c	67) c
8) b	38) b	68) c
9) b	39) d	69) c
10) a	40) c	70) b
11) c	41) b	71) d
12) c	42) c	72) c
13) d	43) c	73) b
14) c	44) b	74) c
15) c	45) c	75) c
16) c	46) b	76) b
17) c	47) b	77) d
18) b	48) c	78) c
19) c	49) d	79) a
20) b	50) c	80) b
21) b	51) a	81) c
22) b	52) c	82) b
23) a	53) a	83) b
24) c	54) d	84) c
25) d	55) c	85) c
26) b	56) a	86) b
27) c	57) d	87) b
28) b	58) c	88) c
29) b	59) b	89) c
30) c	60) d	90) b

PTCB Sim Test 9

1) A new eRx is received: "Xarelto 20 mg tablet. Take 1 tablet PO daily with evening meal. #30." The patient profile shows an active prescription for Eliquis 5 mg BID filled 10 days ago. What is the best next step for the technician?

- a) Enter Xarelto as written and place in pharmacist review for possible duplicate anticoagulation/therapy change
- b) Discontinue Eliquis in the profile and process Xarelto
- c) Process both prescriptions and let the patient decide which to take
- d) Change Xarelto directions to match the Eliquis directions for consistency

2) A prescription reads: "Augmentin 875-125 mg. Take 1 tablet PO BID x 7 days. Disp #14." Which product should be selected?

- a) ampicillin/sulbactam
- b) amoxicillin/clavulanate
- c) amoxicillin
- d) cefdinir

3) A customer requests pseudoephedrine 30 mg tablets OTC. Before the sale is completed, which action is required under the Combat Methamphetamine Epidemic Act?

- a) Submit the purchase to the state PDMP
- b) Require a prescription if the customer is under 21
- c) Verify identification, record the sale in the log, and ensure quantity limits are not exceeded
- d) Provide the patient a medication guide and document acceptance

4) An order reads: "Azithromycin suspension 200 mg/5 mL: 12.5 mL on day 1, then 6.25 mL daily on days 2–5." How many mL should be dispensed?

- a) 40 mL
- b) 43.75 mL
- c) 37.5 mL
- d) 31.25 mL

5) While filling, a barcode scan alerts "NDC mismatch." The label is for hydroxyzine 25 mg, but the stock bottle scanned is hydralazine 25 mg. What should the technician do?

- a) Continue filling; the pharmacist will correct it during verification
- b) Override the alert since the strength matches
- c) Reprint the label with the drug name that matches the scanned bottle
- d) Stop and correct the product selection before proceeding

6) A new prescription is written for "metoprolol succinate ER 50 mg daily." The patient previously received "metoprolol tartrate 50 mg BID." What is the best technician action?

- a) Change the prescription to tartrate since it matches the patient's history
- b) Enter succinate ER as written and queue it for pharmacist review due to formulation change
- c) Enter tartrate once daily to approximate the daily dose
- d) Cancel the prescription as a duplicate therapy

7) A pharmacy discovers a significant loss of controlled substances due to theft. Which DEA form is used to report this?

- a) DEA Form 106
- b) DEA Form 222
- c) DEA Form 41
- d) DEA Form 363

8) A prescription reads: "Prednisone 10 mg tablets: Take 3 tablets daily x 5 days, then 2 tablets daily x 5 days, then 1 tablet daily x 5 days." What quantity should be dispensed?

- a) 25 tablets
- b) 45 tablets
- c) 20 tablets
- d) 30 tablets

9) A patient asks you to tell their employer which medication they picked up last week. There is no written authorization on file. What is the best response?

- a) Provide the medication name only, not the directions or prescriber
- b) Decline to disclose and offer to provide the patient a printout they can share
- c) Provide fill date and price only
- d) Fax the information directly to the employer to avoid discussing it aloud

10) A paper prescription for alprazolam contains the patient name, drug, strength, directions, and prescriber signature but has no date written. What should the technician do?

- a) Fill a limited emergency supply and ask for the date later
- b) Enter today's date and continue processing
- c) Set it aside for pharmacist review so the date can be obtained before dispensing
- d) Ask the patient what date it was written and use that

11) A patient is prescribed "Crestor 20 mg daily." Which generic should be selected?

- a) simvastatin
- b) rosuvastatin
- c) pravastatin
- d) atorvastatin

12) "Insulin aspart U-100: inject 8 units SQ TID with meals." If dispensing one 10 mL vial, what days' supply should be entered?

- a) 41 days
- b) 50 days
- c) 31 days
- d) 125 days

13) A patient presents a Schedule II prescription that appears altered (quantity ink is darker and the handwriting is misaligned). What is the best technician action?

- a) Call the prescriber immediately and request a replacement prescription
- b) Process it as written; the pharmacist will address it at pickup
- c) Return it to the patient and refuse to document anything
- d) Hold the prescription for pharmacist review before any dispensing steps

14) An IV order reads: "Dopamine 400 mg in 250 mL D5W. Infuse at 10 mcg/kg/min. Patient weight 70 kg." What infusion rate in mL/hr is needed?

- a) 13.1 mL/hr
- b) 19.7 mL/hr
- c) 26.3 mL/hr
- d) 32.8 mL/hr

15) Under federal requirements, how must Schedule II prescription records be stored?

- a) They must be kept in a separate Schedule II file
- b) They must be kept in a locked safe at all times
- c) They must be filed with Schedule III–V prescriptions only
- d) They may only be stored electronically with no paper copies

16) A patient picking up nitroglycerin SL asks, "When should I call 911?" What is the best technician response?

- a) Refer the question to the pharmacist for counseling
- b) Advise to call 911 only after all 3 tablets are taken
- c) Read the directions on the label and answer from that
- d) Tell the patient to consult the package insert at home

17) "Amoxicillin 250 mg/5 mL: Give 7.5 mL PO TID x 10 days." How many mL should be dispensed?

- a) 300 mL
- b) 150 mL
- c) 225 mL
- d) 200 mL

18) A DSCSA alert identifies a "suspect product" lot of semaglutide pens due to abnormal labeling. What should the technician do?

- a) Return all semaglutide to the wholesaler immediately without documentation
- b) Relabel the boxes with the correct NDC if it can be identified
- c) Quarantine the affected lot and alert the pharmacist/manager to investigate and document
- d) Continue dispensing until a recall is issued

19) Latanoprost 0.005%: "Instill 1 drop OU QHS." A 2.5 mL bottle is billed. If the plan uses 20 drops/mL, what days' supply should be entered?

- a) 25 days
- b) 10 days
- c) 50 days
- d) 100 days

20) A technician is asked which drug on a profile is an ACE inhibitor. Which is correct?

- a) hydrochlorothiazide
- b) amlodipine
- c) losartan
- d) lisinopril

21) A patient returns a vial of tablets that is "unopened," but the seal is broken and

lot/expiration cannot be verified. What should be done?

- a) Return to stock if the count is correct and tablets look clean
- b) Repackage into a new vial and assign a new expiration date
- c) Do not return to stock; handle as non-returnable and dispose per procedure
- d) Return to stock if the patient confirms it was stored properly

22) An eRx arrives: "Methotrexate 2.5 mg: Take 4 tablets daily." The note indicates rheumatoid arthritis. What should the technician do?

- a) Enter as written because the indication supports use
- b) Change the directions to weekly because that is typical
- c) Delete the eRx and ask the patient to contact the prescriber
- d) Enter and route for pharmacist review due to high-risk dosing concern

23) Which is the generic for Zoloft?

- a) citalopram
- b) fluoxetine
- c) sertraline
- d) paroxetine

24) A patient asks to transfer an Adderall XR prescription to another pharmacy. What is correct under federal law?

- a) Transfers are allowed one time only for Schedule II prescriptions
- b) Schedule II prescriptions cannot be transferred between pharmacies
- c) Transfers are allowed only within the same chain
- d) Transfers are allowed if no fills have occurred

25) A prescription reads: "Warfarin 2 mg: Take as directed by anticoag clinic. #30." The patient reports taking 1.5 tablets daily. What should the technician do?

- a) Change the sig to 1.5 tablets daily to calculate days' supply
- b) Reject the prescription as invalid due to missing specific directions
- c) Leave as written and route for pharmacist review to determine billing/days' supply approach
- d) Automatically bill a 30-day supply for all warfarin prescriptions

26) A prescriber's office calls requesting a patient's complete medication list. No authorization is on file. What should the technician do under HIPAA?

- a) Verify the caller identity and disclose the minimum necessary information for treatment purposes
- b) Refuse all disclosures unless the patient is on the phone
- c) Provide only non-controlled medications
- d) Provide the list without verification since it is a prescriber's office

27) Albuterol HFA directions: "Inhale 2 puffs q4–6h PRN." The inhaler contains 200 puffs. If used at the maximum of 12 puffs/day, what days' supply is it?

- a) 14 days
- b) 17 days
- c) 16 days
- d) 30 days

28) A patient requests a refill for clonazepam (Schedule IV). The prescription was written 7 months ago and refills remain. Under federal law, what is correct?

- a) It may be refilled up to 12 months from the date written, if refills remain
- b) It may be refilled indefinitely until refills run out
- c) It cannot be refilled after 6 months
- d) It can only be refilled if the prescriber re-faxes authorization each time

29) "Furosemide 20 mg: Take 1 tablet PO daily. Dispense a 45-day supply." What quantity is needed?

- a) 60
- b) 30
- c) 90
- d) 45

30) At pickup, there are two will-call bags: "Maria L. Cruz" and "Marian L. Cruz." Their birthdays share the same month/day but different years. What should the technician do to reduce the chance of a wrong-patient error?

- a) Ask for two identifiers (e.g., full DOB and address) before retrieving or handing out the bag
- b) Ask only for address to avoid disclosing DOB in public
- c) Open both bags and compare the medications with the patient
- d) Hand the bag that matches the first name most closely and correct it if needed

31) A prescription is entered for Bactrim DS. The profile lists "sulfonamide—anaphylaxis." What should the technician do?

- a) Substitute doxycycline and continue processing
- b) Remove the allergy since the prescriber ordered the drug
- c) Hold the prescription for pharmacist review due to allergy conflict
- d) Fill it because the prescriber likely considered the allergy

32) Which medication is an inhaled corticosteroid/long-acting beta agonist (ICS/LABA) combination?

- a) Flovent
- b) Ventolin
- c) Advair
- d) Spiriva

33) A compounding worksheet requires 240 mL of a 5% w/v solution made from a 20% w/v stock. How many mL of the 20% stock are needed (then q.s. to 240 mL)?

- a) 75 mL
- b) 30 mL
- c) 60 mL
- d) 45 mL

34) A technician attaches the wrong patient leaflet to a bag, but it is caught before the medication leaves the pharmacy. How should this be classified?

- a) Dispensing error that reached the patient

- b) Near miss (good catch)
- c) Adverse drug event
- d) Counterfeit product incident

35) What is the generic for Norvasc?

- a) amlodipine
- b) diltiazem
- c) lisinopril
- d) atenolol

36) "Cephalexin 500 mg: Take 1 capsule PO QID x 7 days." What quantity should be dispensed?

- a) 30
- b) 21
- c) 28
- d) 14

37) Which statement about Schedule II refills is correct?

- a) Refills are allowed if the patient pays cash
- b) Refills are allowed within 30 days of the original fill
- c) Refills are allowed if the prescriber is contacted by phone
- d) Schedule II prescriptions may not be refilled

38) Timolol 0.5% ophthalmic: "Instill 1 drop OS BID." Bottle size is 10 mL. If 20 drops/mL are used for billing, what days' supply should be entered?

- a) 100 days
- b) 200 days
- c) 50 days
- d) 75 days

39) An eRx reads "Synthroid 75 mcg DAW 1." What does DAW 1 indicate?

- a) Patient requested brand
- b) Brand medically necessary; do not substitute
- c) Substitution allowed; dispense generic
- d) Generic not available

40) DEA Form 222 is used for:

- a) reporting theft or significant loss
- b) ordering Schedule II controlled substances
- c) documenting destruction of controlled substances
- d) registering a pharmacy with the DEA

41) A prescription is written for "Humulin R U-500: inject 50 units BID." The technician is unsure which syringes the patient uses. What is the best action?

- a) Convert the dose to mL and change the patient directions to avoid confusion
- b) Change the product to U-100 regular insulin for safety
- c) Enter and route for pharmacist review due to high-alert concentration and dosing safety concerns

d) Enter as written; syringe type does not affect dosing

42) Mupirocin ointment: "Apply to affected area TID x 10 days." A 22 g tube is dispensed. If the patient uses 1 g per application, what days' supply is the tube?

- a) 10 days
- b) 7 days
- c) 5 days
- d) 2 days

43) Which medication is a proton pump inhibitor (PPI)?

- a) sucralfate
- b) dicyclomine
- c) famotidine
- d) omeprazole

44) A new technician stores varicella vaccine in the refrigerator door where temperatures fluctuate. What is the best immediate corrective action?

- a) Move it to a stable temperature storage area and flag it for pharmacist evaluation of viability
- b) Leave it; door storage is acceptable for vaccines
- c) Dispose of it immediately
- d) Place it in will-call refrigeration to prevent frequent opening

45) "Gabapentin 300 mg: Take 1 capsule in the morning and 2 capsules at bedtime." What quantity is needed for 30 days?

- a) 60
- b) 90
- c) 75
- d) 120

46) What is the generic for Lasix?

- a) torsemide
- b) bumetanide
- c) spironolactone
- d) furosemide

47) A faxed prescription for hydrocodone/APAP is missing the prescriber's signature. What is the best technician action?

- a) Fill it because it came from the prescriber's office fax
- b) Add the signature from a prior scanned copy on file
- c) Ask the patient to sign the prescription
- d) Hold it for pharmacist review; a valid prescription must be obtained before dispensing

48) A prescription says "venlafaxine ER 75 mg capsule daily," but the technician selected venlafaxine IR 75 mg tablet. What should the technician do before filling?

- a) Change the directions to BID and proceed
- b) Re-select the correct ER formulation/dosage form to match the prescription
- c) Print the label and let the pharmacist correct it later

d) Dispense IR once daily because the strength matches

49) Which medication is an SSRI?

- a) bupropion
- b) amitriptyline
- c) sertraline
- d) duloxetine

50) "Clindamycin 150 mg: Take 2 capsules PO TID x 7 days." What quantity should be dispensed?

- a) 56
- b) 84
- c) 42
- d) 28

51) A printed label with PHI is left on the counter visible to customers. What should the technician do immediately?

- a) Throw it in the regular trash to remove it from view
- b) Read it aloud to locate the patient quickly
- c) Continue working; it will be used soon
- d) Turn it face down or move it to a secure area right away

52) A patient requests an early refill for zolpidem 10 mg (Schedule IV) stating the bottle was lost. It was filled 6 days ago for a 30-day supply. What should the technician do?

- a) Process the refill and document "lost medication"
- b) Collect details and route to the pharmacist for evaluation and documentation
- c) Tell the patient early refills are never allowed for any controlled substance
- d) Call the prescriber and request a new prescription without involving the pharmacist

53) Which medication is a direct oral anticoagulant (DOAC)?

- a) enoxaparin
- b) clopidogrel
- c) apixaban
- d) aspirin

54) "Vancomycin 1 g in 200 mL NS infuse over 90 minutes." What is the infusion rate in mL/hr?

- a) 150 mL/hr
- b) 120 mL/hr
- c) 100 mL/hr
- d) 133 mL/hr

55) A prescriber calls in a verbal prescription for amoxicillin. Who may accept a verbal prescription order (federal baseline)?

- a) Only a certified technician
- b) Any technician with at least 6 months experience
- c) Any staff member if it is non-controlled
- d) Only a pharmacist

56) A profile lists "penicillin—hives." A prescription arrives for amoxicillin. What should the technician do?

- a) Route for pharmacist review due to allergy match
- b) Process and add an auxiliary label to take with food
- c) Change to azithromycin and proceed
- d) Process because hives are not a true allergy

57) What is a brand name for metformin?

- a) Januvia
- b) Glucotrol
- c) Glucophage
- d) Glumetza

58) Budesonide/formoterol 160/4.5: "Inhale 2 puffs BID." The inhaler contains 120 actuations. What days' supply should be entered?

- a) 60 days
- b) 30 days
- c) 20 days
- d) 15 days

59) A recall notice is received for specific lot numbers of atorvastatin. What should the technician do first?

- a) Remove all atorvastatin from stock immediately
- b) Call every patient who has ever taken atorvastatin
- c) Quarantine and block the affected lots/NDCs from dispensing
- d) Return all stock to the wholesaler without documentation

60) A prescription says: "Take 1 tablet PO q6h PRN pain. Max 4 tablets/day. Disp #48." What is the days' supply?

- a) 8 days
- b) 10 days
- c) 12 days
- d) 16 days

61) Which medication is a beta blocker?

- a) losartan
- b) metoprolol
- c) atorvastatin
- d) furosemide

62) A cream is labeled: "Apply 2 g topically QID." The tube contains 120 g. What days' supply is the tube?

- a) 7 days
- b) 10 days
- c) 30 days
- d) 15 days

63) While counting, a technician notices tablets in a stock bottle do not match the imprint on the bottle description. What should the technician do?

- a) Continue filling; imprint variations happen
- b) Use the tablets only for established patients
- c) Quarantine the bottle and alert the pharmacist/manager to investigate
- d) Pour the tablets into a bottle with a matching label

64) Which medication is an opioid antagonist used for overdose reversal?

- a) buprenorphine/naloxone
- b) clonidine
- c) naltrexone/bupropion
- d) naloxone

65) "Valacyclovir 1 g: Take 2 tablets PO BID for 1 day." What quantity should be dispensed?

- a) 4 tablets
- b) 6 tablets
- c) 8 tablets
- d) 2 tablets

66) A technician enters "Lamictal 100 mg" and notices the profile already lists "lamotrigine 100 mg" as active. What is the best technician action?

- a) Dispense both because brand and generic are different medications
- b) Confirm it is the same medication and proceed appropriately without duplicating the profile entry
- c) Override because brand and generic must be kept separate
- d) Delete the generic automatically

67) Under federal law, what is the maximum number of refills allowed for a Schedule III prescription within the allowable time period?

- a) 11 refills in 12 months
- b) No refills allowed
- c) 5 refills in 6 months
- d) 3 refills in 6 months

68) Insulin glargine U-100 pens: "Inject 24 units SQ daily." Dispense 1 box (5 pens, 3 mL each). What days' supply should be entered?

- a) 25 days
- b) 62 days
- c) 75 days
- d) 31 days

69) Which is the generic for Keppra?

- a) carbamazepine
- b) lamotrigine
- c) levetiracetam
- d) valproic acid

70) A label prints for diltiazem ER 240 mg, but the technician pulls diltiazem IR 120 mg.

What should the technician do next?

- a) Adjust directions to BID to match the pulled product
- b) Continue; the pharmacist will catch it during final verification
- c) Fill with IR and note "ER" on the label
- d) Stop and pull the correct ER product/strength that matches the prescription

71) A patient requests a printed medication profile to bring to a new provider. What should the technician do?

- a) Print it and hand it to anyone who requests it
- b) Verify identity and provide it to the patient per procedure
- c) Refuse because profiles can't be printed
- d) Fax it to the new provider without patient request

72) "Ciprofloxacin 500 mg: Take 1 tablet PO BID x 10 days." What quantity should be dispensed?

- a) 14
- b) 20
- c) 28
- d) 10

73) A refrigerated insulin shipment arrives with a temperature indicator showing exposure above the acceptable range. What should the technician do?

- a) Freeze the insulin to restore stability
- b) Quarantine the shipment and alert the pharmacist/manager per procedure
- c) Refrigerate and use it first to reduce waste
- d) Dispense only if the patient will finish it within 7 days

74) Which medication is used to treat hypothyroidism?

- a) methimazole
- b) liothyronine
- c) propylthiouracil
- d) levothyroxine

75) A prescription is "Fluconazole 150 mg: Take 1 tablet now; repeat in 72 hours." What days' supply is most appropriate for billing?

- a) 1 day
- b) 2 days
- c) 4 days
- d) 3 days

76) A new eRx is received for lisinopril 10 mg daily. The allergy field shows "ACE inhibitors—angioedema." What should the technician do?

- a) Route for pharmacist review due to contraindication risk
- b) Enter and process; the pharmacist will counsel later
- c) Inactivate the allergy since the prescriber ordered it
- d) Change the medication to losartan 25 mg and process

77) Which medication is a thiazide diuretic?

- a) spironolactone
- b) acetazolamide
- c) furosemide
- d) hydrochlorothiazide

78) A prescription is written for OxyContin 10 mg q12h. The technician selected oxycodone 10 mg immediate-release. What should the technician do?

- a) Change the sig to q6h and proceed
- b) Stop and select the correct extended-release product as prescribed
- c) Continue because both contain oxycodone
- d) Dispense IR and add an "extended release" auxiliary label

79) A patient arrives to pick up isotretinoin. The system shows REMS requirements are not complete for the current fill. What should the technician do?

- a) Substitute topical tretinoin to avoid delay
- b) Complete REMS steps under the technician login to bypass the alert
- c) Dispense and document that counseling will occur later
- d) Hold the prescription until REMS requirements are confirmed complete

80) Metformin ER 500 mg is prescribed "Take 2 tablets BID." The claim rejects for "dose exceeds maximum." What should the technician do first?

- a) Split into two separate prescriptions to force coverage
- b) Route to the pharmacist to review options (e.g., prescriber contact, PA, dose clarification, plan limits)
- c) Change directions to 1 tablet BID and re-bill
- d) Bypass insurance and run as cash automatically

81) A caregiver says at pickup, "This tablet looks different than last time—can they still take it?" What is the best technician response?

- a) Explain that generics are always identical and there is no concern
- b) Open the vial and compare tablets in front of the line
- c) Refer the question to the pharmacist before the medication is released
- d) Provide a detailed counseling explanation about therapeutic equivalence

82) Which medication is a calcium channel blocker?

- a) clonidine
- b) metoprolol
- c) amlodipine
- d) lisinopril

83) "Enoxaparin 40 mg/0.4 mL: Inject 0.4 mL SQ daily x 14 days." How many syringes should be dispensed?

- a) 10
- b) 14
- c) 28
- d) 7

84) A prescription directions field includes "qd." What is the best way to reduce

misinterpretation during order entry?

- a) Enter "qd" exactly as written
- b) Change it to "QID" to avoid confusion
- c) Expand it to "daily" in the patient directions
- d) Reject the prescription as invalid

85) Which medication is an H2 blocker commonly used for GERD?

- a) sucralfate
- b) omeprazole
- c) famotidine
- d) pantoprazole

86) A patient requests a transfer of a non-controlled maintenance medication to another pharmacy. What is the best technician action?

- a) Transfer it over the phone to save time
- b) Route the request appropriately so the pharmacist can complete/authorize the transfer per law and workflow
- c) Cancel the prescription and tell the patient to get a new one
- d) Tell the patient transfers are only for controlled substances

87) "Acetaminophen 325 mg: Take 2 tablets every 4 hours PRN fever." If taken around the clock at the maximum frequency, how many tablets per day is that?

- a) 10 tablets
- b) 12 tablets
- c) 14 tablets
- d) 8 tablets

88) During a controlled substance inventory count, which practice best supports accurate accountability?

- a) Combine partial bottles to reduce clutter
- b) Estimate open bottle counts to save time
- c) Count only when a discrepancy is reported
- d) Maintain a perpetual inventory and document counts promptly

89) Nexium is the brand name for which medication?

- a) pantoprazole
- b) omeprazole
- c) esomeprazole
- d) lansoprazole

90) A caller reports facial swelling and trouble breathing after taking the first dose of amoxicillin today. What should the technician do first?

- a) Recommend diphenhydramine and monitor symptoms
- b) Tell the caller to stop the medication and call back tomorrow
- c) Transfer the call to the pharmacist immediately
- d) Add the allergy to the profile and offer to refill a different antibiotic

Answer Key

1) a	32) c	63) c
2) b	33) c	64) d
3) c	34) b	65) a
4) b	35) a	66) b
5) d	36) c	67) c
6) b	37) d	68) b
7) a	38) a	69) c
8) d	39) b	70) d
9) b	40) b	71) b
10) c	41) c	72) b
11) b	42) c	73) b
12) b	43) d	74) d
13) d	44) a	75) d
14) c	45) c	76) a
15) a	46) d	77) d
16) a	47) d	78) b
17) c	48) b	79) d
18) c	49) c	80) b
19) c	50) b	81) c
20) d	51) d	82) c
21) c	52) b	83) b
22) d	53) c	84) c
23) c	54) d	85) c
24) b	55) d	86) b
25) c	56) a	87) b
26) a	57) c	88) d
27) b	58) b	89) c
28) a	59) c	90) c
29) d	60) c	
30) a	61) b	
31) c	62) b	

PTCB Sim Test 10

1) A new eRx reads: "glipizide 10 mg ER tablet. Take 1 tablet PO daily with breakfast. Qty 90." Which product selection best matches?

- a) Glipizide 10 mg immediate-release tablet
- b) Glipizide XL 10 mg extended-release tablet
- c) Glyburide 10 mg tablet
- d) Metformin ER 500 mg tablet

2) A caller says, "I'm picking up for my spouse. What medication is it and what's it for?" The spouse is not present and there is no authorized representative documented. What is the most appropriate response?

- a) Provide the medication name only, not the indication
- b) Provide the ready time only and offer to contact the patient for permission
- c) Provide the medication name and indication because they're married
- d) Provide the indication but not the medication name

3) A hard copy for a controlled medication has the prescriber signature, patient name, drug, strength, and directions, but no patient address. What should the technician do next?

- a) Enter it as written; address is optional
- b) Ask the patient to write their address on the prescription and proceed
- c) Send it back to the prescriber to rewrite it
- d) Set it aside for pharmacist review to resolve missing required elements

4) An eRx for "metoprolol succinate ER 25 mg daily" is entered, but the product selected is "metoprolol tartrate 25 mg." What is the main concern?

- a) The tartrate form is a different drug class
- b) The release form differs and may change dosing frequency/effect
- c) Succinate is only available as brand-name
- d) Tartrate is contraindicated in hypertension

5) A prescription reads: "cefдинир 250 mg/5 mL. Take 6 mL PO BID x 10 days." How many mL should be dispensed?

- a) 120 mL
- b) 100 mL
- c) 60 mL
- d) 180 mL

6) A tech is pulling a product and notices the NDC on the stock bottle does not match the NDC on the label that just printed. What is the best next step?

- a) Use the stock bottle anyway if the strength matches
- b) Reprint the label and keep filling
- c) Stop and resolve the product/label mismatch before counting
- d) Continue filling and let the pharmacist catch it at final check

7) A patient requests a transfer of their non-controlled prescription to another pharmacy. Which information must be communicated and documented as part of a valid transfer?

- a) Directions, drug/strength, quantity, refills remaining, RX number, prescriber, original date, and last fill date
- b) Diagnosis code and prescriber NPI only
- c) Patient allergies and vaccination history
- d) Wholesaler invoice number and acquisition cost

8) A nurse calls requesting an “emergency” supply of a Schedule II medication for a patient who ran out at a facility. There is no prescription on file. What is the appropriate technician action?

- a) Offer a 3-day loaner supply and bill later
- b) Take a verbal order from the nurse and process it
- c) Tell the nurse to bring the empty bottle and dispense based on that
- d) Refer the call for pharmacist handling; Schedule II emergency dispensing has strict requirements

9) A prescription label direction must be clarified from “Take 1 tab BID PRN” because it lacks an indication. Which field is most appropriate to use if the system allows it?

- a) Patient address field
- b) Sig field with added indication wording if provided on the prescription
- c) Prescriber DEA field
- d) NDC override field

10) A patient is prescribed “warfarin 5 mg daily” and a new eRx arrives for “rivaroxaban 20 mg daily.” What should the technician do during processing?

- a) Enter and fill both because they treat different clots
- b) Hold the new prescription until the patient confirms they stopped warfarin
- c) Enter the new prescription and route for pharmacist review due to therapeutic duplication risk
- d) Delete warfarin from the profile automatically

11) A prescription is for “insulin glargine U-100: inject 18 units SQ nightly. Dispense 1 pen box (5 pens x 3 mL).” What day supply should be billed? (100 units/mL)

- a) 30 days
- b) 83 days
- c) 28 days
- d) 14 days

12) A vaccine shipment arrives. The cold-chain indicator shows an excursion. What should the technician do first?

- a) Put it immediately into the refrigerator and ignore the indicator
- b) Place it in the freezer to stabilize it
- c) Quarantine the shipment and document/escalate per procedure before use
- d) Stock it but mark it “use first”

13) An eRx says: “levothyroxine 0.1 mg tab daily.” Which strength should be selected?

- a) 100 mcg
- b) 10 mcg
- c) 1 mg

d) 0.01 mg

14) A patient asks to buy pseudoephedrine. Under CMEA, which action is required before completing the sale?

- a) Record the sale in the log system after the transaction is finalized
- b) Verify acceptable photo ID and record required sale information
- c) Allow purchase without ID if the patient is known to staff
- d) Restrict sales to prescription-only items

15) A prescription reads: "Take 2.5 mL PO TID." The bottle is 120 mL. How many days will it last?

- a) 8 days
- b) 12 days
- c) 16 days
- d) 24 days

16) A patient presents a printed eRx for a Schedule II stimulant. The prescriber says they "printed it from their system." What is required for validity in most outpatient settings?

- a) The prescription must be handwritten in ink
- b) The prescription must include the clinic's DEA registration number
- c) The prescription must be on tamper-resistant paper with wet signature
- d) The prescription must have an original manual signature by the prescriber

17) A prescription is for "Bactrim DS." Which generic is correct?

- a) Sulfadiazine/pyrimethamine
- b) Sulfamethoxazole/trimethoprim DS
- c) Sulfasalazine DR
- d) Trimethoprim alone

18) A patient's profile shows "penicillin—anaphylaxis." A new prescription is for amoxicillin. What should the technician do?

- a) Process it and add an auxiliary "take with food"
- b) Change it to azithromycin because it's safer
- c) Route the prescription for pharmacist review before dispensing
- d) Remove the allergy because it was reported long ago

19) A hard copy prescription has an obvious alteration to the quantity (different handwriting/ink). What should the technician do?

- a) Fill the smallest quantity that appears on the paper
- b) Continue processing but note "possible alteration"
- c) Ask the patient what quantity the doctor intended
- d) Stop processing and route for pharmacist verification steps

20) The sig reads: "1 gtt OU qHS." What is the correct patient-friendly direction?

- a) Instill 1 drop into each eye at bedtime
- b) Instill 1 drop into the right eye every morning
- c) Instill 2 drops into both ears at bedtime
- d) Apply 1 drop to both eyes every 4 hours

21) A prescription is for “amoxicillin 875 mg: 1 tab PO BID x 7 days.” How many tablets should be dispensed?

- a) 7
- b) 21
- c) 14
- d) 28

22) A technician is performing data entry and notices two patient profiles with the same name and similar DOB. What is the best next step before entering the prescription?

- a) Choose the profile with the most recent activity
- b) Verify using additional identifiers (address/phone) and merge only if appropriate
- c) Create a third profile to avoid mistakes
- d) Enter it under either profile and fix later if rejected

23) A DSCSA request asks for transaction information for a prescription medication. What is the purpose of DSCSA documentation?

- a) Document patient counseling
- b) Ensure correct billing for insurance claims
- c) Track prescription products through the supply chain and help identify suspect product
- d) Provide prescriber credential verification

24) A prescription is for “morphine ER 30 mg” but “morphine IR 30 mg” is selected. What type of error is this most directly?

- a) Look-alike/sound-alike error
- b) Wrong dosage form/release formulation error
- c) Wrong route of administration error
- d) Therapeutic duplication error

25) A controlled substance (Schedule III) prescription is written with 7 refills. What is the correct handling?

- a) Process as written; federal law allows any refills if medically necessary
- b) Reduce to 5 refills and proceed without contacting anyone
- c) Route for pharmacist resolution because it exceeds legal refill limits
- d) Transfer it to another pharmacy to “split” the refills

26) A patient returns an opened bottle of tablets and requests the pharmacy “put it back on the shelf” because it was never used. What is the appropriate technician response in most community settings?

- a) Restock if the tablets appear intact
- b) Restock if the expiration date is not past
- c) Restock after wiping the bottle
- d) Do not restock; handle as return per procedure and regulations

27) A prescription is for “fluoxetine 20 mg daily.” The patient says, “I’m taking Prozac.” What is Prozac?

- a) Brand name for fluoxetine
- b) Generic for sertraline

- c) Brand name for paroxetine
- d) Generic for citalopram

28) A technician is asked to file ordering records for Schedule II controlled substances when paper ordering is used. Which DEA form is used to order Schedule II?

- a) DEA Form 222
- b) DEA Form 106
- c) DEA Form 41
- d) DEA Form 363

29) A prescription reads: "Hydrochlorothiazide 25 mg daily." The system generates an alert due to "sulfa allergy" on file. What should the technician do?

- a) Override the alert because HCTZ is not an antibiotic
- b) Route the alert for pharmacist review
- c) Delete the sulfa allergy
- d) Change the medication to furosemide

30) A prescription is for "albuterol HFA: inhale 2 puffs q4h PRN. Inhaler contains 200 actuations." If used at the maximum frequency, what is the day supply?

- a) 25 days
- b) 16 days
- c) 20 days
- d) 10 days

31) A patient asks, "Can I stop taking my atorvastatin since my cholesterol is better?" What is the best technician response?

- a) Yes, stop immediately if levels are normal
- b) Cut the dose in half for one week
- c) Offer to have the pharmacist speak with the patient
- d) Continue only if taken with grapefruit

32) A prescription for "ondansetron ODT 8 mg" is being entered. Which selection is correct?

- a) Ondansetron immediate-release tablet
- b) Omeprazole delayed-release capsule
- c) Olanzapine ODT tablet
- d) Ondansetron orally disintegrating tablet

33) A technician is preparing to compound a nonsterile oral solution and realizes the balance was not calibrated today. What should the technician do?

- a) Proceed; calibration is only required weekly
- b) Use a volume estimate instead of weighing
- c) Calibrate the balance (or use a verified calibrated balance) before compounding
- d) Ask another tech to eyeball the amount

34) A pharmacist requests you remove a recalled lot of OTC product due to a Class I recall. What is the best technician action?

- a) Keep it on the shelf but place a "do not sell" note
- b) Quarantine the affected lot and document per recall procedure

- c) Return it to the manufacturer without documentation
- d) Sell through remaining stock to avoid waste

35) A prescription is for "tamsulosin 0.4 mg daily." What is the most common indication?

- a) Migraine prevention
- b) Benign prostatic hyperplasia
- c) Hyperthyroidism
- d) Heart failure

36) A technician receives a request to fax a patient's full medication profile to a specialist office. No authorization is documented. What is the best action?

- a) Fax it since it's for treatment and never needs verification
- b) Call the specialist and read the profile over the phone instead
- c) Refuse; profiles can never be shared
- d) Obtain appropriate authorization or route for pharmacist determination of permitted disclosure before sending

37) An eRx for isotretinoin arrives and the system flags REMS requirements. What should the technician do next?

- a) Enter and dispense; REMS is only for prescribers
- b) Ensure required REMS documentation/steps are completed before the prescription can be dispensed
- c) Change to topical tretinoin without contacting anyone
- d) Reduce quantity to avoid REMS

38) A technician is asked to complete a partial fill of a Schedule II in a typical retail setting because not enough stock is available. Under federal rules, when must the remainder generally be filled (if allowed)?

- a) Within 72 hours or the remainder is forfeited
- b) Within 7 days
- c) Within 30 days
- d) Anytime before the prescription expires

39) A prescription is for "clopidogrel 75 mg daily." Which brand name is correct?

- a) Coumadin
- b) Plavix
- c) Paxil
- d) Pravachol

40) During pick-up, a patient says, "These pills look different than last month." The label NDC matches the fill record. What should the technician do?

- a) Tell the patient to take them; appearance changes are irrelevant
- b) Replace with a different manufacturer without checking
- c) Compare imprint/description to the system and route to pharmacist if mismatch or concern remains
- d) Call the prescriber to change the prescription

41) A prescription is for "methotrexate 2.5 mg: take 6 tablets once weekly." Which action

best supports safety during order entry?

- a) Enter it as “6 tablets daily” to avoid missed doses
- b) Enter “once weekly” clearly and route for pharmacist verification due to high-alert weekly dosing
- c) Dispense only 6 tablets total
- d) Change to injectable methotrexate automatically

42) A patient’s prescription says “DAW 1” for a brand product. What should the technician do?

- a) Dispense brand and submit the appropriate DAW code consistent with the prescription
- b) Dispense generic and submit DAW 0
- c) Dispense generic and submit DAW 1 anyway
- d) Call the prescriber to confirm DAW 1 before processing

43) A technician is calculating heparin volume for an IV admixture: Order is 20,000 units. Stock is 5,000 units/mL. How many mL are needed?

- a) 2 mL
- b) 5 mL
- c) 4 mL
- d) 10 mL

44) A prescription reads: “Azithromycin 250 mg: take 2 tablets on day 1, then 1 tablet daily days 2–5.” Total number of tablets is:

- a) 5
- b) 6
- c) 4
- d) 7

45) A patient requests to refill diazepam written 8 months ago with refills remaining. What is correct under federal law?

- a) It can be refilled as long as refills remain
- b) It can be refilled within 1 year
- c) It can be refilled only if the patient pays cash
- d) It generally cannot be refilled because Schedule III–V prescriptions are limited to 6 months from issue date

46) A technician receives an eRx for “losartan 50 mg daily.” Which class is losartan?

- a) ACE inhibitor
- b) ARB
- c) Beta blocker
- d) Thiazide diuretic

47) A technician finds two different strengths of the same drug stored behind one shelf label in the same bin. What is the best action to reduce selection errors?

- a) Keep together to reduce time
- b) Remove the slower-moving strength from inventory
- c) Separate strengths into distinct labeled locations and correct the shelf label/bin
- d) Put the higher strength in front and lower strength behind

48) An Rx reads: "Take ii caps PO TID. Dispense #180." What is the day supply?

- a) 15
- b) 20
- c) 30
- d) 45

49) A technician is entering "Augmentin XR 1000/62.5 mg." The selected product is "amoxicillin/clavulanate 875/125 mg." What should happen?

- a) Continue; they are equivalent
- b) Change directions to match the selected product
- c) Correct the product selection to the XR formulation as written
- d) Reduce the quantity to avoid diarrhea

50) A prescriber's DEA number fails an automated validation during controlled substance entry. What should the technician do?

- a) Override and proceed
- b) Replace DEA with the prescriber's NPI
- c) Enter a default clinic DEA number
- d) Hold for pharmacist verification/correction before dispensing

51) A controlled substance is discovered missing from inventory after counts are short. What is the best immediate step for a technician?

- a) Adjust the perpetual inventory to match the shelf count
- b) Recount, review recent transactions/receiving, and escalate per procedure
- c) Wait until the next shift for a recount
- d) Report to DEA immediately without internal review

52) A prescription is for "carvedilol." Which common indication fits best?

- a) Heart failure/hypertension
- b) Bacterial infection
- c) Hypothyroidism
- d) Osteoporosis

53) A prescription is for "latanoprost 2.5 mL: instill 1 drop OU qHS." Using 20 drops/mL, approximate day supply is:

- a) 12 days
- b) 25 days
- c) 50 days
- d) 100 days

54) A technician notices a printed patient leaflet for one patient was accidentally attached to another patient's bag. It has not left the pharmacy. What should be done?

- a) Cross out the wrong name and keep it
- b) Place it in the correct bag later without reprinting
- c) Securely dispose of the incorrect leaflet and print the correct one
- d) Give both leaflets to the patient and ask them to discard one

55) A prescription is for "tramadol 50 mg." What is the federal schedule?

- a) Schedule II
- b) Schedule III
- c) Schedule IV
- d) Schedule V

56) A prescription is written for "methylphenidate IR 10 mg." What schedule is methylphenidate?

- a) Schedule II
- b) Schedule III
- c) Schedule IV
- d) Schedule V

57) A technician is asked what "DEA Form 106" is used for. The correct use is:

- a) Ordering Schedule II controlled substances
- b) Reporting theft or significant loss of controlled substances
- c) Inventory of controlled substances at opening/closing
- d) Registration renewal for a pharmacy

58) A patient asks to move nitroglycerin SL tablets into a weekly pill organizer. What is the most appropriate technician action?

- a) Confirm it's okay as long as the organizer is amber
- b) Recommend storing in the bathroom cabinet to reduce humidity exposure
- c) Suggest transferring to any vial with a tight cap
- d) Refer to pharmacist due to storage requirements in the original container

59) A prescription reads: "clindamycin 150 mg: take 1 cap PO QID. Qty 40." Day supply is:

- a) 8 days
- b) 10 days
- c) 5 days
- d) 12 days

60) While typing, a technician selects "hydroxyzine 25 mg" instead of "hydralazine 25 mg." Which safeguard best helps prevent this error?

- a) Store the two drugs adjacent to each other
- b) Turn off alerts to reduce distractions
- c) Tall Man lettering and barcode verification at product selection
- d) Use handwriting only instead of eRx

61) A patient requests Medicare Part B billing for diabetic test strips. Which is commonly required for compliance?

- a) Prescriber NPI only
- b) Diagnosis/medical necessity documentation elements along with a valid order/prescription
- c) DEA number on the prescription
- d) Proof of cash purchase in the last 6 months

62) A patient says they have "sulfa allergy—hives." A prescription arrives for sulfamethoxazole/trimethoprim. What should the technician do?

- a) Enter and dispense with an allergy sticker
- b) Change the drug to nitrofurantoin automatically
- c) Route to pharmacist review before dispensing
- d) Tell the patient to take diphenhydramine with each dose

63) A prescription is for "furosemide 20 mg qAM," but the patient says they take it "BID." What should the technician do?

- a) Change the directions to BID and adjust quantity
- b) Process as written and route discrepancy for pharmacist review
- c) Reject the prescription because it conflicts with patient report
- d) Increase strength to 40 mg daily to match total dose

64) A prescription for an inhaled steroid reads: "Flovent HFA." Which generic is correct?

- a) Fluticasone propionate
- b) Fluticasone furoate
- c) Budesonide
- d) Beclomethasone

65) A tech is reconstituting an antibiotic and notices the powder is clumped and the seal appears compromised. What is the best action?

- a) Add extra diluent and shake longer
- b) Use it if it's within expiration
- c) Dispense but shorten the beyond-use date
- d) Quarantine as suspect product and escalate for pharmacist disposition

66) A patient brings in a prescription for a non-controlled medication that is missing the prescriber's signature (not eRx). What is the most appropriate action?

- a) Fill it if the prescriber name is printed
- b) Ask the patient to sign for the prescriber
- c) Hold and obtain a properly authorized prescription before dispensing
- d) Dispense a 3-day supply and call later

67) A new prescription is for "diltiazem ER 120 mg daily," and the profile shows "diltiazem IR 60 mg QID" still active. What should the technician do?

- a) Discontinue the IR product automatically and fill ER
- b) Change ER directions to QID
- c) Enter the ER prescription and route for pharmacist review due to possible therapy change/duplication
- d) Replace ER with IR since total daily mg is similar

68) A prescription says: "Prednisolone 15 mg/5 mL oral solution." The label printed says "prednisone 10 mg tablets." What is the best technician action?

- a) Dispense prednisone; they are interchangeable
- b) Correct the entry to match drug and dosage form before filling
- c) Keep the label and change the hard copy to match
- d) Dispense prednisolone but use the tablet label

69) A prescription is for "allopurinol 300 mg daily." Primary indication is:

- a) COPD maintenance
- b) Acute pain
- c) Gout prophylaxis/hyperuricemia
- d) Bacterial sinusitis

70) A pharmacy receives a request to transfer alprazolam (Schedule IV) with refills remaining. Which statement is correct under federal rules (if state law permits)?

- a) Controlled substances can never be transferred
- b) Schedule IV prescriptions may be transferred with proper documentation
- c) Transfers are allowed only after one refill has been used
- d) Schedule II prescriptions can be transferred between pharmacies

71) A refrigerator log shows temperatures reached 12°C (53.6°F) for 2 hours overnight for refrigerated biologics. What should be done first?

- a) Continue using products; 2 hours is always acceptable
- b) Adjust the thermostat and do not document to avoid waste
- c) Freeze the products to stop degradation
- d) Quarantine affected products and escalate per procedure for stability guidance

72) A prescription for “omeprazole 20 mg daily” is entered, but the product chosen is “omeprazole/sodium bicarbonate.” What is the issue?

- a) Combination product is equivalent to omeprazole alone
- b) Wrong ingredients/formulation; not the same product
- c) Omeprazole cannot be taken daily
- d) Sodium bicarbonate is inactive and irrelevant

73) A patient calls stating a toddler may have swallowed iron tablets. What should the technician do?

- a) Advise inducing vomiting
- b) Recommend milk and monitoring at home
- c) Direct to Poison Control immediately and alert the pharmacist
- d) Tell them to wait for symptoms before acting

74) A prescription is for “clonazepam.” Which common use best fits?

- a) GERD
- b) Anxiety/panic or seizure disorders
- c) Bacterial pneumonia
- d) Type 1 diabetes

75) A prescription is for “acetaminophen 160 mg/5 mL: take 7.5 mL q6h PRN.” How many mg per dose?

- a) 120 mg
- b) 160 mg
- c) 200 mg
- d) 240 mg

76) A technician is preparing to discard expired controlled substances. What is appropriate?

- a) Place in regular trash after removing labels

- b) Pour into sink with water
- c) Combine with coffee grounds and throw away
- d) Follow controlled substance destruction/return procedures with proper documentation and pharmacist oversight

77) A prescription is for "vitamin D 50,000 units weekly." Which product most commonly matches this?

- a) Ergocalciferol 50,000 unit capsules
- b) Calcitriol 0.25 mcg capsules
- c) Cholecalciferol 1,000 unit tablets
- d) Calcium carbonate 600 mg tablets

78) A patient requests the "cheapest version" of levothyroxine and the prescription is DAW 0. What is the best technician action?

- a) Dispense an FDA-approved generic equivalent per payer/inventory rules
- b) Dispense brand only because substitution is not allowed for thyroid meds
- c) Change to liothyronine to reduce cost
- d) Dispense half quantity and refill more often

79) A prescription is for "Cipro 500 mg." Which generic is correct?

- a) Cefdinir
- b) Clarithromycin
- c) Ciprofloxacin
- d) Clindamycin

80) A pharmacy is notified of a significant loss of controlled substances. Which DEA form is typically used to report it?

- a) DEA Form 222
- b) DEA Form 106
- c) DEA Form 41
- d) DEA Form 363

81) A prescription reads: "Use 1 gram topically BID." The tube is 45 g. Approximate day supply is:

- a) 14 days
- b) 18 days
- c) 22 days
- d) 45 days

82) A new eRx includes weight-based vancomycin dosing notes and desired trough goals. What should the technician do?

- a) Adjust the dose to target the trough goal
- b) Enter exactly as written and route clinical dosing notes for pharmacist review
- c) Convert IV vancomycin to oral automatically
- d) Hold the prescription until a fixed dose is provided

83) A prescription is for "sertraline 50 mg daily." A new order arrives for linezolid. What is the technician's best action?

- a) Fill both; antibiotics override interactions
- b) Tell the patient to separate doses by 2 hours
- c) Route for pharmacist review due to serious interaction risk
- d) Remove sertraline from the profile and fill linezolid

84) A technician discovers an unlabeled vial of tablets returned from a nursing unit. What should be done?

- a) Identify by imprint and restock if it seems to match
- b) Quarantine as unknown/suspect medication and follow facility disposition procedure
- c) Combine into a "miscellaneous" bottle for later sorting
- d) Dispense if the color looks right

85) A prescription is for "apixaban." Which brand name is correct?

- a) Xarelto
- b) Eliquis
- c) Lovenox
- d) Coumadin

86) A prescription for "acetaminophen/codeine 300/30 mg" (Tylenol #3) is received. What schedule is this combination under federal law?

- a) Schedule II
- b) Schedule III
- c) Schedule IV
- d) Schedule V

87) A patient has a new prescription with "Do not fill until 07/20/26." Today is 07/15/26. What is the most appropriate action?

- a) Fill now and hold in will-call until 07/20/26
- b) Change the date to today because the prescription is valid
- c) Enter and place on hold to be filled on/after 07/20/26
- d) Transfer to another pharmacy to fill early

88) A prescription reads: "instill 5 gtt AS BID x 7 days" for an otic suspension. What is the correct label translation?

- a) Place 5 drops into the left ear twice daily for 7 days
- b) Place 5 drops into both ears twice daily for 7 days
- c) Place 5 drops into the right ear twice daily for 7 days
- d) Place 5 drops into the left eye twice daily for 7 days

89) A prescription is for "gabapentin 300 mg: take 2 caps PO TID. Qty 180." Day supply is:

- a) 45 days
- b) 60 days
- c) 20 days
- d) 30 days

90) A technician is asked what DEA Form 41 is used for. The correct answer is:

- a) Report significant loss/theft
- b) Order Schedule II controlled substances

- c) Register a new pharmacy location
- d) Document controlled substance destruction

Answer Key

1) b	31) c	61) b
2) b	32) d	62) c
3) d	33) c	63) b
4) b	34) b	64) a
5) a	35) b	65) d
6) c	36) d	66) c
7) a	37) b	67) c
8) d	38) a	68) b
9) b	39) b	69) c
10) c	40) c	70) b
11) b	41) b	71) d
12) c	42) a	72) b
13) a	43) c	73) c
14) b	44) b	74) b
15) c	45) d	75) d
16) d	46) b	76) d
17) b	47) c	77) a
18) c	48) c	78) a
19) d	49) c	79) c
20) a	50) d	80) b
21) c	51) b	81) b
22) b	52) a	82) b
23) c	53) c	83) c
24) b	54) c	84) b
25) c	55) c	85) b
26) d	56) a	86) b
27) a	57) b	87) c
28) a	58) d	88) a
29) b	59) b	89) d
30) b	60) c	90) d

PTCB Sim Test 11

1) A new eRx arrives for “Zoloft 50 mg: take 1 daily.” The prescriber selected “DAW 1” and the patient’s plan requires prior authorization for brand but covers generic. What should the technician enter?

- a) Generic sertraline because it’s equivalent and covered
- b) Brand Zoloft because DAW 1 indicates dispense as written
- c) Generic sertraline and annotate “brand not covered”
- d) Hold entry until the patient agrees to generic

2) A patient asks to buy pseudoephedrine 30 mg tablets, 96-count. The register prompts for logbook completion. Which action is required before completing the sale?

- a) Record the purchaser’s ID information and obtain signature/electronic acknowledgment
- b) Ask the patient to sign a HIPAA form
- c) Require a diagnosis code on a prescription
- d) Limit to one box per visit under federal law

3) A prescription reads: “Augmentin 400 mg/57 mg per 5 mL: 6 mL PO BID x 10 days.” How many mL should be dispensed?

- a) 60 mL
- b) 100 mL
- c) 120 mL
- d) 150 mL

4) While entering a new order, the technician sees two profiles: “Maria L. Santos” DOB 02/03/1982 and “Maria L Santos” DOB 02/03/1983 at the same address. An eRx for metoprolol succinate ER 25 mg arrives. What should the technician do next?

- a) Choose the profile with the most recent pickup activity
- b) Merge the profiles immediately and proceed
- c) Verify additional identifiers (phone, full DOB) before selecting the correct profile
- d) Enter it under either profile and correct it at pickup

5) A vial of insulin glargine pens arrived in a tote with a temperature indicator showing out-of-range exposure during transit. What is the correct technician action?

- a) Refrigerate immediately and dispense first to minimize waste
- b) Quarantine the product per procedure for pharmacist disposition
- c) Dispense if the patient will use within 28 days of opening
- d) Return to shelf stock but note the excursion on the invoice

6) A patient wants to transfer their clonazepam prescription from another pharmacy. What is within a technician’s scope?

- a) Take the transfer by phone and document all required elements
- b) Request the transfer and route it to the pharmacist to complete/authorize the transfer
- c) Refuse because controlled substance transfers are never allowed
- d) Tell the patient to bring in the original hard copy for verification

7) The sig says: “Instill 1 gtt OU qHS.” Which interpretation should print on the label?

- a) Instill 1 drop in both eyes at bedtime
- b) Instill 1 drop in each eye every morning
- c) Instill 2 drops in the right eye at bedtime
- d) Instill 1 drop in the left eye four times daily

8) A prescription is for “amLODIPine 5 mg: take 1 daily.” When selecting an NDC, two bottles are available: amlodipine 5 mg tablets and amlodipine 5 mg/benazepril 10 mg capsules. What is the best verification step before pulling stock?

- a) Verify dosage form and active ingredients match the label exactly before scanning/pulling
- b) Choose the cheapest NDC to reduce copay
- c) Choose whichever is on the same shelf section as “amlodipine”
- d) Pull the combination product because it’s commonly used for BP

9) A patient profile lists “ACE inhibitors—angioedema.” An eRx for lisinopril 10 mg arrives. What should the technician do?

- a) Override the alert if the prescriber ordered it electronically
- b) Change to losartan 50 mg and process
- c) Continue data entry but print with no allergy warnings
- d) Stop the workflow and ensure the pharmacist reviews the allergy conflict

10) A prescription reads: “metformin 500 mg tab: take 1 tab BID.” The patient has filled metformin ER 500 mg BID for the last year. What is the best technician action during entry?

- a) Enter metformin ER to match the patient history
- b) Enter metformin IR as written and proceed without question
- c) Enter metformin IR as written and flag it for pharmacist review due to formulation discrepancy
- d) Change directions to once daily to reduce GI upset

11) A label is printed for Toprol-XL 50 mg, but the technician pulled metoprolol tartrate 50 mg. What is the appropriate next step?

- a) Change the directions to BID to match metoprolol tartrate and continue
- b) Replace with metoprolol succinate ER 50 mg and document the near-miss per QA process
- c) Dispense metoprolol tartrate since the strength matches
- d) Ask the patient which one they were expecting and dispense that

12) A prescription: “amoxicillin 250 mg/5 mL: give 7.5 mL PO TID x 7 days.” How many mL should be dispensed?

- a) 105 mL
- b) 120 mL
- c) 157.5 mL
- d) 175 mL

13) A patient asks at pickup, “Can I drink alcohol on metronidazole?” What should the technician do?

- a) Advise avoiding alcohol for 72 hours after the last dose
- b) Tell the patient a single drink is fine if taken with food
- c) Refer the question to the pharmacist

d) Suggest spacing alcohol 2 hours away from the dose

14) The pharmacy receives a manufacturer recall notice for a specific lot of atorvastatin 20 mg. The shelf has two lots with identical expiration dates. What should the technician do?

- a) Remove all atorvastatin 20 mg from the shelf and return everything
- b) Quarantine only the recalled lot and follow the recall instructions for disposition
- c) Dispense the recalled lot first to reduce inventory loss
- d) Cover the lot number on the recalled bottles and continue using them

15) A C-II paper prescription for oxycodone 5 mg appears altered: the quantity has darker ink and overwritten numerals. What should the technician do?

- a) Fill it as it appears because the prescriber signed it
- b) Ask the patient to initial the change and proceed
- c) Set aside and have the pharmacist evaluate/verify with the prescriber before processing
- d) Call the prescriber and accept a verbal confirmation to proceed

16) A prescription reads: "prednisone 10 mg: take 4 tabs daily x 3 days, then 3 tabs daily x 3 days, then 2 tabs daily x 3 days, then 1 tab daily x 3 days." How many tablets are needed?

- a) 24
- b) 30
- c) 36
- d) 40

17) A controlled substance inventory is being prepared. Under federal requirements, which schedule requires a separate inventory record from all other schedules?

- a) Schedule II
- b) Schedule III
- c) Schedule IV
- d) Schedule V

18) A patient requests an early fill of Adderall due to "lost medication." Which is the most appropriate technician action?

- a) Process as "lost" and bill insurance immediately
- b) Tell the patient early fills are permitted once every 30 days for C-II's
- c) Offer a partial fill without involving the pharmacist
- d) Route the request to the pharmacist for evaluation and prescriber contact if needed

19) A technician spills powder from a crushed hazardous tablet (not in blister packaging) on the counting tray. What should the technician do first?

- a) Wipe with alcohol and continue filling
- b) Put on appropriate PPE and follow hazardous drug spill cleanup procedure
- c) Sweep into regular trash and wash hands
- d) Rinse the tray in the sink and return it to use

20) A prescription for nitroglycerin sublingual tablets is being prepared. Which auxiliary instruction is most appropriate?

- a) Store in original container; protect from light and moisture
- b) Take with food to prevent upset stomach

- c) Refrigerate after opening
- d) Take on a full stomach to reduce headache

21) A prescription reads: "diclofenac 1% gel: apply 2 g to affected area QID." The tube contains 100 g. Approximately how many days will it last?

- a) 5 days
- b) 10 days
- c) 12.5 days
- d) 25 days

22) A hospital technician restocking an ADC notices heparin 1,000 units/mL and 10,000 units/mL are stored next to each other with similar packaging. What is the best safety step?

- a) Place both strengths in the same pocket to reduce selection errors
- b) Separate storage locations and use barcode scanning during stocking and removal
- c) Remove the warning stickers so labels are easier to read
- d) Pre-draw syringes for nursing units to standardize doses

23) A patient's profile shows "sulfonamide antibiotics—hives." A new prescription is for Bactrim DS. What is the technician's best action?

- a) Process normally because hives is a mild reaction
- b) Substitute nitrofurantoin because it's a common alternative
- c) Ensure the pharmacist reviews the allergy before the medication is dispensed
- d) Ask the patient if they still want to fill it and document their answer

24) A prescription reads "levothyroxine 0.075 mg PO daily." The stocked strength is 75 mcg tablets. What should the technician enter?

- a) 0.075 mg daily
- b) 75 mcg daily
- c) 7.5 mcg daily
- d) 750 mcg daily

25) A patient asks for a copy of their medication profile to give to a new doctor. What is the best technician action?

- a) Print and hand it over without verification since it's the patient
- b) Verify identity using appropriate identifiers before releasing the printout
- c) Refuse because profile printouts are PHI that can never be released
- d) Provide only the drug names, not strengths or directions

26) A prescription reads: "Ventolin HFA: inhale 2 puffs q6h PRN." The inhaler contains 200 actuations. Using the maximum frequency, what is the days' supply?

- a) 12 days
- b) 16 days
- c) 25 days
- d) 50 days

27) A technician is entering a ketorolac tablet prescription: "10 mg PO QID for 10 days." What is the appropriate next step?

- a) Enter as written and proceed

- b) Reduce duration to 5 days and proceed
- c) Ensure the pharmacist reviews because the duration exceeds typical limits and may be unsafe
- d) Substitute ibuprofen 800 mg because it's safer

28) A prescriber's office calls asking for the last fill date and quantity of a patient's gabapentin. The caller cannot provide a verifiable callback number. What should the technician do?

- a) Provide the information because it's for continuity of care
- b) Provide only the last fill date, not the quantity
- c) Obtain verified contact information and route to the pharmacist before disclosing PHI
- d) Refuse all requests from prescriber offices under HIPAA

29) An eRx arrives for "Ritalin 10 mg: take 1 tab BID; refills: 3." What is the correct processing action?

- a) Enter as written; Schedule II refills are allowed if eRx
- b) Enter with refills reduced to 0 and continue filling
- c) Ensure the pharmacist addresses the invalid refills because Schedule II prescriptions cannot be refilled
- d) Convert to a Schedule III entry so the system accepts refills

30) A prescription reads: "ondansetron ODT 4 mg: dissolve 1 tab q8h PRN N/V." Quantity is 18 tablets. If taken around-the-clock at the maximum frequency, what is the days' supply?

- a) 2 days
- b) 3 days
- c) 6 days
- d) 9 days

31) A technician is preparing to reconstitute an oral antibiotic suspension. Which step should be done before adding water?

- a) Add all water at once to reduce foaming
- b) Tap the bottle to loosen powder and verify the final volume instruction on the label
- c) Use warm water to dissolve faster
- d) Skip shaking to avoid bubbles

32) A prescription reads "Take ii caps PO TID x 10 days." How many capsules should be dispensed?

- a) 42
- b) 50
- c) 60
- d) 84

33) During data entry, a technician sees an active profile medication for apixaban 5 mg BID and a new prescription for warfarin 5 mg daily. What is the best next step?

- a) Inactivate apixaban and fill warfarin
- b) Fill both because prescribers may overlap therapy intentionally
- c) Split warfarin to 2.5 mg daily to reduce bleeding risk
- d) Ensure the pharmacist reviews for therapeutic duplication/high-risk anticoagulation

overlap

34) A patient on isotretinoin arrives to pick up. The system indicates iPLEDGE requirements are not complete. What should happen?

- a) Dispense because the prescription is valid and the patient is present
- b) Hold the prescription until iPLEDGE requirements are confirmed complete
- c) Dispense only a 3-day emergency supply
- d) Substitute tretinoin cream and dispense instead

35) A pharmacy is asked to provide DSCSA product tracing for a recalled prescription drug. What should the technician do?

- a) Gather the transaction information/transaction statement and route per DSCSA procedure
- b) Delete the receiving record to prevent audit issues
- c) Dispense remaining stock before returning paperwork
- d) Re-label the recalled items with a new NDC

36) A prescription is written "Keflex 500 mg." Which product should be selected?

- a) cefdinir
- b) ceftriaxone
- c) cephalixin
- d) cefuroxime

37) A label prints with the wrong patient name but correct drug and directions. What is the best immediate action?

- a) Cross out the name and write the correct name neatly
- b) Reprint the correct label and document the event per QA procedure
- c) Keep the label and correct it at pickup after ID check
- d) Place in will-call and wait for the patient to notice

38) A technician receives a request for test strips with sig "use as directed." The Medicaid plan rejects because frequency is missing. What is the best next step?

- a) Enter "QD" as a default and resubmit
- b) Reduce quantity to the plan minimum and resubmit
- c) Bypass insurance and charge cash price
- d) Route to obtain/confirm documented testing frequency so the claim can be billed correctly

39) A technician is repackaging tablets into unit-dose packages for a facility. Which information must be on the repackaged label?

- a) Drug name and strength only
- b) Drug name/strength, lot number, and expiration date or assigned BUD per policy
- c) Patient name and prescription number
- d) Prescriber name and DEA number

40) A patient requests to return an opened bottle of hydrocodone/APAP for a refund. What is the correct action?

- a) Accept return and place back into stock if count matches
- b) Accept return only if the bottle is mostly full
- c) Do not accept for re-dispensing; route to pharmacist for disposal guidance if needed

d) Accept and re-dispense only to the same patient later

41) A prescription reads: "fluconazole 150 mg: take 1 now and repeat in 3 days." Quantity to dispense is:

- a) 1 tablet
- b) 2 tablets
- c) 3 tablets
- d) 4 tablets

42) The technician is processing a prescription for "Novolog FlexPen." Which is the correct generic product?

- a) insulin glargine U-100 pen
- b) insulin aspart pen
- c) insulin lispro pen
- d) insulin regular U-100 vial

43) A patient profile shows pregnancy, and a new prescription arrives for rosuvastatin 20 mg daily. What should the technician do?

- a) Process because only high-dose statins are contraindicated
- b) Change to pravastatin and process
- c) Ensure the pharmacist reviews due to contraindication/therapy concern
- d) Ask the patient which trimester and decide based on that

44) A prescription reads: "Take 1 cap PO qHS." Quantity is #90. What is the days' supply?

- a) 30
- b) 45
- c) 60
- d) 90

45) A patient presents a paper prescription for a C-III medication. The prescriber's address is missing from the printed prescription. What should the technician do?

- a) Add the address from a prior prescription and continue
- b) Add the address from an online search and continue
- c) Process as a non-controlled prescription since it's signed
- d) Set aside for pharmacist review to determine whether required elements are met before dispensing

46) A technician is entering an inhaled medication: "inhale 2 puffs QID." The inhaler contains 120 actuations. What is the correct days' supply?

- a) 10 days
- b) 12 days
- c) 15 days
- d) 30 days

47) A prescription reads: "clindamycin 150 mg caps: take 2 caps PO QID x 7 days." How many capsules should be dispensed?

- a) 28
- b) 42

- c) 56
- d) 84

48) A prescription is written for "Azor." Which combination is correct?

- a) amlodipine/olmesartan
- b) amlodipine/valsartan
- c) losartan/hydrochlorothiazide
- d) lisinopril/amlodipine

49) A patient reports "these tablets look different" after a refill. The NDC changed due to a different manufacturer substitution, and the label is correct. What is the best technician action?

- a) Refile the prescription; generics always look different
- b) Replace with the old manufacturer without changing anything in the system
- c) Verify the scanned NDC matches the label and route patient concern to the pharmacist if needed
- d) Open multiple stock bottles to compare tablets by taste and texture

50) While typing an eRx, the technician notices "metoprolol succinate ER 25 mg: take 1 tab BID." What should happen next?

- a) Enter as written and proceed
- b) Change to once daily automatically because it's ER
- c) Change to metoprolol tartrate 25 mg BID automatically
- d) Ensure the pharmacist reviews for appropriateness/clarification due to unusual dosing for ER

51) A patient asks the technician to recommend an OTC product for "new chest tightness and pain that goes to my left arm." What should the technician do?

- a) Recommend omeprazole OTC for 14 days
- b) Recommend ibuprofen and rest
- c) Refer to pharmacist and advise immediate medical evaluation/emergency care
- d) Recommend a decongestant and fluids

52) A prescription reads: "Take 1/2 tab PO daily." Strength is 10 mg tablets. Quantity written is #15 for 30 days. What is the issue?

- a) Quantity is correct
- b) Quantity should be #30
- c) Quantity should be #60
- d) Directions are invalid because tablets cannot be split

53) A prescription is for "eye drops: instill 2 gtt OS q4h while awake x 5 days." A 5 mL bottle contains ~20 gtt/mL. How many bottles are needed (assume 16 waking hours/day)?

- a) 1 bottle
- b) 2 bottles
- c) 3 bottles
- d) 4 bottles

54) A patient asks to partially fill a Schedule II prescription because they can't afford the full

quantity. What is the best technician action?

- a) Split the quantity into two fills under the same prescription number
- b) Tell the patient partial fills are never allowed for C-II
- c) Ensure the pharmacist handles the request because partial-fill rules and documentation are strict
- d) Change the quantity on the label and complete the remainder later without documentation

55) A prescription is for "MS Contin 15 mg." Which is correct?

- a) morphine sulfate extended-release
- b) morphine sulfate immediate-release
- c) oxycodone extended-release
- d) hydromorphone immediate-release

56) A hospital unit calls stating an ADC count for a controlled substance is short and requests "send one more vial now." What is the appropriate technician action?

- a) Deliver a vial immediately to avoid delaying care
- b) Ask the nurse to email a note and then send it
- c) Document the call in a notebook and deliver later
- d) Route per controlled substance discrepancy/ADC procedures for pharmacist oversight

57) A prescriber calls in a verbal prescription for amoxicillin. The state allows trained technicians to receive non-controlled verbal orders, but pharmacist verification is required before dispensing. What should the technician do?

- a) Take the verbal order, document required elements, and route for pharmacist verification
- b) Refuse because technicians can never take any verbal orders
- c) Take the order and dispense once entered without further review
- d) Ask the prescriber to fax it because verbal orders are invalid

58) During will-call cleanup, the technician finds a filled prescription that has been sitting for 16 days. The medication is an antibiotic suspension that was reconstituted with a 14-day BUD. What is the best action?

- a) Return it to stock as long as the original manufacturer expiration date hasn't passed
- b) Replace with a newly reconstituted bottle before dispensing
- c) Dispense but instruct the patient to use it quickly
- d) Change the BUD on the label to match the manufacturer date

59) A prescription is for "Lantus SoloStar." Which storage instruction is most appropriate before first use?

- a) Refrigerate; do not freeze
- b) Store in freezer until needed
- c) Store at room temperature only, even before first use
- d) Refrigerate after opening only

60) A technician is asked to provide the results of a patient's INR test and warfarin dose changes over the phone. What should the technician do?

- a) Provide the INR result and recommend a dose adjustment
- b) Provide only the INR result and tell the patient to decide the dose
- c) Refer the patient to the pharmacist; do not provide clinical interpretation or dosing

instructions

d) Tell the patient to double the dose if the INR is low

61) A prescription is for "Eliquis." Which is the correct generic name?

a) rivaroxaban

b) apixaban

c) dabigatran

d) clopidogrel

62) A prescription label prints: "Take 1 tablet by mouth twice daily" for clopidogrel 75 mg. What is the best action during processing?

a) Change to once daily automatically because it's standard

b) Fill as written; prescribers may intentionally double the dose

c) Ensure the pharmacist reviews the dose directions for verification

d) Ask the patient at pickup which schedule they prefer

63) A technician finds an unlabeled bottle of tablets on the shelf with unknown identity. What should be done?

a) Use a pill identifier app and relabel it

b) Combine with a similar-looking stock bottle

c) Quarantine for pharmacist review and proper disposal procedure

d) Return to the shelf based on color and shape

64) A prescription: "diltiazem 120 mg ER daily." The prescriber calls later and states they meant immediate-release. Who must authorize and make that change?

a) Technician, because it came directly from the prescriber

b) Technician, as long as notes are entered

c) Either technician or pharmacist if the patient agrees

d) Pharmacist, because it is a formulation change requiring professional verification

65) A prescription reads: "Take 5 mL PO TID." Concentration is 250 mg/5 mL. How many mg per dose?

a) 50 mg

b) 125 mg

c) 250 mg

d) 750 mg

66) A prescription for tramadol 50 mg is received electronically. The prescriber's address field is blank in the eRx. What should the technician do?

a) Add the address from an internet search and proceed

b) Proceed because eRx controlled prescriptions do not require an address

c) Set aside for pharmacist review to determine if the prescription contains required elements

d) Convert it to a non-controlled prescription entry to avoid compliance issues

67) A technician is asked to shred old hard copies containing PHI. Which is the best practice?

a) Shred only controlled substance prescriptions

- b) Remove only insurance ID numbers; names may remain
- c) Use approved secure destruction for all PHI-containing documents
- d) Keep all originals intact and discard only copies

68) A prescription is for “omeprazole 20 mg daily,” but the profile shows an active pantoprazole 40 mg daily from another prescriber. What should the technician do?

- a) Inactivate pantoprazole automatically and fill omeprazole
- b) Fill both; the patient can decide which to take
- c) Ensure the pharmacist reviews for therapeutic duplication and appropriateness
- d) Switch omeprazole to famotidine and fill

69) A prescription is written for “hydroCHLOROthiazide 25 mg qAM.” The patient profile lists “sulfonamides—anaphylaxis.” What is the best next step?

- a) Fill because HCTZ is not an antibiotic
- b) Delete the allergy because it’s likely outdated
- c) Change to furosemide automatically and proceed
- d) Ensure the pharmacist reviews the allergy/clinical relevance before dispensing

70) A technician is preparing to compound a nonsterile oral solution. Which must be verified before starting?

- a) Prescriber’s DEA number
- b) Master formulation record and ingredient expiration dates
- c) Patient’s insurance BIN/PCN
- d) Wholesaler acquisition cost

71) A patient’s mother asks for her 16-year-old child’s immunization record printout; the child is not present. What should the technician do?

- a) Provide it because parents always have access
- b) Provide only vaccine dates, not vaccine names
- c) Refer for authorization verification per privacy rules before releasing
- d) Refuse and instruct them to call the pediatrician

72) A prescription is written: “Take 1 tablet PO daily.” Medication is amiodarone 200 mg. The profile shows digoxin 0.25 mg daily and an interaction alert appears. What should the technician do?

- a) Override because both are common cardiac medications
- b) Ensure the pharmacist reviews the interaction before dispensing
- c) Delete digoxin from the profile to clear the alert
- d) Change amiodarone to 100 mg and proceed

73) A patient brings in a C-II hard copy dated 04/01/26. The signature line is signed, but the date next to the signature line is blank. What should the technician do?

- a) Fill because the written date is present
- b) Add today’s date next to the signature
- c) Set aside for pharmacist determination of validity/required elements
- d) Ask the patient to write the date and initial it

74) A prescription is written: “sumatriptan 100 mg: take 1 at onset, may repeat once in 2

hours.” The patient profile note says “uncontrolled hypertension.” What should the technician do?

- a) Continue processing; technicians do not consider conditions
- b) Delete the note to prevent future alerts
- c) Change to acetaminophen 1,000 mg and proceed
- d) Ensure the pharmacist reviews for appropriateness due to condition warning

75) A technician is asked to restock a high-alert medication in a hospital cart fill. Which item is high-alert and commonly requires extra safeguards?

- a) docusate sodium
- b) acetaminophen 325 mg
- c) potassium chloride concentrate
- d) loratadine 10 mg

76) A prescription reads: “Budesonide/formoterol 160/4.5: inhale 2 puffs BID.” The patient asks if it replaces their rescue inhaler. What is the best technician response?

- a) Yes, use it only as needed for sudden symptoms
- b) No, it’s a maintenance inhaler; refer clinical use questions to the pharmacist
- c) Yes, stop albuterol if you have this inhaler
- d) It depends on insurance coverage

77) A patient’s prescription label prints “instill 1 gtt OD BID.” What does OD mean in this context?

- a) Right eye
- b) Left eye
- c) Both eyes
- d) By mouth

78) A prescription is received for “U-500 insulin: inject 50 units daily,” but the order does not specify device/syringe details and the patient is new to therapy. What should the technician do?

- a) Enter as written and select U-100 syringes
- b) Convert the dose to U-100 units and dispense U-100 insulin
- c) Ensure pharmacist review/clarification occurs due to high-concentration insulin safety concerns
- d) Change to insulin glargine because it is safer

79) A prescription is written “brand medically necessary” for Synthroid 88 mcg. Which entry is correct?

- a) Substitute levothyroxine generic automatically and proceed
- b) Dispense brand Synthroid as indicated and bill accordingly
- c) Change to Euthyrox because it is also levothyroxine
- d) Dispense generic and include a note that it is equivalent

80) A prescription: “cephalexin suspension: 250 mg/5 mL; take 10 mL BID x 10 days.” How many mL are needed?

- a) 100 mL
- b) 150 mL

- c) 200 mL
- d) 250 mL

81) A technician is asked to handle a Medication Guide that must be dispensed with certain medications. What is the correct workflow action?

- a) Provide the Medication Guide only on the first fill
- b) Provide only if the patient requests it
- c) Provide only for cash-paying patients
- d) Ensure the Medication Guide is provided when required at dispensing

82) A technician receives a fax requesting a transfer of a Schedule IV prescription with refills remaining. Which is the best next step?

- a) Transfer immediately; Schedule IV transfers never require pharmacist involvement
- b) Deny because controlled substances can't be transferred
- c) Route to the pharmacist to determine transfer eligibility and provide required transfer details
- d) Transfer but omit the original fill date to protect privacy

83) A pharmacy receives two look-alike products in the same bin location due to a stocking error: bupropion SR 150 mg and bupropion XL 150 mg. What is the best corrective action?

- a) Leave them together but add a "look alike" sticker
- b) Separate them into different locations and report the event per QA/near-miss process
- c) Place them in will-call so staff will see them more often
- d) Remove SR and keep only XL since it's more common

84) A prescription label states: "Take 1 tablet PO daily." Quantity #30, directions indicate 1 tablet daily. The claim rejects "refill too soon (75% usage)." The last fill was 20 days ago. What is the earliest day the plan will typically allow the refill?

- a) Day 15
- b) Day 20
- c) Day 23
- d) Day 30

85) A prescription is written for "morphine ER 30 mg." The technician accidentally selects morphine IR 30 mg in the system. What is the best action?

- a) Keep morphine IR and change sig to q12h
- b) Correct to the ER product and document the near-miss per QA process
- c) Dispense IR because the strength matches and patient can take less often
- d) Ask the patient whether they want ER or IR

86) A patient requests a refill for a Schedule III medication, but the prescription has zero refills remaining. What should the technician do?

- a) Add one refill to prevent interruption
- b) Dispense a 3-day emergency supply without pharmacist involvement
- c) Send a renewal/refill request through the appropriate prescriber contact workflow
- d) Transfer the prescription to another pharmacy to renew it

87) A medication label prints for "prednisoLONE acetate ophthalmic suspension," but the

technician pulls "prednisoLONE sodium phosphate ophthalmic solution." What is the best prevention step?

- a) Choose whichever is cheaper; both are prednisolone
- b) Use barcode scanning and verify full drug name/salt form and dosage form before dispensing
- c) Dispense the solution and add "shake well"
- d) Change the directions to match the product pulled

88) A prescription for eye drops is: "latanoprost 0.005%: instill 1 drop OU qPM." The bottle size is 2.5 mL (20 gtt/mL). Approximate days' supply?

- a) 12 days
- b) 25 days
- c) 50 days
- d) 75 days

89) A technician receives a request from a third-party shred vendor to pick up bins containing pharmacy documents. Which is the best action before release?

- a) Allow pickup as long as bins are sealed
- b) Verify vendor authorization and chain-of-custody requirements are met per procedure
- c) Remove patient names but leave other identifiers
- d) Send only controlled-substance records for shredding

90) A prescription reads: "Apply 1 patch to skin daily for 12 hours on, 12 hours off." Quantity #30 patches. What is the days' supply?

- a) 15 days
- b) 30 days
- c) 45 days
- d) 60 days

Answer Key

1) b	32) c	63) c
2) a	33) d	64) d
3) c	34) b	65) c
4) c	35) a	66) c
5) b	36) c	67) c
6) b	37) b	68) c
7) a	38) d	69) d
8) a	39) b	70) b
9) d	40) c	71) c
10) c	41) b	72) b
11) b	42) b	73) c
12) c	43) c	74) d
13) c	44) d	75) c
14) b	45) d	76) b
15) c	46) b	77) a
16) c	47) d	78) c
17) a	48) a	79) b
18) d	49) c	80) c
19) b	50) d	81) d
20) a	51) c	82) c
21) b	52) a	83) b
22) b	53) a	84) c
23) c	54) c	85) b
24) b	55) a	86) c
25) b	56) d	87) b
26) b	57) a	88) b
27) c	58) b	89) b
28) c	59) a	90) b
29) c	60) c	
30) b	61) b	
31) b	62) c	

PTCB Sim Test 12

1. A new eRx reads: "levothyroxine 75 mcg tab. Take 1 tab PO daily. Qty 90." The product list shows levothyroxine 75 mcg and liothyronine 25 mcg. What is the best selection?

- a) Liothyronine 25 mcg tablets
- b) Levothyroxine 75 mcg tablets
- c) Levothyroxine 50 mcg tablets
- d) Levothyroxine 75 mcg capsules

2. A patient brings a paper prescription for hydrocodone/acetaminophen (C-II) with the prescriber signature but no date written. What should the technician do next?

- a) Enter it and dispense; the computer timestamps it
- b) Ask the patient to write today's date on it
- c) Set it aside for pharmacist review before any processing
- d) Call the prescriber and add the date after verbal confirmation

3. While typing an eRx, the technician sees "metoprolol succinate ER 25 mg daily," but the selected item is metoprolol tartrate 25 mg. What is the best action before moving forward?

- a) Change the sig to BID and keep tartrate
- b) Change to succinate ER 25 mg as written
- c) Keep tartrate and add "extended release" in notes
- d) Keep tartrate because it's the same drug

4. A prescription reads: "amoxicillin 500 mg caps. Take 1 cap PO TID x 7 days." What quantity should be dispensed?

- a) 14
- b) 20
- c) 21
- d) 28

5. A customer requests a pseudoephedrine product and presents an unexpired ID. The system indicates the customer has reached the daily limit. What should the technician do?

- a) Decline the sale and offer a nonrestricted alternative
- b) Split the transaction into two smaller purchases
- c) Sell it as an OTC item without logging it
- d) Sell it if the customer pays cash

6. A patient's spouse calls and asks, "Is her prescription ready, and what is it?" The spouse is not listed as authorized and cannot verify any additional identifiers. What can the technician disclose?

- a) The medication name and directions only
- b) Only that a prescription is or is not ready for pickup
- c) The prescriber name and drug name
- d) Full details if the spouse knows the address

7. A new Rx is for "Latuda 40 mg daily." The technician accidentally selects "Lamisil 250 mg." Which safeguard most directly prevents this type of error at the selection step?

- a) NDC barcode scan match of stock bottle to the label
- b) Checking the patient's diagnosis in the chart
- c) Counting the tablets twice
- d) Using amber vials instead of clear vials

8. A vial label prints with another patient's name due to a queue mix-up. No medication has been counted yet. What should the technician do?

- a) Cross out the name and write the correct one
- b) Reprint the correct label and place the incorrect label in secure PHI destruction
- c) Save the label to reuse later for the correct patient
- d) Discard the label in regular trash

9. A prescription reads: "prednisone 10 mg tabs: 4 tabs daily x 3 days, then 3 tabs daily x 3 days, then 2 tabs daily x 3 days, then 1 tab daily x 3 days." What quantity is needed?

- a) 24
- b) 30
- c) 36
- d) 40

10. A patient profile shows "allergy: penicillin—hives." A new prescription arrives for cephalexin. What should the technician do?

- a) Override the alert because cross-reactivity is impossible
- b) Change cephalexin to azithromycin
- c) Delete the penicillin allergy from the profile
- d) Route for pharmacist assessment before dispensing

11. A prescription reads: "Eliquis 5 mg: take 1 tablet PO BID." The profile shows an active warfarin regimen filled 10 days ago. What is the best processing step for the technician?

- a) Deactivate warfarin and process Eliquis
- b) Process Eliquis and note "switch" in the comments
- c) Route for pharmacist review for potential duplicate anticoagulation therapy
- d) Reduce Eliquis to a 7-day supply and process

12. A new eRx says "ondansetron ODT 4 mg." The selected product is ondansetron 4 mg tablets (not ODT). What should the technician do?

- a) Keep tablets; the strength is the same
- b) Keep tablets and add "dissolve on tongue"
- c) Change to ODT as written before filling
- d) Change to oral solution to avoid confusion

13. A prescription reads: "albuterol HFA: inhale 2 puffs q4h PRN. Dispense 1 inhaler (200 actuations)." If used at maximum frequency, what is the day supply?

- a) 16 days
- b) 25 days
- c) 17 days
- d) 8 days

14. A patient requests to return an unopened, dispensed bottle of tablets because therapy

changed. What is the usual community pharmacy handling?

- a) Restock if the seal is intact
- b) Restock after visual inspection
- c) Do not restock; handle through return/disposal process
- d) Restock only if it is noncontrolled

15. A new prescription reads: "insulin glargine 100 units/mL pens. Inject 18 units SQ QHS. Dispense 1 box (5 pens)." Each pen is 3 mL. What day supply should be billed?

- a) 83
- b) 75
- c) 90
- d) 50

16. A technician receives an eRx for isotretinoin. The system requires REMS verification before dispensing. Which REMS program is associated with isotretinoin?

- a) MedWatch
- b) iPLEDGE
- c) VAERS
- d) PDMP

17. A prescription reads: "clindamycin 300 mg caps. Take 1 cap PO QID. Dispense #28." How many days is the course?

- a) 5
- b) 14
- c) 7
- d) 10

18. A technician is asked to take a new verbal prescription from a prescriber over the phone. In most outpatient/community settings, what should the technician do?

- a) Take the verbal if it's noncontrolled
- b) Take the verbal and have the pharmacist sign later
- c) Ask the prescriber to fax it instead of calling
- d) Transfer the call to the pharmacist to receive the verbal order

19. A prescription reads: "fentanyl 25 mcg/hr patch. Apply 1 patch every 72 hours. Dispense #5." What day supply should be billed?

- a) 10 days
- b) 12 days
- c) 15 days
- d) 30 days

20. A patient says, "My bottle was stolen; insurance says refill too soon." What is the best technician next step?

- a) Change the NDC and rebill to force approval
- b) Gather details and route through the proper override/documentation workflow
- c) Tell the patient early refills are never allowed
- d) Run the claim as cash and avoid insurance

21. A prescription is written for "Crestor 20 mg" with DAW 1. What should be dispensed?

- a) Brand Crestor 20 mg
- b) Generic rosuvastatin 20 mg
- c) Rosuvastatin 40 mg half tablets
- d) Any statin equivalent covered by insurance

22. A prescription reads: "Apply 1 patch every 72 hours. Dispense #10." What day supply is this?

- a) 10 days
- b) 20 days
- c) 30 days
- d) 72 days

23. The pharmacy receives a DSCSA tracing request for a lot of atorvastatin received last month. Which record is used to respond?

- a) Patient signature log
- b) Controlled substance perpetual inventory
- c) Transaction information/transaction history from receiving records
- d) Prescription hard copies for that month

24. A technician notices a wholesaler bottle has a broken seal with tablets visible. What should be done?

- a) Stock it if the NDC matches the invoice
- b) Quarantine it as suspect product for return/investigation
- c) Tape it shut and mark "opened"
- d) Use it first to reduce waste

25. A prescription reads: "Azithromycin 250 mg tabs: Take 2 tabs today, then 1 tab daily on days 2–5." What quantity should be dispensed?

- a) 5
- b) 10
- c) 6
- d) 7

26. A hard copy arrives for a Schedule II drug missing the patient address. What is the most appropriate technician action?

- a) Add the address from the patient profile and proceed
- b) Ask the patient to write their address on the prescription
- c) Hold for pharmacist review to confirm legal acceptability before dispensing
- d) Refuse and tell the patient it can never be filled

27. A label reads: "Instill 1 gtt OU BID." What does OU mean?

- a) Right eye
- b) Left eye
- c) Both ears
- d) Both eyes

28. Rx: "enoxaparin 40 mg/0.4 mL syringes. Inject 0.4 mL SQ daily. Dispense #10." What is

the day supply?

- a) 14
- b) 10
- c) 7
- d) 20

29. A refrigerated vaccine shipment arrives; the temperature monitor indicates a possible excursion. What should the technician do first?

- a) Place it in the refrigerator and use it quickly
- b) Freeze it to stabilize it
- c) Quarantine the shipment and alert the pharmacist/manager for guidance and documentation
- d) Discard it immediately without documentation

30. A prescription reads: "diclofenac 1% gel. Apply 2 g to affected area QID." Tube size is 100 g. What is the day supply?

- a) 12 days
- b) 14 days
- c) 10 days
- d) 8 days

31. A patient asks whether grapefruit is safe with amiodarone. What should the technician do?

- a) Advise avoiding grapefruit and document the counseling
- b) Refer the interaction question to the pharmacist
- c) Recommend switching to a different antiarrhythmic
- d) Tell the patient it's safe because it's food, not medicine

32. A noncontrolled prescription is being transferred to another pharmacy. Which information is typically included?

- a) Only the Rx number and total refills authorized
- b) Diagnosis code only
- c) Only patient name and drug name
- d) Original date, drug/strength/directions, quantity, refills remaining, prescriber info, and last fill date

33. A prescription reads: "Bactrim DS: take 1 tab PO BID x 7 days." The prescriber wrote "Dispense #7." What should be done during processing?

- a) Enter quantity 7 because it was written
- b) Enter quantity 14 based on directions
- c) Enter quantity 10 as an intermediate amount
- d) Enter quantity 12 based on typical packaging

34. A Schedule III prescription with refills remaining is being transferred between two pharmacies that do not share a real-time database. Under federal rules, how many transfers are permitted?

- a) Unlimited transfers
- b) 0 transfers

- c) 1 transfer
- d) 2 transfers

35. A technician finds two patient profiles with the same name and similar DOBs. An eRx was attached to the wrong profile, but nothing has been filled. What is the best technician action?

- a) Move the prescription to the correct profile and document the near miss per QA process
- b) Continue processing since it wasn't dispensed
- c) Delete the eRx and request it be resent
- d) Merge the two profiles immediately without review

36. Rx: "valacyclovir 1 g tab. Take 1 tab PO TID x 7 days." What quantity should be dispensed?

- a) 14
- b) 28
- c) 20
- d) 21

37. A technician is preparing 150 mL of a 2% w/v oral solution. How many grams of drug are needed?

- a) 0.3 g
- b) 3 g
- c) 15 g
- d) 30 g

38. A controlled substance inventory is being performed. Which schedule requires an exact count (not an estimate) at inventory?

- a) Schedule II only
- b) Schedule III only
- c) Schedule II–V
- d) Schedule IV–V only

39. An eRx arrives for "diltiazem ER 240 mg daily" with no ER subtype. The system lists ER (CD), ER (LA), and ER (XR), and the patient has no prior history at your pharmacy. What should the technician do?

- a) Pick any ER 240 mg because all are equivalent
- b) Choose the least expensive ER option
- c) Enter diltiazem IR divided doses to match 240 mg/day
- d) Set aside for pharmacist clarification of formulation before dispensing

40. A prescription label prints "hydralazine 25 mg" but the hard copy is for "hydroxyzine 25 mg." What should the technician do?

- a) Continue filling and let the pharmacist catch it at final check
- b) Correct the drug selection/label before filling proceeds
- c) Dispense hydralazine because both are tablets
- d) Change the directions to match hydralazine

41. A prescription reads: "Take 1.5 tablets daily." Quantity dispensed is #45. What day

supply should be billed?

- a) 20 days
- b) 45 days
- c) 25 days
- d) 30 days

42. A prescriber office calls requesting a patient's oxycodone refill history. The caller cannot confirm DOB or address. What should the technician do?

- a) Provide last fill date only
- b) Provide the full refill history because it's a prescriber office
- c) Provide quantity only and omit dates
- d) Decline disclosure until proper verification/authorization is obtained

43. A prescription reads: "KCl oral solution 10 mEq/15 mL. Give 20 mEq PO now." What volume should be measured?

- a) 20 mL
- b) 30 mL
- c) 45 mL
- d) 15 mL

44. The pharmacy receives a manufacturer recall notice for certain lot numbers of losartan. What is the first operational step?

- a) Call every patient who has ever taken losartan
- b) Pull and quarantine the affected lots and follow recall instructions
- c) Return all losartan regardless of lot number
- d) Continue dispensing until replacement arrives

45. A prescription is written: "methylprednisolone dose pack 4 mg. Use as directed." What must the technician ensure during product selection?

- a) The patient's height and weight are documented
- b) The directions and quantity correspond to the specific dose pack product selected
- c) The prescriber DEA is present on the label
- d) The patient has no food allergies

46. An eRx includes "sertraline 50 mg daily." The profile shows an active "fluoxetine 20 mg daily" filled 2 weeks ago. What is the best technician action?

- a) Process both because SSRIs are commonly combined
- b) Inactivate fluoxetine and process sertraline
- c) Change sertraline to citalopram
- d) Route for pharmacist review for potential duplicate SSRI therapy

47. A prescription reads: "ciprofloxacin 0.3% ophthalmic. Instill 2 gtt OS q4h while awake x 7 days." A 5 mL bottle is dispensed. Using 20 gtt/mL, approximately how many total drops are in the bottle?

- a) 50
- b) 200
- c) 100
- d) 400

48. A patient asks what "qid" means on the label. What does it mean?

- a) Four times daily
- b) Once daily
- c) Every other day
- d) At bedtime

49. A technician is asked to process a partial fill of a Schedule II prescription due to insufficient stock. What is the best technician action?

- a) Partially fill and tell the patient the rest can be filled anytime within 6 months
- b) Refer the situation to the pharmacist to ensure timing/documentation rules are met
- c) Transfer the remainder to another pharmacy
- d) Convert the remainder to refills

50. A prescription reads: "brimonidine 0.2% eye drops. Instill 1 drop OU TID." Bottle size is 5 mL and 20 gtt/mL. Approximately what day supply is one bottle?

- a) 11 days
- b) 33 days
- c) 16 days
- d) 22 days

51. A patient requests "Zyrtec-D." Their photo ID is expired. What should the technician do under pseudoephedrine sale requirements?

- a) Complete the sale if the photo is clear
- b) Decline the sale because the ID is not valid for the logbook requirements
- c) Sell one box and document "expired ID" in the notes
- d) Sell it if the patient is a known customer

52. A prescription reads: "Take 5 mL q6h PRN fever." A 120 mL bottle is dispensed. If used at maximum frequency, what is the day supply?

- a) 5 days
- b) 8 days
- c) 4 days
- d) 6 days

53. A patient profile indicates "pregnant." A new prescription arrives for lisinopril. What should the technician do?

- a) Fill as written because the prescriber already considered risks
- b) Remove the pregnancy note since it could be outdated
- c) Change to labetalol per pregnancy guidelines
- d) Route for pharmacist review due to a significant safety concern

54. A technician is asked which element is required on most outpatient prescription labels under federal standards.

- a) Prescriber DEA number on all prescriptions
- b) Patient Social Security number
- c) Pharmacy name and address
- d) Patient diagnosis code

55. A technician is mixing a 1:4 dilution to a final volume of 240 mL. How many mL of stock solution are needed?

- a) 48 mL
- b) 192 mL
- c) 60 mL
- d) 80 mL

56. A prescription reads: "Nitrostat 0.4 mg SL tabs #25." The technician selected nitroglycerin transdermal patches. What step most directly prevents this error?

- a) Reviewing the prescriber specialty
- b) Scanning the stock bottle to match the label/NDC during selection
- c) Using a clean counting tray
- d) Adding auxiliary labels

57. An eRx for "bupropion XL 150 mg daily" is entered as bupropion SR 150 mg. What should the technician do?

- a) Keep SR because both are extended-release
- b) Change directions to once daily and dispense SR
- c) Correct the product to XL as written because SR and XL are not interchangeable
- d) Substitute immediate-release and adjust the sig

58. A patient requests their prescription profile be faxed to a new physician office. What should the technician do?

- a) Fax immediately because it's for care coordination
- b) Provide only the controlled substance portion of the profile
- c) Refuse because profiles cannot be shared
- d) Obtain required patient authorization and verify the destination fax number before sending

59. A technician is restocking a clinic refrigerator and finds an opened multidose vial labeled "opened 45 days ago," not past manufacturer expiration. What should be done?

- a) Keep using it until manufacturer expiration
- b) Use it first to reduce waste
- c) Remove from use per beyond-use dating and facility procedure
- d) Freeze it to extend stability

60. A prescription reads: "warfarin 2.5 mg: take 1 tab Mon/Wed/Fri and 2 tabs Tue/Thu/Sat/Sun. Dispense #50." What is the day supply?

- a) 25 days
- b) 21 days
- c) 28 days
- d) 30 days

61. A new eRx for "Tylenol #3" includes directions but no quantity. What should the technician do?

- a) Enter quantity 20 as a default and process
- b) Enter quantity 30 and process
- c) Clarify the missing quantity before dispensing

d) Dispense a 3-day supply and clarify later

62. A technician is cleaning a counting tray after handling methotrexate tablets and then needs to count a pediatric antibiotic. What is best practice?

- a) Use the same tray without cleaning to avoid delays
- b) Wipe the tray with a dry paper towel only
- c) Rinse the tray under water and continue
- d) Clean the tray using the approved method between products, with extra caution for hazardous drugs

63. A technician receives a "screenshot" on a phone of an electronic prescription for a Schedule II stimulant. What should the technician do?

- a) Treat it as a valid eRx and fill it
- b) Treat it as a fax and fill it
- c) Accept it if the prescriber's DEA is visible
- d) Explain the pharmacy must receive the prescription through a valid transmission method and request it be sent directly

64. A prescription reads: "tacrolimus 0.1% ointment apply BID." The technician selected triamcinolone 0.1% ointment because the strength matches. What should be done?

- a) Dispense triamcinolone because both treat dermatitis
- b) Substitute only if insurance rejects tacrolimus
- c) Dispense both and let the patient choose
- d) Select tacrolimus as written because they are not interchangeable

65. Rx: "amoxicillin/clavulanate 875/125 mg tab. 1 tab PO BID x 10 days. Dispense #20." What quantity should be dispensed?

- a) 10
- b) 14
- c) 20
- d) 28

66. During filling, tablets drop onto a counter that was recently sprayed with disinfectant. What should the technician do?

- a) Wipe the tablets and continue
- b) Return tablets to the stock bottle
- c) Rinse tablets and dry them
- d) Discard the contaminated tablets per procedure

67. A prescription reads: "fluticasone/salmeterol 250/50 Diskus: inhale 1 inhalation BID." The technician selected fluticasone HFA. What is the correct action?

- a) Dispense fluticasone HFA because it contains the steroid
- b) Correct to the combination product and device as prescribed
- c) Change directions to 2 puffs BID to match HFA
- d) Substitute budesonide/formoterol without authorization

68. A technician is processing an eRx with "take 1 tablet daily" in the structured sig field, but free-text notes say "take 2 tablets daily." Quantity is #60. What should be done?

- a) Enter 2 daily because quantity supports it
- b) Enter 1 daily because structured sig is always correct
- c) Clarify the conflicting directions before dispensing
- d) Enter 2 daily and change quantity to #30

69. A prescription reads: "Take 1 tablet PO q8h." How many doses per day is this?

- a) 2
- b) 3
- c) 4
- d) 6

70. A prescription for a Schedule II stimulant is dated 10 months ago and has not been filled. What should the technician do?

- a) Fill it because federal law has no expiration for Schedule II
- b) Fill a partial and void the remainder
- c) Hold for pharmacist review because validity may be limited by state law and other requirements
- d) Transfer it to another pharmacy with stock

71. A prescription reads: "omeprazole 20 mg daily #30." The patient says, "I always take 40 mg—change it." What should the technician do?

- a) Change to 40 mg because the patient reports it
- b) Change directions to take 2 capsules daily and process
- c) Dispense 40 mg but bill as 20 mg
- d) Keep as written and route the request for pharmacist assessment if a change is needed

72. A technician is asked which event is most accurately classified as a near miss.

- a) Wrong drug dispensed and taken by the patient
- b) Wrong label printed but caught before dispensing
- c) Manufacturer recall after the drug was dispensed
- d) Patient declined counseling

73. An order reads: "amoxicillin 400 mg/5 mL suspension. Give 7.5 mL PO BID x 10 days." How many mL should be dispensed?

- a) 75 mL
- b) 100 mL
- c) 150 mL
- d) 200 mL

74. While receiving controlled substances, the pharmacy uses DEA Form 222. Who may sign when executing/authorizing the order?

- a) Any technician with register access
- b) Any pharmacist on duty automatically
- c) The delivery driver after verifying the invoice
- d) The registrant or an authorized person with power of attorney on file

75. A patient requests "some leftover antibiotics from last time" for a sore throat. What is the appropriate technician response?

- a) Sell amoxicillin as a cash item
- b) Suggest taking half-doses to stretch leftovers
- c) Explain antibiotics require a valid prescription and offer to help contact a prescriber if requested
- d) Recommend clindamycin because it is stronger

76. A prescription reads: "Apply 1 gram topically TID. Dispense 30 grams." What is the day supply?

- a) 30 days
- b) 10 days
- c) 7 days
- d) 14 days

77. A technician enters "tamsulosin 0.4 mg daily" but selects "tamoxifen 20 mg daily." What risk factor most contributes to this error?

- a) Similar drug indications
- b) Same controlled schedule
- c) Same manufacturer lot number
- d) Look-alike/sound-alike names

78. A patient profile shows "sulfonamide antibiotics—anaphylaxis." A new prescription arrives for hydrochlorothiazide. What should the technician do?

- a) Override the alert because it's not an antibiotic
- b) Route for pharmacist assessment of allergy relevance and risk
- c) Delete the allergy because it is too broad
- d) Automatically change to furosemide

79. A technician is asked to process an early refill request for a Schedule II medication because the patient is traveling. What is the best technician action?

- a) Override and fill early if the patient provides travel dates
- b) Refuse because early fills are always illegal
- c) Collect details and route to the pharmacist for review
- d) Suggest the patient take less each day until the trip

80. A prescription is written for "gabapentin 300 mg: take 1 cap TID. #90 with 2 refills." The patient asks to transfer it to another pharmacy. Under federal rules (assuming state law allows), which is correct?

- a) Transfer is permitted for noncontrolled prescriptions, including remaining refills
- b) Transfer is never permitted once filled
- c) Transfer is permitted only if no refills remain
- d) Transfer requires a new prescription

81. A patient asks what "OS" means on an eye drop label. What does OS indicate?

- a) Right eye
- b) Both eyes
- c) Left eye
- d) Both ears

82. A technician sees a REMS requirement pop-up for dispensing clozapine. What does REMS stand for?

- a) Risk Evaluation and Mitigation Strategy
- b) Required Electronic Medication System
- c) Regulated Emergency Medication Standard
- d) Retail Electronic Monitoring Service

83. A prescription reads: "clopidogrel 75 mg daily." Which brand name matches clopidogrel?

- a) Plavix
- b) Coumadin
- c) Crestor
- d) Zocor

84. A patient is picking up a new prescription for metronidazole. Which counseling point is most likely associated with metronidazole?

- a) Avoid alcohol during therapy
- b) Take with grapefruit juice for best absorption
- c) Avoid dairy products for 48 hours
- d) Double the dose if you miss one

85. A paper prescription for a Schedule II drug is presented with alterations to the quantity written in ink that appears different from the rest of the prescription. What should the technician do?

- a) Process it as written because it is signed
- b) Ask the patient what quantity they want
- c) Set aside for pharmacist review before any dispensing steps
- d) Fill a partial supply to be safe

86. A prescription reads: "atorvastatin 20 mg daily." Which drug class is atorvastatin?

- a) ACE inhibitor
- b) Proton pump inhibitor
- c) Beta blocker
- d) HMG-CoA reductase inhibitor (statin)

87. A patient's new prescription is for "glipizide 5 mg daily." The patient asks if it is for blood pressure or blood sugar. Which is correct?

- a) Blood pressure
- b) Blood sugar
- c) Acid reflux
- d) Infection

88. A prescription reads: "lisinopril 10 mg daily." Which common indication matches lisinopril?

- a) Hypertension
- b) Bacterial infection
- c) Hypothyroidism
- d) Osteoporosis

89. A patient is prescribed "montelukast 10 mg daily." Which brand name matches montelukast?

- a) Flonase
- b) Singulair
- c) Symbicort
- d) Spiriva

90. A new prescription is for "metformin." Which condition is metformin primarily used to treat?

- a) Type 2 diabetes
- b) Hyperlipidemia
- c) GERD
- d) Acute pain

Answer Key

1) b	32) d	63) d
2) c	33) b	64) d
3) b	34) c	65) c
4) c	35) a	66) d
5) a	36) d	67) b
6) b	37) b	68) c
7) a	38) a	69) b
8) b	39) d	70) c
9) b	40) b	71) d
10) d	41) d	72) b
11) c	42) d	73) c
12) c	43) b	74) d
13) d	44) b	75) c
14) c	45) b	76) b
15) a	46) d	77) d
16) b	47) c	78) b
17) c	48) a	79) c
18) d	49) b	80) a
19) c	50) d	81) c
20) b	51) b	82) a
21) a	52) c	83) a
22) c	53) d	84) a
23) c	54) c	85) c
24) b	55) a	86) d
25) c	56) b	87) b
26) c	57) c	88) a
27) d	58) d	89) b
28) b	59) c	90) a
29) c	60) a	
30) d	61) c	
31) b	62) d	

PTCB Sim Test 13

1) A new e-Rx arrives: "Synthroid 50 mcg take 1 tablet PO daily #30." The patient profile shows an active "levothyroxine 50 mcg take 1 tablet PO daily" filled 10 days ago from the same prescriber. What should the technician do next?

- a) Process the new Rx as a duplicate and inactivate the older one
- b) Change the product to generic automatically to avoid confusion
- c) Enter the prescription and route it for pharmacist review due to potential duplicate therapy/brand-vs-generic concerns
- d) Reject the e-Rx because a refill is too soon

2) A prescriber phones in: "metoprolol succinate ER 25 mg, take 1 daily, #30, 2 refills." The callback number matches the clinic directory, but the prescriber's last name is muffled and unclear. What is the best technician action before entry is completed?

- a) Ask the caller to repeat and spell the prescriber name, then document it accurately
- b) Enter it under the clinic medical director and proceed
- c) Enter it and leave the prescriber field blank for the pharmacist to fix later
- d) Substitute metoprolol tartrate because it is more common

3) A patient requests "Claritin-D 24 hour" at the counter. The log shows they purchased 2.88 g of pseudoephedrine earlier today at another pharmacy. What should the technician do?

- a) Complete the sale because today's purchase was at a different pharmacy
- b) Refuse the sale because the daily limit has already been reached
- c) Sell it but reduce the quantity so it stays under 3.6 g
- d) Sell it if the patient pays cash and shows ID

4) During production, the label prints "hydroxyzine 25 mg take 1 at bedtime." The technician accidentally counted hydralazine 25 mg. Barcode scanning is available. What should happen next?

- a) Continue because both are 25 mg tablets
- b) Place a sticker noting "verify drug name" and send to will-call
- c) Rescan and correct to the product that matches the label before it goes to verification
- d) Change the directions to match hydralazine therapy

5) A prescription is: "amoxicillin/clavulanate 400 mg-57 mg/5 mL. Take 10 mL PO BID x 7 days." How many mL should be dispensed?

- a) 100 mL
- b) 120 mL
- c) 140 mL
- d) 160 mL

6) A parent asks which OTC product is best for a 2-year-old with fever while picking up a prescription. What is the technician's best response within scope?

- a) Recommend aspirin because it works fastest
- b) Recommend ibuprofen only, never acetaminophen
- c) Ask the child's weight and calculate a dose to recommend
- d) Offer to get the pharmacist to help choose an appropriate OTC fever reducer

7) An e-Rx for isotretinoin is in the queue. The iPLEDGE status shows "requirements not met." What is the technician's next step?

- a) Dispense as long as the prescriber is certified
- b) Place on hold and proceed with other work until the patient arrives
- c) Route for pharmacist review and do not dispense until the system requirements are satisfied
- d) Convert to cash to bypass the system rejection

8) A patient asks to transfer their alprazolam (Schedule IV) to another pharmacy. The prescription has refills remaining. Under federal rules, which statement is correct?

- a) It may be transferred one time if permitted by state law
- b) It may be transferred unlimited times until refills run out
- c) It cannot be transferred because it is a controlled substance
- d) It may be transferred only after all refills are used

9) A refill request is received for lisinopril. The original Rx was written 16 months ago with refills, and the system shows it is expired. What should the technician do?

- a) Refill it because refills remain
- b) Change the written date in the system to within 12 months
- c) Request a new prescription authorization from the prescriber
- d) Transfer it to another pharmacy to avoid the expiration rule

10) A label direction reads: "Instill ii gtt OS qid." What does this mean?

- a) Instill 2 drops in the right eye four times daily
- b) Instill 2 drops in the left eye four times daily
- c) Instill 1 drop in both eyes twice daily
- d) Instill 4 drops in the left eye once daily

11) A patient's spouse arrives to pick up and asks, "What is this medicine for?" The spouse is not listed as an authorized pickup/contact. What is the best technician action?

- a) Explain the indication briefly because the spouse is picking up
- b) Provide the full counseling since the patient is not present
- c) Provide only the drug name and directions
- d) Attempt to contact the patient for permission before disclosing medication information

12) A prescription is: "prednisone 20 mg: take 3 tablets daily x 2 days, then 2 tablets daily x 2 days, then 1 tablet daily x 2 days." How many tablets should be dispensed?

- a) 18
- b) 24
- c) 12
- d) 30

13) A prescription is for "metformin ER 500 mg: take 2 tablets once daily #60." What days' supply should be billed?

- a) 15
- b) 60
- c) 30

d) 45

14) A patient's profile has a documented allergy: "sulfonamides—anaphylaxis." A new e-Rx arrives for sulfamethoxazole/trimethoprim DS. What should the technician do?

- a) Enter and dispense because the prescriber ordered it
- b) Put it on hold and route it for pharmacist review of the allergy alert
- c) Change the medication to nitrofurantoin automatically
- d) Delete the allergy because it is likely outdated

15) A bottle of refrigerated insulin is found on the counter at opening. It was left out overnight. What is the best technician action?

- a) Put it back in the refrigerator and mark "use first"
- b) Return it to stock if it is still within expiration
- c) Quarantine it for pharmacist assessment of stability before any use or return-to-stock
- d) Dispense it but shorten the expiration date on the label

16) A new prescription arrives for "Toprol-XL 50 mg daily." Which generic is appropriate to select?

- a) Metoprolol tartrate
- b) Atenolol
- c) Propranolol ER
- d) Metoprolol succinate ER

17) A pharmacy receives a manufacturer recall notice for a specific lot of amlodipine. What should the technician do first?

- a) Pull and quarantine affected lot inventory from shelves and will-call
- b) Call every patient who has ever filled amlodipine
- c) Continue dispensing until replacement arrives
- d) Return all amlodipine regardless of lot number

18) A controlled-substance prescription is presented on paper for a Schedule II stimulant. Which element is required for validity under federal law?

- a) Prescriber DEA number printed on the prescription
- b) Patient's social security number
- c) Prescriber signature
- d) Diagnosis code

19) A prescription reads: "cephalexin 250 mg/5 mL: take 5 mL PO QID x 10 days." How many mL should be dispensed?

- a) 200 mL
- b) 100 mL
- c) 150 mL
- d) 250 mL

20) A patient wants an early refill on oxycodone/APAP stating, "I'm leaving town tomorrow." The claim rejects "refill too soon." What is the best technician action?

- a) Change the days' supply to force approval
- b) Process as a partial fill without documenting

- c) Tell the patient it is illegal to refill early
- d) Gather details and route to the pharmacist to evaluate options such as contacting the prescriber or insurance vacation override

21) A hospital cart-fill list calls for "morphine IR 15 mg." The technician scans "morphine ER 15 mg" and gets a mismatch warning. What should the technician do?

- a) Override the warning because the strength matches
- b) Select the immediate-release product and rescan
- c) Send it anyway and note "ER substituted"
- d) Replace with hydromorphone of an equivalent dose

22) A technician is asked which medication is a proton pump inhibitor (PPI).

- a) Famotidine
- b) Omeprazole
- c) Sucralfate
- d) Dicyclomine

23) A prescription is: "diclofenac 1% gel: apply 4 g to knee QID." If dispensed as a 100 g tube, about how many days will it last?

- a) 6 days
- b) 10 days
- c) 25 days
- d) 3 days

24) A new e-Rx is "Eliquis 5 mg take 1 tablet BID." The profile shows "Xarelto 20 mg daily" filled last week with remaining refills. The prescriber note says "switch anticoagulant." What should the technician do?

- a) Inactivate Xarelto and fill Eliquis immediately without review
- b) Fill both so the patient has options
- c) Route to pharmacist to confirm the switch and avoid duplication
- d) Delete the prescriber note because it is not part of the prescription

25) A prescription direction says: "Take 1 tab PO qod." How should that be interpreted?

- a) Take 1 tablet every other day
- b) Take 1 tablet once daily
- c) Take 1 tablet four times daily
- d) Take 1 tablet as needed for pain

26) A patient's vaccination record must be reported to the state immunization information system (IIS). Which data element is most critical to ensure correct matching?

- a) Patient's favorite pharmacy location
- b) Patient's SSN
- c) Patient's full name and date of birth
- d) Insurance BIN and PCN

27) A faxed prescription for clonazepam is received, but the strength is illegible. What should the technician do?

- a) Fill the lowest strength available to be safe

- b) Request clarification from the prescriber's office before processing
- c) Ask the patient what strength they take and proceed
- d) Dispense a 3-day emergency supply

28) A prescription reads: "fluconazole 150 mg take 1 now and repeat in 72 hours #2." What days' supply should be billed?

- a) 1
- b) 2
- c) 3
- d) 4

29) A prescriber writes for "doxycycline hyclate 100 mg caps." Only doxycycline monohydrate 100 mg caps are on the shelf. What is the best technician action?

- a) Substitute monohydrate automatically because the strength matches
- b) Dispense monohydrate and note "equivalent" on the hard copy
- c) Process a product change only after pharmacist review/approval per state law and pharmacy procedure
- d) Split the fill: half hyclate and half monohydrate

30) A prescription is for "Humalog KwikPen: inject 8 units with meals TID. Dispense 1 box (5 pens, 3 mL each, 100 units/mL)." What is the days' supply?

- a) 93
- b) 62
- c) 75
- d) 50

31) A patient is at drop-off with a prescription from another country for tramadol. What is the correct handling?

- a) Fill it if it has a prescriber signature
- b) Fill a 3-day supply only
- c) Convert it to a U.S. prescription by transcribing it
- d) It is not valid to dispense; the patient needs a prescription from a U.S.-licensed prescriber

32) A prescription for "warfarin 5 mg daily" is received. The profile shows apixaban 5 mg BID filled yesterday. What should the technician do?

- a) Enter and fill warfarin because it was prescribed today
- b) Route to pharmacist due to potential duplicate anticoagulation
- c) Delete apixaban from the profile because warfarin is older therapy
- d) Change warfarin to 2.5 mg daily to reduce risk

33) Which medication is an ACE inhibitor?

- a) Losartan
- b) Amlodipine
- c) Lisinopril
- d) Metoprolol

34) A controlled substance C-II prescription is partially filled at the patient's request due to cost. Under federal law, what is generally required?

- a) The remainder must be dispensed within 72 hours in all cases
- b) The partial fill must be documented, and remaining quantity may be filled within the permitted timeframe depending on the partial-fill situation
- c) Partial fills are never allowed for Schedule II
- d) The remainder can be filled any time within 6 months

35) A technician receives a DUR alert: "Duplicate therapy—NSAIDs: active meloxicam; new ibuprofen Rx." What is the technician's best next step?

- a) Override the alert to keep workflow moving
- b) Route the alert for pharmacist review before dispensing
- c) Tell the patient to stop meloxicam when they pick up
- d) Remove meloxicam from the medication list

36) Which drug is a direct oral anticoagulant (DOAC)?

- a) Clopidogrel
- b) Warfarin
- c) Apixaban
- d) Enoxaparin

37) A prescription reads: "albuterol HFA: inhale 2 puffs q4-6h PRN. 200 actuations." If the pharmacy bills PRN inhalers using a maximum of 12 puffs/day, what days' supply should be submitted?

- a) 25
- b) 16
- c) 17
- d) 30

38) A unit-dose package has the drug name and strength but the lot number is smeared and unreadable. What should the technician do?

- a) Use it because the identity is clear
- b) Write the lot number from a different package of the same drug
- c) Remove from use because lot traceability is needed for recalls
- d) Place it in an amber vial and label it manually

39) A prescription is: "insulin glargine pens 100 units/mL: inject 24 units subcutaneously nightly. Dispense 1 box (5 pens, 3 mL each)." What is the days' supply?

- a) 31
- b) 62
- c) 75
- d) 50

40) A patient asks to return an unopened bottle of hydrocodone/APAP for a refund. What is the correct response?

- a) Accept it and return to stock if sealed
- b) Accept it but send it to the wholesaler for credit
- c) Accept it and redispense to another patient after inspection
- d) Do not take it back for resale; provide disposal options per program availability

41) A prescription comes in with "MSO4 2 mg IV push now." What is the best technician action in a hospital setting?

- a) Enter as morphine sulfate because MSO4 always means morphine
- b) Clarify because MSO4 is a dangerous abbreviation and could be misread
- c) Enter as magnesium sulfate because it's common IV
- d) Substitute hydromorphone to avoid confusion

42) Which medication requires extra safeguards because it is typically dosed weekly and is considered high-alert in outpatient use?

- a) Methotrexate
- b) Metformin
- c) Amoxicillin
- d) Acyclovir

43) A patient profile shows "penicillin—hives." A new prescription arrives for amoxicillin. What should the technician do?

- a) Proceed; hives is not a true allergy
- b) Change to azithromycin automatically
- c) Route for pharmacist review due to allergy risk
- d) Dispense with an allergy warning sticker only

44) A technician is asked to identify the generic for Zoloft.

- a) Sertraline
- b) Citalopram
- c) Fluoxetine
- d) Paroxetine

45) A written prescription for hydromorphone is missing the patient address. What is the best next step?

- a) Have the patient write their address on the prescription
- b) Process it and use the address on the patient profile without clarifying
- c) Fill it and annotate "address missing"
- d) Route for pharmacist review and obtain a corrected prescription/clarification per legal requirements

46) A prescription is: "azithromycin 250 mg: take 2 tablets today, then 1 tablet daily for 4 days." What quantity should be dispensed?

- a) 5
- b) 6
- c) 4
- d) 8

47) A prescriber sends "ondansetron ODT 4 mg dissolve 1 tablet q8h PRN N/V #12." The technician selected ondansetron 4 mg regular tablets. What should be done?

- a) Keep regular tablets because they are the same drug/strength
- b) Change the directions to "swallow" to match selection
- c) Correct the product to the ODT formulation as prescribed
- d) Reduce quantity to #9 because ODT is usually smaller packages

48) A technician sees two open profiles: "Allen, J." and "Allen, J." with different dates of birth. What is the best way to reduce wrong-patient errors before entering the next Rx?

- a) Use the address only since DOB can change
- b) Close one profile and verify at least two identifiers before continuing
- c) Print both leaflets and compare later during pickup
- d) Continue entry and rely on the pharmacist to catch it at verification

49) A prescription is: "levothyroxine 125 mcg take 1 daily #90." The patient asks if they can switch between brands each month based on price. Best technician response?

- a) Yes, thyroid products are always interchangeable without issue
- b) Advise the patient to alternate brands to prevent tolerance
- c) Recommend taking it with calcium to stabilize absorption
- d) Refer to the pharmacist because consistent product/manufacturer may matter for levothyroxine

50) Which controlled substance schedule applies to testosterone cypionate injection federally?

- a) Schedule II
- b) Schedule V
- c) Schedule III
- d) Schedule IV

51) A technician receives a DSCSA notice about a "suspect product" involving a certain NDC/lot of atorvastatin. What should be done first?

- a) Quarantine affected inventory and notify the pharmacist/manager to begin investigation steps
- b) Continue dispensing until the manufacturer confirms the issue
- c) Remove all atorvastatin from shelves regardless of NDC/lot
- d) Repackage it into amber vials to reduce risk

52) A prescription is: "timolol 0.5% ophthalmic solution instill 1 gtt OU BID. Dispense 10 mL." What quantity should be entered for billing if the plan wants mL?

- a) 2
- b) 10
- c) 1
- d) 20

53) A patient is picking up nitroglycerin SL tablets and asks how to store them. Which is correct?

- a) In the original, tightly closed light-resistant container at room temperature
- b) In a weekly pill organizer for convenience
- c) In the refrigerator to maintain potency
- d) In the bathroom medicine cabinet for easy access

54) A technician is asked which medication is a statin.

- a) Atorvastatin
- b) Allopurinol

- c) Albuterol
- d) Atenolol

55) A prescription reads: "Take 2 tsp PO TID." How many mL per dose?

- a) 5 mL
- b) 15 mL
- c) 20 mL
- d) 10 mL

56) A patient calls asking for "a list of everything I've filled this year" to give their new doctor. What should the technician do?

- a) Provide it after verifying identity according to pharmacy procedure
- b) Refuse because profiles are proprietary
- c) Provide it only if the prescriber faxes a request
- d) Provide it but remove drug names to protect privacy

57) A technician is entering: "diltiazem CD 240 mg daily." The system suggests diltiazem IR 120 mg. What should the technician do?

- a) Choose IR and change to twice daily
- b) Enter diltiazem without formulation and dispense what's in stock
- c) Select the extended-release CD product as prescribed
- d) Substitute verapamil ER

58) A prescription is for "clopidogrel 75 mg daily." The patient profile shows a documented clopidogrel allergy: rash. What should the technician do?

- a) Enter and dispense; rash is not serious
- b) Route for pharmacist review due to allergy alert
- c) Remove the allergy from the profile as outdated
- d) Change the medication to aspirin automatically

59) A prescription reads: "amoxicillin 400 mg/5 mL: take 7.5 mL PO BID x 10 days." How many mL should be dispensed?

- a) 120 mL
- b) 200 mL
- c) 150 mL
- d) 90 mL

60) A technician is receiving stock and finds a sealed manufacturer bottle with the outer label torn off; no NDC is visible. What is the best action?

- a) Stock it by identifying tablets via imprint
- b) Quarantine it as suspect/unverifiable and notify the pharmacist/manager for return/verification
- c) Re-label it using the shelf label NDC
- d) Open it and compare to another bottle of the same drug

61) A prescription is: "Bactrim DS 1 tab PO BID x 7 days." The patient states they had "throat swelling" with sulfa in the past. What should the technician do?

- a) Process as normal since it's common for UTIs

- b) Ask the patient to sign a waiver and dispense
- c) Route to pharmacist for allergy assessment before dispensing
- d) Change to cephalexin automatically

62) A technician is asked which medication is an SSRI.

- a) Duloxetine
- b) Sertraline
- c) Venlafaxine
- d) Bupropion

63) A technician is preparing a compounded oral suspension and the formula sheet lists a beyond-use date (BUD) different from the one on a commonly used template. What should be done?

- a) Use the longer BUD so the patient has more time
- b) Omit the BUD since it's compounded
- c) Use the formula sheet BUD and have it checked during pharmacist verification
- d) Use the manufacturer expiration date of the active ingredient

64) A prescription for fluticasone/salmeterol 250/50 reads "inhale 1 puff BID." The inhaler contains 60 inhalations. What days' supply should be billed?

- a) 15
- b) 60
- c) 20
- d) 30

65) Which action best supports a "high-alert medications" safety strategy during data entry?

- a) Turning off clinical alerts to reduce interruptions
- b) Using Tall Man lettering and barcode verification where available
- c) Accepting verbal confirmation from patients instead of scanning
- d) Storing look-alike products next to each other for speed

66) A technician is processing a refill for a Schedule II medication with "0 refills." What is correct?

- a) The technician may refill once if the patient is established
- b) A new prescription is required for a Schedule II refill
- c) The pharmacy can transfer it in and refill it
- d) The prescriber can authorize refills verbally without a new prescription

67) Which medication is a benzodiazepine?

- a) Buspirone
- b) Hydroxyzine
- c) Zolpidem
- d) Lorazepam

68) A transfer-in is being accepted for a Schedule III prescription. Which information must be captured for a compliant transfer record?

- a) Only the drug name and directions
- b) Only the prescriber name and patient phone number

- c) Original Rx number, date written, refills authorized/remaining, and other required identifiers from the transferring pharmacy
- d) Only the last fill date and quantity

69) A prescription direction includes "U-500 insulin." Why is this high risk in dispensing?

- a) It is a Schedule II medication
- b) It is a concentrated insulin where dosing errors can be significant compared to U-100
- c) It is only for IV administration
- d) It must be stored frozen

70) A prescription reads: "Apply sparingly to AA BID." What does "AA" mean?

- a) Affected area
- b) Both arms
- c) After activity
- d) Before meals

71) A pharmacy has two products stocked: Augmentin 875/125 and Augmentin XR 1000/62.5. The Rx is for Augmentin 875/125. What should the technician do?

- a) Use XR because it is extended-release and stronger
- b) Use either product because both are amoxicillin/clavulanate
- c) Change the sig to once daily and use XR
- d) Select the exact formulation/strength prescribed

72) A patient asks the technician to recommend an OTC heartburn product while taking clopidogrel. What should the technician do?

- a) Recommend omeprazole because it is most effective
- b) Recommend bismuth subsalicylate for daily use
- c) Refer to the pharmacist due to potential drug interactions with certain OTC options
- d) Recommend aspirin for chest discomfort

73) A technician is asked which medication is used for gout prophylaxis.

- a) Allopurinol
- b) Colchicine
- c) Prednisone
- d) Indomethacin

74) A prescription is: "vancomycin 125 mg PO QID x 10 days." The only stocked product is vancomycin injection vials used for compounding oral solutions under specific procedures. What is the best technician action?

- a) Select the injection vials and proceed to compound without further steps
- b) Change the route to IV since vials are injectable
- c) Route to pharmacist because formulation selection/compounding requirements must be verified
- d) Substitute metronidazole automatically

75) A patient's insurance requires days' supply for an albuterol inhaler written "2 puffs q4-6h PRN." The pharmacy uses 12 puffs/day for PRN. If the inhaler has 200 puffs, what is the days' supply?

- a) 14
- b) 16
- c) 25
- d) 20

76) Which abbreviation is generally discouraged on outpatient labels due to error risk?

- a) mL
- b) qid
- c) PO
- d) QD

77) A patient's profile shows "torsemide 10 mg daily" active. A new e-Rx comes in: "Lasix 40 mg daily." What should the technician do?

- a) Substitute torsemide because it's the same class
- b) Process Lasix and refill torsemide too
- c) Route for pharmacist review for possible therapeutic duplication/change
- d) Delete torsemide from the profile automatically

78) A technician is asked which medication is an inhaled short-acting beta-agonist (SABA).

- a) Fluticasone
- b) Albuterol
- c) Tiotropium
- d) Budesonide/formoterol

79) A prescription label prints "Take 1 tablet by mouth QD." The patient asks what it means. What should the technician say?

- a) "It means once daily."
- b) "It means four times daily."
- c) "It means every other day."
- d) "It means once weekly."

80) A prescription is for "benzonatate 100 mg TID PRN cough." A caregiver asks if it can be given to a 4-year-old. What should the technician do?

- a) Suggest opening the capsule and giving half
- b) Say it is safe if the child can swallow
- c) Recommend OTC dextromethorphan instead and skip further discussion
- d) Refer to the pharmacist for age-appropriateness/safety counseling

81) A prescription is written: "ibuprofen 200 mg #120. Sig: 1 tab q6h PRN." Using a maximum of 4 doses/day for billing, what is the days' supply?

- a) 20
- b) 24
- c) 30
- d) 10

82) A technician is asked to print a perpetual inventory report for Schedule II drugs. The primary purpose is to:

- a) Identify which patients qualify for automatic refills

- b) Detect diversion and reconcile controlled-substance counts
- c) Generate purchase orders for wholesalers
- d) Document HIPAA disclosures

83) A patient is prescribed "gabapentin 300 mg take 1 capsule TID #90." What is the days' supply?

- a) 45
- b) 90
- c) 15
- d) 30

84) A patient calls asking whether their prescription is ready. They provide name and date of birth but refuse any additional identifiers. What is the best technician response?

- a) Provide the medication name and strength but not directions
- b) Provide full details because DOB is enough
- c) Refuse to say anything at all over the phone
- d) Provide only a general status (e.g., "ready/not ready") without medication details

85) Which medication is a calcium channel blocker (CCB)?

- a) Amlodipine
- b) Lisinopril
- c) Hydrochlorothiazide
- d) Losartan

86) A technician is asked how to properly dispose of outdated Schedule II stock at a retail pharmacy. Which is appropriate?

- a) Flush it down the toilet if it is a controlled substance
- b) Place it in regular trash after removing labels
- c) Use a DEA-compliant method with required records and authorized personnel/witnessing
- d) Give it to a reverse distributor without any documentation

87) A prescription reads: "clotrimazole 1% vaginal cream: insert 1 applicatorful PV qhs x 7 days." Which is the best matching interpretation of PV?

- a) By mouth
- b) In the vagina
- c) In the left eye
- d) Under the tongue

88) A prescription is: "lansoprazole 30 mg take 1 capsule daily." Which brand matches lansoprazole?

- a) Prilosec
- b) Nexium
- c) Prevacid
- d) Protonix

89) While filling, the scanned NDC matches the drug name but the strength on the stock bottle is 10 mg and the label is for 20 mg. What should the technician do?

- a) Continue because the scan matched the drug name

- b) Change the label to 10 mg to match what is available
- c) Stop and reselect the correct strength, then rescan before proceeding
- d) Dispense two 10 mg tablets per dose without changing the prescription

90) A prescription is: "fluoxetine 20 mg: take 1 capsule daily. #30." The patient profile shows paroxetine 20 mg daily filled 5 days ago. What should the technician do?

- a) Fill fluoxetine and remove paroxetine automatically
- b) Route to pharmacist due to potential duplicate SSRI therapy or cross-taper considerations
- c) Fill both and tell the patient to decide which one to take
- d) Change fluoxetine to sertraline because it is commonly used

Answer Key

1) c	31) d	61) c
2) a	32) b	62) b
3) b	33) c	63) c
4) c	34) b	64) d
5) c	35) b	65) b
6) d	36) c	66) b
7) c	37) a	67) d
8) a	38) c	68) c
9) c	39) d	69) b
10) b	40) d	70) a
11) d	41) b	71) d
12) c	42) a	72) c
13) c	43) c	73) a
14) b	44) a	74) c
15) c	45) d	75) b
16) d	46) b	76) d
17) a	47) c	77) c
18) c	48) b	78) b
19) b	49) d	79) a
20) d	50) c	80) d
21) b	51) a	81) c
22) b	52) b	82) b
23) a	53) a	83) d
24) c	54) a	84) d
25) a	55) d	85) a
26) c	56) a	86) c
27) b	57) c	87) b
28) d	58) b	88) c
29) c	59) c	89) c
30) b	60) b	90) b

PTCB Sim Test 14

1) A prescriber e-sends: "Lopressor 50 mg. Take 1 tablet PO BID." Which product should be selected?

- a) Metoprolol succinate ER 50 mg tablet
- b) Metoprolol tartrate 50 mg tablet
- c) Propranolol 40 mg tablet
- d) Atenolol 50 mg tablet

2) A patient's allergy field shows "lisinopril—angioedema." A new eRx arrives for enalapril 10 mg daily. What should the technician do at intake?

- a) Enter and fill because enalapril is a different ACE inhibitor
- b) Add a note and continue processing without delay
- c) Ensure the pharmacist reviews the ACE inhibitor allergy risk before the prescription is filled
- d) Remove the allergy because it is already documented

3) A prescription reads: "Amoxicillin 250 mg/5 mL. Give 10 mL PO TID x 7 days." What total volume should be dispensed?

- a) 140 mL
- b) 210 mL
- c) 200 mL
- d) 180 mL

4) A caregiver asks what "qAM" means on the label. What is the best technician response?

- a) Every night at bedtime
- b) Every afternoon
- c) Every 4 hours as needed
- d) Every morning

5) A new eRx arrives for "Cymbalta 60 mg daily." Which generic should be selected?

- a) Venlafaxine
- b) Bupropion
- c) Duloxetine
- d) Fluoxetine

6) A technician receives a faxed request to transfer refills for alprazolam (Schedule IV). Under federal rules (assuming state law permits), what is true?

- a) It may be transferred one time for the purpose of refill dispensing
- b) It may be transferred unlimited times if pharmacies share a database
- c) It can never be transferred
- d) It may be transferred only after all refills are used

7) A new prescription is written for "warfarin 5 mg daily." Which brand name corresponds to warfarin?

- a) Eliquis
- b) Coumadin

- c) Xarelto
- d) Pradaxa

8) A label prints "Take 1 tablet QID" but the original written directions clearly state "Take 1 tablet QD." What should the technician do before the prescription moves forward?

- a) Cross out the label and initial it
- b) Leave it; the pharmacist will catch it at final check
- c) Reprint the label only and keep the incorrect one in the hardcopy bin
- d) Correct the entry to match the prescription and regenerate labels/documentation per workflow

9) A patient requests to buy pseudoephedrine but left their ID at home. What is the correct action?

- a) Sell it if the patient provides date of birth verbally
- b) Sell a smaller quantity to remain under the daily limit
- c) Decline the sale until acceptable identification is presented
- d) Sell phenylephrine and record it in the logbook instead

10) A prescription reads: "Apply 1 patch every 72 hours. Dispense #8." What is the days' supply?

- a) 12 days
- b) 21 days
- c) 24 days
- d) 8 days

11) A patient says, "My spouse is picking up my meds—can you tell them what's ready?" The spouse is not listed in the patient's communications preferences. What should the technician do?

- a) Provide medication names but not prices
- b) Provide only the total price and nothing else
- c) Provide everything if the spouse knows the patient's address
- d) Avoid disclosing protected information and request the patient add authorization or contact the pharmacy directly

12) An eRx arrives for "metformin ER 500 mg: take 2 tablets daily." The system suggests multiple ER products. Which is the most appropriate product selection?

- a) Metformin immediate-release 500 mg tablet
- b) Metformin ER 500 mg tablet
- c) Glyburide 5 mg tablet
- d) Metformin ER 750 mg tablet

13) A controlled substance shipment arrives with a torn seal and the invoice quantity does not match what's in the tote. What should happen first?

- a) Stock what arrived, then email the wholesaler later
- b) Quarantine the tote and alert the pharmacist/manager immediately
- c) Sign the invoice "received in full" to avoid delays
- d) Return the tote with routine returns at end of day

14) A prescription reads: "Prednisone 20 mg: take 2 tablets daily x 5 days." What quantity should be dispensed?

- a) 20 tablets
- b) 5 tablets
- c) 10 tablets
- d) 15 tablets

15) A patient's profile lists "heparin—HIT." A hospital discharge prescription for enoxaparin is dropped off. What should the technician do?

- a) Process because enoxaparin is not heparin
- b) Ensure the pharmacist reviews the HIT history before dispensing
- c) Tell the patient to take aspirin instead
- d) Remove the heparin allergy because it is not a true allergy

16) A patient is picking up two bags. The names are identical, but one profile is "Maria L. Chen" and the other is "Maria L. Chen (02/14/1978)." What should the technician do before handing out any medication?

- a) Ask which doctor they saw and choose based on that
- b) Ask for at least two identifiers (e.g., DOB and address) and match to the bag label/profile
- c) Open the bag and ask the patient to identify tablets by color
- d) Give the most recently filled bag first and see if the patient objects

17) A refill is requested for a Schedule III prescription originally written 7 months ago with refills remaining. Under federal law, what is correct?

- a) It may be refilled only within 6 months of the date written
- b) It may be refilled for up to 1 year if refills remain
- c) It may be refilled indefinitely with refills remaining
- d) It may be refilled only if the patient pays cash

18) A new eRx: "Levaquin 500 mg daily x 7 days." Which generic should be selected?

- a) Levetiracetam
- b) Linezolid
- c) Levofloxacin
- d) Levothyroxine

19) The pharmacy receives a manufacturer recall notice for a specific lot of amlodipine 10 mg. The lot is found on the shelf. What should the technician do?

- a) Remove it from active stock and place it in quarantine according to recall procedure
- b) Use it first to reduce waste
- c) Put a sticker on it that says "check with pharmacist" and continue dispensing
- d) Return it to the shelf after counting remaining tablets

20) A technician is entering: "Instill 1 gtt OU BID." What does OU mean?

- a) Both eyes
- b) Right eye
- c) Left eye
- d) Both ears

21) A prescription reads: "Humalog U-100 vial. Inject 12 units SQ TID with meals." The vial is 10 mL. What is the days' supply for one vial?

- a) 21 days
- b) 27 days
- c) 18 days
- d) 30 days

22) A prescriber phones in a verbal order for hydromorphone (Schedule II). The technician answers. What is the correct next step?

- a) Take the verbal order and have the pharmacist sign later
- b) Transfer the call to the pharmacist to receive the verbal Schedule II order (if allowed under the circumstance)
- c) Refuse and request a fax because controlled substances cannot be phoned in
- d) Ask the patient to bring the prescriber's written note instead

23) A prescription reads: "Ventolin HFA: inhale 2 puffs q4h PRN wheeze. 1 inhaler (200 actuations)." What is the maximum days' supply that can be billed using the directions as written?

- a) 17 days
- b) 25 days
- c) 12 days
- d) 8 days

24) A technician finds two different-looking tablets mixed in a bottle labeled "clopidogrel 75 mg." What should the technician do?

- a) Continue filling from that bottle if the NDC matches the label
- b) Separate tablets by color and use them accordingly
- c) Remove the bottle from use and escalate for investigation before dispensing
- d) Finish the current fill and then replace the bottle

25) A patient asks for "Zocor 20 mg." Which generic matches?

- a) Rosuvastatin
- b) Simvastatin
- c) Pravastatin
- d) Atorvastatin

26) A prescription reads: "Clindamycin 1% topical solution. Apply to affected area BID. Dispense 60 mL." If the patient uses 1 mL per application, what is the days' supply?

- a) 30 days
- b) 15 days
- c) 60 days
- d) 20 days

27) A patient requests a printed medication list to bring to an appointment. What can the technician do?

- a) Provide it only if the provider faxes a request first
- b) Print and provide it after verifying the patient's identity according to HIPAA/privacy requirements

- c) Refuse because medication lists are confidential
- d) Provide it only if the patient signs it at the register without verification

28) A hospital order: "Heparin 25,000 units in 250 mL D5W. Infuse at 1,250 units/hr." What rate should be entered in mL/hr?

- a) 10 mL/hr
- b) 12.5 mL/hr
- c) 15 mL/hr
- d) 8 mL/hr

29) A new eRx arrives for "Flomax 0.4 mg daily." Which generic should be selected?

- a) Finasteride
- b) Tadalafil
- c) Tamsulosin
- d) Terazosin

30) A patient brings in a photocopy of a Schedule II prescription with no original signature. What is the appropriate action?

- a) Process it if the prescriber's phone number is on it
- b) Decline to treat it as a valid original prescription and escalate per workflow
- c) Fill a partial supply and request the original later
- d) Accept it if the patient shows government ID

31) A prescription reads: "Azithromycin 200 mg/5 mL. Give 12.5 mL on day 1, then 6.25 mL daily on days 2–5." What total volume is needed?

- a) 37.5 mL
- b) 35 mL
- c) 40 mL
- d) 45 mL

32) During pickup, a patient asks: "What does this new carvedilol do, and what side effects should I expect?" What is within technician scope?

- a) Provide detailed counseling and recommend dose changes if dizzy
- b) Provide the medication guide and offer to have the pharmacist counsel the patient
- c) Advise taking it only when blood pressure is high
- d) Recommend stopping it if fatigue occurs

33) A new eRx: "Plavix 75 mg daily." Which generic should be selected?

- a) Clopidogrel
- b) Prasugrel
- c) Ticagrelor
- d) Dipyridamole/aspirin

34) An eRx for oxycodone IR (Schedule II) contains drug, strength, quantity, and prescriber info, but no directions. What should happen before it can be processed?

- a) Enter "Use as directed" and continue
- b) Enter typical directions based on quantity
- c) Obtain complete directions through the appropriate pharmacist workflow before

processing

d) Ask the patient what directions were given and enter that

35) A prescription reads: "Synthroid 50 mcg daily. DAW 1." What should the technician do?

a) Select generic levothyroxine and mark DAW 0

b) Dispense brand Synthroid and submit using the appropriate DAW code

c) Change to generic and write "brand medically necessary" in notes

d) Dispense whichever is in stock and adjust later

36) A patient reports "sulfa allergy—rash." A new prescription is received for TMP-SMX DS. What is the best technician step during intake?

a) Process normally because rash is not serious

b) Change the medication to doxycycline without contacting anyone

c) Delete the allergy because it is nonspecific

d) Ensure pharmacist review occurs before dispensing due to the reported sulfonamide allergy

37) A prescription reads: "Tessalon Perles 100 mg: take 1 capsule TID PRN cough." Which generic should be selected?

a) Dextromethorphan

b) Benzonatate

c) Guaifenesin

d) Codeine

38) A pharmacy receives a DSCSA notice about a "suspect product" for a specific NDC of injectable epinephrine. What should the technician do first?

a) Dispense it only for emergencies

b) Quarantine the product and escalate for investigation steps per DSCSA workflow

c) Relabel it with a warning and keep using it

d) Send it back in the next routine return without documentation

39) A new eRx arrives for "Xarelto 20 mg daily." Which generic should be selected?

a) Dabigatran

b) Warfarin

c) Rivaroxaban

d) Apixaban

40) A patient's insurance rejects "refill too soon." The patient states the medication was lost. What is the best technician action?

a) Change the days' supply to 1 and rebill

b) Collect details and initiate the appropriate override/exception process per plan requirements

c) Immediately run it as cash without discussing options

d) Tell the patient early refills are illegal

41) A prescription reads: "Apply 2 g topically TID. Dispense 60 g." What is the days' supply?

a) 7 days

b) 10 days

- c) 12 days
- d) 14 days

42) A technician receives a phone call from a prescriber's office asking whether a patient picked up buprenorphine/naloxone. The caller provides only the patient name. What should the technician do?

- a) Provide pickup status because it's the prescriber's office
- b) Verify caller identity and confirm appropriate patient identifiers/permissions before disclosing allowed information
- c) Refuse all communication with prescribers about pickups
- d) Provide only the prescription number and nothing else

43) A prescription reads: "Keppra 750 mg: take 1 tablet BID." Which generic should be selected?

- a) Lacosamide
- b) Lamotrigine
- c) Levetiracetam
- d) Levofloxacin

44) A technician is reconstituting an antibiotic. The pharmacy shelf label says "Refrigerate after mixing," but the manufacturer label for the NDC in hand states "Store at room temperature after reconstitution." Which should be followed?

- a) The shelf label because it's pharmacy standard
- b) Either is fine if the patient shakes well
- c) Refrigerate for 24 hours then room temperature
- d) The manufacturer labeling for the specific product being dispensed

45) A prescription reads: "Nitrofurantoin monohydrate/macrocrystals 100 mg BID x 5 days." How many capsules should be dispensed?

- a) 14
- b) 5
- c) 10
- d) 20

46) A prescription reads: "Ciprofloxacin 0.3% eye drops: 2 gtt q2h while awake x 2 days, then 2 gtt q4h while awake x 5 days (affected eye)." Assume awake 16 hours/day and 20 gtt = 1 mL. Minimum mL needed for one eye?

- a) 4.8 mL
- b) 3.2 mL
- c) 6.4 mL
- d) 5.0 mL

47) A technician is entering an order and accidentally selects "hydroxyzine" instead of "hydralazine." Which control most directly reduces this risk at the fill station?

- a) Counting by hand instead of using a tray
- b) Storing look-alike medications next to each other for comparison
- c) Using barcode/NDC scanning to match the stock bottle to the label/order
- d) Turning off alerts to reduce screen clutter

48) A patient asks for "Norvasc." Which generic should be selected?

- a) Amlodipine
- b) Losartan
- c) Lisinopril
- d) Diltiazem

49) A Schedule II eRx is received with an invalid or missing prescriber address in the prescriber section. What is the best technician action?

- a) Enter it and print the label; address is optional for eRx
- b) Substitute the clinic's main address from memory and continue
- c) Ask the patient for the prescriber address and use that
- d) Route through the proper pharmacist workflow to obtain/confirm required prescriber information before dispensing

50) A technician notices a patient profile printout left on a chair in the waiting area. What should the technician do first?

- a) Immediately remove it from public view and secure it appropriately
- b) Leave it in case the patient returns
- c) Put it in will-call so the patient can retrieve it
- d) Read it to identify which patient it belongs to and then file it

51) A prescription reads: "Dopamine 400 mg in 250 mL D5W. Infuse 10 mcg/kg/min. Patient weight 70 kg." What infusion rate should be entered in mL/hr?

- a) 26.3 mL/hr
- b) 31.5 mL/hr
- c) 19.7 mL/hr
- d) 22.5 mL/hr

52) A new prescription is written for "Adderall XR 20 mg daily." Which schedule is correct under the federal Controlled Substances Act?

- a) Schedule I
- b) Schedule II
- c) Schedule III
- d) Schedule IV

53) A prescription reads: "lactulose 10 g/15 mL. Take 30 mL PO BID." Bottle size is 946 mL. What is the days' supply for one bottle?

- a) 12 days
- b) 18 days
- c) 15 days
- d) 21 days

54) A prescription is received for "Augmentin 875/125 mg: take 1 tablet BID x 10 days. Qty #14." What should the technician do?

- a) Dispense #14 as written and change days' supply to 7 days
- b) Change quantity to #20 and proceed
- c) Enter as written and let the patient run out early

d) Route for resolution of the quantity/directions mismatch before dispensing

55) A patient is prescribed "Eliquis 5 mg: take 1 tablet BID." Which generic should be selected?

- a) Apixaban
- b) Rivaroxaban
- c) Dabigatran
- d) Edoxaban

56) During sterile compounding, the technician sees the hood certification sticker expired last month. What is the best action?

- a) Continue compounding and document it
- b) Use the hood only for antibiotics
- c) Stop and escalate; do not compound in an uncertified hood
- d) Wipe the hood and proceed since cleaning replaces certification

57) A technician is asked what "qod" means in directions. Which is correct?

- a) Every other day
- b) Four times daily
- c) Every morning
- d) As needed

58) A prescription reads: "Gentamicin 80 mg IM once." Vials are 40 mg/mL. How many mL should be drawn up?

- a) 1 mL
- b) 2 mL
- c) 0.5 mL
- d) 4 mL

59) A patient asks for "Lasix 20 mg." Which generic should be selected?

- a) Furosemide
- b) Hydrochlorothiazide
- c) Spironolactone
- d) Bumetanide

60) A patient offers to return an unopened prescription bottle dispensed yesterday for a refund and restocking. What is generally correct in most community pharmacies?

- a) Restock if the seal is intact
- b) Restock if it's not a controlled substance
- c) Do not return dispensed prescription medications to stock; handle returns/disposal per procedure
- d) Restock after removing the label and wiping the bottle

61) A new eRx: "Crestor 10 mg daily." Which generic should be selected?

- a) Simvastatin
- b) Rosuvastatin
- c) Atorvastatin
- d) Pravastatin

62) A prescription reads: "Methotrexate 2.5 mg: take 6 tablets daily" for rheumatoid arthritis. What should the technician do?

- a) Enter as written because the prescriber wrote it
- b) Change "daily" to "weekly" and proceed
- c) Ensure pharmacist review occurs due to high-risk weekly vs daily dosing concern
- d) Reduce quantity to 6 tablets and dispense

63) A new eRx arrives for "Zofran ODT 4 mg: take 1 tablet q8h PRN nausea." Which generic should be selected?

- a) Ondansetron
- b) Promethazine
- c) Metoclopramide
- d) Prochlorperazine

64) A patient asks what "SL" means on their nitroglycerin directions. What is the best technician response?

- a) Swallow whole
- b) Place under the tongue
- c) Chew and swallow
- d) Apply to skin

65) A prescription is received for "bupropion XL 150 mg daily." Which is the correct product to select?

- a) Bupropion SR 150 mg tablet
- b) Bupropion XL 150 mg tablet
- c) Buspirone 15 mg tablet
- d) Buprenorphine/naloxone film

66) A technician is asked to place a medication into a compliance package. The tablets are extended-release and the patient wants them crushed into the blister. What is the best technician action?

- a) Crush them if the patient requests it
- b) Split them and crush half only
- c) Keep them intact and refer the question to the pharmacist for appropriateness of crushing ER dosage forms
- d) Substitute with immediate-release automatically

67) A prescription reads: "K-Dur 20 mEq daily." Which product should be selected?

- a) Potassium citrate 20 mEq tablet
- b) Potassium bicarbonate 20 mEq tablet
- c) Potassium chloride ER 20 mEq tablet
- d) Potassium phosphate 20 mEq tablet

68) A prescription is received for "sertraline 50 mg daily." Which brand name corresponds to sertraline?

- a) Zoloft
- b) Prozac

- c) Celexa
- d) Cymbalta

69) A patient presents a Schedule II prescription dated 45 days ago. Under federal law, what is correct regarding whether it can be filled?

- a) It cannot be filled because federal law limits Schedule II to 30 days
- b) It can always be filled within 6 months under federal law
- c) It can be filled only if the prescriber authorizes a new date on the same prescription
- d) Federal law does not set a specific expiration; state law/policy determines validity timeframe

70) During cycle count, the perpetual inventory for oxycodone is short by 3 tablets. What should the technician do?

- a) Adjust the count to match the computer and move on
- b) Recount and review recent transactions, then escalate the discrepancy per controlled-substance procedure
- c) Ignore it until the next delivery
- d) Add 3 tablets from another strength bottle to correct the inventory

71) A patient is switching pharmacies and asks to transfer a Schedule II prescription with refills remaining. What is correct?

- a) Schedule II prescriptions may be transferred once if both pharmacies agree
- b) Schedule II prescriptions may be transferred multiple times
- c) Schedule II prescriptions cannot be transferred for refills because refills are not permitted
- d) Schedule II prescriptions can be transferred only within 72 hours of issue

72) A prescription reads: "Take 1 tsp PO TID." How many mL is 1 tsp?

- a) 2 mL
- b) 10 mL
- c) 15 mL
- d) 5 mL

73) A technician is entering "timolol 0.5% ophthalmic: instill 1 gtt OS BID." What does OS mean?

- a) Right eye
- b) Left eye
- c) Both eyes
- d) Both ears

74) An eRx arrives for "insulin glargine U-100 pen: inject 20 units nightly." Several glargine pens exist in the system. What is most important during selection?

- a) Any U-100 long-acting insulin pen is acceptable
- b) Select the least expensive pen and annotate "OK to substitute"
- c) Match the exact prescribed product/device (specific glargine product and pen form)
- d) Substitute NPH if the patient used it previously

75) A prescription reads: "fluconazole 150 mg x 1 dose." The patient profile shows simvastatin 40 mg nightly. What should happen?

- a) Process routinely; single-dose therapy has no interaction concerns
- b) Hold simvastatin automatically and add counseling directions to the label
- c) Ensure pharmacist review occurs due to potential interaction considerations
- d) Change to nystatin tablets and process

76) A technician is asked which item requires a permanent bound logbook or electronic log under federal law for sales restrictions.

- a) Pseudoephedrine products
- b) Dextromethorphan products
- c) Guaifenesin products
- d) Acetaminophen products

77) A prescription reads: "Docusate sodium 100 mg: take 1 capsule BID." The pharmacy has docusate calcium 240 mg. What is correct?

- a) Substitute because both are docusate
- b) Substitute if the patient agrees
- c) Substitute if the quantity is changed to match mg
- d) Do not substitute; different salt/strength requires appropriate authorization

78) A new eRx arrives for "Lantus SoloStar." Which generic/ingredient is insulin glargine?

- a) Insulin lispro
- b) Insulin glargine
- c) Insulin aspart
- d) Insulin detemir

79) A prescription reads: "Vancomycin solution concentration 50 mg/mL." How many mL contains 750 mg?

- a) 10 mL
- b) 12 mL
- c) 18 mL
- d) 15 mL

80) A technician is processing a prescription and selects pantoprazole because it's "the same class" as omeprazole on the eRx. What is correct?

- a) Acceptable because both are PPIs
- b) Acceptable if the patient has taken pantoprazole before
- c) Not acceptable; the dispensed drug must match the prescription unless substitution is properly authorized
- d) Acceptable if documented in notes

81) A prescription is received for "atorvastatin 40 mg daily." Which brand name corresponds to atorvastatin?

- a) Lipitor
- b) Zocor
- c) Pravachol
- d) Crestor

82) A hospital order reads: "D5W 250 mL with 1 g ceftriaxone." Vials are labeled 1 g per vial.

How many vials are needed to prepare one dose?

- a) 2 vials
- b) 1 vial
- c) 3 vials
- d) 0.5 vial

83) A prescription reads: "Linezolid 600 mg BID" for a patient taking sertraline 100 mg daily. What should the technician do during processing?

- a) Process; this combination is routinely safe
- b) Tell the patient to stop sertraline immediately
- c) Ensure pharmacist review occurs due to potential serious interaction risk
- d) Automatically change linezolid to doxycycline

84) A prescription reads: "Insulin aspart U-100 pen: inject 6 units with meals TID." One box contains five 3 mL pens. What is the days' supply for one box?

- a) 28 days
- b) 42 days
- c) 50 days
- d) 83 days

85) A new eRx arrives for "Neurontin 300 mg TID." Which generic should be selected?

- a) Pregabalin
- b) Gabapentin
- c) Carbamazepine
- d) Lamotrigine

86) A patient requests an early fill of Adderall XR 12 days early due to travel. The claim rejects as "too soon." What should the technician do?

- a) Change the fill date in the system and rerun
- b) Dispense as cash without discussing options
- c) Tell the patient early fills are always illegal
- d) Collect travel/plan details and initiate the appropriate early-fill/vacation override workflow for review

87) A prescription reads: "Morphine ER 30 mg: take 1 tablet q12h. Qty 60. Refills 2." What is correct under federal law?

- a) Refills are allowed if the prescriber authorizes
- b) Refills are not permitted for Schedule II prescriptions
- c) Only one refill is allowed
- d) Refills are allowed for ER products only

88) A prescription reads: "metronidazole 500 mg BID x 7 days. Qty #7." What should the technician do?

- a) Dispense #7 and change directions to once daily
- b) Dispense #7 and keep directions as BID
- c) Route for resolution of quantity/directions mismatch before dispensing
- d) Change quantity to #14 and proceed without review

89) A high-alert medication that commonly requires extra verification steps in many settings is:

- a) Amoxicillin
- b) Levothyroxine
- c) Metformin
- d) Heparin

90) A patient calls and asks to change the mailing address for prescriptions. The profile note states: "Do not change address without identity verification." What should the technician do?

- a) Update the address immediately to meet the request
- b) Ship to the old address and update it later
- c) Verify identity using required methods, document, then update the address before shipment
- d) Cancel mail service permanently for the patient

Answer Key

1) b	31) d	61) b
2) c	32) b	62) c
3) b	33) a	63) a
4) d	34) c	64) b
5) c	35) b	65) b
6) a	36) d	66) c
7) b	37) b	67) c
8) d	38) b	68) a
9) c	39) c	69) d
10) c	40) b	70) b
11) d	41) b	71) c
12) b	42) b	72) d
13) b	43) c	73) b
14) c	44) d	74) c
15) b	45) c	75) c
16) b	46) a	76) a
17) a	47) c	77) d
18) c	48) a	78) b
19) a	49) d	79) d
20) a	50) a	80) c
21) b	51) a	81) a
22) b	52) b	82) b
23) b	53) c	83) c
24) c	54) d	84) d
25) b	55) a	85) b
26) a	56) c	86) d
27) b	57) a	87) b
28) b	58) b	88) c
29) c	59) a	89) d
30) b	60) c	90) c

PTCB Sim Test 15

1) While entering a new eRx, the system suggests two patient profiles with the same name. One profile matches DOB, the other matches address. What is the best next step before continuing?

- a) Select the most recently used profile and proceed
- b) Merge the two profiles immediately
- c) Confirm at least two identifiers with the patient/prescriber information and link to the correct profile before processing
- d) Create a new profile to avoid choosing incorrectly

2) A prescription is written for "Toprol-XL 25 mg daily." During product selection, the screen shows metoprolol tartrate 25 mg as the first match. What should you select?

- a) Metoprolol succinate ER 25 mg
- b) Metoprolol tartrate 25 mg
- c) Carvedilol 3.125 mg
- d) Atenolol 25 mg

3) A prescriber's office calls in a verbal prescription for alprazolam. The pharmacist is administering a vaccine. What should the technician do?

- a) Take the full verbal order and leave it for the pharmacist to sign later
- b) Ask the caller to fax it because controlled substances cannot be called in
- c) Place the caller on hold and wait until the pharmacist is available to take the call
- d) Tell the caller to leave the order on voicemail

4) Rx: "amoxicillin 875 mg: take 1 tablet PO BID x 10 days." Quantity?

- a) 14
- b) 20
- c) 21
- d) 28

5) A patient picking up warfarin asks which OTC pain reliever they should choose for a headache. What is the most appropriate response by a technician?

- a) Recommend ibuprofen because it works best for headaches
- b) Suggest aspirin because it supports anticoagulation
- c) Recommend naproxen because it lasts longer
- d) Refer the patient to the pharmacist for selection advice

6) An eRx arrives for "Crestor 20 mg daily." No DAW is indicated. What should be dispensed if the plan allows substitution?

- a) Crestor only because brand name was used
- b) Simvastatin 20 mg because it is a statin
- c) Rosuvastatin 20 mg
- d) Atorvastatin 20 mg

7) While filling hydralazine 25 mg, you notice hydroxyzine 25 mg is stored adjacent and the bottles look similar. Which step most directly reduces the risk of dispensing the wrong drug?

- a) Count tablets in multiples of 10
- b) Barcode scan the stock bottle and verify it matches the prescription/NDC in the system
- c) Use the bottle with the newest expiration date
- d) Place both bottles on the counter to compare labels

8) Rx: "azithromycin 250 mg: take 2 tablets today, then 1 tablet daily for 4 more days."

Quantity?

- a) 5
- b) 6
- c) 7
- d) 8

9) A patient's profile shows "levothyroxine—same manufacturer only." The previous NDC is out of stock, but another manufacturer is available. What is the best technician action?

- a) Substitute the available manufacturer and document after the fact
- b) Change to the closest strength available and proceed
- c) Check wholesaler/other locations for the same NDC and route for pharmacist decision if substitution is needed
- d) Convert the order to "dispense as written" to force the same manufacturer

10) A patient returns an unopened vial of tablets and asks the pharmacy to restock it for credit. What is the correct response?

- a) Restock it if the seal is intact
- b) Restock it only if it is not a controlled substance
- c) Restock it only if the patient stored it correctly
- d) Explain that dispensed medications generally cannot be returned to stock and proceed with the pharmacy's return/refund/disposal process

11) During entry, you see a new prescription for apixaban 5 mg BID. The patient filled warfarin yesterday and it is still active on the profile. What should you do during processing?

- a) Inactivate warfarin automatically and fill apixaban
- b) Fill both to ensure therapeutic overlap
- c) Continue processing but ensure this is flagged for pharmacist review before dispensing
- d) Delete the apixaban prescription to avoid duplication

12) Insulin lispro U-100 vial 10 mL (100 units/mL). Sig: "Inject 8 units before meals TID."

Days' supply for 1 vial?

- a) 28
- b) 41
- c) 56
- d) 83

13) A patient allergy list shows "sulfonamides—hives." An eRx arrives for sulfamethoxazole/trimethoprim DS. What should be done?

- a) Process normally because hives is not life-threatening
- b) Remove the allergy because it is nonspecific
- c) Change to nitrofurantoin and continue
- d) Ensure the allergy is highlighted and route the order for pharmacist review before

dispensing

14) A patient presents a written prescription for oxycodone 5 mg dated 9 months ago. What is the best handling step?

- a) Process it because it is within one year
- b) Refuse and shred it immediately
- c) Route for pharmacist review due to controlled-substance validity requirements and safety concerns
- d) Call the prescriber and request a verbal authorization to fill

15) A claim rejects "refill too soon." The patient says the prescriber increased the dose yesterday, but no new prescription is on file. What should you do next?

- a) Decrease the days' supply until it pays
- b) Submit the claim repeatedly to force an approval
- c) Gather the updated prescription directions/order information and route for pharmacist review for appropriate resolution
- d) Bypass insurance and bill cash without documenting

16) Rx: "prednisone 10 mg: take 4 tabs daily x 3 days, then 3 tabs daily x 3 days, then 2 tabs daily x 3 days, then 1 tab daily x 3 days." Quantity?

- a) 24
- b) 30
- c) 36
- d) 40

17) A customer requests pseudoephedrine. The sale is within gram limits, but the customer presents an expired driver's license. What should happen?

- a) Complete the sale and note "expired" in the log
- b) Decline the sale and offer a non-regulated alternative
- c) Accept it if the photo matches the customer
- d) Accept it if the ID number is readable

18) An isotretinoin prescription appears "patient not qualified today" in the REMS workflow. What should you do?

- a) Process and dispense because REMS is only for counseling
- b) Dispense after obtaining the patient's signature
- c) Place the prescription on hold until REMS requirements are met and the system allows dispensing
- d) Substitute tretinoin cream and dispense

19) Rx: "Advair Diskus 250/50: inhale 1 puff BID." Inhaler contains 60 inhalations. Days' supply?

- a) 15
- b) 20
- c) 30
- d) 60

20) While reviewing the label preview, the directions print "take 1 tablet twice daily," but the

eRx clearly states "take 1 tablet daily." What should you do?

- a) Dispense and tell the patient verbally at pickup
- b) Correct the sig in the system and reprint the label before filling continues
- c) Handwrite the corrected directions on the label and initial it
- d) Ignore it because pharmacist verification will catch it

21) A manufacturer recall is issued for a specific lot of atorvastatin 20 mg. Which action is correct?

- a) Remove all atorvastatin products from stock
- b) Quarantine only the affected lot(s) and follow the recall instructions for disposition
- c) Continue dispensing until the supply runs out
- d) Re-label the bottles with a new expiration date and keep dispensing

22) A prescriber writes clindamycin 1% topical solution "Disp 60 mL," but the system defaults to 30 mL. What should you do?

- a) Keep 30 mL because it is standard for topicals
- b) Change the quantity to 60 mL as written
- c) Change the directions to TID to justify 60 mL
- d) Substitute clindamycin capsules

23) Another pharmacy requests a transfer of a remaining refill for a Schedule II prescription. What is correct under federal law?

- a) It can be transferred once if both pharmacies are in the same state
- b) It can be transferred if the patient requests it
- c) It can be transferred only within the same chain
- d) Schedule II prescriptions cannot be transferred

24) Rx: gabapentin 300 mg "take 1 capsule TID." Disp #90. Days' supply?

- a) 15
- b) 30
- c) 45
- d) 90

25) A new prescription arrives for lisinopril 10 mg daily. The patient profile indicates pregnancy. What is the best technician action?

- a) Fill because it is a common antihypertensive
- b) Route for pharmacist review prior to dispensing
- c) Remove the pregnancy note because it may be outdated
- d) Change to labetalol and continue

26) A caller asks for the medication name and directions for an adult patient. The caller is not listed as an authorized representative. What should you do?

- a) Provide the name but not directions
- b) Provide directions but not the name
- c) Provide information if the caller confirms DOB and address
- d) Decline to disclose and advise the patient to contact the pharmacy or add authorization

27) Rx: fentanyl patch "apply 1 patch every 72 hours." Disp #10. Days' supply?

- a) 10
- b) 20
- c) 30
- d) 72

28) You find nitroglycerin sublingual tablets removed from the original container and stored without the desiccant. What should you do?

- a) Repackage in an amber vial and use first
- b) Return them to the original bottle and keep using
- c) Remove from dispensing stock and quarantine for pharmacist determination
- d) Keep them for internal use only

29) Rx: latanoprost 0.005% "instill 1 drop OU QHS." Disp 2.5 mL. Assume 20 drops/mL. Approximate days' supply?

- a) 12
- b) 25
- c) 50
- d) 100

30) Which record must be readily retrievable for DEA inspection at a retail pharmacy?

- a) Cash register deposit slips
- b) Employee vaccination schedules
- c) Controlled substance purchase records (invoices) with drug, quantity, date, and supplier information
- d) Patient driver's license copies for controlled pickups

31) A prescriber's office requests a one-year medication fill history for a patient. No authorization is on file, and the office cannot accept secure communications. What should you do?

- a) Provide it because it is for treatment
- b) Provide only non-controlled medications
- c) Read it over the phone without quantities
- d) Do not release the information without proper authorization and an appropriate secure method

32) Rx: ondansetron ODT 4 mg "take 1 tablet q8h PRN nausea." Disp #12. If billing requires max daily use, what days' supply should be billed?

- a) 2
- b) 3
- c) 4
- d) 12

33) A refrigerated medication is found on the counter with unknown time out of refrigeration. What should you do?

- a) Return it to the refrigerator and continue dispensing
- b) Mark it "use first" and return to stock
- c) Discard immediately without documentation
- d) Quarantine it for pharmacist evaluation using stability/temperature excursion guidance

34) Rx: albuterol HFA "inhale 2 puffs q4h PRN." Canister contains 200 actuations. For max use, assume 6 doses/day. Days' supply?

- a) 10
- b) 12
- c) 16
- d) 25

35) A prescription is for diltiazem ER 240 mg daily, but the robot contains only diltiazem IR 120 mg and 240 mg. What should you do?

- a) Use IR 240 mg because the strength matches
- b) Use IR and change the sig to BID
- c) Select the ER product or route for pharmacist resolution before filling
- d) Substitute verapamil ER 240 mg

36) A prescription is entered for spironolactone 25 mg daily. The profile shows lisinopril and potassium chloride. What is the best next step?

- a) Continue because this combination is common
- b) Remove potassium chloride from the active profile and proceed
- c) Route for pharmacist review due to potential hyperkalemia risk
- d) Change spironolactone to hydrochlorothiazide

37) A system interaction alert appears for sertraline and linezolid on a new prescription. What should you do?

- a) Override and proceed because it is prescriber-directed
- b) Route the order for pharmacist review before dispensing
- c) Substitute doxycycline for linezolid and proceed
- d) Tell the patient to stop sertraline while on the antibiotic

38) Unit-dose delivery arrives with warfarin cards missing lot numbers on the receiving label, and your site requires lot/expiration for high-alert traceability. What should you do?

- a) Stock them because the expiration date is present
- b) Write the lot number from another carton of the same drug
- c) Quarantine and obtain correct traceability information before stocking
- d) Stock and document lot as "unknown"

39) Rx: cephalexin 500 mg "take 1 capsule q6h x 10 days." Quantity?

- a) 30
- b) 40
- c) 60
- d) 120

40) During a scheduled C-II cycle count, what is an appropriate technician role?

- a) Estimate the count based on dispensing history
- b) Count and document per procedure and report discrepancies per workflow
- c) Adjust the perpetual inventory to match what "should be" on hand
- d) Delay counting until after closing so no one is watching

41) A prescription states "take 1½ tablets daily." Each tablet is 80 mg. How many mg per day?

- a) 100 mg
- b) 120 mg
- c) 140 mg
- d) 160 mg

42) A hard-copy noncontrolled prescription is dropped off without a prescriber signature. What should you do?

- a) Fill it because the prescriber name is printed
- b) Have the patient sign for the prescriber
- c) Enter it as a verbal order
- d) Set it aside for pharmacist review to obtain a valid prescription before processing

43) A vaccine shipment arrives with a temperature indicator showing a possible excursion in transit. What should you do?

- a) Stock immediately to prevent further warming
- b) Refrigerate and use first
- c) Quarantine and follow excursion procedures to determine usability before stocking
- d) Discard immediately without contacting anyone

44) A patient requests a partial fill of a Schedule II due to cost. What should you do?

- a) Process a partial fill and tell them they can pick up the remainder anytime
- b) Inform them partial fills are never allowed for Schedule II
- c) Convert to a Schedule III equivalent so refills are allowed
- d) Route to the pharmacist to determine whether a partial fill is permissible and to ensure proper documentation/timing

45) Humalog U-100: "inject 10 units before meals TID." Disp 1 vial (10 mL; 100 units/mL). Days' supply?

- a) 25
- b) 30
- c) 33
- d) 50

46) The sig reads: "Apply sparingly to affected area TID–QID." What does TID–QID mean?

- a) Every 3 to 4 hours
- b) 3 to 4 times daily
- c) Twice daily as needed
- d) Once daily at bedtime

47) While pulling stock, you scan amlodipine 10 mg but the prescription is for amiodarone 200 mg. What should you do?

- a) Override because both are cardiovascular medications
- b) Continue counting; pharmacist will catch it
- c) Change the prescription to amlodipine to match what was scanned
- d) Stop and select the correct medication that matches the prescription before continuing

48) Rx: fluticasone nasal spray "2 sprays in each nostril daily." Bottle contains 120 sprays. Days' supply?

- a) 15
- b) 20
- c) 30
- d) 60

49) You are entering an otic drop prescription and notice the system defaulted to an ophthalmic product with a similar name. What is the best prevention step?

- a) Use more abbreviations to speed entry
- b) Confirm route/dosage form from the original prescription during entry and again during product selection
- c) Choose ophthalmic because it can be used in the ear
- d) Skip barcode scanning for small bottles

50) Lantus SoloStar box contains five 3 mL pens (100 units/mL). Sig: "Inject 28 units nightly." Days' supply?

- a) 25
- b) 45
- c) 53
- d) 75

51) A Schedule II eRx is missing required prescriber information due to a system mapping error. What should you do?

- a) Dispense because the prescriber NPI is present
- b) Find the prescriber DEA online and enter it
- c) Convert it to a verbal order
- d) Hold and route for pharmacist verification/correction before dispensing

52) A patient brings in a bottle of unknown tablets with no label for disposal. What should you do?

- a) Place it in regular trash
- b) Put it back into stock if tablets look intact
- c) Put it in the pharmacy's designated medication disposal process and keep it separate from inventory
- d) Flush the tablets

53) Rx: Augmentin 875/125 mg "1 tablet BID x 7 days." Quantity?

- a) 7
- b) 10
- c) 14
- d) 21

54) Rx: atorvastatin 10 mg nightly #90. Only 20 mg tablets are available. What is the best technician action?

- a) Dispense 20 mg and instruct patient to take 1/2 tablet
- b) Dispense 20 mg and change quantity to 45
- c) Substitute simvastatin 10 mg

d) Route for pharmacist review to determine authorization/appropriate product selection

55) A patient asks whether cetirizine or loratadine is more likely to cause drowsiness. What should the technician do?

- a) Recommend cetirizine because it is stronger
- b) Recommend loratadine because it never causes drowsiness
- c) Tell the patient to take both and see what works
- d) Refer the patient to the pharmacist for product selection counseling

56) A C-II hard copy is presented with the patient address missing. What should happen?

- a) Add the address from the patient profile and process
- b) Have the patient write the address on the prescription
- c) Route for pharmacist review to determine if and how the missing element can be resolved before dispensing
- d) Call the prescriber and take a verbal correction and fill

57) A new prescription for clopidogrel is entered and an interaction alert appears with the patient's active omeprazole. What should you do?

- a) Override because it is common
- b) Switch omeprazole to pantoprazole automatically
- c) Delete omeprazole from the profile and proceed
- d) Route for pharmacist review before dispensing

58) During sterile compounding, a technician touches a door handle with gloved hands and returns to the hood to continue. What should occur next?

- a) Continue because gloves are worn
- b) Disinfect the IV bag only and continue
- c) Stop, perform appropriate decontamination steps, and reglove before continuing
- d) Finish the preparation and document the touch afterward

59) Rx: metformin ER 500 mg "take 2 tablets with evening meal." Disp #60. Days' supply?

- a) 15
- b) 30
- c) 45
- d) 60

60) Patient profile shows methotrexate 2.5 mg weekly. New eRx says methotrexate 2.5 mg "take 1 tablet daily." What should you do?

- a) Enter as written because it is electronic
- b) Change to weekly because that is typical
- c) Fill daily but reduce quantity
- d) Route for pharmacist review due to high-alert dosing frequency concern

61) A new order comes in for "Toprol-XL 50 mg daily." The patient recently filled metoprolol tartrate 50 mg BID. What should you do?

- a) Fill Toprol-XL and inactivate tartrate automatically
- b) Fill both because they are both metoprolol
- c) Change Toprol-XL to tartrate to match the profile

d) Ensure the pharmacist reviews for formulation change/duplicate therapy risk before dispensing

62) Rx: gentamicin ophthalmic "instill 2 drops OU q4h x 7 days." Assume 6 doses/day and 20 drops = 1 mL. Approximately how many mL are needed?

- a) 4.2 mL
- b) 6.0 mL
- c) 8.4 mL
- d) 12.6 mL

63) While counting, you find one tablet is chipped and powdery. What should you do?

- a) Dispense it; chips do not matter
- b) Set it aside and replace it with an intact tablet to maintain the correct count
- c) Crush it and include the powder to keep count accurate
- d) Add an extra tablet and keep the chipped one in the vial

64) A friend arrives to pick up a controlled medication for a patient. What should you do?

- a) Refuse all third-party pickups for controlled substances
- b) Release it if they know the patient's DOB
- c) Release it only if the person is a family member
- d) Verify identity and authorization per pharmacy requirements and document as required before release

65) Rx: clindamycin 300 mg "take 1 capsule q8h x 10 days." Quantity?

- a) 20
- b) 30
- c) 40
- d) 60

66) Rx: fentanyl patch "apply 1 patch every 48 hours." Disp #5. Days' supply?

- a) 5
- b) 10
- c) 12
- d) 15

67) A patient asks you to transfer gabapentin from another pharmacy. What is the best technician action?

- a) Complete the transfer independently without pharmacist involvement
- b) Tell the patient transfers are never allowed
- c) Ask the patient to bring the old bottle so you can copy the label and dispense
- d) Collect required transfer information and proceed according to law/policy, involving the pharmacist as required

68) Rx: Victoza 1.2 mg SQ daily. One pen contains 18 mg total. Days' supply for 1 pen?

- a) 10
- b) 12
- c) 15
- d) 30

69) A patient requests an early refill for oxycodone/APAP stating it was stolen; last fill was 9 days ago for a 30-day supply. What should you do first?

- a) Refuse because controlled substances are never replaced
- b) Route the request for pharmacist evaluation and documentation steps
- c) Immediately call the prescriber and request an emergency replacement
- d) Process a partial fill as cash to avoid insurance blocks

70) Which generic name matches Xarelto?

- a) Dabigatran
- b) Rivaroxaban
- c) Apixaban
- d) Edoxaban

71) During will-call staging, the bag label shows the correct patient name but a different DOB than the profile. What should you do?

- a) Release if the address matches
- b) Correct the DOB with pen and initial
- c) Leave it because DOB is not used at pickup
- d) Pull it from will-call for correction/verification before it can be sold

72) An ear drop prescription says "instill 4 drops in left ear BID." Which abbreviation indicates left ear?

- a) OD
- b) AU
- c) AS
- d) OS

73) A DSCSA inquiry is opened for a suspect product. What should the technician help locate?

- a) Patient copay receipts for the last 30 days
- b) The prescriber's state license number
- c) Transaction information/history and related tracing statements for the product
- d) The pharmacy's controlled substance perpetual inventory report

74) A prescription for losartan 50 mg daily arrives. The profile note indicates pregnancy (2nd trimester). What should you do?

- a) Continue because it is only a profile note
- b) Change to valsartan because it is a different ARB
- c) Delete the pregnancy note as outdated
- d) Route for pharmacist review due to contraindication risk

75) Rx: clotrimazole 1% cream "apply BID," Disp 45 g. The computer default is 30 g. What should you enter?

- a) Dispense 30 g because that is standard
- b) Dispense 45 g as written
- c) Change directions to TID so 45 g makes sense
- d) Substitute clotrimazole/betamethasone to justify 45 g

76) Warfarin eRx sig field says "take as directed," and the prescriber note states "start 1 tablet daily; adjust per INR." What is the best way to enter for the label?

- a) Enter "take as directed" only
- b) Enter "take 1 tablet by mouth daily" and include titration language in the appropriate notes/auxiliary area per system capability
- c) Enter "take 1 tablet twice daily"
- d) Enter "take 1 tablet daily PRN"

77) A caller claims to be the patient's spouse and requests the patient's medication list. The caller is not listed as authorized but confirms DOB. What should you do?

- a) Provide the list because spouse implies authorization
- b) Provide drug names only
- c) Provide the information if they also know the prescriber name
- d) Decline disclosure until authorization is documented or the patient contacts the pharmacy

78) Rx: doxycycline 100 mg "take 1 capsule BID x 10 days." Quantity?

- a) 10
- b) 14
- c) 20
- d) 28

79) A patient asks whether an antibiotic will interfere with birth control. What should the technician do?

- a) Tell them yes for all antibiotics
- b) Tell them no for all antibiotics
- c) Recommend doubling the contraceptive dose
- d) Refer the question to the pharmacist

80) Rx: bupropion XL 150 mg daily. Stock includes bupropion SR 150 mg and bupropion XL 150 mg. Which should be dispensed?

- a) SR 150 mg
- b) XL 150 mg
- c) Either SR or XL is acceptable once daily
- d) IR 75 mg, two tablets daily

81) You see morphine sulfate ER 30 mg and morphine sulfate IR 30 mg on the shelf. The prescription is for ER. What is the safest product-selection step?

- a) Select IR because it works faster
- b) Select either because the strength matches
- c) Select ER and verify with barcode/NDC match before filling
- d) Select IR and change the directions to q12h

82) Same-day eRx orders arrive for omeprazole 20 mg daily and esomeprazole 20 mg daily from different prescribers for the same patient. What should you do?

- a) Fill both because they are different drugs
- b) Route for pharmacist review due to potential duplicate therapy
- c) Delete the older prescription automatically

d) Change one to famotidine and proceed

83) Under federal law, which schedules may be refilled up to 5 times within 6 months of the date written (if authorized)?

a) Schedule II

b) Schedule III and IV

c) Schedule V only

d) Schedule I

84) Insulin glargine U-100: inject 34 units nightly. Disp 1 box (5 pens, each 3 mL). Days' supply?

a) 30

b) 39

c) 44

d) 53

85) You notice an IV label prints a concentration that does not match the compounding worksheet. What should you do?

a) Hand-correct the label and initial it

b) Compound using the worksheet and ignore the label

c) Stop and ensure the discrepancy is resolved and correct labeling is produced before continuing

d) Make a second bag as backup

86) A patient asks you to teach them how to use a new inhaler. What should you do?

a) Provide full device counseling because it is not a medication

b) Demonstrate technique and recommend timing changes

c) Refer the patient to the pharmacist for counseling

d) Tell the patient to read the package insert

87) Rx: cefdinir suspension "take 5 mL BID x 10 days." How many mL should be dispensed?

a) 50 mL

b) 75 mL

c) 100 mL

d) 120 mL

88) A claim for triamcinolone 0.1% cream rejects "days' supply required." Sig: "apply to affected area BID." Quantity: 30 g. No area specified. What is the most appropriate days' supply entry without further clarification?

a) 7 days

b) 14 days

c) 30 days

d) 90 days

89) During entry for digoxin 0.125 mg daily, 0.25 mg was accidentally selected. Which workflow control most directly reduces the chance of dispensing the wrong strength?

a) Store all cardiac drugs together

b) Print labels in larger font

- c) Count by fives
- d) Barcode/NDC verification of drug and strength during filling

90) A prescription is for "Eliquis 5 mg BID." Which generic name matches Eliquis?

- a) Apixaban
- b) Edoxaban
- c) Rivaroxaban
- d) Dabigatran

Answer Key

1) c	32) c	63) b
2) a	33) d	64) d
3) c	34) c	65) b
4) b	35) c	66) b
5) d	36) c	67) d
6) c	37) b	68) c
7) b	38) c	69) b
8) b	39) b	70) b
9) c	40) b	71) d
10) d	41) b	72) c
11) c	42) d	73) c
12) b	43) c	74) d
13) d	44) d	75) b
14) c	45) c	76) b
15) c	46) b	77) d
16) c	47) d	78) c
17) b	48) c	79) d
18) c	49) b	80) b
19) c	50) c	81) c
20) b	51) d	82) b
21) b	52) c	83) b
22) b	53) c	84) c
23) d	54) d	85) c
24) b	55) d	86) c
25) b	56) c	87) c
26) d	57) d	88) b
27) c	58) c	89) d
28) c	59) b	90) a
29) c	60) d	
30) c	61) d	
31) d	62) c	

PTCB Sim Test 16

1) A new eRx shows: "levothyroxine 75 mcg PO daily. #90." During product selection, the technician accidentally selects levothyroxine 75 mg. What is the best technician action before continuing?

- a) Continue and reduce the quantity to #9
- b) Correct the strength to 75 mcg and proceed through normal verification workflow
- c) Change the sig to "take 1/1000 tablet daily"
- d) Dispense 75 mg because it is the brand-equivalent strength

2) A person at the counter asks to pick up a prescription for their roommate and asks, "What is it for?" They can provide the roommate's name and address but not DOB. What should the technician do?

- a) Provide only the medication name but not the directions
- b) Provide the indication only because it isn't the same as the drug name
- c) Ask for an additional identifier before discussing any prescription details
- d) Provide the price only and confirm the drug and directions

3) A new eRx reads: "metformin ER 500 mg: take 2 tablets with evening meal. #60." The profile shows metformin IR 500 mg 1 tablet BID with refills remaining. What is the best next step?

- a) Fill the ER and automatically discontinue the IR in the profile
- b) Fill both to avoid interruption until the patient decides
- c) Change ER directions to 1 tablet BID to match the old Rx
- d) Enter the ER prescription and send for pharmacist review for therapy change/duplication

4) The refrigerator log shows vaccines at 11°C (51.8°F) overnight. What is the best immediate action for the vaccines?

- a) Quarantine and label "Do Not Use" while excursion guidance is obtained
- b) Return them to the fridge and continue using if vials look normal
- c) Move them to the freezer to restore proper temperature faster
- d) Discard immediately because any excursion requires disposal

5) A paper prescription for alprazolam 0.5 mg includes patient name, drug, directions, date, and prescriber signature, but no prescriber address. What should the technician do?

- a) Add the pharmacy address as the prescriber address
- b) Process it because prescriber address is never required
- c) Ask the patient to write the prescriber address from memory
- d) Hold for pharmacist review to confirm required elements per law/state rules

6) Azithromycin suspension is prescribed: "12.5 mL PO day 1, then 6.25 mL PO daily days 2–5." How many mL should be dispensed?

- a) 35 mL
- b) 31.25 mL
- c) 37.5 mL
- d) 42.5 mL

- 7) A patient requests a transfer of a clonazepam prescription from another pharmacy. The receiving pharmacist is on a call. What can the technician do to help appropriately?
- a) Accept the verbal transfer and document it as the transferring agent
 - b) Tell the patient transfers of Schedule IV are not allowed
 - c) Collect the other pharmacy's information and prepare the patient profile/notes for pharmacist-to-pharmacist transfer
 - d) Use a photo of the old label to create a new prescription
- 8) While stocking, the technician finds cefdinir capsules that expired last month. What is the correct action?
- a) Place them behind in-date stock and dispense only if needed
 - b) Relabel the expiration one year out if the bottle is unopened
 - c) Remove from active stock and place in the designated expired/return area with required documentation
 - d) Keep them for cash patients to reduce waste
- 9) Prednisone 10 mg directions: "Take 2 tablets PO BID x 5 days, then 1 tablet PO BID x 5 days." How many tablets are needed?
- a) 40
 - b) 20
 - c) 30
 - d) 60
- 10) A patient asks if morphine ER tablets can be crushed. What is the best technician response?
- a) "Yes, if mixed with soft food it's fine"
 - b) "No—let me get the pharmacist to discuss safe options"
 - c) "Yes, but only if you take it with water"
 - d) "Crush it and take half now and half later"
- 11) A customer requests pseudoephedrine 30 mg tablets, quantity 96. Under federal rules, what must occur before completing the sale?
- a) Verify photo ID, complete the required log entry, and confirm sales limits are not exceeded
 - b) Require a written prescription
 - c) Obtain a signed HIPAA authorization
 - d) Have the pharmacist perform mandatory counseling
- 12) Isotretinoin is ready, but the system shows REMS steps incomplete. What should the technician do?
- a) Dispense if the patient states they completed requirements
 - b) Dispense a partial supply and complete the rest later
 - c) Ensure REMS dispensing authorization is completed and requirements are met before release
 - d) Bypass REMS because it is not a controlled substance
- 13) A technician counts a sulfonamide antibiotic and then begins counting warfarin on the same tray without cleaning. What is the best immediate action?

- a) Continue because tablets are solid and cross-contact is minimal
- b) Only clean when switching between controlled and noncontrolled medications
- c) Clean the tray/spatula per procedure before continuing, and re-count as needed
- d) Spray the tray with alcohol and immediately count the next drug while wet

14) An Rx is entered for Latuda 40 mg 1 tablet daily. The technician pulled lurasidone 20 mg. What should the technician do?

- a) Change the prescription to 20 mg to match the bottle
- b) Dispense two 20 mg tablets daily but leave the label as 40 mg 1 tablet daily
- c) Put a note on the bag saying "take 2 tablets"
- d) Pull the correct strength product (or correct entry per authorization) before proceeding

15) A new Rx for cephalexin arrives. The profile lists "penicillin—anaphylaxis." What should the technician do?

- a) Fill as normal because cephalexin is not a penicillin
- b) Route to pharmacist for allergy risk assessment before dispensing
- c) Delete the allergy because it's unrelated
- d) Substitute azithromycin and process

16) A new eRx says: "warfarin 5 mg daily (dose change)." The patient filled warfarin 2.5 mg yesterday. What should the technician do?

- a) Enter the new Rx and send for pharmacist review to confirm dose-change plan and discontinue/clarify prior directions
- b) Fill the 5 mg and delete the 2.5 mg from the record entirely
- c) Fill both strengths and let the patient decide which to take
- d) Dispense 5 mg tablets and instruct the patient to take half daily

17) An insurance claim rejects "Refill too soon." The patient says the medication was lost. What should the technician do?

- a) Change the last fill date so it will adjudicate
- b) Collect key details (fill date, days' supply, circumstances) and route for appropriate override/plan steps
- c) Override with any code to keep workflow moving
- d) Tell the patient early refills are never allowed

18) A hospital order prints potassium chloride 20 mEq tablets, but the patient is listed as NPO. What is the best technician action?

- a) Send the tablets because nursing will decide administration
- b) Convert it to IV potassium automatically
- c) Hold and bring the mismatch to the pharmacist for resolution
- d) Crush the tablets and send with "via NG" handwritten

19) A controlled-substance shipment arrives with a broken outer seal and no explanation. What should the technician do?

- a) Accept it if the invoice matches the contents
- b) Quarantine and escalate per suspect shipment procedure
- c) Reseal it and place directly into the C-II safe
- d) Discard the shipment to prevent diversion

20) Diltiazem CD 240 mg daily is prescribed. The technician selected diltiazem IR 120 mg tablets. What should be done next?

- a) Change directions to 120 mg BID and fill
- b) Fill IR once daily because total daily dose matches
- c) Correct to the CD formulation and correct strength as written before continuing
- d) Substitute another calcium channel blocker

21) A manufacturer recall notice lists a specific lot of losartan currently on the shelf. What should the technician do?

- a) Keep dispensing until inventory is depleted, then remove it
- b) Remove the affected lot from dispensing stock and quarantine/return per recall instructions
- c) Relabel with a different lot number to avoid confusion
- d) Ignore it unless a patient complains

22) Diclofenac 1% gel directions: "Apply 1 g topically BID." The tube contains 100 g. What is the days' supply?

- a) 100 days
- b) 75 days
- c) 50 days
- d) 25 days

23) During entry, the system flags a major interaction: simvastatin + clarithromycin. What is the correct technician action?

- a) Override because alerts are common
- b) Change clarithromycin to azithromycin automatically
- c) Reduce simvastatin quantity to a 5-day supply and proceed
- d) Route the interaction for pharmacist review before dispensing

24) A pharmacist received a verbal tramadol prescription, but the documentation hardcopy is missing the prescriber phone number. What should occur before dispensing?

- a) Dispense because prescriber name is enough for a verbal order
- b) Add any phone number found online and initial it
- c) Ensure required verbal order documentation is completed/validated before processing to fill
- d) Ask the patient for the prescriber phone number at pickup

25) A label prints for hydroxyzine 25 mg, but the technician accidentally grabbed hydralazine 25 mg. Which workflow control best prevents this error?

- a) Store look-alike drugs adjacent to compare labels
- b) Rely on cap color differences among manufacturers
- c) Use barcode/NDC scanning at product selection and again at verification points
- d) Remove Tall Man lettering to reduce confusion

26) Ventolin HFA: "Inhale 2 puffs q4–6h PRN." Inhaler contains 200 actuations. If used at maximum of 2 puffs every 4 hours, what is the days' supply?

- a) 20 days

- b) 16 days
- c) 14 days
- d) 12 days

27) A patient requests an early fill of Adderall due to travel. The Rx is Schedule II. What is the best technician action?

- a) Process early if the patient pays cash
- b) Directly change the days' supply to help insurance pay
- c) Transfer the prescription to another pharmacy
- d) Refer the request for pharmacist review of timing, payer/state rules, and whether a new Rx is needed

28) A new Rx arrives for atorvastatin 40 mg daily. The profile shows simvastatin 40 mg filled 7 days ago. What should the technician do?

- a) Enter atorvastatin and send for pharmacist review for therapeutic duplication vs therapy change
- b) Delete atorvastatin as a duplicate
- c) Discontinue simvastatin in the system and fill atorvastatin without review
- d) Dispense atorvastatin 20 mg to reduce duplication risk

29) A DSCSA package will not scan and the human-readable lot number is smeared. What should the technician do?

- a) Type the closest-looking lot number and accept it
- b) Remove the label and print a replacement
- c) Set aside as suspect product and escalate for verification with supplier/manufacturer
- d) Dispense first so it doesn't remain in inventory

30) Augmentin 875/125 mg: "1 tablet PO BID x 10 days." What quantity should be dispensed?

- a) 28 tablets
- b) 14 tablets
- c) 10 tablets
- d) 20 tablets

31) A patient returns an opened bottle of lisinopril stating it "made them tired" and wants it put back for later. What should the technician do?

- a) Restock if tablets appear intact
- b) Restock if the bottle is more than half full
- c) Do not restock returned patient medication; process return/disposal per procedure and route clinical concern appropriately
- d) Repackage into a new vial and restock

32) The sig reads: "1 gtt OU BID." What does this mean?

- a) Instill 1 drop in right eye twice daily
- b) Instill 1 drop in left eye twice daily
- c) Instill 1 drop in both eyes twice daily
- d) Instill 2 drops in one eye twice daily

33) An eRx reads: "methotrexate 2.5 mg: take 6 tablets daily." Notes indicate rheumatoid arthritis. What should the technician do?

- a) Fill as written because RA dosing can be daily early on
- b) Change "daily" to "weekly" and continue
- c) Fill but add an auxiliary label "weekly"
- d) Hold and route for pharmacist review due to high-risk dosing frequency concern

34) A patient presents a Schedule II paper prescription for oxycodone dated today, but the prescriber signature is missing. What should the technician do?

- a) Process it and ask the patient to return later for signature
- b) Do not process; route because a C-II must be signed to be valid
- c) Call the prescriber for a verbal authorization to dispense
- d) Have the patient sign as the prescriber's agent

35) Apixaban 5 mg BID is prescribed, and the profile shows rivaroxaban 20 mg daily filled 5 days ago. What is the best technician action?

- a) Discontinue rivaroxaban and fill apixaban automatically
- b) Fill both because overlap is always required
- c) Change apixaban to 2.5 mg BID to reduce bleeding risk
- d) Route for pharmacist review of duplicate anticoagulant therapy/transition plan

36) A prescription for furosemide is missing the patient's address. It is not controlled. What is the best action?

- a) Refuse to fill until the prescriber rewrites it
- b) Fill only a 72-hour emergency supply
- c) Process the prescription; address is not required for validity of a non-controlled prescription under federal rules
- d) Substitute a different diuretic that does not require address

37) Clindamycin 150 mg: "Take 2 capsules PO QID x 7 days." How many capsules are needed?

- a) 56
- b) 42
- c) 84
- d) 28

38) A prescriber's office calls requesting a patient's full 12-month fill history. No release is on file. Caller provides patient name and DOB but cannot confirm a callback number. What should the technician do?

- a) Provide it because it's for treatment
- b) Provide only noncontrolled medications
- c) Refuse all medication-history requests from prescribers
- d) Use callback/verification procedure through authenticated contact information before disclosure

39) A patient is newly prescribed Suboxone film. What are the active ingredients?

- a) Buprenorphine/naloxone
- b) Buprenorphine/naltrexone

- c) Methadone/naloxone
- d) Naloxone/naltrexone

40) A child weighs 44 lb. Order: amoxicillin 40 mg/kg/day divided BID. What is the total daily dose in mg?

- a) 600 mg/day
- b) 1,200 mg/day
- c) 800 mg/day
- d) 400 mg/day

41) Two patient profiles have the same name and similar DOBs. A refill is queued for processing. What should the technician do first?

- a) Use the profile with the most recent activity
- b) Merge the profiles to prevent confusion
- c) Verify at least two identifiers (DOB plus address/phone) before processing
- d) Process the refill and correct later if rejected

42) A prescription for hydrocodone/APAP 5/325 mg shows "refills: 3." What is the correct action?

- a) Process with 3 refills if the prescriber authorized it
- b) Change refills to 0 and fill without escalation
- c) Route because Schedule II prescriptions cannot have refills
- d) Transfer to a pharmacy that can allow refills

43) A patient asks, "Will this antibiotic affect my warfarin?" What should the technician do?

- a) Recommend a vitamin K supplement while on the antibiotic
- b) Tell the patient to stop warfarin until the antibiotic is finished
- c) Tell the patient to separate doses by 2 hours
- d) Refer the question for pharmacist counseling

44) In a hospital setting, a vancomycin label prints "infuse over 30 minutes," but the verified order states "infuse over 2 hours." What should the technician do?

- a) Cross out the label and handwrite "2 hours" without updating system records
- b) Send it because nursing will follow the MAR
- c) Stop distribution and route for correction so the label matches the verified order
- d) Change the order to 30 minutes to match the label default

45) The sig says: "Take 1 tablet PO qAM." What does qAM mean?

- a) At bedtime
- b) Every other morning
- c) Every morning
- d) Every evening

46) The profile lists "TMP-SMX—rash." A new Rx arrives for sulfadiazine. What should the technician do?

- a) Process normally because sulfa allergies only apply to penicillins
- b) Route for pharmacist review due to possible sulfonamide allergy cross-reactivity concern
- c) Delete the allergy because it is not anaphylaxis

d) Automatically switch to doxycycline

47) Omeprazole 20 mg is billed to insurance. The technician selected an OTC omeprazole NDC, resulting in rejection. What is the best next step?

- a) Select the correct covered product/NDC and rebill per plan requirements
- b) Bill cash automatically without asking the patient
- c) Change therapy to pantoprazole without authorization
- d) Split the claim into two 14-day supplies

48) A Schedule III prescription is dated 1/10/2026 and presented 8/15/2026. Under federal law, can it be filled?

- a) Yes, if any refills remain it is valid indefinitely
- b) No, all controlled prescriptions expire after 30 days
- c) Yes, controlled prescriptions are valid for 12 months
- d) No, Schedule III–V prescriptions cannot be filled more than 6 months after date written

49) A patient requests a printed medication history for taxes and presents valid ID. What should the technician do?

- a) Refuse because medication histories cannot be provided to patients
- b) Provide it after verifying identity and completing the pharmacy's release process
- c) Provide only controlled-substance fills
- d) Email it unsecured for convenience

50) Label: "Take 0.5 mL PO daily." Concentration is 250 mg/5 mL. How many mg is each dose?

- a) 25 mg
- b) 12.5 mg
- c) 50 mg
- d) 125 mg

51) A prescription is for phenytoin ER 100 mg capsules, but the technician selected phenytoin injection 50 mg/mL. What should happen next?

- a) Dispense injection with oral syringes
- b) Continue because both forms are equivalent
- c) Correct the dosage form/route selection before proceeding
- d) Substitute carbamazepine 100 mg capsules

52) A patient requests non-child-resistant caps due to arthritis. What is the best technician action?

- a) Provide non-safety caps only for controlled substances
- b) Refuse because federal law always requires child-resistant caps
- c) Open the bottle at pickup and keep the safety cap for home
- d) Provide a non-safety cap upon request and document per procedure

53) A faxed clonazepam prescription arrives and the signature appears copied. What should the technician do?

- a) Fill it because fax signatures often look copied
- b) Partially fill and complete later

- c) Call the patient to confirm authenticity
- d) Route for pharmacist validation of prescription authenticity before processing

54) The prescription is for erythromycin ophthalmic ointment, but the technician selected erythromycin topical (skin) ointment. What is the correct step?

- a) Dispense topical ointment and add "for eye use" on the label
- b) Correct to the ophthalmic dosage form before filling/labeling
- c) Dispense topical because ointments are interchangeable
- d) Dilute topical ointment with sterile saline

55) A patient asks whether sertraline can be taken with St. John's wort. What should the technician do?

- a) Advise taking them 6 hours apart
- b) Recommend doubling sertraline to offset the herb
- c) Refer the question for pharmacist counseling
- d) Assure the patient herbs do not interact with prescriptions

56) A refrigerated biologic carton is damp and torn; the vial appears intact. What should the technician do?

- a) Tape the carton and continue processing
- b) Dispense because only vial integrity matters
- c) Quarantine and escalate for storage/integrity determination per workflow
- d) Replace the carton with another lot and dispense the same vial

57) Directions: "Take 1.5 tablets PO daily." Quantity dispensed: 45 tablets. What is the days' supply?

- a) 20 days
- b) 45 days
- c) 30 days
- d) 60 days

58) A patient requests home delivery to a new address and asks the technician to update it. What is the best action?

- a) Update only after pharmacist counseling
- b) Update after verifying identity and document per workflow
- c) Refuse because only pharmacists can change demographics
- d) Ship to the old address and change later

59) Zofran ODT 4 mg is which medication?

- a) Oseltamivir 4 mg capsule
- b) Olanzapine ODT 4 mg
- c) Ondansetron orally disintegrating tablet 4 mg
- d) Omeprazole delayed-release tablet 4 mg

60) A patient returns an unopened eye-drop bottle that was dispensed yesterday and stored at home; the seal is intact. What should the technician do?

- a) Restock because the seal is intact
- b) Restock after wiping the bottle with alcohol

- c) Repackage into a new bottle and relabel with a new beyond-use date
- d) Do not restock returned medication; handle disposal/return per procedure

61) A new Rx for lisinopril 10 mg daily is entered; the profile indicates pregnancy (third trimester). What should the technician do?

- a) Process as usual; pregnancy does not affect ACE inhibitors
- b) Switch to labetalol automatically
- c) Route for pharmacist review due to contraindication concern
- d) Add an auxiliary "avoid in pregnancy" label and dispense

62) A patient wants a transfer of a noncontrolled refill from a pharmacy that permanently closed. No one answers the phone and there is no shared database. What is the best next step?

- a) Create a new prescription with remaining refills and dispense
- b) Route for pharmacist review; a new prescription may be needed if transfer information cannot be obtained
- c) Dispense a 30-day emergency supply without escalation
- d) Deny because transfers are never allowed

63) Sig: "Take 2 tsp PO TID." How many mL is each dose?

- a) 15 mL
- b) 5 mL
- c) 10 mL
- d) 30 mL

64) A sterile compounding worksheet shows volumes added that do not match the documented final volume. What should the technician do?

- a) Adjust the final volume on paper to match the recipe
- b) Proceed if the final concentration looks correct
- c) Stop and route for pharmacist verification/correction before proceeding
- d) Discard everything immediately without documenting the discrepancy

65) A patient asks whether a Schedule II prescription can be partially filled today and the remainder next week. What should the technician do?

- a) Tell the patient partial filling is never allowed for C-II
- b) Partially fill and store the remainder in will-call without notes
- c) Change to a Schedule III alternative so it can be refilled
- d) Refer the question for pharmacist review of legality, documentation, and time limits

66) Spironolactone 25 mg daily is prescribed, but the technician selected spironolactone/HCTZ 25/25. What should the technician do?

- a) Dispense the combination to improve blood pressure control
- b) Correct to single-ingredient spironolactone as prescribed
- c) Dispense combination but cut quantity in half
- d) Add "take half tablet daily" and dispense combination

67) A patient asks which vaccine they should receive today based on their medical history. What should the technician do?

- a) Recommend the most common adult vaccine
- b) Ask another technician what they would recommend
- c) Tell the patient to decide based on online reviews
- d) Refer the patient for pharmacist clinical assessment

68) Sig: "i tab po qd." The technician entered "1 tablet BID." What is the correct next step?

- a) Leave it because most maintenance meds are BID
- b) Change quantity to double so the patient won't run out
- c) Correct the directions to once daily before processing further
- d) Replace directions with "use as directed" to avoid an entry error

69) Stock bottle is metoprolol succinate ER 50 mg. The Rx is metoprolol tartrate 50 mg BID. What should the technician do?

- a) Dispense succinate ER with BID directions
- b) Select metoprolol tartrate to match the prescription exactly
- c) Dispense succinate ER and change directions to daily
- d) Dispense either because both are metoprolol

70) A patient profile lists latex allergy. The ordered medication comes in a vial with a latex-containing stopper. What should the technician do?

- a) Ignore because latex isn't a drug ingredient
- b) Remove the stopper and proceed
- c) Add a "latex allergy" sticker and dispense
- d) Route for pharmacist review of product options and risk

71) DUR flags "duplicate antiplatelet therapy" when entering clopidogrel for a patient already taking aspirin 81 mg daily. What should the technician do?

- a) Override because aspirin is OTC
- b) Reduce clopidogrel dose to half tablet and continue
- c) Route the alert for pharmacist review of appropriateness
- d) Delete clopidogrel and continue aspirin only

72) Mupirocin ointment: apply TID x 7 days. Estimated use is 1 g per application. How many grams are needed?

- a) 28 g
- b) 21 g
- c) 14 g
- d) 7 g

73) Another pharmacy calls to transfer a Schedule IV prescription. The pharmacist is administering immunizations. What can the technician do?

- a) Complete the transfer with the other pharmacy's technician
- b) Ask for a fax of the original and treat it as transferred
- c) Cancel the prescription to prevent duplicates
- d) Take the request details and arrange pharmacist-to-pharmacist transfer communication when available

74) An eRx arrives for Cymbalta 60 mg daily. The profile lists "duloxetine—nausea." What

should the technician do?

- a) Delete the prescription because any listed reaction blocks dispensing
- b) Process without review because nausea is not a true allergy
- c) Route for pharmacist assessment of intolerance vs allergy and appropriateness
- d) Automatically change to venlafaxine XR 75 mg

75) The prescriber writes "brand medically necessary" for Keppra 500 mg. What should the technician do during product selection?

- a) Select any levetiracetam generic; bioequivalent is always acceptable
- b) Select levetiracetam ER to justify brand necessity
- c) Select brand Keppra NDC and process using correct DAW/brand coding
- d) Dispense generic but print "brand necessary" on the label

76) Prepare 120 mL of a 5% w/v dextrose solution. How many grams of dextrose are needed?

- a) 3 g
- b) 24 g
- c) 12 g
- d) 6 g

77) A technician sees a coworker scanning a stock bottle with an NDC mismatch versus the one on the screen, though the drug name appears similar. What should happen next?

- a) Override if strength matches
- b) Stop and resolve the mismatch by confirming the intended product and selecting the correct NDC
- c) Continue if tablet color matches expected
- d) Change the label to match the scanned bottle

78) Xarelto 20 mg rejects for prior authorization. The patient insists they must start tonight and asks for a few tablets. What should the technician do?

- a) Substitute warfarin and dispense
- b) Dispense a few tablets without billing and fix later
- c) Tell the patient there is nothing the pharmacy can do
- d) Route for pharmacist review of options (PA steps, coupons, cash options, emergency supply rules)

79) Law enforcement submits a written request for a patient profile but provides no subpoena/court order and asks for an immediate fax. What should the technician do?

- a) Release it if the officer shows a badge
- b) Fax it immediately because it's law enforcement
- c) Provide only controlled-substance records without a court order
- d) Escalate to pharmacist/management and require appropriate legal documentation before releasing PHI

80) Timolol 0.5% ophthalmic solution directions: "Instill 1 drop OS BID." What does OS mean?

- a) Left ear
- b) Left eye

- c) Both eyes
- d) Right eye

81) A prescription is written for 8 fl oz of cough syrup. Approximately how many mL is this?

- a) 180 mL
- b) 120 mL
- c) 300 mL
- d) 240 mL

82) DUR flags “max daily acetaminophen exceeded” for a patient receiving hydrocodone/APAP while also having OTC APAP documented in the profile. What should the technician do?

- a) Delete OTC APAP from the profile to clear the alert
- b) Override because OTC documentation is unreliable
- c) Route for pharmacist review of total APAP exposure and safety
- d) Reduce the hydrocodone/APAP quantity and proceed

83) Flonase 50 mcg/spray: “2 sprays in each nostril once daily.” Bottle contains 120 sprays. What is the days’ supply?

- a) 30 days
- b) 45 days
- c) 15 days
- d) 60 days

84) A patient asks the technician to change amlodipine 10 mg to amlodipine/benazepril because they “want fewer pills.” What should the technician do?

- a) Make the change if doses seem comparable
- b) Swap to the combination and label “as directed”
- c) Tell the patient to add OTC benazepril
- d) Explain that a medication change requires a new prescription and route for pharmacist follow-up

85) During final scan, the barcode indicates a different strength than the prescription label, and the NDC differs by one digit. What should the technician do?

- a) Override because it’s likely packaging
- b) Dispense and make a note for next time
- c) Stop and correct the mismatch before dispensing
- d) Change the label strength to match the scanned bottle

86) A patient requests an early refill of a Schedule V codeine-containing cough product due to “lost bottle.” What should the technician do?

- a) Refill early automatically because it is Schedule V
- b) Require a police report before any refill
- c) Transfer the prescription to another pharmacy to bypass limits
- d) Route for pharmacist review of appropriateness, documentation, and applicable legal/payer limits

87) A patient picking up a methylprednisolone dose pack asks how to take it. What should

the technician do?

- a) Provide the full taper schedule from memory
- b) Ensure the dosing card/package directions are included and refer counseling questions for pharmacist consultation
- c) Tell the patient to take all tablets on day 1
- d) Tell the patient to call the prescriber for instructions

88) The profile lists glipizide. A new Rx arrives for glyburide. What is the best next step?

- a) Process both; dual sulfonylureas improve control
- b) Delete glyburide as an obvious error
- c) Route for pharmacist review for therapeutic duplication/therapy change
- d) Automatically change glyburide to metformin

89) Valacyclovir 1 g: "Take 1 tablet PO TID x 7 days." What quantity should be dispensed?

- a) 14 tablets
- b) 30 tablets
- c) 28 tablets
- d) 21 tablets

90) A long-term care facility requests a patient's full medication profile by unsecured email "because it's urgent." What is the best action?

- a) Email it because urgency overrides privacy requirements
- b) Email only the medication names without directions
- c) Use an approved secure method (verified fax/secure portal) for transmission or route for an acceptable alternative
- d) Email it if the facility agrees to delete it after receipt

Answer Key

1

) b

2) c

3) d

4) a

5) d

6) b

7) c

8) c

9) a

10) b

11) a

12) c

13) c

14) d

15) b

16) a

17) b

18) c

19) b

20) c

21) b

22) d

23) d

24) c

25) c

26) d

27) d

28) a

29) c

30) d

31) c

32) c

33) d

34) b

35) d

36) c

37) a

38) d

39) a

40) c

41) c

42) c

43) d

44) c

45) c

46) b

47) a

48) d

49) b

50) a

51) c

52) d

53) d

54) b

55) c

56) c

57) c

58) b

59) c

60) d

61) c

62) b

63) c

64) c

65) d

66) b

67) d

68) c

69) b

70) d

71) c

72) b

73) d

74) c

75) c

76) d

77) b

78) d

79) d

80) b

81) d

82) c

83) a

84) d

85) c

86) d

87) b

88) c

89) d

90) c

PTCB Sim Test 17

1) A new eRx is received: "levETIRAcetam 500 mg tab: take 1 tab PO BID." During product selection, the technician selects levETIRAcetam 750 mg because the strengths are adjacent in the list. What is the best next step before labels print?

- a) Keep 750 mg and reduce directions to once daily
- b) Change the quantity so the total daily mg matches 1000 mg
- c) Select levETIRAcetam 500 mg with the correct dosage form and confirm the NDC
- d) Continue and rely on counseling to catch the error

2) A patient's spouse calls asking, "What did you fill for my husband today?" The caller is not listed as authorized and cannot verify the patient's DOB. What should the technician do?

- a) Provide only the medication names without directions
- b) Ask for the patient's address and then provide the information
- c) Provide the total price only, not the medication information
- d) Decline to disclose PHI and offer to speak when the patient is available or properly verified

3) A prescription reads: "cephalexin 500 mg caps: 1 cap PO QID x 10 days." How many capsules should be dispensed?

- a) 28
- b) 40
- c) 30
- d) 20

4) While putting away stock, a technician finds a bottle of tablets with the label smeared and no readable NDC/lot. The tablets look like a common generic. What is the safest action?

- a) Place it in a separate bin and use it only after comparing to an image
- b) Quarantine it and remove it from dispensing inventory until it can be positively identified
- c) Create a new label using another bottle's NDC and keep it in stock
- d) Use it only for cash patients to avoid third-party audits

5) A prescription is for "Humalog KwikPen U-100." The technician pulls "Humulin R U-100 vial" because both are insulin. What should the technician do?

- a) Replace with the prescribed Humalog KwikPen and confirm NDC and dosage form
- b) Dispense the vial and include pen needles
- c) Dispense the vial and instruct the patient to draw up the correct dose
- d) Substitute Novolog FlexPen because it is the same class

6) A customer purchases pseudoephedrine 30 mg tablets, quantity 96. Under CMEA limits, which is the most relevant check to determine if the sale is permissible?

- a) Maximum 6 g per day and 18 g per 30 days
- b) Maximum 3.6 g per day and 9 g per 30 days
- c) Maximum 9 g per day and 3.6 g per 30 days
- d) Maximum 10 g per week and 30 g per year

7) A delivery of Schedule II medications arrives. The invoice lists the correct drugs and

quantities, but the pharmacy's DEA number on the invoice is incorrect. What should the technician do?

- a) File the invoice; DEA numbers are not required on invoices
- b) Correct the DEA number in ink and initial the correction
- c) Set the invoice aside and ensure the discrepancy is addressed before routine filing
- d) Discard the invoice after the medications are stocked

8) A prescription reads: "azithromycin 200 mg/5 mL: 12.5 mL PO day 1, then 6.25 mL PO daily days 2–5." What total volume should be dispensed?

- a) 37.5 mL
- b) 40 mL
- c) 45 mL
- d) 42.5 mL

9) During entry, the patient profile shows "allergy: ACE inhibitors—angioedema." A new prescription is for lisinopril 10 mg daily. What is the best technician action?

- a) Enter it and add a warning label for swelling
- b) Enter it and remove the allergy since it may be old
- c) Ensure the allergy conflict is addressed through the pharmacist's clinical check before dispensing
- d) Change the medication to losartan and continue processing

10) A label prints for "hydroxyzine 25 mg." The technician accidentally counts hydralazine 25 mg because the bottles are stored nearby. Which workflow control best prevents this from reaching the patient?

- a) Counting twice by the same technician
- b) Barcode scanning that matches the stock bottle NDC to the prescription label/NDC
- c) Reading the first three letters of the drug name on the shelf tag
- d) Placing both drugs in the same bin to reduce searching

11) A prescription is "Lantus SoloStar: inject 22 units SQ daily." One box contains 5 pens, each 3 mL, U-100. What days' supply should be billed?

- a) 68 days
- b) 75 days
- c) 60 days
- d) 45 days

12) A patient presents a paper prescription for a Schedule II medication dated 5 months ago. What should the technician do?

- a) Fill it if it has not been filled before
- b) Refuse automatically because Schedule II expires in 30 days federally
- c) Edit the written date to today if the prescriber confirms by phone
- d) Route it for pharmacist review because validity depends on required elements and applicable state law

13) Which product is commonly stored refrigerated prior to dispensing in many pharmacies?

- a) Latanoprost ophthalmic solution (unopened)
- b) Clopidogrel tablets

- c) Omeprazole capsules
- d) Nitroglycerin SL tablets

14) While processing a refill for warfarin 5 mg daily, the profile shows a recent fill for apixaban 5 mg BID at another location in the same chain. What is the most appropriate technician step?

- a) Continue; both are anticoagulants and can be used together
- b) Discontinue apixaban on the patient profile to prevent confusion
- c) Make sure the pharmacist sees the potential duplicate anticoagulant therapy concern
- d) Ask the patient which one they prefer and process that one

15) A prescription reads: "prednisone 10 mg: 4 tabs daily x 3 days, then 3 tabs daily x 3 days, then 2 tabs daily x 3 days, then 1 tab daily x 3 days." What quantity should be dispensed?

- a) 30
- b) 24
- c) 36
- d) 40

16) A patient asks for a copy of the pharmacy's Notice of Privacy Practices. What should the technician do?

- a) Provide it or direct the patient to where it is available in the pharmacy
- b) Provide it only after the patient signs a HIPAA release
- c) Provide it only at the first prescription pickup
- d) Tell the patient it must be requested from their insurer

17) An IV order: "Heparin 25,000 units in 250 mL D5W. Infuse 18 units/kg/hr. Weight 70 kg." What pump rate should be entered in mL/hr?

- a) 12.6 mL/hr
- b) 11.8 mL/hr
- c) 10.1 mL/hr
- d) 14.4 mL/hr

18) The prescription is written for "Toprol-XL 50 mg daily." Which product should be selected?

- a) Atenolol 50 mg
- b) Metoprolol tartrate 50 mg
- c) Metoprolol succinate ER 50 mg
- d) Propranolol ER 60 mg

19) At pickup, a patient asks, "Is this safe to take with ibuprofen?" What is the best technician response?

- a) Suggest taking ibuprofen with food
- b) Recommend spacing doses by 2 hours
- c) Ask the ibuprofen dose and provide a maximum daily dose
- d) Offer to get the pharmacist to discuss interactions and safety

20) A claim rejects "Refill too soon." The patient states the medication was lost. Which is the

most appropriate next step for the technician?

- a) Use a lost-medication override code immediately to speed pickup
- b) Gather details and route for pharmacist/payer-guided decision and documentation
- c) Tell the patient no early refills are allowed for any medication
- d) Transfer the prescription to a different pharmacy to bypass the reject

21) A prescription: "albuterol HFA: inhale 2 puffs q4h PRN." The inhaler contains 200 actuations. Using maximum directions, what days' supply should be billed?

- a) 14
- b) 17
- c) 25
- d) 10

22) A technician finds a will-call bag labeled for Patient A, but inside is a vial labeled for Patient B. What is the best immediate action?

- a) Switch the vials into the correct bags and continue
- b) Place both vials into one bag and leave a note for counseling
- c) Separate and reprocess through the correct verification/matching steps before any pickup
- d) Leave it until end-of-day cleanup to avoid delaying workflow

23) A prescription is entered for "metformin ER 500 mg: take 2 tablets PO daily with evening meal." The technician selected metformin IR 500 mg. What should be done?

- a) Keep IR and change directions to 1 tablet BID
- b) Switch to ER as written and confirm correct product selection before proceeding
- c) Keep IR and reduce dose to 250 mg to reduce GI effects
- d) Dispense both ER and IR and let the patient decide

24) A clinic faxes a prescription for alprazolam with drug, directions, quantity, and prescriber info, but it has no prescriber signature. What is the correct handling?

- a) Fill it because it came from the clinic's fax number
- b) Hold it as invalid as received and route for pharmacist follow-up
- c) Convert it to a verbal order and enter it immediately
- d) Ask the patient to sign the faxed prescription at pickup

25) A manufacturer recall notice is received for a specific lot of atorvastatin tablets. What should the technician do first?

- a) Pull the affected lot from inventory and quarantine it per recall procedure
- b) Return all atorvastatin in stock to the wholesaler
- c) Continue dispensing until replacement stock arrives
- d) Call every patient who has ever taken atorvastatin

26) A prescription reads: "cefdinir 250 mg/5 mL: take 6 mL PO BID x 10 days." How many mL are needed?

- a) 100 mL
- b) 110 mL
- c) 120 mL
- d) 90 mL

27) A prescription arrives for "Eliquis 5 mg: take 1 tablet PO BID." Which generic should be selected?

- a) Apixaban
- b) Rivaroxaban
- c) Dabigatran
- d) Edoxaban

28) A perpetual inventory shows 250 tablets of a controlled medication, but the physical count is 246. What should the technician do?

- a) Adjust the count to 246 and continue work
- b) Recount and initiate the discrepancy process used for suspected loss/diversion
- c) Ignore it because small variances are expected
- d) Add 4 tablets from another bottle without documentation

29) A patient requests to transfer an unfilled Adderall XR (Schedule II) prescription from another pharmacy that is out of stock. What is the correct response?

- a) Transfer it like a non-controlled prescription if both pharmacies are in-state
- b) Accept a fax transfer from the other pharmacy for documentation
- c) Accept a verbal transfer between pharmacists
- d) The prescriber must issue a new prescription to the receiving pharmacy

30) A prescription is "diclofenac 1% gel: apply 4 g to knee QID." Tube size is 100 g. What is the days' supply if used exactly as directed?

- a) 5 days
- b) 6 days
- c) 7 days
- d) 10 days

31) The hard copy directions are "Take 1 tablet PO BID," but the printed label reads "Take 1 tablet PO daily." What should the technician do before the prescription proceeds?

- a) Correct the SIG to match the prescription and reprint the label
- b) Handwrite "BID" on the label and initial it
- c) Keep the label and adjust the days' supply only
- d) Dispense as daily to reduce side effects

32) A prescriber writes "start sertraline 50 mg daily." Which class is sertraline?

- a) SNRI
- b) SSRI
- c) Tricyclic antidepressant
- d) MAOI

33) A patient asks the technician to transfer nitroglycerin SL tablets into a standard amber vial because the original bottle is hard to open. What is the best response?

- a) Repackage into an amber vial and add a "protect from light" label
- b) Repackage only if the patient signs a waiver
- c) Do not repackage into a standard vial; maintain appropriate container requirements for stability
- d) Crush tablets and package as unit-dose for easier use

34) A prescription is "levothyroxine 0.05 mg daily." Which strength should be selected?

- a) 5 mcg
- b) 500 mcg
- c) 0.5 mcg
- d) 50 mcg

35) A prescription is "nystatin oral suspension 100,000 units/mL: swish and swallow 5 mL QID x 7 days." What quantity should be dispensed?

- a) 200 mL
- b) 140 mL
- c) 120 mL
- d) 150 mL

36) A patient requests a printout of their prescription history. What is the best initial technician step before printing?

- a) Print immediately; it's their information
- b) Verify identity and confirm the specific timeframe/details requested
- c) Provide only drug names without dates
- d) Refuse because printouts are not allowed under HIPAA

37) A prescriber sends three separate e-prescriptions for a Schedule II stimulant with "do not fill before" dates for the next two months. What is the key technician task during entry?

- a) Combine them into one prescription for the total quantity
- b) Enter each prescription separately and apply the correct earliest fill date for each
- c) Hold them and only enter into the system on each fill date
- d) Convert them to Schedule III since they are monthly supplies

38) A sterile compounding refrigerator temperature log shows the unit was out of range for 4 hours, and several batches were stored there during that time. What should be done with those batches?

- a) Release them because temperature does not affect sterility
- b) Extend the beyond-use dates to compensate for the deviation
- c) Quarantine/hold and route for quality review per the compounding program
- d) Dispense them but add "refrigerate" labels

39) A prescription is "Crestor 20 mg daily" with DAW 0. If substitution is permitted, what should be dispensed?

- a) Brand Crestor 20 mg
- b) Atorvastatin 40 mg as a therapeutic equivalent
- c) Generic rosuvastatin 20 mg
- d) Any statin at an equivalent dose

40) A prescriber's office calls asking for the last fill date of a patient's clonazepam. What should the technician do?

- a) Provide it immediately since it is the prescriber
- b) Request the patient's SSN, then provide it
- c) Refuse because prescribers cannot receive fill information

d) Confirm appropriate verification/authorization and route through the pharmacist/workflow for disclosure and documentation

41) A patient profile lists “sulfa—hives.” A new prescription arrives for sulfamethoxazole/trimethoprim DS. What is the best technician action?

- a) Process it; allergies can be discussed at pickup
- b) Delete the allergy since it is nonspecific
- c) Ensure the allergy conflict is reviewed before dispensing proceeds
- d) Substitute nitrofurantoin and continue processing

42) A hospital order reads: “dopamine 400 mg in 250 mL D5W. Infuse 5 mcg/kg/min. Weight 80 kg.” What is the infusion rate in mL/hr?

- a) 12 mL/hr
- b) 15 mL/hr
- c) 18 mL/hr
- d) 9 mL/hr

43) A technician is restocking look-alike/sound-alike items: clonidine and clonazepam. Which action best reduces selection error?

- a) Store them next to each other so staff learns the difference
- b) Separate storage locations and apply Tall Man/alert labeling practices used at the site
- c) Place both in the same bin to reduce walking time
- d) Remove one from inventory to eliminate confusion

44) Which medication is a proton pump inhibitor?

- a) Dicyclomine
- b) Famotidine
- c) Sucralfate
- d) Omeprazole

45) A prescription reads: “fluconazole 150 mg: take 1 tablet today and repeat in 72 hours.” What quantity should be dispensed?

- a) 1
- b) 2
- c) 3
- d) 4

46) A new prescription arrives: “methotrexate 2.5 mg: take 6 tablets daily.” The patient’s previous directions on profile show “6 tablets once weekly.” What should the technician do?

- a) Enter as written; dosing is the prescriber’s responsibility
- b) Change to weekly because that is typical
- c) Fill it and counsel the patient to follow the old directions
- d) Pause processing and route for pharmacist review due to high-risk dosing concern

47) A third-party claim rejects: “NDC not covered—use preferred product.” What is the best technician next step?

- a) Change to a preferred NDC of the same drug/strength/dosage form and reprocess
- b) Change to a different medication in the same class

- c) Bill cash without telling the patient
- d) Change the prescriber field to match the plan

48) Which is a high-alert medication commonly targeted for extra safeguards in hospitals?

- a) Docusate
- b) Loratadine
- c) Potassium chloride concentrated injection
- d) Amoxicillin

49) A prescription is written for "Advair Diskus." Which combination is correct?

- a) Budesonide/formoterol
- b) Fluticasone/salmeterol
- c) Fluticasone/vilanterol
- d) Tiotropium/olodaterol

50) While counting, a technician drops several tablets onto the floor. What should be done?

- a) Return them to stock if they appear clean
- b) Wipe with alcohol and dispense
- c) Place them aside for later use as "waste" doses
- d) Discard per procedure and recount using clean stock

51) Under federal law, alprazolam is which schedule?

- a) II
- b) III
- c) IV
- d) V

52) A prescription is "insulin aspart: inject 8 units SQ TID with meals." One box contains 5 pens, each 3 mL, U-100. What days' supply should be billed?

- a) 50 days
- b) 56 days
- c) 62 days
- d) 46 days

53) A wholesaler requests DSCSA transaction information for a specific lot and NDC dispensed last month due to an investigation. What should the technician do?

- a) Provide the pharmacy's DSCSA transaction documentation using the established retrieval process
- b) Provide the patient's entire medication profile to prove dispensing
- c) Ignore it; DSCSA applies only to controlled substances
- d) Provide only the invoice total without identifiers

54) At pickup, the patient asks, "Which one is for blood pressure?" The bag contains amlodipine and metformin. What should the technician do?

- a) Tell them amlodipine is for blood pressure and discuss side effects
- b) Tell them metformin is for blood pressure because it improves circulation
- c) Read the indication from the stock bottle and answer
- d) Offer to get the pharmacist for medication counseling

55) A prescription reads: "tramadol 50 mg: take 1 tablet q6h PRN pain, qty 24." Using maximum directions, what days' supply should be billed?

- a) 3
- b) 4
- c) 5
- d) 6

56) A prescription is "diltiazem ER 240 mg daily." The technician entered diltiazem IR 120 mg BID because the total daily dose matches. Once discovered, what should be done?

- a) Leave it since total daily dose is the same
- b) Correct to the ER formulation as prescribed before it proceeds
- c) Keep IR and add an auxiliary label "extended-release"
- d) Dispense IR and counsel the patient to take once daily

57) Under CMEA, which element is required at the point of sale for pseudoephedrine products?

- a) Purchaser's name and verification using a government-issued photo ID
- b) Prescriber's DEA number
- c) Purchaser's diagnosis code
- d) Purchaser's credit card number

58) Which medication is an ARB?

- a) Losartan
- b) Lisinopril
- c) Metoprolol
- d) Amlodipine

59) A new eRx arrives and the date of birth on the prescription does not match the patient profile. What is the best first step?

- a) Ignore it if the address matches
- b) Change the prescription DOB to match the profile
- c) Verify patient identity and correct demographics before continuing processing
- d) Fill it and fix the profile at pickup

60) Directions: "Test blood glucose up to 4 times daily." Quantity is 200 test strips. What days' supply should be billed?

- a) 30
- b) 60
- c) 45
- d) 50

61) A patient requests that a Schedule III prescription with refills be transferred to another pharmacy. What is the correct approach?

- a) Transfer is never allowed for controlled substances
- b) The transfer must be handled according to controlled-substance transfer rules, typically pharmacist-to-pharmacist
- c) A technician may transfer it if both pharmacies are in the same chain

d) Transfer is allowed only after all refills are used

62) The prescription is "Zoloft 50 mg daily." Which generic corresponds?

- a) Fluoxetine
- b) Venlafaxine
- c) Sertraline
- d) Citalopram

63) A patient returns an opened bottle of gabapentin requesting a refund and asks if the pharmacy will reuse the medication. What is the most appropriate technician response?

- a) Accept and return to stock if tablets look intact
- b) Accept and pour into a stock bottle, then document quantity
- c) Refuse all returns, even for disposal
- d) Do not return it to stock; handle per return/disposal procedures and offer take-back options if available

64) An IV label reads: "vancomycin 1 g in 250 mL NS; infuse over 2 hours." What rate is this in mL/hr?

- a) 100 mL/hr
- b) 125 mL/hr
- c) 150 mL/hr
- d) 200 mL/hr

65) Under federal law, which record type must generally be retained for at least 2 years?

- a) Prescription and refill records
- b) Technician timecards
- c) Patient counseling documentation for every prescription
- d) OTC planograms

66) In a hospital unit-dose cart, a blister pack shows a lot number and expiration but the medication name is missing. What should the technician do?

- a) Use it if the barcode scans and the dose matches
- b) Handwrite the medication name on the blister and continue using it
- c) Remove it from use and replace it with properly labeled product
- d) Place it in the controlled-substance drawer for extra security

67) A prescription reads: "enoxaparin 40 mg/0.4 mL: inject 0.4 mL SQ daily x 14 days." Syringes are packaged in boxes of 10. What quantity should be dispensed?

- a) 10
- b) 14
- c) 20
- d) 28

68) Which medication is a sulfonylurea?

- a) Sitagliptin
- b) Empagliflozin
- c) Glipizide
- d) Metformin

69) A hard copy for a Schedule II medication is missing the patient address. What should happen next?

- a) Add the address from the patient profile and process
- b) Fill it and note "address on file" in the system
- c) Hold for pharmacist review because the prescription may be missing required elements
- d) Call the prescriber's office staff and take the address verbally

70) A prescription reads: "amoxicillin/clavulanate 875/125 mg: 1 tab PO BID x 10 days." What quantity should be dispensed?

- a) 14
- b) 18
- c) 28
- d) 20

71) A patient requests "my cholesterol med—Lipitor." Which generic is Lipitor?

- a) Simvastatin
- b) Atorvastatin
- c) Rosuvastatin
- d) Pravastatin

72) The refrigerator temperature log shows 10°C for 8 hours overnight where unopened insulin glargine was stored. What should the technician do first?

- a) Put it back and use it first to prevent waste
- b) Freeze it briefly to re-stabilize
- c) Quarantine the affected product for stability review and disposition
- d) Relabel it with a shorter expiration date and dispense

73) A prescription is "Synthroid 125 mcg daily" with DAW 1. What should be dispensed?

- a) Generic levothyroxine 125 mcg
- b) Any levothyroxine product as long as it is 125 mcg
- c) Liothyronine 25 mcg
- d) Brand Synthroid 125 mcg

74) A claim message reads: "Coupon requires primary paid." What should the technician do?

- a) Bill the coupon as primary with an override
- b) Bill cash and manually reduce the price
- c) Bill the primary insurance first, then submit the coupon as secondary if allowed
- d) Call the manufacturer for a universal authorization code

75) A prescription is for "allopurinol 300 mg daily." What is the primary use?

- a) Acute relief of gout pain
- b) Gout prophylaxis by lowering uric acid
- c) Treatment of rheumatoid arthritis flares
- d) Prevention of kidney stones by increasing calcium excretion

76) Directions read: "Take 2 tablets PO TID." Quantity dispensed is 180 tablets. What is the days' supply?

- a) 45
- b) 20
- c) 30
- d) 15

77) A prescription for triamcinolone 0.1% cream states: "apply thin layer to affected area BID for 7 days." Which SIG entry approach is best?

- a) Enter "apply BID" only to keep it short
- b) Enter "use as directed" to reduce liability
- c) Include the key elements as written (site if provided, frequency, and duration when possible)
- d) Omit duration because creams do not require it

78) A patient has isotretinoin on their profile and the system indicates REMS requirements are incomplete for today's fill. What should the technician do?

- a) Dispense anyway; REMS is the prescriber's responsibility
- b) Hold the fill until required REMS steps are completed before dispensing
- c) Substitute topical tretinoin and dispense instead
- d) Change to a non-REMS acne product automatically

79) Which medication is a benzodiazepine?

- a) Buspirone
- b) Hydroxyzine
- c) Propranolol
- d) Lorazepam

80) Clopidogrel 75 mg daily is best described as which class?

- a) Statin
- b) Proton pump inhibitor
- c) Antiplatelet (P2Y12 inhibitor)
- d) ACE inhibitor

81) A patient requests a claim reversal stating they did not pick up the medication. The will-call log shows a signature from yesterday. What is the best technician action?

- a) Reverse immediately; signatures are unreliable
- b) Review pickup documentation and route for investigation per workflow before reversing
- c) Rebill under a different patient profile
- d) Delete the signature entry and reverse the claim

82) A prescription is "acetaminophen 160 mg/5 mL: take 7.5 mL PO q6h PRN." Quantity dispensed is 240 mL. Using maximum frequency, what days' supply should be billed?

- a) 8
- b) 6
- c) 10
- d) 7

83) Which medication is used to treat hypothyroidism?

- a) Methimazole

- b) Levothyroxine
- c) Propranolol
- d) Prednisone

84) A retail pharmacy completes a partial fill of a Schedule II prescription due to insufficient stock. Under federal rules for typical outpatient partial fills, the remainder generally must be dispensed within:

- a) 72 hours
- b) 7 days
- c) 30 days
- d) 24 hours

85) A prescription reads: "Augmentin ES-600 600 mg/5 mL: 7.5 mL PO BID x 10 days." What total volume should be dispensed?

- a) 150 mL
- b) 180 mL
- c) 120 mL
- d) 160 mL

86) Which medication is a calcium channel blocker commonly used for hypertension?

- a) Losartan
- b) Metoprolol
- c) Hydrochlorothiazide
- d) Amlodipine

87) During verification prep, a technician notices refills were entered as 6, but the prescription shows "0 refills." What should be done?

- a) Leave it and fix it at the next refill request
- b) Correct the refill entry to match the prescription and reprint any affected labels/documentation
- c) Cross out the refills on the label and initial it
- d) Keep the refills but add a note "no refills allowed"

88) Which medication is a direct-acting oral anticoagulant (DOAC)?

- a) Clopidogrel
- b) Warfarin
- c) Apixaban
- d) Aspirin

89) Clonazepam is which federal controlled substance schedule?

- a) II
- b) III
- c) V
- d) IV

90) A prescription is entered for sertraline 100 mg daily, but the technician pulled sertraline 50 mg tablets. What is the best next step?

- a) Dispense 50 mg and change directions to 2 tablets daily

- b) Dispense 50 mg and add "take as directed"
- c) Reduce quantity to keep the same days' supply and dispense
- d) Pull the correct strength (or correct the entry if it was entered incorrectly) before continuing

Answer Key

1)		
c	31) a	61) b
2) d	32) b	62) c
3) b	33) c	63) d
4) b	34) d	64) b
5) a	35) b	65) a
6) b	36) b	66) c
7) c	37) b	67) b
8) d	38) c	68) c
9) c	39) c	69) c
10) b	40) d	70) d
11) a	41) c	71) b
12) d	42) a	72) c
13) a	43) b	73) d
14) c	44) d	74) c
15) a	45) b	75) b
16) a	46) d	76) c
17) a	47) a	77) c
18) c	48) c	78) b
19) d	49) b	79) d
20) b	50) d	80) c
21) b	51) c	81) b
22) c	52) b	82) a
23) b	53) a	83) b
24) b	54) d	84) a
25) a	55) b	85) a
26) c	56) b	86) d
27) a	57) a	87) b
28) b	58) a	88) c
29) d	59) c	89) d
30) b	60) d	90) d

PTCB Sim Test 18

1) While entering an eRx for levothyroxine, the system auto-populates "levothyroxine 125 mcg" from the last patient searched, but the new eRx clearly states 112 mcg. The label has not printed yet. What should you do next?

- a) Keep 125 mcg and add a note for the pharmacist
- b) Change to 112 mcg, then re-verify patient, drug, and directions before proceeding
- c) Print the label and correct it at pickup if needed
- d) Delete the eRx and request a new one

2) A patient asks for the generic of Crestor at the register. Which is correct?

- a) Atorvastatin
- b) Pravastatin
- c) Rosuvastatin
- d) Simvastatin

3) A C-II eRx arrives for oxycodone. The prescriber name is present, but the DEA number field is blank in the pharmacy system display. What is the most appropriate next step?

- a) Continue processing; DEA number is not required on an eRx
- b) Add a DEA number from a previous hardcopy and proceed
- c) Convert it to a verbal order and document the call
- d) Hold the prescription for pharmacist review before dispensing steps continue

4) Rx: amoxicillin/clavulanate 600 mg/42.9 mg per 5 mL; take 7 mL PO BID x 10 days. How many mL should be dispensed?

- a) 120 mL
- b) 140 mL
- c) 150 mL
- d) 200 mL

5) A patient requests an early refill for lorazepam saying the tablets were "lost." The last fill was 9 days ago for a 30-day supply. What should a technician do?

- a) Process the refill and let the claim reject if too soon
- b) Tell the patient it is illegal to replace lost controlled substances
- c) Create the request and route it for pharmacist decision before processing
- d) Transfer the prescription to a different pharmacy immediately

6) Which brand/generic pair is correct?

- a) Xarelto — apixaban
- b) Eliquis — apixaban
- c) Pradaxa — rivaroxaban
- d) Brilinta — prasugrel

7) During product selection, the barcode scan reads "clonidine 0.1 mg," but the stock bottle label says "clozapine 100 mg." What is the best immediate action?

- a) Use the bottle label and override the scan
- b) Select it anyway if the prescriber wrote "clo—"

- c) Stop and resolve the mismatch before continuing the fill
- d) Change the entered drug to match the scanned bottle

8) You are counting tablets and notice two different labels on the counter for two patients with the same last name. What is the safest next step?

- a) Finish counting both and separate later
- b) Stop, clear the workstation, and match each label to the correct prescription before continuing
- c) Put both vials in will-call and let pickup sort it out
- d) Reprint both labels to avoid confusion

9) A customer purchases pseudoephedrine behind the counter. Which federal requirement applies?

- a) The prescriber's NPI must be documented
- b) The sale must be logged and quantity limits enforced
- c) The customer must sign a HIPAA consent form
- d) The customer must present a prescription

10) Rx: prednisone 10 mg tablets: take 4 tablets daily x 3 days, then 2 tablets daily x 3 days, then 1 tablet daily x 3 days. What quantity should be dispensed?

- a) 18
- b) 21
- c) 24
- d) 27

11) A new eRx is for metoprolol succinate ER 25 mg daily. The patient profile shows an active metoprolol tartrate 25 mg BID started two days ago. What should the technician do?

- a) Enter the new prescription and inactivate the old one automatically
- b) Change succinate to tartrate because strengths match
- c) Enter as written and flag for pharmacist review due to formulation/therapy overlap risk
- d) Place the new eRx on hold and ignore it

12) Which medication is a proton pump inhibitor?

- a) Famotidine
- b) Sucralfate
- c) Omeprazole
- d) Dicyclomine

13) A patient asks for the generic of Norvasc. Which is correct?

- a) Amlodipine
- b) Losartan
- c) Lisinopril
- d) Hydrochlorothiazide

14) A Schedule III prescription was written 7 months ago and still has refills remaining. Under federal law, can it be refilled?

- a) Yes, until all refills are used
- b) Yes, for 1 year from the date written

- c) Yes, but only once more
- d) No, it is not refillable after 6 months

15) Rx directions: "Take 1 tablet by mouth every 6 hours as needed for pain." Quantity #24. For insurance, what days' supply should be billed using maximum daily use?

- a) 4
- b) 6
- c) 8
- d) 12

16) The refrigerator temperature log shows 50°F at opening. Several insulin and antibiotic suspensions are stored inside. What should happen next before any are dispensed?

- a) Continue using them; 50°F is close to normal
- b) Move them to room temperature for 30 minutes and continue
- c) Quarantine the affected stock and alert the pharmacist for stability assessment and documentation
- d) Put the items in the freezer to bring the temperature down quickly

17) Which brand/generic pair is correct?

- a) Ozempic — dulaglutide
- b) Trulicity — semaglutide
- c) Jardiance — empagliflozin
- d) Januvia — canagliflozin

18) A caller asks for a medication list for their spouse. The spouse is not listed as authorized and is not on the line. What should the technician do?

- a) Provide only medication names but not directions
- b) Provide the list if the caller confirms date of birth and address
- c) Provide only prescriber names and pharmacy phone number
- d) Decline to disclose and offer options to add authorization or have the patient call

19) A patient has an active warfarin prescription. A new eRx arrives for apixaban 5 mg BID from another prescriber. What is the best technician action?

- a) Fill both because they treat clotting
- b) Enter the new prescription and automatically discontinue warfarin
- c) Enter and route for pharmacist review before dispensing due to therapeutic duplication risk
- d) Substitute warfarin for apixaban to reduce cost

20) Rx: albuterol HFA inhaler (200 actuations): inhale 2 puffs QID. What days' supply is one inhaler?

- a) 12
- b) 25
- c) 50
- d) 100

21) A hardcopy prescription for Synthroid 75 mcg is marked DAW 1. Which product should be selected?

- a) Any AB-rated levothyroxine 75 mcg
- b) Brand Synthroid 75 mcg (no substitution)
- c) Liothyronine 25 mcg
- d) Any thyroid medication with 75 mcg strength

22) Which is the correct generic for Lasix?

- a) Torsemide
- b) Bumetanide
- c) Spironolactone
- d) Furosemide

23) In a hospital, an order reads: heparin 25,000 units in 250 mL. Infuse at 1,600 units/hour. What is the pump rate in mL/hour?

- a) 14 mL/hr
- b) 16 mL/hr
- c) 18 mL/hr
- d) 20 mL/hr

24) A prescriber's office calls in a new prescription while the pharmacist is counseling. The prescriber asks the technician to take the verbal order. What should the technician do?

- a) Take the order and have the pharmacist sign later
- b) Ask the caller to fax it instead because verbals are never allowed
- c) Transfer the call to the pharmacist to receive the verbal prescription
- d) Take only the patient name and medication, then fill in the rest from the profile

25) During will-call checks, you find a vial with tablets and an auxiliary label but no patient label. What should you do?

- a) Put it back; the pharmacist will handle it at pickup
- b) Add the patient label for the last prescription you filled
- c) Remove it from will-call and send it back through reprocessing/correction workflow
- d) Write the patient name on tape and attach it to the vial

26) Which medication is a benzodiazepine?

- a) Buspirone
- b) Hydroxyzine
- c) Alprazolam
- d) Bupropion

27) A manufacturer recall notice lists a specific lot of metformin ER. You locate that lot on the shelf. What is the appropriate technician action?

- a) Keep dispensing until replacement stock arrives
- b) Segregate/quarantine the lot and follow the recall process for disposition
- c) Re-label with a shorter beyond-use date and continue dispensing
- d) Move it behind other stock to reduce chance of selection

28) Rx: valacyclovir 1 g: take 1 tablet TID x 7 days. What quantity should be dispensed?

- a) 14
- b) 21

- c) 28
- d) 30

29) The patient profile lists "anaphylaxis to penicillin." A new eRx arrives for cephalexin. What should the technician do?

- a) Fill normally; cephalexin is not a penicillin
- b) Delete the allergy because it is not a current medication
- c) Substitute azithromycin without prescriber approval
- d) Flag for pharmacist review before dispensing due to allergy history

30) Under the Combat Methamphetamine Epidemic Act, what is the maximum pseudoephedrine base allowed per day for a customer?

- a) 1.8 g
- b) 2.4 g
- c) 3.6 g
- d) 9.0 g

31) An eRx arrives for omeprazole 20 mg with directions "UAD" only. What is the best next step?

- a) Enter "take 1 daily" because it is typical
- b) Enter "use as directed" and proceed to fill
- c) Route for clarification so complete directions can be obtained/confirmed
- d) Convert it to an OTC purchase and bypass prescription processing

32) Which medication is an ACE inhibitor?

- a) Losartan
- b) Metoprolol
- c) Lisinopril
- d) Amlodipine

33) Rx: insulin lispro U-100 vial. Inject 10 units before meals TID. One 10 mL vial is dispensed. What days' supply should be billed?

- a) 28
- b) 33
- c) 30
- d) 24

34) Which abbreviation means "both eyes"?

- a) OD
- b) OU
- c) OS
- d) AD

35) A perpetual inventory for a controlled substance shows 500 tablets on hand, but a physical count shows 497. What should be done first?

- a) Adjust the count to 497 and move on
- b) Document "breakage" to balance the count
- c) Wait for the next shipment and recheck

d) Recount and review recent transactions per process, then report discrepancy as required

36) Which brand/generic pair is correct?

- a) Cymbalta — duloxetine
- b) Seroquel — sertraline
- c) Neurontin — pregabalin
- d) Lyrica — gabapentin

37) Rx: “apply 1 gram topically BID.” You dispense a 45-gram tube. What days’ supply should be billed?

- a) 21
- b) 30
- c) 45
- d) 90

38) A patient brings loose tablets and asks you to identify them. What is within technician scope?

- a) Identify based on color/shape and confirm for the patient
- b) Use an approved identifier tool using the imprint code and refer to the pharmacist before confirming
- c) Tell the patient to take one and see if it works
- d) Refuse to discuss pills not filled at your pharmacy

39) A pharmacy orders a Schedule II medication from the primary wholesaler. Which is required?

- a) DEA Form 106
- b) DEA Form 41
- c) DEA Form 222 or CSOS
- d) DEA Form 363

40) A Schedule II prescription is partially filled due to insufficient stock. What documentation is most critical to allow completion later?

- a) Patient’s allergy list
- b) Prescriber specialty
- c) Date and quantity dispensed and quantity remaining owed
- d) Technician initials only

41) Which is the correct generic for Plavix?

- a) Prasugrel
- b) Clopidogrel
- c) Ticagrelor
- d) Dipyridamole

42) Rx: fentanyl 25 mcg/hr patches: apply 1 patch every 72 hours. A box contains 5 patches. What is the days’ supply for one box?

- a) 12
- b) 15
- c) 18

d) 30

43) Two eRxs arrive on the same day: levothyroxine 75 mcg daily and levothyroxine 88 mcg daily, same patient. What should the technician do?

- a) Fill both and let the patient decide
- b) Fill the higher strength because it is likely an increase
- c) Enter both and route for pharmacist review to determine intended therapy
- d) Delete the older one automatically

44) A patient asks what "BID" means on the label. What is an appropriate response?

- a) Twice daily
- b) Every 6 hours
- c) Once daily
- d) At bedtime

45) Under HIPAA, what is an appropriate identity check before releasing a printed medication history to a patient in person?

- a) Ask for last name only
- b) Ask for insurance card only
- c) Ask for prescriber name only
- d) Use at least two identifiers (e.g., full name and date of birth) per workflow

46) Which medication is a statin?

- a) Ropinirole
- b) Ranitidine
- c) Rosuvastatin
- d) Rosiglitazone

47) A prescription is written for "Bactrim DS." Which product is correct?

- a) Trimethoprim/sulfamethoxazole 160 mg/800 mg
- b) Amoxicillin/clavulanate 875 mg/125 mg
- c) Nitrofurantoin 100 mg
- d) Doxycycline 100 mg

48) In sterile compounding, a technician notices a glove tear after sanitizing but before starting compounding. What should the technician do?

- a) Continue; tear is minor
- b) Put on a second glove over the tear and proceed
- c) Replace gloves and re-sanitize as required before proceeding
- d) Tape the tear closed and proceed

49) The label prints "take 1 tablet daily," but the eRx image states "take 1 tablet BID." The medication has been counted but not verified. What should be done next?

- a) Dispense as entered; counseling will correct it
- b) Update the sig to match the prescription, adjust quantity if needed, and reprint the label before moving forward
- c) Handwrite "BID" on the label and send to verification
- d) Keep the sig and double the quantity

50) A Schedule IV prescription is presented with 6 refills authorized. Under federal law, what is the maximum number of refills allowed within 6 months?

- a) 5
- b) 6
- c) 3
- d) 0

51) A patient profile lists "sulfa—rash." A new prescription arrives for celecoxib. What should the technician do?

- a) Fill normally; celecoxib is unrelated
- b) Remove the sulfa allergy because reaction was mild
- c) Substitute naproxen without approval
- d) Route for pharmacist review due to potential sulfonamide concern

52) Which is the correct generic for Flonase?

- a) Mometasone
- b) Fluticasone
- c) Budesonide
- d) Triamcinolone

53) Rx: azithromycin 200 mg/5 mL: 12.5 mL on day 1, then 6.25 mL daily on days 2–5. How many mL should be dispensed?

- a) 37.5 mL
- b) 40 mL
- c) 45 mL
- d) 50 mL

54) A paper prescription for a controlled substance has the quantity overwritten in a different ink. What should the technician do?

- a) Ask the patient to initial the change
- b) Enter it as written and proceed
- c) Bring it to the pharmacist promptly for evaluation before processing
- d) Call the prescriber and change it in the system, then discard the paper

55) A patient asks for the generic of Ventolin HFA. Which is correct?

- a) Levalbuterol
- b) Albuterol
- c) Ipratropium
- d) Budesonide

56) A patient is on sertraline. A new prescription arrives for linezolid. What should the technician do?

- a) Process normally; interaction is minimal
- b) Add an auxiliary label "avoid sunlight"
- c) Dispense but tell the patient to separate doses by 2 hours
- d) Flag for pharmacist review due to serious interaction risk

57) Which is the correct generic for Zofran?

- a) Oseltamivir
- b) Omeprazole
- c) Ondansetron
- d) Olanzapine

58) Directions: "Take 2 tablets PO TID." Quantity #180. What days' supply should be billed?

- a) 20
- b) 30
- c) 45
- d) 90

59) Under DSCSA, a returned product from another store arrives with a damaged 2D barcode and no readable product identifier. What should be done?

- a) Return to stock if the NDC is readable
- b) Dispense it first to avoid waste
- c) Print and apply a new product identifier label
- d) Segregate as suspect product until verification/disposition is completed

60) An eRx shows DOB 04/06/1986, but the existing profile shows 06/04/1986 for the same name. Addresses also differ. What should the technician do?

- a) Assume it is the same patient and proceed
- b) Confirm with additional identifiers and correct/merge only after verification
- c) Change the eRx DOB to match the profile automatically
- d) Create a new profile and dispense under both profiles

61) Which medication is used for hypothyroidism?

- a) Methimazole
- b) Propranolol
- c) Levothyroxine
- d) Prednisone

62) A patient wants to return an opened bottle of gabapentin for a refund. What is the best technician response?

- a) Accept it and restock if tablets look intact
- b) Accept it but relabel for another patient
- c) Explain it generally cannot be returned to stock; offer disposal options per services available
- d) Pour tablets into a disposal bin at the counter without documentation

63) While entering metformin 500 mg, the technician accidentally selects metformin ER 500 mg. No label has printed. What should be done?

- a) Leave it; ER is preferred
- b) Correct to the prescribed formulation before printing and filling
- c) Change the directions to match ER and proceed
- d) Dispense ER and tell the patient to cut tablets in half

64) Trulicity is the brand name for which medication?

- a) Dulaglutide
- b) Liraglutide
- c) Semaglutide
- d) Exenatide

65) A technician observes a coworker repeatedly bypassing barcode scanning to save time. What is the most appropriate action?

- a) Ignore; it is not your role
- b) Do the same so the workflow stays even
- c) Report the safety concern through the appropriate supervisor/incident channel
- d) Confront the coworker loudly so they stop immediately

66) The sig reads: "instill 1 gtt AD BID." What does AD mean?

- a) Right ear
- b) Left ear
- c) Right eye
- d) Both ears

67) Keflex is which generic medication?

- a) Cefuroxime
- b) Cefdinir
- c) Cephalexin
- d) Ceftriaxone

68) Which medication is an ARB?

- a) Propranolol
- b) Losartan
- c) Diltiazem
- d) Lisinopril

69) Rx: doxycycline 100 mg caps; take 1 cap BID x 7 days; quantity #14. The claim rejects for days' supply mismatch. What days' supply is correct?

- a) 5
- b) 7
- c) 10
- d) 14

70) A prescription is flagged "Medication Guide required." What should the technician ensure happens at dispensing?

- a) Provide the Medication Guide with the prescription
- b) Provide it only if the patient requests it
- c) Replace it with a pharmacy-generated monograph
- d) Skip it if the patient has received it before

71) A patient requests "Cymbalta." Which generic should be selected?

- a) Citalopram
- b) Duloxetine
- c) Fluoxetine

d) Venlafaxine

72) Directions: "Take 1 tablet every 8 hours." Quantity #90. What is the days' supply?

- a) 15
- b) 20
- c) 30
- d) 45

73) A prescriber's office requests confirmation of whether a patient picked up a medication. No patient authorization is on file. What should be done?

- a) Provide pickup status because it is part of care coordination
- b) Provide only the fill date, not pickup date
- c) Decline and route to the pharmacist/manager to determine if disclosure is permitted
- d) Provide it if the caller knows the patient's DOB

74) In an inpatient facility, a partially used vial of a Schedule II injectable must be wasted. What is the correct approach?

- a) Dispose in regular trash if capped
- b) Waste per controlled-substance procedure with required documentation and witnessing
- c) Pour into the sink and note it later
- d) Send it back to the wholesaler for credit

75) Which medication is a calcium channel blocker?

- a) Losartan
- b) Furosemide
- c) Lisinopril
- d) Amlodipine

76) A prescription is for tamsulosin 0.4 mg daily. The patient states they are female and asks if it is a mistake. What should the technician do?

- a) Refuse to fill because it is only for men
- b) Change it to oxybutynin because it is for bladder issues
- c) Continue processing without documenting the concern
- d) Refer the question and prescription for pharmacist assessment before dispensing

77) During filling, you find the only stock bottle of nitroglycerin SL tablets is expired. What should you do?

- a) Dispense it; expiration is not important for nitroglycerin
- b) Relabel with a new expiration date based on opening
- c) Remove/quarantine expired product and obtain in-date stock before filling
- d) Dispense and advise the patient to replace it soon

78) Another pharmacy requests a transfer of a noncontrolled prescription. Which information is typically required in the transfer record?

- a) Diagnosis code and lab values
- b) Patient employer and insurance group
- c) Original Rx number, drug/strength, directions, quantity, refills remaining, and identifiers of both pharmacies

d) Only patient name and drug name

79) Carbidopa/levodopa 25/100 is associated with which brand?

- a) Sinemet
- b) Spiriva
- c) Seroquel
- d) Cymbalta

80) A prescription is for "apply 1 patch twice weekly." One carton contains 8 patches. What is the days' supply for one carton?

- a) 14
- b) 21
- c) 28
- d) 56

81) Which situation requires DEA Form 106?

- a) Ordering Schedule II medications
- b) Reporting theft or significant loss of controlled substances
- c) Transferring a controlled substance between pharmacies
- d) Destroying expired noncontrolled medications

82) Which is the correct generic for Augmentin?

- a) Amoxicillin
- b) Amoxicillin/clavulanate
- c) Ampicillin
- d) Cefdinir

83) A hospital order reads "MSO4 2 mg IV q2h PRN pain." Why is this problematic for safe order processing?

- a) The dose is too low to work
- b) IV PRN medications are not permitted
- c) MSO4 is error-prone and can be confused with magnesium sulfate
- d) Morphine cannot be ordered in mg

84) Humalog KwikPen: inject 8 units with meals TID. One box contains five 3 mL pens (100 units/mL). What days' supply is one box?

- a) 56
- b) 62
- c) 46
- d) 75

85) A patient requests naloxone under a statewide standing order. What is within technician scope?

- a) Select the best product based on opioid risk scoring
- b) Provide counseling on when to administer based on respiratory rate
- c) Recommend dose titration after the first spray
- d) Help locate/process the product and refer clinical questions to the pharmacist

86) Which statement is correct about Schedule II refills under federal law?

- a) Refills are allowed if "PRN" is written
- b) Refills are allowed up to 5 within 6 months
- c) Schedule II prescriptions cannot be refilled; a new prescription is required for more
- d) Refills are allowed for hospice patients only

87) A patient profile indicates asthma. A new prescription arrives for timolol ophthalmic drops. What should the technician do?

- a) Fill normally since it is topical
- b) Add an auxiliary label "may cause drowsiness"
- c) Change to latanoprost without contacting anyone
- d) Flag for pharmacist review due to potential beta-blocker effects in asthma

88) Quetiapine is which brand?

- a) Zyprexa
- b) Seroquel
- c) Celexa
- d) Neurontin

89) A stock bottle label is torn and the expiration date cannot be read. What is the best action?

- a) Use it first to avoid waste
- b) Estimate expiration based on when it was ordered
- c) Use it only for noncontrolled prescriptions
- d) Remove it from active stock and follow procedure to identify/confirm expiration before use

90) Rx: hydrocodone/acetaminophen 5/325: take 1 tablet q6h PRN pain; quantity #56. Using maximum daily use, what days' supply should be billed?

- a) 7
- b) 14
- c) 10
- d) 28

Answer Key

1) b	31) c	61) c
2) c	32) c	62) c
3) d	33) c	63) b
4) b	34) b	64) a
5) c	35) d	65) c
6) b	36) a	66) a
7) c	37) a	67) c
8) b	38) b	68) b
9) b	39) c	69) b
10) d	40) c	70) a
11) c	41) b	71) b
12) c	42) b	72) c
13) a	43) c	73) c
14) d	44) a	74) b
15) a	45) d	75) d
16) c	46) c	76) d
17) c	47) a	77) c
18) d	48) c	78) c
19) c	49) b	79) a
20) b	50) b	80) c
21) b	51) d	81) b
22) d	52) b	82) b
23) b	53) a	83) c
24) c	54) c	84) a
25) c	55) b	85) d
26) c	56) d	86) c
27) b	57) c	87) d
28) b	58) b	88) b
29) d	59) d	89) d
30) c	60) b	90) b

PTCB Sim Test 19

- 1) During data entry, an eRx shows “metoprolol succinate ER 25 mg: take 1 tablet daily.” The patient’s last fill was metoprolol tartrate 25 mg BID. What is the best technician action?
- a) Change the prescription to tartrate 25 mg BID to match the profile
 - b) Enter succinate ER as written and note the previous formulation for pharmacist review
 - c) Enter tartrate 25 mg daily because the strengths match
 - d) Split the difference and enter tartrate 25 mg BID with ER selected
- 2) A patient asks to purchase pseudoephedrine 30 mg tablets, quantity 96. Which step is required at the point of sale under federal law?
- a) Record the transaction in the required log after confirming a valid photo ID
 - b) Sell only 24 tablets per day to any purchaser
 - c) Require a prescription if the purchaser is under age 21
 - d) Provide a Medication Guide and document refusal if declined
- 3) An eRx reads: “Augmentin 875/125 mg: take 1 tablet PO BID for 10 days.” How many tablets should be dispensed?
- a) 10
 - b) 14
 - c) 20
 - d) 28
- 4) A prescriber’s office calls to give a verbal order for hydromorphone. The pharmacist is counseling and cannot come to the phone. What should the technician do?
- a) Take the verbal order and label it “pending RPh signature”
 - b) Ask the caller to repeat the order slowly and you will write it down for later
 - c) Obtain a callback number and have the pharmacist take the call when available
 - d) Accept the verbal order only if it is for a 72-hour supply
- 5) During filling, the technician pulls “hydroxyzine 25 mg” from the shelf for a label that reads “hydralazine 25 mg.” Which step most directly prevents this error at the fill station?
- a) Counting on a separate tray for each patient
 - b) Comparing tablet color and shape to the computer image
 - c) Placing both drugs in the same shelf section for faster picking
 - d) Scanning the stock bottle barcode/NDC to match the label
- 6) A patient presents a paper Schedule II prescription that is signed but not dated. What should the technician do?
- a) Enter it and leave the date blank; the pharmacist will decide
 - b) Ask the patient what date it was written and enter that date
 - c) Have the patient write today’s date next to the signature
 - d) Set it aside for pharmacist review because a required element is missing
- 7) A prescription is for albuterol HFA: “inhale 2 puffs q4h while awake.” The patient reports using it 4 times daily. The inhaler contains 200 actuations. What days’ supply should be billed?

- a) 25
- b) 50
- c) 75
- d) 100

8) A profile lists "sulfonamides—anaphylaxis." A new eRx arrives for sulfamethoxazole/trimethoprim. What is the correct technician response?

- a) Fill it because the prescriber sent it electronically
- b) Override the allergy alert if the patient used it in the past
- c) Hold processing and route the allergy conflict for pharmacist evaluation
- d) Substitute nitrofurantoin to avoid a delay

9) A tote from the wholesaler arrives with broken tamper tape and no matching invoice inside. What is the best technician action?

- a) Put items away but keep the tote for documentation
- b) Quarantine the tote contents as suspect and alert the pharmacist/manager
- c) Return the tote to the wholesaler driver immediately without documenting
- d) Place the tote in will-call so it is separated from inventory

10) While entering, the technician selects "liothyronine 5 mcg" instead of "levothyroxine 75 mcg" due to adjacent search results. What is the best immediate correction?

- a) Keep the selection and rely on counseling to catch it
- b) Re-select the correct drug/strength and continue through normal verification steps
- c) Change the sig to match the drug selected
- d) Substitute another thyroid medication to avoid re-entry

11) The profile lists "warfarin—hives." A new prescription for warfarin 5 mg is received. What should the technician do?

- a) Remove the allergy since warfarin is commonly tolerated
- b) Proceed and add a label: "Stop if rash occurs"
- c) Call the prescriber directly to request a different drug
- d) Stop and route to the pharmacist to assess the allergy history

12) A patient requests an early refill for oxycodone because they are traveling tomorrow. What is the best technician step?

- a) Change the pickup date so insurance will pay
- b) Gather travel details and route for pharmacist decision and documentation
- c) Dispense a partial now and the remainder after the due date without documentation
- d) Advise the patient to take half doses until the refill date

13) Amoxicillin 400 mg/5 mL: "take 6 mL PO BID for 10 days." What total quantity should be dispensed?

- a) 60 mL
- b) 100 mL
- c) 120 mL
- d) 150 mL

14) A patient returns an unopened vial of tablets that left the pharmacy yesterday and asks

to have it restocked. What is the correct action?

- a) Restock if the seal is intact and the medication is not refrigerated
- b) Restock only if it is not a controlled substance
- c) Do not restock once it has left pharmacy control; follow return procedure
- d) Restock after verifying the lot number and expiration date

15) A prescription is for Lantus SoloStar 100 units/mL. The technician accidentally selects Toujeo SoloStar 300 units/mL because both are insulin glargine pens. What should the technician do?

- a) Dispense Toujeo but adjust the days' supply
- b) Re-select the exact prescribed product and verify with barcode/NDC during fill
- c) Add directions to inject one-third the units
- d) Dispense either product since both contain insulin glargine

16) A law office delivers a subpoena requesting a patient's complete pharmacy profile. What should the technician do first?

- a) Print the profile and provide it immediately
- b) Decline all subpoenas due to HIPAA
- c) Route the request for pharmacist/management review before any release
- d) Release only controlled substance history to limit PHI

17) Nystatin 100,000 units/mL oral suspension: "swish and swallow 5 mL QID for 10 days." What quantity should be dispensed?

- a) 100 mL
- b) 150 mL
- c) 200 mL
- d) 250 mL

18) A prescriber writes "1 tab PO qd." Your pharmacy discourages "qd" due to misread risk. How should a technician enter it?

- a) Enter "qd" exactly as written
- b) Enter "daily"
- c) Enter "qod" to reduce confusion
- d) Reject the prescription and require rewrite

19) Which federal recordkeeping practice is required for Schedule II purchase invoices?

- a) Keep only electronically; paper copies are not allowed
- b) File separately from non-controlled invoices
- c) Keep in the will-call area until fully dispensed
- d) Maintain only until the next controlled substance inventory

20) An isotretinoin prescription is ready to fill, but the patient is new and has not completed required program steps. What can legally prevent dispensing even if the label is correct?

- a) The patient must be at least 21 years old
- b) The prescription must be filled within 24 hours of being written
- c) REMS requirements must be satisfied prior to dispensing
- d) The drug can only be dispensed in 90-day supplies

- 21) A claim rejects "refill too soon." The patient reports the medication was stolen. What is the best technician action?
- a) Reduce days' supply to 7 to push it through insurance
 - b) Change the fill date to yesterday and resubmit
 - c) Tell the patient nothing can be done until due
 - d) Collect details (date, report/plan requirements) and route for authorized override handling
- 22) Gabapentin 300 mg: "take 1 capsule TID." Dispense #90. What is the days' supply?
- a) 15
 - b) 30
 - c) 45
 - d) 90
- 23) Another pharmacy requests a transfer for a Schedule II prescription because they have stock and you do not. What is the correct response?
- a) Transfer is allowed once with pharmacist-to-pharmacist communication
 - b) Transfer is allowed only within the same chain
 - c) Schedule II prescriptions generally cannot be transferred; a new prescription is needed
 - d) Transfer is allowed if the prescriber authorizes by phone
- 24) The label prints "sertraline 50 mg," but the bottle pulled is "sildenafil 50 mg." Which workflow step best prevents this?
- a) Using a larger vial so the label is clearer
 - b) Scanning the stock bottle barcode/NDC before counting
 - c) Comparing tablet color to the on-screen image only
 - d) Storing similar names adjacent to speed picking
- 25) Latanoprost 0.005%: "instill 1 drop OU qhs." Dispense 2.5 mL. Assume 20 drops = 1 mL. What days' supply should be billed?
- a) 25
 - b) 30
 - c) 50
 - d) 75
- 26) DUR shows "Max daily acetaminophen exceeded." Rx A: acetaminophen 500 mg, 2 tablets QID. Active Rx B: hydrocodone/APAP 5/325 mg, 1 tablet QID. What should the technician do?
- a) Bypass the alert and proceed to fill both
 - b) Route for pharmacist DUR resolution before dispensing continues
 - c) Change acetaminophen directions to TID and proceed
 - d) Delete the hydrocodone/APAP because it is controlled
- 27) Prednisone 20 mg taper: "3 tablets daily x 3 days, then 2 tablets daily x 3 days, then 1 tablet daily x 3 days." How many tablets are needed?
- a) 12
 - b) 18
 - c) 24
 - d) 27

28) A spouse calls asking, "Did my husband pick up his HIV medicine today?" The spouse is not listed as an authorized contact. What should the technician do?

- a) Confirm pickup status but not the drug name
- b) Confirm only if the spouse verifies DOB and address
- c) Decline to disclose without patient authorization
- d) Confirm the copay amount only

29) An opened bottle of nitroglycerin SL tablets is found with no open date and unknown beyond-use status. What should be done?

- a) Keep it and use it first to avoid waste
- b) Assign a new 6-month date from today
- c) Tape the cap and keep it in active stock
- d) Remove from active inventory and quarantine for proper disposition

30) The correct drug and directions are printed, but the label has the wrong patient name due to selecting the wrong profile. What is the best technician action?

- a) Reprint with the correct name and dispense
- b) Cross out the name and write the correct one
- c) Stop and correct patient selection per QA workflow before dispensing
- d) Dispense if the address matches

31) Which generic corresponds to Plavix?

- a) clopidogrel
- b) clonidine
- c) clozapine
- d) clarithromycin

32) A patient asks whether a new antibiotic is safe with their cholesterol medication. What should the technician do?

- a) Suggest separating doses by 2 hours
- b) Tell the patient to stop the cholesterol medicine until finished
- c) State that antibiotics do not interact with statins
- d) Refer the patient to the pharmacist for interaction assessment

33) A vaccine shipment arrives with an indicator showing the package exceeded temperature limits during transit. What should the technician do?

- a) Refrigerate and stock it immediately
- b) Quarantine and alert the pharmacist for disposition guidance
- c) Put it in will-call to keep it separate
- d) Use it today only to avoid further warming

34) Triamcinolone 0.1% cream: "apply to both hands BID." Quantity 30 g. Billing rule: both hands together use 1 g per application. What days' supply should be billed?

- a) 10
- b) 15
- c) 20
- d) 30

35) For an initial controlled substance inventory at a new pharmacy, what is required for Schedule II drugs?

- a) Exact count for all Schedule II substances
- b) Estimated counts are allowed for all Schedule II bottles
- c) Exact counts only for Schedule II narcotics, not stimulants
- d) Inventory must be submitted weekly to the DEA

36) Dorzolamide/timolol ophthalmic: "instill 1 drop OS BID." Dispense 10 mL. Assume 20 drops = 1 mL. What days' supply should be billed?

- a) 50
- b) 75
- c) 100
- d) 200

37) Which generic corresponds to Zofran ODT?

- a) omeprazole
- b) ondansetron
- c) oxybutynin
- d) olanzapine

38) An eRx says "Synthroid 25 mcg DAW." Generic levothyroxine 25 mcg is in stock. What should be dispensed?

- a) Generic levothyroxine because it is AB-rated
- b) Brand Synthroid as written
- c) Any levothyroxine product; DAW is optional
- d) Liothyronine because it is another thyroid hormone

39) Methotrexate 2.5 mg arrives with directions "Take 6 tablets daily" for rheumatoid arthritis. What is the correct technician action?

- a) Fill as written because it is a common regimen
- b) Change "daily" to "weekly" because that is typical
- c) Stop and route to the pharmacist due to high-alert dosing concern
- d) Reduce quantity to limit risk and dispense

40) Estradiol patch 0.05 mg/day: "apply 1 patch twice weekly." Dispense 8 patches. What is the days' supply?

- a) 14
- b) 21
- c) 28
- d) 56

41) A manufacturer recall is issued for a specific lot of losartan due to contamination risk. What should the technician do with affected stock?

- a) Continue dispensing until a replacement arrives
- b) Quarantine the lot and follow recall instructions for return/credit documentation
- c) Re-label the lot with a new expiration date
- d) Transfer it to another store to reduce waste

42) A prescription direction includes "bid ac." What does "ac" mean?

- a) after meals
- b) before meals
- c) at bedtime
- d) with meals

43) The prescription is for glipizide XL 10 mg daily, but only glipizide IR 10 mg is on the shelf. What should the technician do?

- a) Substitute IR and change directions to BID
- b) Dispense IR with the same directions
- c) Substitute IR because strength matches
- d) Put on hold and route for pharmacist review due to non-interchangeable dosage forms

44) A patient asks to buy insulin syringes without a prescription. What is the best technician response?

- a) Federal law prohibits sales without a prescription
- b) It is always allowed with no restrictions
- c) It depends on state law and store policy; confirm what applies here
- d) It is allowed only if the patient has insulin on file in all states

45) The stock bottle expires next month, but the system prints a 1-year date on the label. What should the technician do?

- a) Dispense because the label date controls
- b) Use the manufacturer's expiration if earlier and ensure the label reflects the earliest applicable date
- c) Cross out the manufacturer date on the bottle to match the label
- d) Dispense if the patient will finish before next month

46) Humalog KwikPen 100 units/mL: "inject 8 units before meals TID." A box has 5 pens, each 3 mL. What days' supply is 1 box?

- a) 56
- b) 62
- c) 75
- d) 90

47) A neighbor asks whether a patient picked up medication today and is not authorized on the patient's profile. What should the technician do?

- a) Confirm pickup status only
- b) Confirm if the neighbor can state DOB
- c) Decline to disclose and offer to contact the patient for authorization
- d) Provide only the copay amount

48) Diclofenac 1% gel: "apply 2 g topically BID." Dispense 100 g. What days' supply should be billed?

- a) 10
- b) 25
- c) 50

d) 100

49) Which brand corresponds to rivaroxaban?

- a) Eliquis
- b) Pradaxa
- c) Xarelto
- d) Savaysa

50) A scanned prescription is too blurry to tell whether the dose is “10 mg” or “100 mg.” What should the technician do before completing entry?

- a) Enter 10 mg because it is more common
- b) Enter 100 mg and let verification catch it
- c) Use the last fill to choose the dose
- d) Obtain a clearer copy or contact the prescriber’s office for clarification

51) A patient has an active rosuvastatin prescription. A new prescription for simvastatin is received from a different prescriber. What should the technician do?

- a) Fill both because they are different statins
- b) Cancel the older statin automatically
- c) Enter both and let the patient decide which to take
- d) Route to the pharmacist to assess duplication vs intended switch

52) A prescription is written for “Keflex 500 mg.” Which generic should be selected?

- a) cephalexin
- b) cefdinir
- c) ceftriaxone
- d) ceftazidime

53) Metformin ER 500 mg is prescribed. Which auxiliary label is most appropriate?

- a) “Take on an empty stomach”
- b) “Do not crush or chew”
- c) “May cause drowsiness”
- d) “Avoid sunlight”

54) Under HIPAA, which disclosure is generally permitted without patient authorization?

- a) Sharing PHI with an employer to verify adherence
- b) Sharing PHI with a spouse who can verify DOB
- c) Sharing PHI with another provider for treatment purposes
- d) Sharing PHI with a neighbor providing transportation

55) Clindamycin 75 mg/5 mL: “take 10 mL PO TID for 7 days.” What quantity should be dispensed?

- a) 140 mL
- b) 200 mL
- c) 210 mL
- d) 280 mL

56) An eRx for Adderall XR #30 arrives with 0 refills. The patient asks you to “add two refills

like last time.” What should the technician do?

- a) Add refills if the patient has been stable
- b) Add refills and document “per patient request”
- c) Leave as written and route if the patient requests pharmacist assistance contacting the prescriber
- d) Convert to Adderall IR with refills

57) A label direction says “take 1 capsule by mouth q8h.” How many doses per day is that?

- a) 2
- b) 3
- c) 4
- d) 8

58) A prescription is written for morphine ER 30 mg with directions “take 1 tablet q4–6h PRN pain.” What should the technician do?

- a) Enter as written; ER can be used at that frequency
- b) Change to morphine IR because PRN implies IR
- c) Change the directions to BID automatically
- d) Route for pharmacist clarification due to mismatch between formulation and directions

59) A profile shows omeprazole daily. A new prescription arrives for pantoprazole daily. What is the best technician action?

- a) Fill both because they are different PPIs
- b) Route to the pharmacist to assess duplication vs switch
- c) Substitute famotidine automatically
- d) Cancel omeprazole automatically and fill pantoprazole

60) Under federal law, how many times may a Schedule III–V prescription be transferred between pharmacies for refill dispensing (traditional rule)?

- a) 0
- b) 1
- c) 3
- d) Unlimited until refills are used

61) A technician prepares a nonsterile compounded oral suspension from a master formulation record. What must occur before it can be dispensed?

- a) It can be dispensed once labeled; documentation can be finished later
- b) Any beyond-use date up to 90 days is acceptable
- c) Ensure compounding documentation is complete and the product proceeds through pharmacist final verification
- d) Only the active ingredient lot number needs to be recorded

62) A patient has “penicillin—anaphylaxis” documented. A prescription arrives for cephalexin. What should the technician do?

- a) Process normally because it is not penicillin
- b) Override the alert because cephalosporins are always safe
- c) Substitute azithromycin automatically
- d) Pause and route to the pharmacist due to allergy severity and cross-reactivity concern

63) A patient asks what time of day to take a medication labeled "take 1 tablet once daily." What is the best technician response?

- a) Recommend morning for best absorption
- b) Recommend bedtime to avoid side effects
- c) Ask what other medications they take and decide for them
- d) Offer pharmacist counseling for individualized timing guidance

64) Toujeo SoloStar 300 units/mL: inject 24 units daily. Dispense 3 pens. Each pen contains 1.5 mL. What is the days' supply?

- a) 47
- b) 56
- c) 70
- d) 90

65) What ingredient makes Augmentin different from amoxicillin alone?

- a) sulbactam
- b) clavulanate
- c) tazobactam
- d) probenecid

66) For a Schedule III–V prescription transfer, which statement best reflects federal requirements?

- a) Transfers are never allowed
- b) A technician may complete the transfer independently if documented
- c) Transfers may occur with proper documentation and required pharmacist involvement
- d) Transfers are allowed only if no refills remain

67) Ciprofloxacin 0.3% ophthalmic: "instill 2 drops OU q4h for 7 days." Dispense 5 mL. Assume 20 drops = 1 mL. Is 5 mL sufficient?

- a) Yes
- b) No, dispense 2.5 mL
- c) No, dispense 7.5 mL
- d) No, dispense 15 mL

68) During unit-dose cart fill, which practice best reduces administration errors?

- a) Pre-opening all blisters to save time
- b) Combining tablets for multiple patients in one cup
- c) Using barcode scanning to match product to the order/MAR
- d) Skipping expiration checks for fast-moving items

69) A written prescription for amoxicillin is presented without a prescriber signature. What should the technician do?

- a) Enter it and leave the signature area blank
- b) Ask the patient to sign for the prescriber
- c) Fill a 3-day supply and request a signed copy later
- d) Hold and route for pharmacist review because the prescription is not valid as written

70) Which generic corresponds to Eliquis?

- a) rivaroxaban
- b) dabigatran
- c) apixaban
- d) edoxaban

71) While restocking vaccines, you find a box that expires tomorrow. What is the best technician action?

- a) Remove the expiration sticker so it can still be used
- b) Move it to the front and alert the pharmacist so it can be prioritized appropriately
- c) Freeze it to extend dating
- d) Place it in will-call to keep it separate

72) Acetaminophen 500 mg: "take 2 tablets QID." What is the total daily dose in mg?

- a) 2,000 mg
- b) 3,000 mg
- c) 4,000 mg
- d) 5,000 mg

73) A Schedule II prescription is partially filled because the pharmacy does not have enough stock. Which action best supports compliance?

- a) Add refills for the remainder once stock arrives
- b) Convert the remaining quantity to a new Schedule III prescription
- c) Document the partial fill and ensure the remainder is handled within applicable time limits and requirements
- d) Void the original and issue a new prescription for the remaining amount

74) A reconstituted refrigerated antibiotic suspension has been sitting in will-call at room temperature for 15 days. What should the technician do?

- a) Refrigerate it now and dispense
- b) Add a "refrigerate" label and dispense
- c) Quarantine and route for pharmacist evaluation; it may require reversal and re-preparation
- d) Dispense if the patient agrees to refrigerate at home

75) A prescription is for bupropion XL 150 mg daily, but the shelf has bupropion SR 150 mg. What should the technician do?

- a) Dispense SR with the same directions since strength matches
- b) Dispense SR and change the sig to BID
- c) Dispense SR if the patient used it previously
- d) Select the correct XL formulation and proceed through normal verification

76) A patient asks for an OTC recommendation for new chest pain with shortness of breath. What should the technician do?

- a) Recommend an antacid and advise follow-up if it persists
- b) Recommend ibuprofen and rest
- c) Refer immediately for pharmacist assistance and urgent evaluation as appropriate
- d) Recommend aspirin only and send the patient home

77) Promethazine syrup: "take 5 mL PO QID." How many mL will be used in 6 days?

- a) 60 mL
- b) 90 mL
- c) 120 mL
- d) 150 mL

78) Under federal controlled substance recordkeeping requirements, which records must be maintained for at least 2 years and be readily retrievable?

- a) Only Schedule II dispensing records
- b) Only prescriber DEA certificates
- c) Only controlled substance return-to-stock logs
- d) Schedule II and Schedule III–V records (separately filed or readily retrievable)

79) A prescription indicates "DAW 1." What does this mean for product selection?

- a) Generic substitution is required
- b) Dispense brand as written (no substitution)
- c) Dispense any AB-rated generic only
- d) Patient may choose brand or generic regardless of prescriber

80) A patient shows a photo of an old label for a Schedule II stimulant and requests a refill, but there is no active prescription on file. What should the technician do?

- a) Dispense a 3-day emergency supply
- b) Use the label photo as authorization to refill
- c) Transfer the prescription from another pharmacy
- d) Explain a new prescription is required and route to the pharmacist for next steps

81) Which medication is the generic for Ventolin HFA?

- a) budesonide
- b) albuterol
- c) fluticasone
- d) salmeterol

82) Two NSAIDs appear active: ibuprofen 800 mg TID and naproxen 500 mg BID from different prescribers. A duplicate therapy alert appears. What should the technician do?

- a) Override because combining NSAIDs improves pain control
- b) Cancel one automatically
- c) Route to the pharmacist for duplication review before dispensing
- d) Tell the patient to alternate them every other day

83) A prescription reads "morphine sulfate 15 mg tablets" with no IR/ER specified. The patient previously filled MS Contin. What should the technician do?

- a) Dispense MS Contin because it's on the profile
- b) Dispense morphine IR because "tablets" implies IR
- c) Route to the pharmacist for clarification of formulation before processing
- d) Split the quantity between IR and ER

84) A prescription arrives for furosemide 40 mg daily. The profile shows "sulfa allergy—mild rash." What should the technician do?

- a) Remove the allergy because it is mild
- b) Ignore the alert because furosemide is not an antibiotic
- c) Substitute ethacrynic acid automatically
- d) Route to the pharmacist to evaluate allergy relevance and risk

85) Levetiracetam 500 mg: "1 tablet BID #60." The patient requests a 90-day supply today. What should the technician do?

- a) Change quantity to #180 and process
- b) Change refills to cover 90 days
- c) Tell the patient 90-day supplies are never allowed for seizure medications
- d) Route for pharmacist review of authorization/plan rules and prescriber intent

86) Diltiazem CD 240 mg daily is prescribed, but only diltiazem immediate-release 120 mg is available. What should the technician do?

- a) Substitute IR and enter TID directions
- b) Dispense IR with the same directions because it's the same drug
- c) Route to the pharmacist because dosage form substitution is not automatic
- d) Change CD to SR automatically

87) While entering a controlled substance prescription, the prescriber profile is missing the DEA number. The paper prescription shows a DEA number. What should the technician do?

- a) Use a placeholder DEA number to proceed
- b) Enter the DEA from the prescription and proceed through the normal verification workflow
- c) Copy a DEA from another prescriber with a similar name
- d) Leave DEA blank because it is optional if the prescriber is licensed

88) A prescription is for EpiPen 0.3 mg auto-injector. Which generic name matches?

- a) epinephrine
- b) phenylephrine
- c) norepinephrine
- d) diphenhydramine

89) A prescription arrives for lamotrigine 100 mg. The patient profile lists "lamotrigine—rash" as an allergy. What should the technician do?

- a) Override and proceed because rash is common
- b) Dispense with an auxiliary label "may cause rash"
- c) Route to the pharmacist due to documented allergy conflict
- d) Change to levetiracetam automatically

90) A new eRx is for sertraline 100 mg daily. The patient has an active sertraline 50 mg daily with refills remaining. The patient says the dose was increased. What should the technician do?

- a) Cancel the new prescription because refills remain on the old one
- b) Fill both strengths so the patient can adjust dosing
- c) Enter the new prescription as written and route for pharmacist review for dose-change/duplication handling
- d) Change to 75 mg daily as a compromise

Answer Key

1) b	31) a	61) c
2) a	32) d	62) d
3) c	33) b	63) d
4) c	34) b	64) b
5) d	35) a	65) b
6) d	36) c	66) c
7) b	37) b	67) a
8) c	38) b	68) c
9) b	39) c	69) d
10) b	40) c	70) c
11) d	41) b	71) b
12) b	42) b	72) c
13) c	43) d	73) c
14) c	44) c	74) c
15) b	45) b	75) d
16) c	46) b	76) c
17) c	47) c	77) c
18) b	48) c	78) d
19) b	49) c	79) b
20) c	50) d	80) d
21) d	51) d	81) b
22) b	52) a	82) c
23) c	53) b	83) c
24) b	54) c	84) d
25) c	55) c	85) d
26) b	56) c	86) c
27) b	57) b	87) b
28) c	58) d	88) a
29) d	59) b	89) c
30) c	60) b	90) c

PTCB Sim Test 20

1) A patient asks to pick up “my antibiotic” for Jordan Lee. The will-call has two bags: Jordan Lee (DOB 02/14/1990) and Jordan Lee (DOB 02/14/2009). What is the best next step before handing over any bag?

- a) Ask which medication they’re expecting and match to the answer
- b) Ask for DOB and one additional identifier, then match to the label
- c) Hand both bags over and let the patient sort them out
- d) Ask for the address only, since names and DOB match

2) A prescription reads: “metFORMIN ER 500 mg tab. Take 2 tabs po qpm with dinner. #60.” What days’ supply should be entered?

- a) 15
- b) 30
- c) 45
- d) 60

3) A new eRx: “Synthroid 75 mcg daily #30” arrives. The profile shows levothyroxine 75 mcg daily filled 5 days ago from the same prescriber with 25 days remaining. The patient says, “My doctor increased it today.” What is the best workflow step for the technician?

- a) Process as a duplicate and delete the new eRx
- b) Put it on hold until the patient runs out
- c) Continue processing but flag as possible therapy change for pharmacist review
- d) Reverse the previous claim immediately and rebill the new one

4) A prescription is written: “Augmentin 875/125 mg: 1 tab po bid x 7 days.” What quantity should be dispensed?

- a) 7
- b) 10
- c) 14
- d) 21

5) A patient’s profile lists “ACE inhibitors—angioedema.” A new prescription is received for lisinopril 10 mg daily. What should the technician do?

- a) Enter it and print labels; allergy checks are pharmacist-only
- b) Remove the allergy since it’s a class allergy, not a drug allergy
- c) Proceed but add “take with food” to reduce reaction risk
- d) Ensure the allergy alert is visible and route for pharmacist review before filling

6) A patient drops off a prescription for “Adderall 20 mg tablet #60 take 1 tab bid,” but it is missing the prescriber’s address. What is the most appropriate action?

- a) Call the office to obtain the missing required information and document per pharmacist direction
- b) Add the prescriber address from Google Maps and continue
- c) Fill it because the DEA number is present
- d) Convert it to an emergency supply until the address is provided

7) A technician is selecting stock for hydrALAZINE 25 mg but accidentally scans hydrOXYzine 25 mg. The system displays an NDC mismatch alert. What should be done next?

- a) Override since the dose strength matches
- b) Stop, reselect the correct product, and rescan before proceeding
- c) Continue and leave a note for the pharmacist to verify
- d) Change the drug in the system to match the scanned bottle

8) A prescription for warfarin is being processed. The patient started trimethoprim-sulfamethoxazole yesterday per profile history. What is the best technician action?

- a) Add an auxiliary label "separate from dairy"
- b) Enter normally; interactions are covered at counseling
- c) Flag for pharmacist review due to high-risk interaction potential
- d) Tell the patient to stop warfarin until the antibiotic is finished

9) A prescription reads: "predniSONE 10 mg tabs: 4 tabs daily x 3 days, then 3 tabs daily x 3 days, then 2 tabs daily x 3 days, then 1 tab daily x 3 days." How many tablets should be dispensed?

- a) 24
- b) 30
- c) 36
- d) 40

10) A patient is purchasing pseudoephedrine. Under federal CMEA requirements, which must occur before the sale is completed?

- a) Verify the patient is at least 21 years old
- b) Obtain a prescription from a prescriber
- c) Patient presents valid photo ID and signs the log (paper or electronic)
- d) Pharmacist must provide mandatory counseling

11) A prescription reads: "cephalexin 250 mg/5 mL. Take 10 mL po bid x 10 days." What total volume should be dispensed?

- a) 100 mL
- b) 150 mL
- c) 200 mL
- d) 250 mL

12) A patient asks to return an unopened bottle of tablets that was dispensed yesterday because "the doctor changed my mind." What is the correct action in most community pharmacies?

- a) Return to stock if the seal is intact
- b) Accept the return and relabel for another patient
- c) Accept and place into regular trash to prevent diversion
- d) Accept for disposal per policy, but do not return to stock for redispensing

13) A hospital order reads: "Heparin infusion per protocol." No patient weight or protocol details are attached. What should the technician do first?

- a) Prepare the standard concentration used most often on the unit

- b) Ask nursing what rate they want to start at
- c) Route for pharmacist verification/clarification before compounding
- d) Substitute enoxaparin because it's safer

14) A prescription reads: "dorzolamide-timolol: instill i gtt OU bid." Which is correct?

- a) Instill 1 drop in each eye twice daily
- b) Instill 2 drops in each eye twice daily
- c) Instill 1 drop in the right eye twice daily
- d) Instill 1 drop in both eyes once daily

15) A vial label shows: "Take 1 tab po tid." Quantity is 90 tablets. What days' supply should be billed?

- a) 15
- b) 20
- c) 30
- d) 45

16) A patient requests a printed list of all prescriptions filled in the last 12 months. What is the best approach?

- a) Provide the printout after verifying identity with at least two identifiers
- b) Provide it only if the patient signs a notarized request
- c) Refuse because fill history is pharmacy property
- d) Give it to anyone who can state the patient's DOB

17) A prescription is written for "Lantus Solostar: inject 18 units subQ nightly. Dispense 15 mL." Using 100 units/mL, what days' supply is this?

- a) 75
- b) 83
- c) 90
- d) 100

18) A patient asks which OTC product to take for chest pressure that started an hour ago and radiates to the left arm. What is the technician's best response?

- a) Recommend aspirin and have them sit down for 10 minutes
- b) Recommend omeprazole in case it is reflux
- c) Direct them to seek immediate medical attention and alert the pharmacist immediately
- d) Recommend ibuprofen and avoid caffeine

19) A prescription for isotretinoin is ready, but the system shows the patient's REMS requirements are incomplete for the current month. What should be done?

- a) Dispense and remind the patient to complete requirements next month
- b) Hold the prescription until REMS is complete and involve the pharmacist
- c) Dispense a smaller quantity to bypass the restriction
- d) Substitute tretinoin cream without contacting the prescriber

20) A technician finds a stock bottle on the shelf containing tablets but no label, no NDC, and no lot/expiration. What is the best action?

- a) Use a pill identifier and relabel the bottle for use

- b) Quarantine the bottle and alert the pharmacist; do not use for dispensing
- c) Put it aside and use it only for cash patients
- d) Repackage into unit-dose packaging to improve safety

21) A patient asks for a refill of a Schedule II medication and says, "The label says no refills, but can you just refill it one time?" What is correct under federal law?

- a) Schedule II prescriptions may be refilled once within 30 days
- b) Schedule II prescriptions cannot be refilled; a new prescription is required
- c) Schedule II prescriptions can be refilled up to 5 times in 6 months
- d) Schedule II prescriptions can be refilled if the patient pays cash

22) A prescription reads: "amLODIPine 5 mg tab i po qd #30." Which SIG entry is correct?

- a) Take 1 tablet by mouth daily
- b) Take 5 tablets by mouth daily
- c) Take 1 tablet by mouth twice daily
- d) Take 1 tablet by mouth daily as needed

23) A technician accidentally selects bupropion XL 150 mg when the prescription is for bupropion SR 150 mg, 1 tablet twice daily. What is the best correction?

- a) Keep XL and change directions to once daily
- b) Change to SR and keep directions as written
- c) Keep XL and keep twice-daily directions
- d) Change to IR and adjust to three times daily

24) A prescription reads: "nystatin 100,000 units/mL: swish and swallow 5 mL qid x 14 days." How many mL should be dispensed?

- a) 140 mL
- b) 200 mL
- c) 240 mL
- d) 280 mL

25) A controlled substance prescription is presented with the prescriber's DEA number present, but the date is missing. What is the best action?

- a) Ask the patient what date it was written and use that
- b) Use today's date as the written date
- c) Set aside and obtain a valid prescription with required elements before dispensing
- d) Fill it if the patient has filled from the prescriber before

26) During data entry, a prescription is entered for sertraline 50 mg daily, but the technician accidentally selects paroxetine 20 mg daily from the dropdown. Which step is most likely to catch this before the patient receives it?

- a) Cashier review at pickup
- b) Pharmacist product verification against the original prescription and stock bottle/NDC
- c) Patient counseling refusal documentation
- d) Bin location verification by another technician

27) A prescription reads: "Take 2.5 mL po q12h." Concentration is 250 mg/5 mL. How many mg per dose?

- a) 62.5 mg
- b) 100 mg
- c) 125 mg
- d) 250 mg

28) A technician receives a manufacturer recall notice for specific lots of losartan. The technician finds matching lot numbers on the shelf. What should be done?

- a) Remove from shelf, quarantine per recall procedure, and notify the pharmacist/manager
- b) Continue dispensing until replacement arrives
- c) Place a sticker warning on the bottle and keep dispensing
- d) Re-label with a different lot number and segregate

29) A prescription for hydrocodone/acetaminophen 5/325 mg is written: "1 tab q6h prn pain. Qty 18." What is the days' supply (assume max use)?

- a) 3 days
- b) 4 days
- c) 5 days
- d) 6 days

30) A patient requests transfer of a prescription for buprenorphine/naloxone from another pharmacy. What is the best next step?

- a) Transfer like any legend drug
- b) Tell the patient controlled substances are never transferable
- c) Transfer only if the original pharmacy is in the same chain
- d) Confirm schedule and transfer eligibility under federal/state rules and route to pharmacist to complete the transfer

31) A prescription reads: "qod" in the SIG. What does this mean?

- a) Every day
- b) Every other day
- c) Four times daily
- d) Once weekly

32) A prescription is written: "valACYclovir 1 g: take 2 tabs po bid for 1 day." What quantity should be dispensed?

- a) 2
- b) 4
- c) 6
- d) 8

33) A technician is asked to sell insulin syringes U-100, 0.5 mL, 31G. The patient uses insulin 4 times daily. Quantity is 100 syringes. What days' supply is this?

- a) 20
- b) 25
- c) 30
- d) 50

34) An eRx for clonazepam arrives missing the prescriber's phone number in the message.

What is the best action?

- a) Fill it because eRx's do not require a phone number
- b) Add the phone number from an internet search and proceed
- c) Hold and obtain the missing required prescriber information through proper channels before dispensing
- d) Convert it to a verbal order and document it as such

35) At pickup, a patient asks how to store nitroglycerin SL tablets. Which response is appropriate within technician scope?

- a) Store in the original container, tightly closed, away from heat and moisture
- b) Store in a weekly pill organizer for convenience
- c) Store in the bathroom medicine cabinet
- d) Store in the refrigerator to prevent breakdown

36) A prescription reads: "ciprofloxacin 0.3% eye drops: 2 gtt q2h while awake x 2 days, then 2 gtt q4h while awake x 5 days." Total duration is:

- a) 5 days
- b) 7 days
- c) 10 days
- d) 14 days

37) A patient's spouse calls asking for the patient's medication list "to bring to the hospital." There is no documented authorization. What should the technician do?

- a) Provide it if the spouse knows the patient's DOB
- b) Provide only medication names, not directions
- c) Request to speak directly with the patient or obtain documented authorization before releasing PHI
- d) Fax it to the spouse's work fax number marked "confidential"

38) An order entry screen shows "Humulin R U-100" selected, but the eRx says "Humalog U-100." What is the best technician action?

- a) Keep the selection; both are insulins and interchangeable
- b) Change the prescription to match the selected product
- c) Stop and correct the product selection to match the prescription before filling
- d) Dispense and ask the patient which one they use

39) A prescription is written for "Eliquis 5 mg" but the SIG line is blank. What is the best next step?

- a) Enter the standard SIG "1 tab bid" and proceed
- b) Enter "use as directed" and dispense
- c) Contact the patient to guess the dose based on indication
- d) Obtain complete directions before dispensing

40) A patient's profile shows anaphylaxis to penicillin. A prescription is received for amoxicillin/clavulanate. What is the best technician action?

- a) Process normally since clavulanate reduces allergy risk
- b) Ask the patient if they've taken it before and proceed if yes
- c) Ensure the allergy is flagged and route for pharmacist assessment before dispensing

d) Change to cephalexin automatically

41) A pharmacy receives refrigerated semaglutide that arrived with a temperature indicator showing an excursion. What should be done?

- a) Refrigerate immediately and dispense first
- b) Quarantine the shipment and notify the pharmacist/manager per procedure
- c) Return to stock if the carton is sealed
- d) Freeze it to restore potency

42) A prescription reads: "fluticasone nasal spray: 2 sprays each nostril daily." Bottle contains 120 sprays. What days' supply should be billed?

- a) 15
- b) 30
- c) 45
- d) 60

43) A patient asks if they can drink alcohol while taking metronidazole. What should the technician do?

- a) Provide counseling on safe limits
- b) Tell them it is okay if they drink only beer
- c) Refer the question to the pharmacist
- d) Suggest skipping doses when drinking

44) A prescription reads: "omeprazole 20 mg daily before breakfast." The technician accidentally selects omeprazole/sodium bicarbonate due to similar names. What is the best next step when the product image and description don't match expectations?

- a) Ignore images; they are often outdated
- b) Stop and verify the drug and formulation match the prescription before continuing
- c) Change the SIG to match the selected product
- d) Dispense and tell the patient to call if it looks different

45) A prescription reads: "ondansetron ODT 4 mg: dissolve 1 tab on tongue q8h prn nausea #12." The technician selected ondansetron 4 mg tablets (non-ODT). What should be done?

- a) Dispense tablets; patient can chew them
- b) Change directions to "swallow whole"
- c) Select the correct dosage form (ODT) before filling
- d) Substitute promethazine tablets

46) A technician drops one tablet on the floor while counting. What is the best action?

- a) Place it back if it appears clean
- b) Discard per policy and recount to ensure accurate quantity
- c) Wipe with alcohol and return it
- d) Put it in a separate vial and let the patient decide

47) A medication label prints for budesonide-formoterol 80/4.5 inhaler, but the selected stock is 160/4.5. What should the technician do?

- a) Dispense the stronger one and instruct fewer puffs
- b) Reprint the label to match the stronger inhaler

- c) Stop and select the correct strength/NDC before continuing
- d) Dispense and annotate "strength substituted" in the profile

48) An IV order is for 1,000 mL to infuse over 8 hours. What is the rate in mL/hr?

- a) 80
- b) 100
- c) 125
- d) 150

49) A patient requests an early refill of alprazolam saying it was lost. Insurance rejects as "too soon." What is the best technician step?

- a) Process as cash automatically since insurance is the issue
- b) Refuse because early refills are never allowed for controls
- c) Offer to transfer to another pharmacy for immediate filling
- d) Gather details and route to the pharmacist for decision-making and documentation

50) A prescriber's office leaves a voicemail for a new Schedule II stimulant prescription. What is the best next step?

- a) Transcribe and fill as a verbal order
- b) Fill a partial supply and request the original later
- c) Use the voicemail as a fax equivalent if caller ID matches
- d) Route to the pharmacist; voicemail is not an acceptable method for issuing a Schedule II prescription

51) A prescription for atorvastatin 40 mg daily is received. The profile shows an active simvastatin 40 mg daily from last month. What should the technician do?

- a) Fill both because dual statins work better
- b) Discontinue simvastatin in the system automatically
- c) Flag for pharmacist review for therapeutic duplication/therapy change
- d) Change atorvastatin to 20 mg to avoid overlap

52) A prescription reads: "metoprolol succinate 50 mg. Take 1 tab bid." The patient previously took metoprolol tartrate 50 mg bid. What is the best technician action?

- a) Enter as written and ignore the history
- b) Change to tartrate to match prior fills
- c) Flag for pharmacist review because formulation/dosing may be inconsistent
- d) Dispense succinate but label it as "tartrate" for clarity

53) A DSCSA inquiry is received requesting transaction history for a suspect bottle of rivaroxaban. What is the best action?

- a) Ignore because DSCSA applies only to controlled substances
- b) Provide required transaction documentation through the pharmacy's DSCSA process and notify the pharmacist/manager
- c) Dispense the bottle quickly to remove it from inventory
- d) Replace the lot number on the bottle with another lot

54) A patient returns from the counter saying they received the wrong strength last week and brought it back unopened. What should be done with the returned bottle?

- a) Remove the patient label and return to stock immediately
- b) Return to stock if sealed to prevent waste
- c) Quarantine and route to the pharmacist for disposition and error documentation
- d) Place it in the will-call bin to be re-dispensed later

55) Which medication is an HMG-CoA reductase inhibitor (statin)?

- a) Losartan
- b) Atorvastatin
- c) Clopidogrel
- d) Pantoprazole

56) A prescription reads: "levETIRAcetam 500 mg: 1 tab po bid." The technician accidentally selects lamotrigine 100 mg. What is the most reliable prevention strategy during product selection?

- a) Verify the patient's diagnosis before filling
- b) Use NDC/barcode verification and match the stock bottle label to the prescription
- c) Store all seizure medications together for easy access
- d) Choose whichever is preferred by the insurance plan

57) A patient with COPD asks what "Spiriva" is on their med list. Which generic corresponds to Spiriva?

- a) Tiotropium
- b) Salmeterol
- c) Ipratropium
- d) Fluticasone

58) Which medication is a proton pump inhibitor?

- a) Famotidine
- b) Omeprazole
- c) Dicyclomine
- d) Hydrochlorothiazide

59) A prescription is written for "clopidogrel." Which brand name matches?

- a) Plavix
- b) Eliquis
- c) Crestor
- d) Coumadin

60) A patient asks which medication is an SSRI. Which is an SSRI?

- a) Sertraline
- b) Bupropion
- c) Duloxetine
- d) Amitriptyline

61) Which medication is a calcium channel blocker commonly used for hypertension?

- a) Lisinopril
- b) Amlodipine
- c) Metoprolol

d) Hydrochlorothiazide

62) Which medication is an ARB?

- a) Losartan
- b) Propranolol
- c) Amlodipine
- d) Furosemide

63) A prescription is for "albuterol HFA." Which brand is albuterol HFA?

- a) Ventolin HFA
- b) Symbicort
- c) Spiriva
- d) Flonase

64) Which medication is a sulfonylurea?

- a) Metformin
- b) Sitagliptin
- c) Glipizide
- d) Acarbose

65) Which medication is a direct oral anticoagulant (factor Xa inhibitor)?

- a) Apixaban
- b) Warfarin
- c) Heparin
- d) Clopidogrel

66) Which medication is a macrolide antibiotic?

- a) Doxycycline
- b) Azithromycin
- c) Amoxicillin
- d) Ciprofloxacin

67) Which medication is a benzodiazepine?

- a) Buspirone
- b) Hydroxyzine
- c) Alprazolam
- d) Zolpidem

68) Which medication is an opioid antagonist used for overdose reversal?

- a) Naloxone
- b) Buprenorphine
- c) Naltrexone
- d) Methadone

69) Which medication is a loop diuretic?

- a) Spironolactone
- b) Furosemide
- c) Hydrochlorothiazide

d) Losartan

70) Which medication is commonly used for hypothyroidism?

- a) Liothyronine
- b) Levothyroxine
- c) Methimazole
- d) Propranolol

71) Which medication is a potassium supplement?

- a) Potassium chloride
- b) Calcium carbonate
- c) Magnesium hydroxide
- d) Sodium bicarbonate

72) Which medication is an H2 blocker?

- a) Pantoprazole
- b) Omeprazole
- c) Famotidine
- d) Sucralfate

73) Which medication is a beta blocker?

- a) Metoprolol
- b) Losartan
- c) Amlodipine
- d) Hydrochlorothiazide

74) Which medication is a tetracycline antibiotic?

- a) Cephalexin
- b) Doxycycline
- c) Azithromycin
- d) Clindamycin

75) Which medication is an antiepileptic?

- a) Levetiracetam
- b) Sertraline
- c) Atorvastatin
- d) Metformin

76) Which medication is a basal insulin?

- a) Insulin lispro
- b) Insulin aspart
- c) Insulin glargine
- d) Regular insulin

77) Which medication is a long-acting anticholinergic inhaler (LAMA)?

- a) Tiotropium
- b) Albuterol
- c) Budesonide-formoterol

d) Montelukast

78) Under federal law, Schedule III and IV controlled substances may be refilled:

- a) Unlimited times within 1 year
- b) Up to 5 refills within 6 months
- c) Up to 3 refills within 90 days
- d) Only with a new prescription each time

79) A prescriber writes for a Schedule III medication on 1/2. The patient presents it for the first time on 7/20 of the same year. Under federal law, it is:

- a) Valid because it is within 1 year
- b) Invalid because it is beyond 6 months
- c) Valid only if the prescriber authorizes by phone
- d) Valid only for a partial fill

80) A technician receives a request to transfer a prescription for clonazepam (Schedule IV) that still has refills remaining. Which is correct under federal rules?

- a) Transfers are never allowed for controlled substances
- b) Schedule IV transfers are allowed one time between pharmacies (unless pharmacies share a real-time online database, where multiple transfers are permitted)
- c) Schedule IV transfers are allowed unlimited times between any pharmacies
- d) Schedule IV prescriptions can only be transferred if no refills remain

81) A patient asks to buy dextromethorphan-containing cough syrup in a state with age restrictions for OTC DXM sales. What is the technician's appropriate action?

- a) Complete the sale without restrictions since it is OTC
- b) Verify age and follow the applicable state restriction requirements
- c) Require a prescription for any DXM product
- d) Refuse all sales of DXM products regardless of age

82) A technician is asked to accept a patient's unused oxycodone for disposal. The pharmacy does not have a DEA-registered collection receptacle and does not run take-back events. What is the best action?

- a) Accept it and place in the hazardous waste bin
- b) Accept it and store in the safe until reverse distribution
- c) Decline acceptance and offer information on authorized disposal options per policy
- d) Accept it only if the patient removes the label first

83) A faxed Schedule II prescription is received from a prescriber's office. Under federal rules, the fax may serve as the original prescription only in:

- a) Any case if the prescriber's signature is on the fax
- b) Any case if the patient has filled there before
- c) Limited situations such as for a patient in a LTCF or for hospice, when requirements are met
- d) No situations; a fax is never allowed for Schedule II

84) A patient requests a copy of their medication profile be emailed to them. What is the best response?

- a) Email it immediately if the address on file matches
- b) Email is never permitted under HIPAA
- c) Provide it only in person; remote requests are prohibited
- d) Verify identity and follow secure communication procedures; if insecure email is used, obtain acknowledgment/consent as required

85) A technician is reconciling a controlled-substance inventory count and finds a discrepancy. What is the best technician action?

- a) Adjust the count to what should be on hand to match records
- b) Set aside the record and recheck later if time allows
- c) Immediately report the discrepancy to the pharmacist/manager per procedure
- d) Call the wholesaler to ask what quantity was shipped

86) A prescription for a controlled substance is presented by an out-of-state prescriber. The technician is unsure if it can be filled. What is the best next step?

- a) Fill it because out-of-state controlled prescriptions are always valid
- b) Refuse immediately; out-of-state controls are never valid
- c) Ask the patient to have the prescriber rewrite it on local prescription paper
- d) Route to the pharmacist to assess validity and state-specific requirements

87) Two similar patient names appear in the queue: "Maria Gonzalez" and "Mariana Gonzalez," same street address, different DOBs. What is the best way to reduce wrong-patient selection during data entry?

- a) Use the address as the primary identifier since it matches
- b) Confirm DOB and compare the prescriber name and medication history before selecting the profile
- c) Select whichever profile has the most recent activity
- d) Merge the profiles to prevent future confusion

88) A high-alert medication warning appears: methotrexate entered as "10 mg daily," but the hard copy says "10 mg once weekly." What should the technician do?

- a) Override the warning because prescribers sometimes use daily dosing
- b) Correct the SIG to once weekly per the prescription and route for pharmacist verification
- c) Dispense with daily directions but verbally tell the patient weekly
- d) Change to 2.5 mg daily to reduce toxicity risk

89) A prescription is written for "lidocaine 5% patch: apply 1 patch daily" with no on/off schedule. What should the technician do?

- a) Default to 24 hours on and proceed
- b) Default to 12 hours on/12 hours off and proceed without asking
- c) Obtain clarification (through pharmacist workflow) before dispensing if required by pharmacy policy/clinical appropriateness
- d) Substitute OTC lidocaine 4% patches and proceed

90) A patient's insurance rejects a claim for "plan limit exceeded." The technician notices the day supply was submitted as 90, but the prescription is for #30 once daily. What should be done first?

- a) Change quantity to 90 to match the submitted day supply

- b) Confirm correct quantity/day supply per the prescription and rebill with corrected values
- c) Split into three separate 30-day claims on the same date
- d) Bypass insurance and run the claim as cash

Answer Key

1) b	31) b	61) b
2) b	32) b	62) a
3) c	33) b	63) a
4) c	34) c	64) c
5) d	35) a	65) a
6) a	36) b	66) b
7) b	37) c	67) c
8) c	38) c	68) a
9) c	39) d	69) b
10) c	40) c	70) b
11) c	41) b	71) a
12) d	42) b	72) c
13) c	43) c	73) a
14) a	44) b	74) b
15) c	45) c	75) a
16) a	46) b	76) c
17) b	47) c	77) a
18) c	48) c	78) b
19) b	49) d	79) a
20) b	50) d	80) b
21) b	51) c	81) b
22) a	52) c	82) c
23) b	53) b	83) c
24) d	54) c	84) d
25) c	55) b	85) c
26) b	56) b	86) d
27) c	57) a	87) b
28) a	58) b	88) b
29) b	59) a	89) c
30) d	60) a	90) b

PTCB Sim Test 21

1) While typing a new eRx, the technician sees “metformin ER 500 mg: take 2 tablets BID.” The patient profile shows metformin IR 500 mg: “take 2 tablets BID,” last filled 5 days ago. What is the best next step?

- a) Enter the ER prescription and discontinue the IR in the profile
- b) Enter the ER prescription but keep the IR active so the patient can choose
- c) Hold the order for pharmacist review due to formulation change/duplication risk
- d) Change the ER to IR to match the patient's existing therapy

2) A customer requests pseudoephedrine 30 mg tablets OTC. Before completing the sale, what must the technician do under federal requirements?

- a) Ask the patient to sign a waiver acknowledging risks
- b) Verify acceptable photo ID and complete the required logbook/electronic log entry
- c) Obtain a written prescription from a prescriber
- d) Limit the sale to one box regardless of package size

3) During filling, the label is for “Augmentin 875/125,” but the technician accidentally grabbed “amoxicillin 875.” Which workflow tool most directly prevents this from reaching the patient?

- a) Barcode scan match of the stock bottle/NDC to the prescription label
- b) Patient signature at pickup
- c) Counting the tablets twice
- d) Printing a Medication Guide

4) Rx: cephalexin 250 mg/5 mL. Sig: “Give 10 mL PO QID x 10 days.” How many mL should be dispensed?

- a) 350 mL
- b) 450 mL
- c) 300 mL
- d) 400 mL

5) Insurance asks for days supply. Albuterol HFA 200 actuations. Sig: “Inhale 2 puffs q4h PRN.” Using maximum frequency, what days supply should be submitted?

- a) 12 days
- b) 8 days
- c) 17 days
- d) 25 days

6) A paper prescription for oxycodone 5 mg has patient and prescriber identifiers, date, drug, and quantity, but directions are missing. What should the technician do?

- a) Enter “use as directed” and proceed
- b) Ask the patient what directions were intended and enter them
- c) Send it back to the patient to obtain a new prescription
- d) Contact the prescriber for directions and document per workflow

7) An isotretinoin prescription is in process and the system blocks dispensing until REMS steps are satisfied. What is the best technician action?

- a) Verify required REMS authorization/steps are complete in the system before proceeding
- b) Override the alert because prescribers handle REMS
- c) Switch to tretinoin topical because it is not restricted
- d) Dispense with a Medication Guide and proceed to sale

8) At pickup, the patient asks: "Can I take ibuprofen with lisinopril?" What should the technician do?

- a) Suggest acetaminophen instead and refuse to sell ibuprofen
- b) Tell the patient to separate doses by at least 2 hours
- c) Refer the question to the pharmacist for counseling
- d) Recommend taking ibuprofen with food to prevent interaction

9) The printed label says "take 1 tablet daily," but the eRx directions read "take 1 tablet twice daily." What should the technician do before filling?

- a) Fill using the label to avoid delays
- b) Correct the sig to match the prescription directions and reprint the label per workflow
- c) Add an auxiliary label "take twice daily" and continue
- d) Fill as is and rely on counseling to correct it

10) The profile lists "sulfa—hives." A new eRx arrives for sulfamethoxazole/trimethoprim. What should the technician do?

- a) Process normally because hives is mild
- b) Change to nitrofurantoin and proceed
- c) Place in pharmacist review due to documented allergy
- d) Ask the patient to sign an allergy waiver

11) At opening, the refrigerator temperature log shows 12°C overnight. Vaccines were stored in this unit. What is the best immediate technician action?

- a) Lower the temperature and continue using the vaccines
- b) Quarantine affected vaccines and alert the pharmacist/manager for excursion evaluation
- c) Move vaccines to the freezer to stabilize them
- d) Discard all vaccines immediately

12) A manufacturer coupon is valid only for brand Crestor. The prescription is written for rosuvastatin 20 mg. What should the technician do?

- a) Keep generic selected and attach the coupon to the bag for pickup
- b) Select brand Crestor in the system and bill according to coupon/coverage requirements, then proceed through verification
- c) Change to atorvastatin 20 mg because it is equivalent
- d) Decline because coupons cannot be used on statins

13) Insulin glargine U-100 pens: "Inject 24 units daily." Dispense 1 box (5 pens × 3 mL each). What days supply should be billed?

- a) 50 days
- b) 75 days
- c) 62 days
- d) 45 days

14) A patient requests a transfer of lorazepam (Schedule IV) with refills remaining from another pharmacy. Under federal law, the transfer must be completed by:

- a) A technician trained on transfers
- b) A pharmacist (or authorized intern where allowed)
- c) The prescriber's office only
- d) The patient through the pharmacy app

15) A patient asks for a refill of alprazolam, but the profile shows 0 refills remaining. What should the technician do next?

- a) Dispense a short emergency supply
- b) Create a new prescription number and send to fill
- c) Send a refill request to the prescriber per workflow
- d) Transfer the prescription to another pharmacy to obtain refills

16) Prednisone 10 mg taper: "Take 2 tablets daily x 5 days, then 1 tablet daily x 5 days." Quantity is missing. How many tablets should be dispensed?

- a) 10
- b) 20
- c) 30
- d) 15

17) DSCSA notice: "Stop distribution" for a specific lot of losartan 50 mg. The technician finds that lot on the shelf. What should be done?

- a) Dispense remaining stock first to prevent waste
- b) Quarantine the lot and follow recall/return instructions
- c) Relabel with a new lot number and keep dispensing
- d) Remove shelf tags but keep the product available

18) During entry, a technician selected clonidine instead of clonazepam from a dropdown list. Which safeguard best reduces this risk?

- a) Store both products next to each other
- b) Rely on counseling at pickup to catch errors
- c) Use Tall Man lettering/alerts and verify name/strength during entry and product selection
- d) Use larger label font only for controlled substances

19) Dorzolamide-timolol ophthalmic: "Instill 1 drop OU BID." Bottle size 10 mL. If 20 drops = 1 mL, what is the days supply for 1 bottle?

- a) 25 days
- b) 50 days
- c) 40 days
- d) 100 days

20) A severe interaction alert appears: patient on warfarin; new Rx for TMP-SMX. What should the technician do?

- a) Override the alert because the prescriber ordered it
- b) Tell the patient to hold warfarin while on the antibiotic
- c) Change to azithromycin and proceed
- d) Route to pharmacist review before dispensing

21) Under federal law, which records must be maintained separately or be readily retrievable?

- a) Schedule II records and Schedule III–V records must be readily retrievable from other records
- b) Only Schedule II records must be separate; Schedule III–V may be mixed
- c) Only Schedule III–V records must be separate
- d) All prescription records must be stored in separate files by class

22) A stock bottle of nitroglycerin SL tablets was opened 8 months ago; the bottle label states “discard 6 months after opening.” Manufacturer expiration is next year. What should the technician do?

- a) Continue using it until manufacturer expiration
- b) Transfer tablets to a new bottle and assign a new discard date
- c) Remove from dispensing stock for proper disposition per workflow
- d) Keep using it but add an auxiliary label on the next fill

23) A hard copy for hydrocodone/acetaminophen is missing the prescriber’s address. What should the technician do?

- a) Fill because the prescriber signature is present
- b) Add the address from an online search without documentation
- c) Have the patient write in the prescriber address
- d) Route for pharmacist handling because required information is incomplete

24) Azithromycin 200 mg/5 mL: “Give 12.5 mL on day 1, then 6.25 mL daily on days 2–5.” Total volume needed:

- a) 37.5 mL
- b) 31.25 mL
- c) 43.75 mL
- d) 50 mL

25) A friend attempts to pick up a patient’s prescription and knows the patient’s DOB and address but is not listed as an authorized pickup person. What should the technician do?

- a) Release the prescription because identifiers match
- b) Release only if the friend pays with the patient’s card
- c) Do not release without proper authorization/verification per workflow
- d) Remove the label from the bag and release the medication

26) While pulling levothyroxine 25 mcg, the technician grabs 125 mcg due to similar packaging. What is the best immediate check before counting?

- a) Compare bottle color to the shelf tag
- b) Verify NDC/barcode match between stock bottle and label
- c) Count tablets first, then compare imprint
- d) Ask another technician to initial the selection

27) Insurance rejects an Adderall XR refill: “refill too soon; next fill date in 5 days.” The patient is traveling tomorrow. Best first technician step:

- a) Change the day supply to force approval

- b) Offer cash price and bypass insurance without documenting
- c) Tell the patient early fills are never allowed
- d) Gather travel details and route to the pharmacist for options (vacation override/plan requirements)

28) The sig reads "Take 1 tablet PO qhs." qhs means:

- a) Every morning
- b) At bedtime
- c) Every hour
- d) Four times daily

29) A label with PHI is left on the counter visible to customers. What should the technician do immediately?

- a) Leave it until the pharmacist notices
- b) Throw it in the regular trash
- c) Remove it promptly and report/document the privacy incident per workflow
- d) Reprint and staple both labels to the hard copy

30) For a nonsterile compound, which action best supports quality and traceability?

- a) Use approximate measurements when close
- b) Document ingredients, lot numbers, quantities, and beyond-use date; clean the area before starting
- c) Skip documentation for small batches
- d) Reuse a returned patient vial if it looks clean

31) An eRx for bupropion SR 150 mg BID is received. The profile shows bupropion XL 150 mg daily from another prescriber. The technician should:

- a) Substitute SR for XL and proceed
- b) Inactivate XL and proceed with SR
- c) Change SR to IR to avoid duplication
- d) Route to pharmacist review due to duplication/formulation concerns

32) A prescriber writes "Norvasc 5 mg daily." What product should be selected?

- a) Omeprazole 5 mg
- b) Lisinopril 5 mg
- c) Amlodipine 5 mg
- d) Atorvastatin 5 mg

33) Lidocaine 5% patches: "Apply 1 patch for 12 hours on, 12 hours off." Dispense #30. What day supply should be billed?

- a) 30 days
- b) 60 days
- c) 45 days
- d) 15 days

34) Under federal law, controlled substance records must generally be kept for at least:

- a) 1 year
- b) 2 years

- c) 3 years
- d) 5 years

35) A Schedule II prescription is partially filled because the pharmacy is out of stock. What must be documented for the partial fill?

- a) Only the quantity owed
- b) Date of partial fill, quantity dispensed, and quantity remaining authorized
- c) Patient diagnosis code and prescriber NPI
- d) Patient signature at the time of partial fill

36) Heparin infusion: 25,000 units in 250 mL. Order: 18 units/kg/hr for a 70 kg patient. What rate (mL/hr) should be set?

- a) 12.6 mL/hr
- b) 18.9 mL/hr
- c) 10.1 mL/hr
- d) 15.2 mL/hr

37) An Augmentin prescription is entered for a patient whose allergy list includes "penicillin— anaphylaxis." The technician should:

- a) Continue because clavulanate reduces allergy risk
- b) Change to cephalexin and proceed
- c) Queue for pharmacist review due to severe allergy risk
- d) Ask the patient if they are sure it was anaphylaxis and proceed if unsure

38) A patient picking up sertraline asks, "Is this for anxiety or depression?" The technician should:

- a) Recommend which condition it treats based on the patient's symptoms
- b) Provide a brief explanation and skip pharmacist counseling since it's common
- c) State it is only for depression
- d) Offer pharmacist counseling for medication questions

39) Docusate 100 mg: "Take 1 capsule PO BID." Quantity for 30 days:

- a) 45
- b) 30
- c) 90
- d) 60

40) Under federal law (assuming state law allows), a Schedule IV prescription may be transferred between pharmacies:

- a) Unlimited times until refills are exhausted
- b) 0 times
- c) 2 times
- d) 1 time

41) A patient requests the same manufacturer for metformin ER due to tolerance, but that NDC is unavailable. What should the technician do?

- a) Dispense metformin IR instead
- b) Select an available equivalent NDC and note the manufacturer change per workflow

- c) Substitute any generic and say nothing
- d) Tell the patient manufacturer requests are never allowed

42) Fluconazole 150 mg: "Take 1 tablet today and repeat in 72 hours." Quantity:

- a) 1
- b) 4
- c) 2
- d) 3

43) A prescriber's office calls in a new verbal prescription for tramadol while the pharmacist is unavailable. The technician should:

- a) Take the order and have the pharmacist sign later
- b) Accept if the caller provides the DEA number
- c) Ask the patient to relay the prescription details
- d) Ask the office to call back when the pharmacist is available or send via permitted transmission method

44) Valacyclovir 1 g: "Take 1 tablet PO BID x 7 days." Quantity:

- a) 21
- b) 14
- c) 7
- d) 10

45) Two will-call bags have similar patient names. Best practice at pickup to reduce wrong-patient errors:

- a) Use two identifiers (e.g., full name and DOB) before release
- b) Confirm last name only
- c) Ask what medication they're picking up and match the bag
- d) Use payment method as the primary identifier

46) A patient requests an early refill for oxycodone stating the bottle was lost. What should the technician do?

- a) Promise an early fill if the patient pays cash
- b) Tell the patient to file a police report and it will be replaced
- c) Offer a lower-strength opioid without prescriber involvement
- d) Collect details and route to the pharmacist without promising an outcome

47) Order: KCl 20 mEq in 100 mL NS. Stock KCl is 2 mEq/mL. How many mL of KCl are needed?

- a) 10 mL
- b) 15 mL
- c) 5 mL
- d) 20 mL

48) The sig reads "Take 1 tablet PO qod." qod means:

- a) Four times daily
- b) Twice daily
- c) Every other day

d) Once weekly

49) Plavix is best described as:

- a) ACE inhibitor used for hypertension
- b) Opioid analgesic for severe pain
- c) Proton pump inhibitor for GERD
- d) Antiplatelet used to reduce risk of stroke/MI

50) Humalog U-100 vial: "Inject 8 units SQ before meals TID." Vial is 10 mL. What days supply for 1 vial?

- a) 25 days
- b) 41 days
- c) 31 days
- d) 125 days

51) A patient asks, "Is carvedilol the same as Coreg?" The best response is:

- a) No, Coreg is a cholesterol medication
- b) Yes, Coreg is the brand name for carvedilol
- c) No, carvedilol is a diuretic
- d) Coreg is generic and carvedilol is brand

52) DUR alert: patient currently taking apixaban. New eRx: rivaroxaban. The technician should:

- a) Continue because dual anticoagulants are commonly combined
- b) Inactivate apixaban and proceed
- c) Route to pharmacist review due to duplication/bleeding risk
- d) Change rivaroxaban to aspirin 81 mg

53) Losartan 50 mg: "Take 1 tablet daily." The plan requests a 90-day fill. Rx is #30 with 2 refills. Quantity to dispense for 90 days (if allowed):

- a) 120
- b) 60
- c) 30
- d) 90

54) A Schedule II prescription is presented dated 2 months ago. Federal law does not set a specific expiration, but state law may. The technician should:

- a) Fill it because federal law allows it
- b) Route to the pharmacist to determine validity under state law/policy before processing
- c) Refuse immediately without review
- d) Call the DEA for approval

55) A shipping tote with refrigerated insulin is found warm after sitting out for several hours. The technician should:

- a) Refrigerate immediately and dispense first to reduce waste
- b) Return to stock if the seal is intact
- c) Quarantine and alert the pharmacist/manager for excursion assessment
- d) Use only for cash-paying patients

56) Enoxaparin 40 mg/0.4 mL syringes: "Inject 0.4 mL SQ daily x 14 days." How many syringes should be dispensed?

- a) 10
- b) 28
- c) 7
- d) 14

57) A written prescription uses "QD." The pharmacy avoids error-prone abbreviations on labels. Best entry approach:

- a) Enter "QD" exactly on the label
- b) Convert to "daily" on the label while preserving the original in the record/image per workflow
- c) Convert to "QID" to avoid confusion
- d) Refuse to fill prescriptions containing abbreviations

58) Which medication commonly has an FDA-required Medication Guide at dispensing?

- a) Metformin
- b) Hydrochlorothiazide
- c) Lamotrigine
- d) Amoxicillin

59) A morphine ER prescription was entered as IR, but the error is caught before the medication is dispensed. This event is best classified as:

- a) Adverse drug event
- b) Drug diversion
- c) Near miss
- d) Therapeutic duplication

60) Pantoprazole corresponds to which brand name?

- a) Protonix
- b) Prevacid
- c) Nexium
- d) Prilosec

61) The pharmacy is returning outdated Schedule II inventory to a reverse distributor. Which task can a technician perform?

- a) Destroy the drugs on-site without witnesses
- b) Assist with pulling items and preparing documentation under pharmacist supervision
- c) Decide which items can be returned without documentation
- d) Adjust controlled inventory counts without review

62) Profile shows simvastatin 40 mg active. New eRx from a different prescriber: atorvastatin 40 mg. The technician should:

- a) Fill both; dual statins are common
- b) Switch atorvastatin to simvastatin to match the profile
- c) Discontinue simvastatin in the system and proceed with atorvastatin
- d) Route to pharmacist review due to potential therapy change/duplication

63) Oral solution order: "Take 2.5 mL PO TID x 10 days." Concentration 100 mg/5 mL. Total mg taken over 10 days:

- a) 3,000 mg
- b) 750 mg
- c) 1,500 mg
- d) 1,000 mg

64) Which statement about Schedule II prescriptions is correct under federal law?

- a) May be refilled up to 5 times in 6 months
- b) May be refilled once if authorized verbally
- c) May not be refilled; a new prescription is required
- d) May be refilled only for established patients

65) A patient requests a printed prescription history to give to a new provider. The technician should:

- a) Provide it to anyone who requests it if they know the DOB
- b) Verify identity and obtain required authorization/consent per workflow before releasing
- c) Refuse because patients cannot receive their records
- d) Fax it to the new provider without consent to speed care

66) Gabapentin Rx: 300 mg "Take 1 capsule TID #90." Patient insists the prescriber said "Take 2 capsules TID." The technician should:

- a) Change directions to match the patient's statement
- b) Enter "1-2 capsules TID" so both are covered
- c) Fill as written and tell the patient to follow the doctor's verbal directions
- d) Route to pharmacist review for possible clarification with prescriber

67) An eRx arrives with the patient's last name off by one letter, but DOB and phone match an existing profile. The technician should:

- a) Merge profiles immediately
- b) Ignore the mismatch if the claim adjudicates
- c) Verify identity with additional identifiers and correct demographics per workflow
- d) Create a new profile and avoid changing anything

68) Which medication is a beta-1 selective blocker commonly used for HTN and rate control?

- a) Hydrochlorothiazide
- b) Metoprolol
- c) Amlodipine
- d) Losartan

69) Amiodarone is most commonly used for:

- a) Arrhythmias
- b) Hypothyroidism
- c) GERD
- d) Bacterial sinusitis

70) In ophthalmic directions, "OS" means:

- a) Both eyes
- b) Left eye
- c) Right eye
- d) Left ear

71) Antibiotic Rx: "Take 1 capsule PO TID x 7 days. Dispense #21." Correct days supply:

- a) 14
- b) 21
- c) 5
- d) 7

72) Which brand/generic pairing is correct?

- a) Celebrex — cephalexin
- b) Flonase — fluoxetine
- c) Zofran — ondansetron
- d) Keppra — cephalexin

73) Profile shows potassium chloride ER. New Rx: spironolactone. The technician should:

- a) Process as normal because they are often used together
- b) Discontinue potassium chloride and fill spironolactone
- c) Route to pharmacist review due to hyperkalemia risk
- d) Change spironolactone to furosemide and proceed

74) During a DEA inspection, controlled substance records must be:

- a) Stored only in a locked cabinet
- b) Destroyed after 12 months to protect privacy
- c) Sent to the DEA monthly
- d) Maintained so they are readily retrievable for inspection

75) A patient presents an out-of-state Schedule II prescription. The technician should:

- a) Refuse immediately because out-of-state CII is always invalid
- b) Process normally without additional review
- c) Ask the prescriber to phone it in instead
- d) Route to the pharmacist to confirm legality and validity under applicable state law before processing

76) A stock bottle has a torn label and the lot number is unreadable; the tablets cannot be confidently identified. The technician should:

- a) Use it if the tablet image in the computer seems to match
- b) Quarantine the bottle and alert the pharmacist; do not dispense from it
- c) Copy the lot number from another bottle of the same medication
- d) Dispense it only for refills to established patients

77) "Synthroid 75 mcg DAW" rejects because the plan prefers generic. The technician should:

- a) Switch to generic levothyroxine and dispense
- b) Call the prescriber to remove DAW without pharmacist involvement

- c) Dispense generic but keep "Synthroid" as the label name
- d) Process as brand/DAW and route per workflow for required authorization/override steps

78) Coreg is the brand name for:

- a) Clonazepam
- b) Carvedilol
- c) Colchicine
- d) Clopidogrel

79) Nasal spray directions: "Use 2 sprays in each nostril daily." Bottle contains 120 sprays.
Days supply:

- a) 15 days
- b) 20 days
- c) 60 days
- d) 30 days

80) In a unit-dose cart fill, hydroxyzine and hydralazine are both stocked. Best method to reduce selection errors:

- a) Store them next to each other for easier comparison
- b) Rely on dosage form differences to prevent errors
- c) Separate storage locations and use barcode scanning during picking
- d) Handwrite labels to force careful reading

81) A technician is asked to "fix" an invalid prescriber DEA by changing digits until the check digit passes. The correct action is to:

- a) Adjust digits to make it pass so the claim can be billed
- b) Use the pharmacy's DEA number instead
- c) Leave the DEA blank and continue
- d) Do not alter identifiers; route to pharmacist to resolve using accurate prescriber information

82) A patient on warfarin asks about starting an OTC herbal product. The technician should:

- a) Recommend avoiding all herbals without exception
- b) Tell the patient it is safe if taken at a different time
- c) Suggest switching to aspirin instead
- d) Offer pharmacist consultation due to interaction potential

83) A patient returns tablets saying they look different and smell odd; the label indicates a manufacturer change. The technician should:

- a) Tell the patient all generics vary and no action is needed
- b) Replace immediately with a different manufacturer without review
- c) Tell the patient to stop the medication immediately
- d) Gather product details and route to the pharmacist to assess possible defect or dispensing error

84) Furosemide is best described as:

- a) SSRI used for depression
- b) Alpha-1 blocker used for BPH

- c) Loop diuretic used for edema and hypertension
- d) Inhaled corticosteroid for asthma maintenance

85) Metronidazole 500 mg BID x 7 days, #14. The technician enters quantity #28 but catches it before the label prints. Best action:

- a) Correct quantity to #14 and continue processing
- b) Leave #28 and rely on final verification to catch it
- c) Change directions to "take QID" to match #28
- d) Dispense #28 and adjust days supply on the claim

86) A patient returns a patient-labeled vial that left the pharmacy yesterday because they "don't need it." In most community settings, the correct handling is:

- a) Return to stock if the seal is intact
- b) Do not return to stock; follow procedure for returns/disposal
- c) Return to stock if tablets appear clean
- d) Repackage into unit dose and return to stock

87) Which prescription may be refilled as written under federal law (assume all other elements are valid)?

- a) Schedule IV zolpidem with 5 refills
- b) Schedule II methylphenidate with 1 refill
- c) Schedule III medication with 6 refills
- d) Schedule V medication with 12 refills

88) IV admixture requires 1.5 g vancomycin. Stock vials are 1 g each. Minimum number of vials needed:

- a) 3 vials
- b) 1 vial
- c) 4 vials
- d) 2 vials

89) New eRx: amlodipine 10 mg with directions "Take 10 tablets daily." The technician should:

- a) Enter as written because technicians cannot question directions
- b) Change to "Take 1 tablet daily" because it is standard dosing
- c) Ask the patient what was intended and enter that
- d) Route for pharmacist review due to likely prescribing/entry error

90) A technician is unsure whether a codeine-containing cough syrup is controlled because formulations differ. Best next step:

- a) Assume it is noncontrolled because it is a cough syrup
- b) Confirm the schedule in the drug file/system; if unclear, route for pharmacist confirmation before processing
- c) Process as Schedule II to be safe
- d) Refuse all codeine cough syrup prescriptions

Answer Key

1) c	31) d	61) b
2) b	32) c	62) d
3) a	33) a	63) c
4) d	34) b	64) c
5) b	35) b	65) b
6) d	36) a	66) d
7) a	37) c	67) c
8) c	38) d	68) b
9) b	39) d	69) a
10) c	40) d	70) b
11) b	41) b	71) d
12) b	42) c	72) c
13) c	43) d	73) c
14) b	44) b	74) d
15) c	45) a	75) d
16) d	46) d	76) b
17) b	47) c	77) d
18) c	48) c	78) b
19) b	49) d	79) c
20) d	50) b	80) c
21) a	51) b	81) d
22) c	52) c	82) d
23) d	53) d	83) d
24) a	54) b	84) c
25) c	55) c	85) a
26) b	56) d	86) b
27) d	57) b	87) a
28) b	58) c	88) d
29) c	59) c	89) d
30) b	60) a	90) b

PTCB Sim Test 22

- 1) An eRx reads: "metoprolol succinate ER 50 mg—take 1 tablet twice daily." What is the most appropriate technician action during data entry?
- a) Enter exactly as written and route for pharmacist verification due to unusual ER directions
 - b) Change to metoprolol tartrate 50 mg twice daily because that's the usual product for BID
 - c) Change directions to once daily because ER is always once daily
 - d) Cancel the prescription and tell the patient it cannot be filled
- 2) A prescription is written for "levothyroxine 0.075 mg daily." Which product should be selected?
- a) Levothyroxine 75 mcg tablets
 - b) Levothyroxine 7.5 mcg tablets
 - c) Liothyronine 75 mcg tablets
 - d) Levothyroxine 0.75 mg tablets
- 3) A patient picking up apixaban (Eliquis) says, "I missed yesterday—should I double up today?" What should the technician do?
- a) Tell the patient to take two tablets now
 - b) Tell the patient to skip today and restart tomorrow
 - c) Offer to have the pharmacist speak with the patient now
 - d) Suggest taking one extra dose tonight
- 4) Prednisone 5 mg tablets: "Take 8 tablets daily x 3 days, then 6 daily x 3 days, then 4 daily x 3 days, then 2 daily x 3 days." How many tablets should be dispensed?
- a) 42
 - b) 54
 - c) 60
 - d) 72
- 5) A customer requests pseudoephedrine 30 mg tablets. Under CMEA, what must be completed before the sale?
- a) Verify government-issued photo ID and obtain required logbook/signature
 - b) Provide counseling and document refusal/acceptance
 - c) Require a prescription if buying more than one box
 - d) Record the buyer's insurance information
- 6) Insulin glargine (Lantus) SoloStar: "Inject 20 units SQ once daily. Dispense 15 mL." How many pens should be dispensed? (Each pen = 3 mL)
- a) 3 pens
 - b) 4 pens
 - c) 5 pens
 - d) 6 pens
- 7) A paper Schedule II prescription is missing the patient's address. What is the best technician action?
- a) Ask the patient to write the address on the prescription

- b) Route to the pharmacist because a required element is missing
- c) Add the address from the patient profile and proceed
- d) Fill it because address is not required for Schedule II

8) During product verification by barcode scan, the NDC scanned does not match the NDC on the label. What should the technician do first?

- a) Override the scan if the drug name matches
- b) Stop and correct the product selection to match the label/prescription
- c) Continue and let the pharmacist fix it at final check
- d) Reprint the label to match the scanned NDC

9) Metformin ER 500 mg: "Take 2 tablets daily with evening meal. Qty 180." What day supply should be entered?

- a) 45 days
- b) 60 days
- c) 75 days
- d) 90 days

10) An eRx is received for amoxicillin/clavulanate. The patient profile shows "Penicillin—
anaphylaxis." What should the technician do?

- a) Process normally because clavulanate reduces allergy risk
- b) Hold in the problem/clarification queue for pharmacist review of allergy risk
- c) Ask the patient if the allergy is "still active" and proceed if unsure
- d) Substitute azithromycin per common practice

11) A nearby pharmacy calls to transfer an active lorazepam prescription with refills. What is the correct technician action?

- a) Provide the transfer information if both stores are in the same chain
- b) Provide transfer information after confirming the caller's license number
- c) Route the request to the pharmacist to complete the transfer
- d) Refuse because controlled substances cannot be transferred

12) A recall notice is received for a specific lot of atorvastatin on your shelf. What should the technician do?

- a) Pull the affected lot from stock and quarantine it for pharmacist/manager handling
- b) Pull all statins from the shelf until the manufacturer clarifies
- c) Keep dispensing until patients report adverse effects
- d) Return all open bottles to the wholesaler immediately

13) Nitroglycerin 0.4 mg SL tablets are dispensed. Which auxiliary label is most appropriate?

- a) "Do not drink grapefruit juice"
- b) "Swallow whole; do not crush"
- c) "Take with food to prevent stomach upset"
- d) "Place under the tongue; do not swallow"

14) A person claiming to be the patient's spouse asks for the patient's full medication list by phone. They are not listed as an authorized representative. What should the technician do?

- a) Provide it after verifying date of birth

- b) Provide only drug names, not directions
- c) Offer to contact the patient directly or obtain documented authorization before releasing PHI
- d) Provide it if the caller knows the patient's address

15) Carvedilol 25 mg: "Take 1 tablet twice daily. Qty 120." What is the day supply?

- a) 30 days
- b) 45 days
- c) 60 days
- d) 120 days

16) A patient asks, "Is Toprol-XL the same as metoprolol tartrate?" What is the best technician response?

- a) "Yes, they're the same—just different brands."
- b) "No, they're different forms; I can have the pharmacist explain."
- c) "Yes, tartrate is the extended-release form."
- d) "No, Toprol-XL is propranolol."

17) A patient presents a Schedule II prescription dated 5 months ago. What is the best technician action?

- a) Process it because federal law has no expiration date for Schedule II
- b) Reject it because Schedule II expires after 30 days everywhere
- c) Route to the pharmacist to determine validity based on state law and professional judgment
- d) Change the written date to today if the patient states it was never filled

18) While stocking, you find unopened latanoprost that has been left at room temperature since yesterday morning. What should the technician do?

- a) Refrigerate and label "use first" without further action
- b) Quarantine and bring to pharmacist/manager to determine if it can be used
- c) Dispense it first before it expires
- d) Discard immediately because any room temperature exposure ruins it

19) Triamcinolone 0.1% cream: "Apply 2 grams topically twice daily. Dispense 80 g." What day supply should be entered?

- a) 10 days
- b) 14 days
- c) 20 days
- d) 40 days

20) An eRx for zolpidem 10 mg arrives with "Refills: 6." What should the technician do?

- a) Enter as written because it is electronic
- b) Change refills to 0 and process the fill
- c) Enter and route for pharmacist review due to legal refill limits for Schedule IV
- d) Split into two prescriptions to make the refills legal

21) Your state restricts OTC dextromethorphan sales to age 18+. A 17-year-old attempts to purchase it. What should the technician do?

- a) Complete the sale because it is OTC
- b) Refuse the sale if age requirement is not met after checking ID
- c) Complete the sale if a parent is present at pickup
- d) Allow the sale if the patient signs a misuse acknowledgment

22) A label prints for hydroxyzine 25 mg, but the product pulled is hydralazine 25 mg. Which safety practice best reduces this risk?

- a) Stock both drugs next to each other for easy comparison
- b) Use Tall Man lettering and separate/storage safeguards for LASA medications
- c) Use only brand-name products for look-alike names
- d) Disable sound-alike alerts to reduce interruptions

23) Ibuprofen 200 mg: "Take 2 tablets every 4 hours as needed." What is the maximum tablets per day based on this schedule?

- a) 10
- b) 12
- c) 8
- d) 6

24) Timolol 0.5% ophthalmic: "Instill 1 drop in each eye twice daily. Dispense 10 mL." If 1 mL = 20 drops, what is the closest day supply?

- a) 25 days
- b) 40 days
- c) 50 days
- d) 100 days

25) Breo Ellipta 100/25: "Inhale 1 puff once daily. Dispense 1 inhaler (30 doses)." What day supply should be billed?

- a) 15 days
- b) 30 days
- c) 45 days
- d) 60 days

26) Which filing method most appropriately meets typical controlled-substance recordkeeping?

- a) File Schedule II prescriptions separately from noncontrolled prescriptions
- b) File all prescriptions together alphabetically by patient last name only
- c) File all prescriptions together by prescriber DEA number
- d) Shred all hard copies after scanning to the system

27) A patient asks to pay cash for an oxycodone prescription "so insurance won't know." What is the best technician action?

- a) Refuse because Schedule II cannot be paid in cash
- b) Process as cash if allowed and refer concerns to the pharmacist if the request seems suspicious
- c) Require a diagnosis code before proceeding
- d) Report the patient to the DEA immediately

28) "Toprol-XL 100 mg once daily" is prescribed. Which generic should be selected?

- a) Metoprolol tartrate 100 mg
- b) Metoprolol succinate ER 100 mg
- c) Propranolol ER 100 mg
- d) Atenolol 100 mg

29) A patient calls and reports a new rash after starting TMP-SMX yesterday. What should the technician do?

- a) Recommend continuing and adding an OTC antihistamine
- b) Recommend stopping immediately and switching to doxycycline
- c) Transfer the call to the pharmacist promptly
- d) Tell the patient mild rash is expected and will resolve

30) Albuterol HFA: "Inhale 2 puffs every 4 hours as needed. Dispense 1 inhaler (200 actuations)." If used at maximum frequency, what day supply is closest?

- a) 12 days
- b) 17 days
- c) 25 days
- d) 8 days

31) A fax from a prescriber office says "OK refill x1" but does not identify the medication or patient and has no signature. What should the technician do?

- a) Add one refill to the most recent medication on the profile
- b) Treat it as invalid and send a proper refill request/clarification
- c) Dispense a 3-day emergency supply and document "office OK"
- d) Use the prescriber's DEA number to validate and proceed

32) Directions: "Take 1/2 tablet twice daily." Tablet strength is 20 mg. How many mg per dose?

- a) 20 mg
- b) 5 mg
- c) 10 mg
- d) 40 mg

33) During a C-II cycle count, the perpetual count is short by 3 tablets. What should the technician do?

- a) Recount and report the discrepancy to the pharmacist/manager for investigation and documentation
- b) Adjust the perpetual count to match the physical count and move on
- c) Borrow tablets from another bottle to balance the count
- d) Wait until the next shipment arrives and recheck then

34) Methotrexate 2.5 mg: "Take 8 tablets once weekly." What is the best technician action during entry?

- a) Enter as daily to avoid missed doses
- b) Enter as written and ensure it is highlighted for high-alert verification
- c) Change to injectable methotrexate because it is safer
- d) Reduce to 1 tablet weekly to lower toxicity risk

35) A clinic requests DSCSA transaction documentation for a bottle of dulaglutide your pharmacy sold to them. What should the technician do?

- a) Provide DSCSA transaction information through the pharmacy's established DSCSA process
- b) Refuse because pharmacies never have DSCSA obligations
- c) Provide the patient's identifying information to prove where it went
- d) Recreate the transaction history from memory and send it

36) Fluconazole 150 mg: "Take 1 tablet today and repeat in 72 hours. Dispense #2." What day supply is typically entered?

- a) 1 day
- b) 6 days
- c) 2 days
- d) 4 days

37) A patient wants to return an unopened bottle of reconstituted amoxicillin suspension dispensed yesterday. What is the correct action?

- a) Accept and restock if it looks sealed
- b) Accept and relabel for the next patient
- c) Accept for disposal/credit if allowed, but do not restock dispensed medication
- d) Convert it back to powder and restock

38) Vancomycin order: 1 g IV. Available: 500 mg/10 mL. How many mL are needed?

- a) 10 mL
- b) 15 mL
- c) 25 mL
- d) 20 mL

39) A prescription says "Azithromycin Z-Pak: use as directed." What should the technician do?

- a) Enter "use as directed" and process
- b) Enter standard Z-Pak directions without clarification
- c) Obtain/confirm complete directions that will appear on the label before processing
- d) Bill as 30 days because it's "as directed"

40) An eRx is for ondansetron ODT 4 mg. ODT is out of stock, and tablets are available. What is the best technician next step?

- a) Substitute tablets and add "dissolve on tongue"
- b) Hold and route for pharmacist review regarding substitution/availability options
- c) Dispense tablets because strength is the same
- d) Split fill: some ODT now, some tablets later

41) The pharmacy must dispose of an expired partial bottle of morphine oral solution from an emergency kit. Which action is correct?

- a) Pour into the sink with water running
- b) Place in regular trash mixed with cat litter
- c) Dispose using the controlled-substance waste procedure with required

documentation/witnessing

d) Return to the wholesaler for credit

42) The label prints "prednisone 10 mg: take 1 tablet daily," but the eRx sig is "take 4 tablets daily for 5 days." What should the technician do?

- a) Dispense and tell the patient to follow the doctor's instructions
- b) Correct the label to match the prescription and route for verification
- c) Change only the quantity and keep the printed directions
- d) Leave it because prednisone regimens vary

43) Which item commonly requires refrigeration while unopened in the pharmacy?

- a) Omeprazole DR capsules
- b) Latanoprost ophthalmic solution
- c) Clopidogrel tablets
- d) Amoxicillin capsules

44) Amoxicillin/clavulanate 400 mg/5 mL: "Give 7.5 mL twice daily for 10 days." What minimum quantity should be dispensed?

- a) 150 mL
- b) 180 mL
- c) 120 mL
- d) 200 mL

45) Why is "QD" discouraged in prescription directions?

- a) It is not billable to insurance
- b) It can be misread as "QID"
- c) It is only permitted for controlled substances
- d) It is only allowed in inpatient orders

46) New eRx: sertraline 100 mg daily. Profile shows active fluoxetine 20 mg daily filled last week. What should the technician do?

- a) Enter sertraline and inactivate fluoxetine
- b) Enter sertraline and route for pharmacist review due to potential duplication/transition issue
- c) Enter both and schedule automatic refills for both
- d) Change sertraline to 50 mg and process

47) Nitrofurantoin monohydrate/macrocrystals 100 mg is prescribed. The technician selects nitrofurantoin macrocrystals 100 mg. What should the technician do?

- a) Continue because both are nitrofurantoin 100 mg
- b) Correct the selection because formulations are not automatically interchangeable
- c) Change the sig to once daily to match the chosen product
- d) Substitute TMP-SMX because it's equivalent for UTI

48) A nurse requests a patient's controlled-substance fill history for med reconciliation, but the caller cannot be verified with approved procedures. What should the technician do?

- a) Provide it because it is for patient care
- b) Provide only the last fill date

- c) Use approved verification/callback procedures before releasing any information
- d) Refuse all requests from healthcare facilities under HIPAA

49) A patient requests an early refill of alprazolam stating it was stolen, with no police report. What should the technician do?

- a) Process an early refill override automatically
- b) Inform the patient early refills are never allowed for any controlled substance
- c) Gather details and route to the pharmacist for evaluation/documentation
- d) Transfer the prescription to another pharmacy

50) Lantus SoloStar: "Inject 24 units at bedtime. Dispense 5 pens." Each pen is 3 mL of 100 units/mL. What day supply is closest?

- a) 50 days
- b) 62 days
- c) 75 days
- d) 90 days

51) A patient asks the pharmacy to email their prescription history to a personal email address. What should the technician do?

- a) Email it after verifying two patient identifiers
- b) Email it if the patient consents verbally
- c) Provide a printed copy or use an approved secure method per workflow
- d) Refuse because patients cannot request their own records

52) Several vials of refrigerated insulin were left out overnight. The temperature log shows 78°F. What is the best action?

- a) Put back in the refrigerator and use first
- b) Quarantine and notify the pharmacist/manager to assess usability per product/storage guidance
- c) Dispense only to patients who use insulin quickly
- d) Return to stock because room temperature is acceptable indefinitely

53) Omeprazole 20 mg: "Take 1 capsule twice daily. Qty 60." Insurance rejects "max daily dose exceeded—PA required." What should the technician do next?

- a) Change directions to once daily so the claim pays
- b) Split into two claims of 30 each
- c) Inform the patient of the rejection and route for prescriber follow-up/PA or alternatives
- d) Bypass insurance and process cash without telling the patient

54) "Wellbutrin SR 150 mg" is prescribed. Which is the correct generic selection?

- a) Buspirone 15 mg
- b) Buprenorphine/naloxone 8 mg/2 mg
- c) Bupropion XL 150 mg
- d) Bupropion SR 150 mg

55) A liquid medication: "Take 5 mL three times daily for 7 days." How many mL should be dispensed?

- a) 120 mL

- b) 90 mL
- c) 105 mL
- d) 75 mL

56) A technician is unit-dose packaging an oral hazardous medication on the facility's NIOSH list. Which action is appropriate?

- a) Package without special precautions because it is oral
- b) Use required PPE/engineering controls and follow the hazardous-drug workflow
- c) Crush tablets to reduce breakage and speed packaging
- d) Remove it from the hazardous list if it is commonly used

57) A patient asks to transfer an unfilled Schedule II prescription from another pharmacy because they are out of stock. Which is correct?

- a) It can be transferred once between pharmacies
- b) It can be transferred within the same chain only
- c) It cannot be transferred; a new prescription must be issued to the receiving pharmacy
- d) It can be transferred if the patient pays cash

58) A patient on rivaroxaban (Xarelto) asks whether they should keep taking aspirin 81 mg they started on their own. What should the technician do?

- a) Tell them to stop aspirin immediately
- b) Tell them to continue aspirin with Xarelto
- c) Suggest taking aspirin every other day
- d) Refer the question to the pharmacist for counseling

59) Potassium chloride oral solution is prescribed. Which technician step best prevents a dosage-form error?

- a) Select tablets because they are cheaper and equivalent
- b) Confirm dosage form during entry and selection; do not substitute dosage forms without pharmacist involvement
- c) Change directions to "dissolve tablets in water" and proceed
- d) Reduce dose because solution absorbs faster

60) A patient brings a sealed sharps container to the pharmacy for disposal. What is the safest appropriate technician action?

- a) Open it to confirm contents
- b) Place it in regular trash if sealed
- c) Use the pharmacy's approved sharps disposal process if the site accepts them
- d) Recap loose needles and transfer to a new container

61) Profile lists "codeine—severe nausea/vomiting." A new prescription arrives for hydrocodone/APAP. What should the technician do?

- a) Fill without any action because nausea is not an allergy
- b) Delete the codeine reaction from the profile
- c) Route for pharmacist review due to opioid intolerance history and counseling considerations
- d) Substitute tramadol automatically

62) Depakote ER 500 mg daily is prescribed. The patient previously used divalproex DR 500 mg. What should the technician do?

- a) Substitute DR for ER because they are equivalent
- b) Change to valproic acid immediate-release to save money
- c) Process as written and ensure pharmacist verification because ER vs DR are not automatically interchangeable
- d) Change the sig to twice daily to match prior DR dosing

63) A prescription reads "diltiazem 120 mg CD daily." Which entry approach is best?

- a) Enter diltiazem IR 30 mg four times daily because CD is unclear
- b) Enter an appropriate extended-release "CD" formulation matching intent and strength
- c) Substitute verapamil SR 120 mg because it's the same class
- d) Enter diltiazem LA 240 mg because it is once daily

64) Isotretinoin is at pickup. What must be confirmed before dispensing?

- a) Patient signs a counseling waiver
- b) Prior authorization is on file
- c) REMS/iPLEDGE requirements are satisfied for patient, prescriber, and pharmacy
- d) Medication is dispensed in amber glass only

65) A shipment arrives with several cracked glass ampules. What should the technician do?

- a) Stock the unbroken ones and discard the cracked ones in regular trash
- b) Tape cracked ampules and set aside to use first
- c) Quarantine damaged items and notify pharmacist/manager to arrange credit/replacement
- d) Wipe ampules with alcohol and stock them

66) A technician accidentally selects digoxin 0.25 mg instead of 0.125 mg during entry. Which safeguard most directly helps prevent this from reaching the patient?

- a) Disabling DUR alerts to reduce interruptions
- b) Barcode scanning during product verification plus pharmacist final check
- c) Storing all strengths together to speed selection
- d) Printing labels in large batches

67) A prescription for promethazine with codeine syrup is presented for a 16-year-old. What should the technician do?

- a) Fill as written if the dose seems appropriate
- b) Substitute promethazine-DM automatically
- c) Route to the pharmacist due to age-related safety/appropriateness concerns
- d) Ask the parent to sign a sedation waiver and dispense

68) Rosuvastatin 40 mg daily is received. Profile shows active atorvastatin 80 mg daily filled recently. What should the technician do?

- a) Fill both because they are different statins
- b) Automatically discontinue atorvastatin and fill rosuvastatin
- c) Change rosuvastatin to 10 mg to reduce duplication
- d) Route for pharmacist review due to likely therapy change/duplication risk

69) An inpatient order says "heparin infusion per protocol." The chart lists a history of HIT.

What should the technician do?

- a) Enter and proceed because it says "per protocol"
- b) Hold and escalate promptly for pharmacist review due to documented HIT history
- c) Substitute enoxaparin per protocol
- d) Delete the HIT entry because it is not an allergy

70) A patient picking up gabapentin asks if it can cause drowsiness while driving. What should the technician do?

- a) Tell them it does not affect driving
- b) Recommend taking all doses at bedtime
- c) Suggest caffeine with each dose
- d) Refer to the pharmacist for counseling

71) A technician is restocking medications returned from will-call. What is best practice?

- a) Restock everything to reduce waste
- b) Restock controlled substances without documentation
- c) Verify integrity and storage conditions; restock only eligible items per workflow
- d) Re-label returned items with a new patient label

72) A prescription is for clonidine 0.1 mg patch weekly, but tablets were pulled. Which process best catches this earlier?

- a) Rely on tablet color and imprint
- b) Compare dosage form at data entry and product selection; use barcode scanning if available
- c) Store patches with tablets to reduce walking time
- d) Skip image verification to improve speed

73) A prescription reads: "Take 1 capsule PO QD. Qty 30." How should "QD" be interpreted?

- a) Once daily
- b) Every other day
- c) Four times daily
- d) Twice daily

74) Amiodarone 200 mg: "Take 1 tablet twice daily for 14 days, then 1 tablet daily. Dispense #42." What is the primary issue?

- a) Quantity is too high for directions
- b) Quantity is too low for directions
- c) Directions and quantity match perfectly
- d) Duration of maintenance phase is unclear and affects total day supply

75) A patient asks for a copy of their own prescription profile information. What is an appropriate technician action?

- a) Refuse because PHI belongs to the pharmacy
- b) Provide access through permitted channels using the pharmacy's records request process
- c) Provide only if requested by an attorney
- d) Provide only after a mandatory HIPAA fee is paid upfront

76) Cephalexin 500 mg: "1 capsule four times daily x 7 days." A profile note indicates "renal dosing adjustments commonly needed—pharmacist to review." What should the technician do?

- a) Process and fill as written without delay
- b) Change to twice daily to be safer
- c) Route for pharmacist review before dispensing
- d) Dispense half now and half later

77) DEA Form 222 is used for ordering which of the following?

- a) Schedule I and II substances
- b) Schedule II substances
- c) Schedule III–V substances
- d) Any controlled substance when ordering from a wholesaler

78) The label prints "morphine ER 15 mg," but the prescription is for "morphine IR 15 mg." What should the technician do?

- a) Dispense ER because it lasts longer
- b) Add "do not crush" and proceed
- c) Stop, correct the selection, and send for verification before continuing
- d) Change directions to every 12 hours and dispense ER

79) While preparing a nonsterile compound, the concentration on the label does not match the master formula record. What should the technician do first?

- a) Compound according to the label since that is what the patient sees
- b) Stop and bring the discrepancy to the pharmacist before compounding
- c) Average the two concentrations and proceed
- d) Prepare both concentrations and let the pharmacist choose later

80) A Medicare Part D claim shows the patient is in the coverage gap and the copay increased. What is the best technician response?

- a) Reverse and rebill until the copay changes
- b) Explain generally what it means and offer pharmacist help for plan-specific questions
- c) Tell the patient insurance is not discussed in the pharmacy
- d) Rebill as cash automatically to lower cost

81) Lorazepam 1 mg: "Take 1 tablet three times daily as needed. Qty 90. Refills 4." What is the best technician action?

- a) Enter as written because Schedule IV can have unlimited refills
- b) Change refills to 0 because benzodiazepines are Schedule II
- c) Route for pharmacist review because refills may exceed legal limits for Schedule IV
- d) Refuse to process any PRN benzodiazepine prescription

82) New prescription: lisinopril 20 mg daily. Profile shows active enalapril 10 mg daily filled recently. What should the technician do?

- a) Fill both to improve BP
- b) Inactivate enalapril and fill lisinopril without review
- c) Route for pharmacist review due to possible duplicate ACE inhibitor therapy/transition
- d) Change enalapril to 20 mg to match lisinopril

83) Patient reports discharge medication Brilinta last week, but today's eRx is clopidogrel 75 mg daily. What should the technician do?

- a) Fill clopidogrel and tell the patient to stop Brilinta
- b) Fill both because dual therapy is always required
- c) Route for pharmacist review to assess transition and duplication risk
- d) Automatically discontinue Brilinta and fill clopidogrel

84) In sterile compounding, a technician realizes hand hygiene documentation is incomplete before entering the cleanroom. What should the technician do?

- a) Enter anyway if gloves are worn
- b) Complete required hand hygiene/garbing steps and documentation before entering
- c) Ask another technician to sign the log
- d) Use extra alcohol on gloves instead of documenting

85) A paper prescription for lisdexamfetamine (Vyvanse) is missing the prescriber's phone number but includes name, address, DEA, and signature. What should the technician do?

- a) Fill it because the phone number is optional
- b) Add the phone number from an online directory and proceed
- c) Route for pharmacist review because the prescription may be missing required elements per state rules
- d) Ask the patient to supply the prescriber phone number and write it on the prescription

86) The prescription says "take 1 tablet every other day," but the label prints "take 1 tablet daily." What should the technician do?

- a) Dispense as daily because it is clearer
- b) Correct the label to match the prescription and route for verification
- c) Change to "take 1 tablet three times weekly" and dispense
- d) Leave it and counsel the patient at pickup

87) A patient says their tablets look different than last month. NDC shows a different manufacturer but same drug/strength. What should the technician do?

- a) Tell the patient it must be counterfeit
- b) Confirm the manufacturer/NDC change and offer pharmacist counseling if concerns remain
- c) Report to the DEA before dispensing
- d) Replace with a different strength that matches old appearance

88) A prescription is written for furosemide 40 mg daily, but the profile shows torsemide 20 mg daily filled recently with no discontinuation note. What should the technician do?

- a) Fill furosemide and inactivate torsemide automatically
- b) Cancel the new prescription as a duplicate
- c) Convert to torsemide 40 mg and fill as equivalent
- d) Route for pharmacist review due to possible therapy change/duplication

89) A patient on Bactrim DS asks if they can drink alcohol while taking it. What should the technician do?

- a) Tell them alcohol is always contraindicated with every antibiotic

- b) Tell them it is always safe with antibiotics
- c) Refer the question to the pharmacist for counseling
- d) Recommend skipping doses when drinking

90) A prescription reads: "phenytoin 100 mg capsules: take 3 capsules daily. DAW." The patient asks for the least expensive generic. What should the technician do?

- a) Substitute the generic automatically because it is bioequivalent
- b) Change the DAW code to allow substitution
- c) Route to the pharmacist to discuss DAW, cost options, and possible prescriber contact
- d) Tell the patient DAW means insurance will not cover it

Answer Key

1) a	31) b	61) c
2) a	32) c	62) c
3) c	33) a	63) b
4) d	34) b	64) c
5) a	35) a	65) c
6) c	36) d	66) b
7) b	37) c	67) c
8) b	38) d	68) d
9) d	39) c	69) b
10) b	40) b	70) d
11) c	41) c	71) c
12) a	42) b	72) b
13) d	43) b	73) a
14) c	44) b	74) d
15) c	45) b	75) b
16) b	46) b	76) c
17) c	47) b	77) b
18) b	48) c	78) c
19) c	49) c	79) b
20) c	50) b	80) b
21) b	51) c	81) c
22) b	52) b	82) c
23) b	53) c	83) c
24) c	54) d	84) b
25) b	55) c	85) c
26) a	56) b	86) b
27) b	57) c	87) b
28) b	58) d	88) d
29) c	59) b	89) c
30) d	60) c	90) c

PTCB Sim Test 23

1) A new eRx arrives for metformin ER 500 mg: "Take 2 tablets daily with evening meal. #180." The patient profile shows metformin IR 500 mg "1 tab BID" with refills remaining. What is the most appropriate technician action?

- a) Discontinue the IR prescription in the profile and process the ER prescription
- b) Enter the ER prescription as written and flag as a formulation change for pharmacist review
- c) Change ER to IR because the total daily dose is the same
- d) Fill both prescriptions to avoid therapy interruption

2) A prescriber sends "levothyroxine 0.1 mg PO daily." Which product strength should be selected?

- a) 0.01 mg
- b) 10 mcg
- c) 100 mcg
- d) 1 mg

3) During data entry, an allergy alert shows "Sulfa—hives." The new prescription is for sulfamethoxazole/trimethoprim DS. What should the technician do next?

- a) Continue processing; sulfa allergy does not apply to antibiotics
- b) Change to nitrofurantoin and proceed
- c) Remove the allergy since it is nonspecific
- d) Stop processing and route for pharmacist evaluation of the allergy conflict

4) A patient requests an early refill on zolpidem 10 mg (Schedule IV). The refill is 12 days early and the patient states it was "lost." What is the best technician action?

- a) Process the refill because it is not Schedule II
- b) Offer a weekend supply without documentation
- c) Document the request details and route to the pharmacist for a decision
- d) Transfer the prescription to another pharmacy for early fill

5) Order: "Infuse 1,000 mL NS over 8 hours." What is the infusion rate in mL/hr?

- a) 80 mL/hr
- b) 125 mL/hr
- c) 160 mL/hr
- d) 1,000 mL/hr

6) A patient asks for the generic for Crestor. Which medication should the technician select?

- a) atorvastatin
- b) rosuvastatin
- c) simvastatin
- d) pravastatin

7) A refrigerated vaccine shipment arrives. The temperature-monitor tag indicates it exceeded the acceptable range during transit. What should the technician do?

- a) Put it in the refrigerator immediately and use it first

- b) Return it to stock if the vials look intact
- c) Quarantine the shipment and follow the excursion/return process for disposition
- d) Discard it immediately without documenting lot numbers

8) A paper prescription for alprazolam 1 mg has the prescriber's signature but is missing the patient's address. What is the best next step?

- a) Add the address from the patient profile and proceed
- b) Hold and obtain the missing required information through appropriate verification before dispensing
- c) Fill it; address is required only for Schedule II
- d) Replace the missing information using a previous prescription image

9) Amoxicillin suspension is prescribed: "Take 7.5 mL PO BID x 10 days." How many mL should be dispensed?

- a) 75 mL
- b) 100 mL
- c) 150 mL
- d) 200 mL

10) The label prints "metoprolol succinate ER 50 mg daily." The technician pulled metoprolol tartrate 50 mg. What should the technician do?

- a) Dispense tartrate; both are metoprolol
- b) Change directions to "BID" to match tartrate
- c) Ask the patient which one they prefer
- d) Stop and obtain the correct ER formulation before continuing

11) A patient asks to transfer a Schedule II prescription for amphetamine salts from a different pharmacy to yours. Under federal rules, which is correct?

- a) It may be transferred once if both pharmacies use the same software
- b) It may be transferred only if it has not been filled
- c) It cannot be transferred between pharmacies for dispensing
- d) It can be transferred if the patient signs a release form

12) A label reads: "Azithromycin 250 mg: take 2 tablets on day 1, then 1 tablet daily days 2–5." How many tablets are needed?

- a) 5
- b) 6
- c) 7
- d) 8

13) A prescriber's office calls to authorize 2 refills for lisinopril 20 mg. What is the best technician action?

- a) Add refills and document without any pharmacist involvement
- b) Tell the caller to resend as an eRx only
- c) Record the authorization details and route for pharmacist approval per workflow/state rules
- d) Refuse because refills cannot be authorized by phone

14) The prescription says "Latuda 40 mg daily." Which generic is Latuda?

- a) lurasidone
- b) lamotrigine
- c) lithium carbonate
- d) duloxetine

15) A new prescription for warfarin is being entered. The system flags a new prescription for trimethoprim-sulfamethoxazole on the same profile. What is the most appropriate technician action?

- a) Proceed; interaction checks are handled at pickup
- b) Add a note and route for pharmacist review due to high-risk interaction
- c) Tell the patient to avoid leafy greens while on the antibiotic
- d) Change the antibiotic to cephalexin and proceed

16) A cream is prescribed: "Apply 2 g topically BID." Tube size is 60 g. What is the days' supply?

- a) 10 days
- b) 12 days
- c) 15 days
- d) 30 days

17) An eRx is received for clonazepam 0.5 mg #30 with 6 refills. What is the federal issue?

- a) Schedule IV prescriptions cannot have any refills
- b) Refills are limited to 5 within 6 months for Schedule III–IV
- c) Quantity for Schedule IV is limited to #15
- d) Clonazepam is Schedule II

18) A customer buys pseudoephedrine 30 mg tablets. Which requirement applies under CMEA?

- a) A prescription is required for all pseudoephedrine products
- b) Only a signature is required; ID is optional
- c) Verify acceptable photo ID and complete required sales log documentation
- d) Record the prescriber's DEA number in the log

19) A patient asks whether morphine and oxycodone are "the same strength." What should the technician do?

- a) Provide a conversion using a standard equianalgesic chart
- b) Tell the patient oxycodone is always stronger than morphine
- c) Refer the question to the pharmacist for counseling
- d) Recommend cutting the dose in half until they know

20) A LASA alert appears for hydroXYzine vs hydrALAZINE. The eRx is "hydroxyzine 25 mg TID prn itching." What is the best technician verification step?

- a) Confirm the indication and directions align with the selected product before proceeding
- b) Ignore the alert if the patient has filled hydroxyzine before
- c) Substitute hydralazine if hydroxyzine is out of stock
- d) Verify only the prescriber's phone number matches the clinic website

21) At pickup, someone in line asks, "What medicine is she getting?" What is the best technician response?

- a) Answer quietly so only the bystander hears
- b) Provide only the drug class, not the name
- c) Decline to disclose any information and maintain privacy
- d) Confirm the name only if the bystander knows the patient's DOB

22) A patient wants two bottles of nitroglycerin SL because "I lose them." The prescription is for one bottle, no refills. What should the technician do?

- a) Increase quantity and document "patient request"
- b) Split one bottle into two vials and label both
- c) Dispense a second bottle cash without changing the prescription quantity
- d) Route the request appropriately because dispensing additional quantity changes what is being provided

23) A vaccine diluent labeled "Refrigerate. Do not freeze." is found frozen overnight. What should the technician do?

- a) Thaw and use today
- b) Quarantine and escalate for manufacturer stability guidance/disposition
- c) Return it to the refrigerator and use first
- d) Relabel with a shorter beyond-use date

24) Prednisone 10 mg taper: 4 tabs daily x 3 days, then 3 tabs daily x 3 days, then 2 tabs daily x 3 days, then 1 tab daily x 3 days. What quantity is needed?

- a) 24
- b) 30
- c) 36
- d) 40

25) Which statement is correct about federal recordkeeping for Schedule II inventory records?

- a) They must be maintained separately from all other records
- b) They may be filed with other controlled-substance records if readily retrievable
- c) They may be filed with noncontrolled records if readily retrievable
- d) They are required only for pharmacies that dispense more than 100 controlled prescriptions per day

26) An eRx arrives for glipizide XL 10 mg daily. The profile shows glimepiride 2 mg daily active. What should the technician do?

- a) Enter and fill both; they are different medications
- b) Hold and route for pharmacist review for possible therapeutic duplication
- c) Change glipizide XL to metformin XR and proceed
- d) Inactivate glimepiride automatically and continue

27) A patient calls: "My antibiotic suspension says refrigerate. I left it out 10 hours." What should the technician do?

- a) Tell them it is always fine as long as it was capped
- b) Tell them to discard it immediately and refill

- c) Refer to the pharmacist because stability varies by product and time/temperature
- d) Tell them to reconstitute it again with water

28) Albuterol HFA contains 200 puffs. Sig: "Inhale 2 puffs q4h prn." If used at the maximum frequency around the clock, what is the days' supply?

- a) 8.3 days
- b) 10 days
- c) 12.5 days
- d) 16.7 days

29) An isotretinoin prescription is dropped off by a patient who states she is pregnant. What is the best technician action?

- a) Proceed; pregnancy counseling occurs at pickup
- b) Ask the patient to sign a waiver and continue
- c) Stop processing and route immediately for pharmacist handling due to REMS/pregnancy risk
- d) Dispense a 7-day supply only

30) A prescription is written for cephalexin 500 mg capsules, but the technician pulled 500 mg tablets due to adjacent stock placement. What should be done?

- a) Dispense tablets because strength is the same
- b) Change the label to "tablets" and proceed
- c) Stop and correct to the prescribed dosage form; obtain authorization if a change is needed
- d) Open capsules and repackage into a vial marked "tablets"

31) A patient requests a printed copy of their refill history. What is the best technician action?

- a) Print and hand it over without verification
- b) Verify identity and release only through the authorized process
- c) Refuse because refill history is always confidential to prescribers
- d) Fax it to the prescriber without patient authorization

32) Two oral morphine liquids are stocked: 20 mg/mL concentrate and 10 mg/5 mL solution. Which practice best reduces selection errors?

- a) Store next to each other to keep morphine together
- b) Separate and clearly label concentrations; use barcode scanning during filling
- c) Remove barcode scanning to avoid nuisance alerts
- d) Keep both in will-call bins to reduce stock selection errors

33) An eRx product field shows "sertraline oral concentrate," but the directions say "take 1 tablet daily," and the patient has only used tablets. What should the technician do?

- a) Change to tablets and process because the sig says tablet
- b) Dispense concentrate and keep "tablet" in the directions
- c) Proceed with concentrate and adjust the directions to 5 mL daily
- d) Hold for clarification because product and directions conflict

34) A returned-to-stock vial from last week has a broken safety seal and appears opened. What is appropriate?

- a) Restock after recounting tablets
- b) Restock if tablets look intact
- c) Do not restock; treat as nonreturnable once dispensed/opened
- d) Restock only if the patient signs a return form

35) A claim rejects “refill too soon” for a maintenance inhaler. The patient reports improper priming and early depletion. What should the technician do?

- a) Reduce the days’ supply and resubmit without notes
- b) Use any override code to force approval
- c) Use the appropriate override/prior authorization process and document the reason as required
- d) Tell the patient to wait until the next fill date; no options exist

36) A patient asks, “Is clopidogrel the same as Plavix?” What is correct?

- a) Yes, clopidogrel is the generic for Plavix
- b) No, Plavix is the generic for clopidogrel
- c) They are different drugs but both are antiplatelets
- d) Only certain manufacturers are equivalent

37) A patient asks the pharmacy to “post-date” a Schedule II prescription so it can be filled next week. What should the technician do?

- a) Change the fill date in the computer and process
- b) Add “do not fill until” on the hardcopy and file it
- c) Route for pharmacist handling; only the prescriber can issue proper instructions for future fill dates
- d) Partially fill now without documenting and finish next week

38) In unit-dose packaging, a prepared blister card label is missing the lot number. What should the technician do?

- a) Use it; lot numbers are optional if expiration is present
- b) Write a guessed lot number based on similar stock
- c) Use it only for internal floor stock
- d) Set it aside and obtain properly labeled product to preserve traceability

39) An eRx for losartan 50 mg daily arrives. The profile shows lisinopril 20 mg active with a note “ACE inhibitor cough.” What should the technician do?

- a) Inactivate lisinopril and process losartan without review
- b) Hold and route because it likely represents a therapy change needing pharmacist assessment/documentation
- c) Delete the cough note to prevent future alerts
- d) Change losartan to lisinopril to match the profile

40) Eye drops: “Instill 1 drop OU BID.” Bottle size 10 mL. Assume 20 drops/mL. What is the days’ supply?

- a) 25 days
- b) 50 days
- c) 75 days
- d) 100 days

41) Fentanyl 25 mcg/hr patches: "Apply 1 patch every 72 hours. Dispense #10." What is the days' supply?

- a) 10
- b) 20
- c) 30
- d) 72

42) An established patient's new eRx has the correct name and address, but DOB differs by one year from the profile. What should the technician do?

- a) Update the profile DOB to match the eRx and continue
- b) Create a second profile to avoid delays
- c) Pause and verify the correct DOB using reliable identifiers before processing
- d) Ignore it because address matches

43) Which is the generic for Eliquis?

- a) rivaroxaban
- b) apixaban
- c) dabigatran
- d) edoxaban

44) During a safe audit, an OTC bottle of acetaminophen is found in the controlled-substance safe. What should the technician do?

- a) Leave it since the safe is more secure
- b) File a DEA Form 106 because it was in the safe
- c) Quarantine it until the next DEA inspection
- d) Remove it and store correctly; no controlled-substance reporting is required

45) A Class I recall is issued for specific lots of a medication due to contamination. What should the technician do first with affected inventory?

- a) Continue dispensing until replacement arrives
- b) Quarantine affected lots and follow recall instructions for returns/credit
- c) Repackage into smaller vials to reduce waste
- d) Dispense only to established patients

46) A label prints for potassium chloride IVPB. Which is the safest technician action?

- a) Bypass checks because IVPB labels are standardized
- b) Ensure required high-alert verification steps are completed before it leaves the pharmacy
- c) Send the vials to the unit to mix without pharmacy verification
- d) Increase the diluent volume to reduce irritation without an order

47) A patient asks, "Is omeprazole for heartburn or blood pressure?" What should the technician do?

- a) Tell them it is for blood pressure and take at bedtime
- b) Tell them it is for heartburn and take with grapefruit juice
- c) Refer to the pharmacist for counseling on indication and proper use
- d) Tell them to stop it if symptoms improve

48) A prescription for hydrocodone/acetaminophen says "1–2 tabs q4–6h prn pain," quantity #30, with no maximum daily dose. What is the best technician action?

- a) Enter as "1 tab q6h" to create a clear day supply
- b) Enter as written and flag/route because ambiguity affects safety, billing, and controlled-substance compliance
- c) Dispense and let the patient decide how to take it
- d) Change to "1–2 tabs q6h" since that is common

49) While restocking a crash cart, the technician finds an epinephrine syringe expired yesterday. What should be done?

- a) Keep it because it is only one day expired
- b) Move it behind the newer syringes and leave it available
- c) Remove it from use and replace with in-date product
- d) Send it to will-call to avoid waste

50) Sig: "Take 1.5 tablets daily." Quantity dispensed is #45. What is the days' supply?

- a) 20
- b) 25
- c) 30
- d) 45

51) A spouse calls and asks, "Did my wife pick up her antidepressant today?" The spouse is not listed as authorized. What should the technician do?

- a) Confirm pickup status since it is not a diagnosis
- b) Provide only the pickup time
- c) Decline to disclose and request the patient contact the pharmacy or add authorization
- d) Provide the information if the spouse can verify address

52) While filling amlodipine 10 mg, the technician accidentally selects amlodipine/benazepril 10/20 mg due to similar names. Which safeguard best helps prevent this during filling?

- a) Rely on tablet color and shape memory
- b) Use barcode scanning of the stock bottle to match the label/NDC selection
- c) Skip scanning for combination products to save time
- d) Always use the cheapest NDC to reduce confusion

53) An out-of-state prescription is presented for promethazine/codeine syrup. The hardcopy lacks the patient address and the prescriber phone number is incomplete. What should the technician do?

- a) Add missing information from the profile and proceed
- b) Refuse all out-of-state controlled-substance prescriptions
- c) Fill promethazine-only to avoid codeine rules
- d) Hold and route for pharmacist determination of legal sufficiency and next steps

54) A sealed bottle is returned from another pharmacy location, but no DSCSA transaction documentation accompanies it. What should the technician do?

- a) Stock it if the seal is intact
- b) Quarantine and follow the process for missing/invalid traceability documentation
- c) Relabel it with your pharmacy's NDC and stock it

d) Dispense it only for cash transactions

55) An eRx shows gabapentin 300 mg “tablet” in the product field, but the sig says “Take 1 capsule TID.” The pharmacy stocks both forms. What should the technician do?

- a) Dispense either; strength is the same
- b) Select capsules and proceed since the sig says capsule
- c) Convert to 600 mg tablets to reduce pill count
- d) Hold for clarification because product selection conflicts with directions

56) Two patients with similar names are waiting at pickup. What is the best step to prevent releasing the wrong prescription?

- a) Ask which medication they are picking up
- b) Use two patient identifiers before handing over any bag
- c) Call out the patient’s address loudly to confirm
- d) Hand both bags and ask them to check labels

57) Enoxaparin is prescribed: “40 mg/0.4 mL syringe; inject 0.4 mL daily.” The technician pulled 30 mg/0.3 mL syringes. What should the technician do?

- a) Dispense 30 mg/0.3 mL and instruct to inject the whole syringe
- b) Dispense 30 mg/0.3 mL and instruct to inject 0.4 mL
- c) Dispense two 30 mg syringes daily to approximate 60 mg
- d) Stop and obtain the correct strength/volume; do not alter the ordered dose

58) A noncontrolled refill authorization call is received from a prescriber office. Who may document it, depending on state law and pharmacy workflow?

- a) Only a pharmacist
- b) Only the prescriber
- c) Only the patient
- d) An appropriately authorized technician or pharmacist, with required pharmacist oversight

59) A prescription for amoxicillin/clavulanate is entered for a patient with “penicillin— anaphylaxis.” What should the technician do?

- a) Process it; allergy issues are handled at counseling
- b) Ask the patient if they still have the allergy and proceed if they say no
- c) Route for pharmacist review because it is a high-risk allergy conflict
- d) Substitute azithromycin and fill immediately

60) A compounding record includes ingredient lot numbers and expiration dates. What is the primary purpose?

- a) Calculate pricing and markup
- b) Enable traceability and quality assurance if a recall or error occurs
- c) Replace cleaning logs
- d) Avoid the need for pharmacist verification

61) Sig: “Take 2 tsp PO TID.” How many mL is each dose?

- a) 5 mL
- b) 10 mL
- c) 15 mL

d) 30 mL

62) Insulin glargine U-100 pens: 3 mL per pen, 5 pens per box. Sig: "Inject 18 units daily." What is the days' supply for one box?

- a) 41 days
- b) 50 days
- c) 60 days
- d) 83 days

63) A prescription reads: "Methotrexate 2.5 mg: take 1 tablet daily." The patient has rheumatoid arthritis. What should the technician do?

- a) Process it because daily methotrexate is standard for RA
- b) Change the directions to weekly and proceed
- c) Hold and route immediately due to high-risk weekly vs daily dosing concern
- d) Dispense 7 tablets and assume it means weekly

64) Which medication is most associated with the auxiliary warning "Avoid grapefruit"?

- a) simvastatin
- b) amoxicillin
- c) levothyroxine
- d) metformin

65) Under federal rules, which statement is most accurate about transferring a Schedule III prescription?

- a) It may be transferred once for refill dispensing if allowed by state law
- b) It cannot be transferred under any circumstances
- c) It can be transferred unlimited times between any pharmacies
- d) It can be transferred only if written on special controlled-substance paper

66) In a hospital, a nurse calls a verbal order: "Hydromorphone 0.5 mg IV now." The nurse asks the technician to enter it. What is the best technician action?

- a) Enter and verify it independently because it is STAT
- b) Enter it per workflow so a pharmacist can verify before the order becomes active
- c) Refuse to enter any verbal orders under any circumstances
- d) Change the route to IM to reduce risk

67) A patient presents a manufacturer coupon for a brand inhaler. The technician bills it as primary and gets "coverage terminated." What is the best next step?

- a) Delete the patient's insurance profile
- b) Rebill using correct coordination of benefits (primary insurance first, coupon as secondary if applicable)
- c) Bill cash and close the transaction
- d) Switch to a different medication without contacting anyone

68) A Schedule II prescription hardcopy is discovered filed with Schedule IV records. What should the technician do?

- a) Leave it; it is still on file
- b) Move it to the correct filing system to meet controlled-substance recordkeeping

requirements

- c) Shred it and reprint later
- d) Staple it to the register tape and discard at month end

69) A patient asks: "Is Cymbalta the same as duloxetine?" What is correct?

- a) Cymbalta is the generic for duloxetine
- b) Duloxetine is the generic for Cymbalta
- c) They are different drugs in the same class and not interchangeable
- d) Duloxetine is the generic for citalopram

70) During filling, barcode scanning returns "NDC mismatch," but the strength appears the same. What should the technician do?

- a) Override the mismatch if the drug name looks similar
- b) Continue and let final verification catch it
- c) Stop and re-verify the selected drug/NDC; correct the product before proceeding
- d) Change the label to match the scanned bottle

71) A patient asks how to dispose of leftover opioid tablets. What is the best technician response?

- a) Flush them to prevent diversion
- b) Recommend an authorized take-back/collection site and offer pharmacist help if needed
- c) Dissolve in water and pour down the sink
- d) Save them for future pain

72) A new prescription is for carvedilol 12.5 mg BID. The profile shows asthma and frequent albuterol fills. What should the technician do?

- a) Process normally; carvedilol has no respiratory effects
- b) Route for pharmacist review due to beta-blocker respiratory risk concerns
- c) Change carvedilol to metoprolol succinate and fill
- d) Tell the patient to use albuterol before taking carvedilol

73) A technician prints a MAR for a long-term care facility that includes patient identifiers and diagnoses. What is the best privacy practice?

- a) Leave it on the printer for nursing pickup
- b) Secure it immediately and release only to authorized personnel
- c) Email it to any requesting nurse
- d) Post it at the nurses' station for easy access

74) A prescription is for diltiazem ER 240 mg daily. The technician selected diltiazem IR 120 mg because it appears in history. What should the technician do?

- a) Dispense IR since the patient used it before
- b) Dispense two IR 120 mg daily to equal 240 mg
- c) Put the prescription on hold until the patient calls
- d) Stop and select the ER formulation as prescribed; route if clarification is needed

75) During expiration checks, a stock bottle shows an expiration date that appears altered with marker. What should the technician do?

- a) Use the altered date if it seems reasonable

- b) Ignore it if tablets look normal
- c) Quarantine/remove from inventory for investigation due to product integrity concerns
- d) Replace it with a new sticker showing a later date

76) An eRx is selected as bupropion XL 150 mg, but the directions entered read "Take 1 tablet twice daily." What should the technician do?

- a) Dispense XL and keep twice-daily directions
- b) Change to immediate-release automatically
- c) Change directions to once daily and process
- d) Hold and route for pharmacist review because formulation and directions conflict

77) A prescription is written for insulin lispro U-100 vial. The patient says they use pens and have no syringes. What should the technician do?

- a) Switch to pens because the insulin is the same
- b) Dispense the vial and suggest using pen needles
- c) Request a new prescription for pens per workflow and route appropriately
- d) Convert to insulin aspart pens to match availability

78) While typing directions, a technician selects "qod" instead of "qd." Which practice best reduces this risk?

- a) Free-type all directions instead of using templates
- b) Use standardized sig selections and compare the final directions to the original order before submitting
- c) Focus on speed because pharmacists will fix it later
- d) Convert all "qd" instructions to "weekly" to prevent confusion

79) A person picks up for a patient and shows ID. Last names match but the person is not listed in the patient profile. What should the technician do?

- a) Release it since last names match
- b) Refuse all third-party pickups
- c) Confirm required identifiers and follow the pharmacy's third-party pickup process before release
- d) Ask them to open the bag at the counter to prove ownership

80) A patient says, "I already take Flomax, and this one says tamsulosin." What should the technician do?

- a) Tell them to take both for better symptom control
- b) Confirm they are brand/generic equivalents and ensure pharmacist counseling is offered/available as required
- c) Tell them to stop Flomax immediately
- d) Dispense both brand and generic to avoid confusion

81) The technician notices two patient profiles were merged incorrectly and medications from two people are combined. What should the technician do?

- a) Delete one profile to fix it quickly
- b) Continue and ask the patient which meds are theirs at pickup
- c) Add a note "verify name" and proceed
- d) Stop processing and escalate immediately to correct patient identity and prevent misfills

82) A faxed refill request for a noncontrolled medication is received, but the fax header number does not match the prescriber's known office number. What should the technician do?

- a) Process it because the directions match prior fills
- b) Call the number on the fax to confirm
- c) Verify using a trusted contact method (known office number) and route as needed
- d) Ignore all faxed refill requests

83) An eRx lists metronidazole vaginal gel, but the directions say "Take 1 tablet by mouth twice daily." What should the technician do?

- a) Dispense the gel and keep oral directions
- b) Change to tablets and continue without clarification
- c) Dispense both gel and tablets
- d) Hold and route for clarification because dosage form and directions conflict

84) Which medication commonly requires providing an FDA Medication Guide when dispensed?

- a) isotretinoin
- b) amoxicillin
- c) calcium carbonate
- d) loratadine

85) A patient asks whether aztreonam is safe with a penicillin anaphylaxis history. What should the technician do?

- a) Assure them it is always safe with any penicillin allergy
- b) Tell them it is a penicillin and should never be used
- c) Recommend premedicating with diphenhydramine
- d) Refer to the pharmacist for allergy/cross-reactivity counseling

86) Pediatric dosing: 12 mg/kg/day divided BID. Child weighs 26 lb. What is the total mg per day?

- a) 72 mg/day
- b) 96 mg/day
- c) 120 mg/day
- d) 156 mg/day

87) A pharmacy dispenses naloxone nasal spray under a statewide standing order. Which technician action is appropriate?

- a) Provide full patient counseling on overdose response
- b) Dispense without documentation because naloxone is OTC
- c) Collect required information, process the order per protocol, and ensure pharmacist counseling is offered/available as required
- d) Substitute injectable naloxone to reduce cost without authorization

88) A patient returns stating the label has another patient's name, though the tablets look correct. What should the technician do?

- a) Remove the label and reprint a new one without reporting

- b) Tell the patient to discard it and come back later
- c) Ask the patient to keep it and swap labels at home
- d) Retrieve the medication immediately and process as a dispensing error/incident for correction and documentation

89) Doxycycline 100 mg capsules: "Take 1 capsule BID. Dispense #28." What is the days' supply?

- a) 7 days
- b) 10 days
- c) 14 days
- d) 28 days

90) A new eRx is received for buprenorphine/naloxone 8/2 mg films "Dissolve 1 film SL daily #30" from a prescriber new to the pharmacy. What should the technician do before it can be dispensed?

- a) Fill immediately because it is not a controlled substance
- b) Verify diagnosis and add it to the patient profile
- c) Convert to tablets because they are less expensive
- d) Complete required validity checks and route through controlled-substance workflow for pharmacist verification

Answer Key

1) b	31) b	61) b
2) c	32) b	62) b
3) d	33) d	63) c
4) c	34) c	64) a
5) b	35) c	65) a
6) b	36) a	66) b
7) c	37) c	67) b
8) b	38) d	68) b
9) c	39) b	69) b
10) d	40) b	70) c
11) c	41) c	71) b
12) b	42) c	72) b
13) c	43) b	73) b
14) a	44) d	74) d
15) b	45) b	75) c
16) c	46) b	76) d
17) b	47) c	77) c
18) c	48) b	78) b
19) c	49) c	79) c
20) a	50) c	80) b
21) c	51) c	81) d
22) d	52) b	82) c
23) b	53) d	83) d
24) c	54) b	84) a
25) a	55) d	85) d
26) b	56) b	86) c
27) c	57) d	87) c
28) a	58) d	88) d
29) c	59) c	89) c
30) c	60) b	90) d

PTCB Sim Test 24

1) A new eRx reads: "Glucophage XR 500 mg: Take 2 tablets by mouth with evening meal daily. Qty 180." Which product should be selected if substitution is permitted?

- a) Metformin IR 500 mg tablets
- b) Metformin ER 500 mg tablets
- c) Glyburide 5 mg tablets
- d) Metformin ER 1000 mg tablets

2) A patient asks for a copy of last month's receipt showing medication names for "my spouse's taxes." No authorization is on file. What should the technician do?

- a) Print it because they are married
- b) Provide only total dollars paid, not medication names, without authorization
- c) Release it if they can state the spouse's DOB
- d) Decline to release and route to the pharmacist/manager for appropriate authorization options

3) Rx: "Amoxicillin/clavulanate 600 mg/42.9 mg per 5 mL. Take 7.5 mL PO BID x 10 days." How many mL should be dispensed?

- a) 150 mL
- b) 120 mL
- c) 75 mL
- d) 90 mL

4) A will-call bag is being handed out. The patient states only, "I'm here for my refill," and several patients are waiting. What is the best next step?

- a) Ask for the patient's name and date of birth, then match to the will-call bag
- b) Ask for the patient's last name only to keep the line moving
- c) Ask what the medication is for, then select the bag
- d) Hand over the bag that matches the current time stamp on the receipt

5) A prescriber's office calls in a new prescription for alprazolam (Schedule IV). The technician answers the phone. What should happen next?

- a) The technician takes the verbal order and processes it
- b) The technician collects callback information and transfers the call to the pharmacist
- c) The technician refuses because controlled substances cannot be phoned in
- d) The technician tells the caller to e-prescribe only and hangs up

6) A new eRx is for "Toprol-XL 50 mg daily." Which generic should be selected?

- a) Metoprolol tartrate 50 mg
- b) Atenolol 50 mg
- c) Metoprolol succinate ER 50 mg
- d) Propranolol ER 60 mg

7) A refrigerator delivery arrives with a min/max indicator showing the shipment reached 12°C. What should the technician do with the affected refrigerated medications?

- a) Place into stock immediately because they are still cold

- b) Quarantine the shipment for pharmacist/manager assessment and documentation
- c) Move to the freezer for 30 minutes to rapidly cool, then stock
- d) Dispense first to avoid waste

8) Rx: "Humalog KwikPen U-100. Inject 12 units SQ with meals TID." Box contains 5 pens (3 mL each). What days' supply should be billed for one box?

- a) 31 days
- b) 25 days
- c) 28 days
- d) 37 days

9) A paper prescription for a Schedule II medication is presented. The date and prescriber signature are present, but the patient address is missing. What is the best technician action?

- a) Accept and fill; patient address is optional for CII
- b) Ask the patient to write in their address on the prescription
- c) Route to the pharmacist to determine if it meets applicable requirements before dispensing
- d) Add the address from the patient profile and proceed without review

10) A patient profile lists "anaphylaxis to sulfa drugs." A new prescription arrives for sulfamethoxazole/trimethoprim. What should the technician do?

- a) Enter and proceed because the prescriber chose it
- b) Change to nitrofurantoin and process
- c) Remove the sulfa allergy because it is nonspecific
- d) Stop processing and route for pharmacist review before dispensing

11) The technician selects "hydroxyzine 25 mg" but the label prints "hydralazine 25 mg." What is the best immediate action before the prescription moves forward?

- a) Leave as is and rely on counseling at pickup
- b) Correct the product selection to match the prescription and regenerate labels as needed
- c) Dispense whichever is in stock and note it on the label
- d) Change the directions to "take as directed" to reduce confusion

12) A patient requests pseudoephedrine 30 mg tablets, quantity 96. The log shows they purchased 2.4 g earlier today. What is the appropriate outcome under federal limits?

- a) Deny the sale because it would exceed the daily limit
- b) Sell the full quantity because only the 30-day limit matters
- c) Sell half now and half later without logging the second sale
- d) Sell only if the patient pays cash

13) A prescription is entered with "Take 1 tab PO qd." For error reduction, what should the patient label display?

- a) qd
- b) daily
- c) once daily
- d) q.i.d.

14) A patient is prescribed warfarin. A new eRx arrives for metronidazole. What should the

technician do during processing?

- a) Process normally; interactions are only discussed at pickup
- b) Route for pharmacist review due to significant interaction risk
- c) Tell the patient to stop warfarin while on the antibiotic
- d) Change metronidazole to amoxicillin and continue

15) Order: "Dexamethasone 10 mg IV now." Stock is 4 mg/mL. How many mL should be prepared?

- a) 2 mL
- b) 2.5 mL
- c) 1 mL
- d) 4 mL

16) A patient presents a manufacturer copay card to use with Medicaid for a brand medication. What is the best technician response?

- a) Apply the coupon because it lowers the patient cost
- b) Split-bill: Medicaid pays part and the coupon pays the rest
- c) Explain coupons generally cannot be used with government-funded plans and offer allowed alternatives
- d) Run the coupon first, then rebill Medicaid after pickup

17) The pharmacy receives a Class I recall notice for a specific lot of losartan 50 mg. What should the technician do?

- a) Continue dispensing until the lot is depleted
- b) Pull only opened bottles; sealed bottles may remain
- c) Remove all strengths of losartan from stock to be safe
- d) Quarantine the affected lot(s) immediately and follow the recall instructions for documentation/disposition

18) Rx: "Azithromycin 200 mg/5 mL: 12.5 mL PO day 1, then 6.25 mL daily days 2–5." Total volume to dispense?

- a) 37.5 mL
- b) 40 mL
- c) 45 mL
- d) 31.25 mL

19) A prescriber leaves a voicemail prescription for clonazepam (Schedule IV). The technician retrieves the message. What is the correct next step?

- a) Transcribe and process it as a valid verbal order taken by the technician
- b) Delete the message because controlled substances cannot be phoned in
- c) Forward the message to the pharmacist to determine acceptance/documentation requirements
- d) Call the patient to confirm the directions and process

20) A new prescription is written: "Eliquis 5 mg PO BID." Which generic is correct?

- a) Rivaroxaban
- b) Dabigatran
- c) Apixaban

d) Edoxaban

21) Barcode scan shows “bupropion XL 150 mg,” but the printed label is for “bupropion SR 150 mg BID.” What should the technician do?

- a) Proceed because bupropion is the same drug
- b) Stop and resolve the mismatch by selecting the correct formulation and/or correcting entry before verification
- c) Change the label to “daily” so SR matches XL
- d) Dispense and rely on patient counseling to correct it

22) A technician notices multiple prescriptions on the counter with visible patient names and medication names where other customers can see them. What is the best immediate action?

- a) Leave them; they are needed for workflow
- b) Turn them face down or move them to a secure area to reduce PHI exposure
- c) Shred them immediately even if still needed
- d) Highlight the names so staff can read them faster

23) Rx: “Prednisone 10 mg tabs: 4 tabs daily x 3 days, then 3 tabs daily x 3 days, then 2 tabs daily x 3 days, then 1 tab daily x 3 days.” Quantity?

- a) 30
- b) 36
- c) 42
- d) 24

24) An insurance claim rejects: “Invalid prescriber NPI.” What is the best technician action?

- a) Use another provider’s NPI from the same clinic
- b) Replace with the pharmacist’s NPI to get a paid claim
- c) Remove the prescriber field and submit as “clinic”
- d) Contact the prescriber office (or follow workflow) to obtain/verify the correct NPI and resubmit

25) A patient asks which medication is the generic for Norvasc. What is correct?

- a) Amlodipine
- b) Valsartan
- c) Lisinopril
- d) Atenolol

26) A controlled-substance cycle count is short by 1 tablet. What should the technician do first?

- a) Adjust the count to match the shelf quantity to close the discrepancy
- b) Recount carefully and review recent activity/transactions, then escalate per discrepancy process
- c) Wait until the next shift because it may self-correct
- d) Add 1 tablet to the next fill to rebalance inventory

27) Rx: “Nitroglycerin 0.4 mg SL tablets.” The technician selected nitroglycerin 0.4 mg/hr patches. Best action?

- a) Dispense patches because the drug matches

- b) Change directions to patch directions and proceed
- c) Select the correct dosage form before it goes to verification
- d) Add an auxiliary "remove at bedtime" and proceed

28) A new eRx arrives for losartan 50 mg daily. Profile shows lisinopril 20 mg daily active with refills. Patient says, "My doctor switched me." Best technician step?

- a) Discontinue lisinopril and fill losartan
- b) Fill both and let the patient decide
- c) Reject the losartan because ACE inhibitors and ARBs cannot be switched
- d) Process the losartan and route for pharmacist review of possible therapy change/duplication

29) A prescription label reads: "Instill 1 gtt OU QID." What does OU mean?

- a) Right eye
- b) Left eye
- c) Both eyes
- d) Both ears

30) A DSCSA trading partner requests transaction information/history/statement for a product under investigation. What should the technician do?

- a) Provide required documentation through the pharmacy's established process and escalate appropriately
- b) Refuse because only manufacturers keep those records
- c) Provide only the total invoice amount without item detail
- d) Delete the records after responding to reduce liability

31) Clindamycin oral solution 75 mg/5 mL: "Take 10 mL PO TID x 7 days." How many mL should be dispensed?

- a) 140 mL
- b) 200 mL
- c) 210 mL
- d) 280 mL

32) A prescription for methylphenidate IR 10 mg lists "Refills: 2." What is the best action?

- a) Process as written; refills are allowed for ADHD medications
- b) Process only the original fill and remove the refills before dispensing
- c) Convert to methylphenidate ER so refills can be used
- d) Route for pharmacist review because Schedule II prescriptions cannot be refilled

33) A patient asks, "What is atorvastatin for?" What is the appropriate technician response?

- a) Explain it lowers LDL and list common side effects
- b) Describe how it works in the liver and discuss monitoring
- c) Refer the patient to the pharmacist for counseling
- d) Tell them it is a blood thinner used for clots

34) Rx: "Ventolin HFA: Inhale 2 puffs q4–6h PRN." The technician selected Advair Diskus. What should happen?

- a) Dispense because both treat breathing problems

- b) Change sig to "1 inhalation BID" and dispense
- c) Stop and select the correct medication/device before verification
- d) Dispense and tell patient to use it as needed

35) A nonsterile compounded oral suspension containing water is assigned a BUD of 180 days. What should the technician do?

- a) Dispense as labeled because BUDs are optional
- b) Remove the BUD and use only the earliest ingredient expiration date
- c) Change automatically to 14 days without asking
- d) Route to the pharmacist for BUD assessment per applicable standards before dispensing

36) Rx: "Cefdinir 250 mg/5 mL. Take 6 mL PO daily x 10 days." How many mL should be dispensed?

- a) 60 mL
- b) 80 mL
- c) 90 mL
- d) 120 mL

37) Two patient profiles match on last name and share the same street address. What is the best way to prevent selection/entry errors?

- a) Use address alone as the identifier
- b) Confirm at least two identifiers such as full name and DOB
- c) Ask the patient what the medication is for
- d) Use phone number only because it is unique

38) A prescriber's office sends a signed patient authorization requesting the patient's medication profile by fax. What should the technician do?

- a) Fax it to any number provided to avoid delays
- b) Refuse because HIPAA prohibits sharing profiles
- c) Email the profile because fax is not secure
- d) Fax to a verified office fax number and document the disclosure per procedure

39) The technician is restocking and sees "DOBUtamine" stored directly next to "DOPamine" in similar packages. What is the best immediate step within technician scope?

- a) Separate the medications and apply/maintain tall-man or shelf alerts per system
- b) Remove one drug from inventory permanently
- c) Open both cartons and store syringes together in one bin
- d) Rely on color differences and keep them adjacent

40) Rx: "Wellbutrin XL 150 mg daily." The technician selected bupropion SR 150 mg. What is the best action?

- a) Substitute SR because the strength matches
- b) Change directions to once daily and dispense SR
- c) Select the correct XL formulation as prescribed before verification
- d) Dispense SR and add "do not crush" auxiliary label

41) A technician receives a request to transfer a Schedule III prescription with refills remaining to another pharmacy. What should the technician do?

- a) Perform the transfer over the phone and document it
- b) Tell the patient controlled substances can never be transferred
- c) Print a copy and hand it to the patient to take to the other pharmacy
- d) Collect destination information and route to the pharmacist to complete the transfer per rules

42) A patient returns an opened bottle of tablets dispensed yesterday and says they no longer need it. What should the technician do?

- a) Return to stock if the label is intact
- b) Pour back into the stock bottle if tablets look clean
- c) Do not redispense returned medication; handle per return/disposal procedure
- d) Repackage into unit-dose and return to inventory

43) A prescription is written "MSO4 2 mg IV q2h PRN pain." What should be done during entry?

- a) Enter as morphine sulfate and proceed
- b) Enter as magnesium sulfate because it is safer
- c) Hold for clarification because the abbreviation is error-prone
- d) Convert to an oral morphine equivalent for safety

44) A technician sees a DUR alert flagged: "Sertraline + linezolid: interaction." What is the best technician action?

- a) Override the alert because it is likely not clinically significant
- b) Tell the patient to stop sertraline and start linezolid
- c) Change linezolid to doxycycline and continue
- d) Route for pharmacist review before dispensing

45) A box of Lantus SoloStar contains 5 pens (3 mL each), U-100. Sig: "Inject 24 units SQ daily." What is the days' supply for one box?

- a) 50 days
- b) 62 days
- c) 75 days
- d) 90 days

46) A technician is asked to dispose of expired Schedule II stock bottles. What is the appropriate action?

- a) Place in regular trash after removing labels
- b) Follow the controlled-substance disposal/reverse distribution process with secure handling and required documentation
- c) Send back to wholesaler in the next regular return tote without paperwork
- d) Pour tablets into a hazardous waste bin to prevent diversion

47) A prescription for tramadol is missing the prescriber signature on a hardcopy. What should the technician do?

- a) Process it; signature is optional for non-CII
- b) Ask the patient to sign to acknowledge it
- c) Route for pharmacist resolution before dispensing
- d) Call the prescriber and take authorization as the technician

- 48) A patient has a documented severe latex allergy. The selected vaccine syringe packaging states "contains natural rubber latex." What should the technician do?
- a) Proceed; latex allergy applies only to gloves
 - b) Have the patient sign a waiver and proceed
 - c) Remove the allergy from the profile since it is not medication related
 - d) Hold and route for pharmacist review to select a latex-free option
- 49) A patient asks if escitalopram is "the same as citalopram." What should the technician do?
- a) Confirm they are identical and interchangeable
 - b) Refer to the pharmacist for counseling about similarity/interchangeability
 - c) Tell the patient they treat completely different conditions
 - d) Change the prescription to citalopram to save money
- 50) Rx: "Albuterol HFA: Inhale 2 puffs q6h PRN. Dispense 1 inhaler (200 puffs)." Using maximum frequency, what days' supply should be billed?
- a) 12
 - b) 16
 - c) 25
 - d) 50
- 51) A prescription is for "Synthroid 75 mcg" with DAW 1. What is correct?
- a) Dispense generic levothyroxine and bill DAW 0
 - b) Dispense brand Synthroid and bill as DAW 1
 - c) Dispense generic and print "Synthroid" on the label
 - d) Change DAW to 0 if patient requests lower cost
- 52) During data entry, the image shows "Take 1 tablet daily," but the typed sig prints "Take 1 tablet BID." What is the best technician action?
- a) Keep BID since it is already typed
 - b) Change to "as directed" to reduce rework
 - c) Correct the directions to match the prescription before sending to verification
 - d) Fill BID and add a note "verify at pickup"
- 53) A customer asks for "the strongest OTC ibuprofen." What should the technician do?
- a) Recommend 800 mg tablets and explain dosing
 - b) Recommend doubling naproxen with ibuprofen for more relief
 - c) Tell them to take 4 tablets every 4 hours
 - d) Help locate products and offer pharmacist assistance for dosing/selection questions
- 54) A CII prescription is presented with signs of alteration: quantity appears overwritten in darker ink. What should the technician do?
- a) Fill the smallest common quantity to reduce risk
 - b) Ask the patient to initial the change and proceed
 - c) Set aside and route to the pharmacist for evaluation and verification steps
 - d) Call the prescriber and confirm as the technician, then fill

55) A technician is entering "ondansetron ODT 4 mg." Which selection best matches?

- a) Ondansetron 4 mg swallow tablets
- b) Ondansetron ODT 4 mg orally disintegrating tablets
- c) Promethazine 12.5 mg tablets
- d) Prochlorperazine 10 mg tablets

56) A patient profile note states "difficulty swallowing; prefers liquid." New prescription: doxycycline 100 mg capsules. What should the technician do?

- a) Change to a liquid doxycycline product and process
- b) Add "open capsule and sprinkle" to the label
- c) Route for pharmacist review for appropriate dosage-form options
- d) Substitute doxycycline monohydrate tablets automatically

57) Rx: "Metformin ER 500 mg: Take 2 tablets daily with evening meal. Dispense #180." What is the days' supply?

- a) 45
- b) 60
- c) 90
- d) 180

58) A pharmacy receives a request from law enforcement for a patient's prescription records without a warrant or patient authorization. What should the technician do?

- a) Provide the records immediately to cooperate
- b) Provide only controlled-substance records
- c) Decline and route to the pharmacist/manager for appropriate legal handling
- d) Print the profile but remove diagnosis information and provide the rest

59) Rx: "Augmentin 875/125 mg: Take 1 tab PO BID x 7 days." Quantity?

- a) 7
- b) 10
- c) 14
- d) 21

60) A patient's controlled-substance prescription is rejected by the system due to missing prescriber DEA number on file. What is the best technician action?

- a) Use a DEA number from an old hardcopy for that clinic
- b) Bill without DEA because it is required only for CII
- c) Ask the patient to provide the DEA number
- d) Obtain/verify required prescriber information through appropriate workflow before processing

61) A technician is asked by a patient to explain why their Medicare Part D is in the coverage gap and what plan they should choose. Best action?

- a) Provide a full plan recommendation based on costs
- b) Refer to appropriate staff/pharmacist for questions beyond basic claim status
- c) Tell the patient Medicare will not answer those questions
- d) Enroll the patient in a new plan using the pharmacy computer

62) Label reads: "Instill 1 drop OS qhs." What does OS mean?

- a) Right eye
- b) Left eye
- c) Both eyes
- d) Left ear

63) A technician accidentally counted digoxin 0.25 mg instead of digoxin 0.125 mg. Which safeguard is designed to catch this before the patient receives it?

- a) Alphabetical will-call organization
- b) Patient signature at pickup
- c) Pharmacist final product verification
- d) Drug manufacturer bottle color consistency

64) An outpatient prescription for isotretinoin is processed, and the system displays: "REMS: dispense blocked—no authorization." What should the technician do?

- a) Rebill as cash to bypass the block
- b) Substitute tretinoin cream and dispense
- c) Print a REMS handout and dispense anyway
- d) Hold the prescription and route for pharmacist REMS resolution before dispensing

65) A patient is currently taking simvastatin. A new prescription arrives for clarithromycin. What is the best technician action?

- a) Process normally; short antibiotic courses do not matter
- b) Route for pharmacist review due to interaction risk
- c) Tell the patient to halve simvastatin while on clarithromycin
- d) Change clarithromycin to azithromycin and proceed

66) A prescription is entered with "Take 0.5 tablet daily." The tablet is scored. What should the technician do?

- a) Change to "1 tablet every other day" for clarity
- b) Enter as written for pharmacist verification
- c) Round to 1 tablet daily to avoid splitting
- d) Reject because tablets cannot be split

67) An order requires 25 g dextrose. Stock is D50W (50 g/100 mL). How many mL are needed?

- a) 25 mL
- b) 40 mL
- c) 50 mL
- d) 75 mL

68) A patient is picking up a new prescription and asks to waive counseling to save time. What is the correct technician action?

- a) Document refusal and bypass counseling without pharmacist involvement
- b) Provide counseling yourself to prevent delays
- c) Offer counseling and ensure pharmacist is available if the patient has questions per workflow
- d) Require counseling for every prescription with no exceptions

69) A CII prescription is partially fillable due to out-of-stock quantity. The patient requests a partial fill today. What should the technician do?

- a) Dispense a partial fill without documentation to speed pickup
- b) Refer to the pharmacist to determine if partial filling is allowed and ensure required documentation/timelines
- c) Transfer the CII to another pharmacy to complete the remainder
- d) Substitute with a Schedule III alternative and dispense

70) Rx: "Nystatin oral suspension: Swish and swallow 5 mL QID." Bottle size: 120 mL. Days' supply?

- a) 4
- b) 6
- c) 8
- d) 10

71) A technician is entering an inpatient order that reads "U" for units (e.g., "Regular insulin 10U"). What is the safest entry practice?

- a) Enter "10U" exactly as written
- b) Enter "10 units" to avoid misinterpretation
- c) Enter "10.0 units" to show precision
- d) Change dose to 1 unit to avoid overdose risk

72) The technician selects levothyroxine 25 mcg, but the NDC scanned is liothyronine 25 mcg. What should the technician do?

- a) Continue because thyroid medications are interchangeable
- b) Dispense and let the patient's physician adjust if needed
- c) Stop and correct the product selection before verification
- d) Change the label to "take as directed" and proceed

73) A patient profile lists "serious rash with lamotrigine." A new prescription arrives for Lamictal. What should the technician do?

- a) Process it because brand may be tolerated better
- b) Remove the allergy since it is in the past
- c) Substitute levetiracetam and fill to avoid risk
- d) Route for pharmacist review due to documented reaction history

74) A technician is reconciling a controlled-substance invoice. What is most critical to verify?

- a) Bottle shape matches prior inventory
- b) Correct drug/NDC and quantity received match the invoice
- c) Manufacturer marketing description matches the wholesaler site
- d) That today's patient copays cover the invoice amount

75) Rx: "Take 2 tsp PO BID." Concentration is 160 mg/5 mL. How many mg per dose?

- a) 160 mg
- b) 240 mg
- c) 320 mg
- d) 640 mg

76) A patient has penicillin anaphylaxis documented. A new prescription arrives for cephalexin 500 mg. What should the technician do?

- a) Fill because cephalosporins never cross-react
- b) Fill and add an allergy sticker for the patient
- c) Change to clindamycin and proceed
- d) Route for pharmacist review to assess allergy risk before dispensing

77) A prescription for a high-alert medication is being filled. Which medication is considered high-alert and commonly requires extra safeguards?

- a) Heparin
- b) Amoxicillin
- c) Docusate
- d) Loratadine

78) A technician is asked to print a patient's full medication profile for a third party at pickup. The patient is not present, and no permission is documented. What should the technician do?

- a) Print it if the third party knows the patient's DOB
- b) Decline to release and route for appropriate authorization handling
- c) Print it but black out controlled substances
- d) Read the list aloud instead of printing it

79) A prescription is for "Zestril 10 mg daily." No DAW is indicated. Which product should be selected?

- a) Enalapril 10 mg
- b) Lisinopril 10 mg
- c) Benazepril 10 mg
- d) Losartan 10 mg

80) A patient previously took metoprolol tartrate 25 mg BID. Today an eRx arrives for metoprolol succinate ER 25 mg daily, and the patient is confused at pickup. Best technician action?

- a) Tell the patient to take both until the next appointment
- b) Change the new prescription to tartrate BID to match history
- c) Route for pharmacist counseling and verification of therapy change
- d) Cancel the new prescription to prevent duplication

81) Rx: "Fluconazole 150 mg: Take 1 tablet now and repeat in 72 hours." Quantity #2. What days' supply is typically billed?

- a) 2
- b) 3
- c) 4
- d) 7

82) A patient requests an early refill for a controlled substance; the system indicates it is "too soon." What is the best technician action?

- a) Override the reject and process to avoid conflict

- b) Tell the patient the prescriber must authorize every refill
- c) Change the days' supply to a smaller number to get it paid
- d) Gather needed information (e.g., reason, travel) and route per workflow for pharmacist/plan handling

83) A prescription is for tamsulosin 0.4 mg daily. The technician selected terazosin 1 mg. Best action?

- a) Dispense terazosin because both are alpha blockers
- b) Change directions to bedtime and dispense terazosin
- c) Select the correct medication; do not substitute within class without proper authorization
- d) Dispense terazosin and note "therapeutic interchange" on label

84) A label prints an auxiliary "May cause drowsiness" for fexofenadine. What should the technician do?

- a) Leave it; all antihistamines cause sedation
- b) Remove it and dispense without telling anyone
- c) Replace it with "Avoid alcohol" to be safer
- d) Correct the auxiliary labeling through the established process so final labeling is accurate before dispensing

85) An IV order reads: "Vancomycin 1 g in 250 mL NS over 2 hours." The technician is asked to calculate infusion rate in mL/hr. What is it?

- a) 125 mL/hr
- b) 250 mL/hr
- c) 100 mL/hr
- d) 150 mL/hr

86) A patient asks, "Can I drink alcohol with metronidazole?" What should the technician do?

- a) Tell them yes if it's only one drink
- b) Tell them to stop metronidazole if they plan to drink
- c) Refer the patient to the pharmacist for counseling
- d) Tell them alcohol is fine if taken with food

87) A paper prescription for acetaminophen/codeine includes drug, strength, directions, date, and prescriber signature, but no quantity. What should the technician do?

- a) Dispense #20 as a standard quantity
- b) Ask the patient what quantity they want and document it
- c) Dispense #30 because that is typical
- d) Route for pharmacist action to obtain quantity from prescriber before dispensing

88) A prescription is for insulin glargine U-100 vial (10 mL). Sig: "Inject 18 units SQ at bedtime." Days' supply for one vial?

- a) 55 days
- b) 50 days
- c) 40 days
- d) 60 days

89) During filling, the technician notices the technician-entered directions say "Take 1 tablet

PO QID" but the prescriber's directions are "Take 1 tablet PO daily." What is the best action?

- a) Leave QID because it provides better coverage
- b) Change to "as directed" and proceed
- c) Correct to match the prescription and reprint the label before verification
- d) Dispense and ask the patient which is correct

90) A Medication Guide is required for a medication being dispensed, but the printer is down and none are available. What should be done?

- a) Dispense without it because counseling is offered at pickup
- b) Provide the guide verbally and document it
- c) Hold dispensing until the Medication Guide can be provided through the workflow
- d) Substitute a different medication that does not require a guide

Answer Key

1) b	31) c	61) b
2) d	32) d	62) b
3) a	33) c	63) c
4) a	34) c	64) d
5) b	35) d	65) b
6) c	36) b	66) b
7) b	37) b	67) c
8) a	38) d	68) c
9) c	39) a	69) b
10) d	40) c	70) c
11) b	41) d	71) b
12) a	42) c	72) c
13) c	43) c	73) d
14) b	44) d	74) b
15) b	45) b	75) c
16) c	46) b	76) d
17) d	47) c	77) a
18) a	48) d	78) b
19) c	49) b	79) b
20) c	50) b	80) c
21) b	51) b	81) c
22) b	52) c	82) d
23) b	53) d	83) c
24) d	54) c	84) d
25) a	55) b	85) a
26) b	56) c	86) c
27) c	57) c	87) d
28) d	58) c	88) b
29) c	59) c	89) c
30) a	60) d	90) c

PTCB Sim Test 25

1) A new eRx reads: "Crestor 10 mg PO daily #30." The patient requests generic. The prescriber did not indicate DAW. What should the technician dispense?

- a) atorvastatin 10 mg
- b) rosuvastatin 10 mg
- c) simvastatin 10 mg
- d) pravastatin 10 mg

2) A written prescription for alprazolam is presented. The date written is 9 months ago. What should the technician do?

- a) Process it because it is a Schedule IV
- b) Fill it but remove refills
- c) Route it for pharmacist review due to controlled-substance validity rules
- d) Ask the patient to have the prescriber backdate a new copy

3) A prescription reads: "azithromycin 200 mg/5 mL. Take 12.5 mL PO day 1, then 6.25 mL PO daily days 2–5." What total volume should be dispensed?

- a) 37.5 mL
- b) 50 mL
- c) 62.5 mL
- d) 75 mL

4) During filling, the barcode scan flags the selected NDC as "omeprazole 20 mg" but the label is for "pantoprazole 40 mg." What should the technician do next?

- a) Override the scan because both are PPIs
- b) Replace the product with the correct NDC and rescan
- c) Change the label to match the scanned product
- d) Bag it for the pharmacist to correct later

5) A patient's caregiver requests the status of a prescription pickup. The caregiver provides the patient's name and DOB, but there is no authorization on file. What can the technician do?

- a) Confirm medication name and readiness
- b) Confirm only that "the pharmacy has prescriptions on file," without details
- c) Provide the copay and the medication name but not directions
- d) Provide full details if they know the address on file

6) A prescription reads: "clopidogrel 75 mg PO daily #90." Which brand name corresponds to clopidogrel?

- a) Brilinta
- b) Effient
- c) Plavix
- d) Coumadin

7) A prescriber calls in a verbal order to the pharmacy voicemail after hours for oxycodone IR 5 mg #20. Which is correct?

- a) The technician may transcribe and fill it if the prescriber left a DEA number
- b) The technician may enter it but must wait for the patient to sign a consent
- c) Schedule II prescriptions may not be authorized by voicemail; route to the pharmacist for handling
- d) The technician may fill it as an emergency oral C-II without pharmacist involvement

8) A prescription for insulin degludec U-100 pens: "Inject 22 units SQ daily." The box contains 3 pens of 3 mL each. What days' supply should be billed?

- a) 30
- b) 34
- c) 40
- d) 45

9) A new prescription is received for metoprolol succinate ER 25 mg daily. The patient profile shows metoprolol tartrate 25 mg BID filled yesterday. What should the technician do?

- a) Process the new one and discontinue the old automatically
- b) Change succinate to tartrate to match the profile
- c) Hold for pharmacist review due to formulation and dosing differences
- d) Fill both because they are the same medication

10) A prescription reads: "Apply 1 gram topically to affected area TID." The tube contains 30 grams. What is the days' supply?

- a) 7
- b) 10
- c) 14
- d) 30

11) A prescription reads: "instill 2 gtt AD q12h." What is the correct label instruction?

- a) Place 2 drops in left ear every 12 hours
- b) Place 2 drops in right ear every 12 hours
- c) Place 2 drops in each ear every 12 hours
- d) Place 2 drops in right eye every 12 hours

12) A shipment arrives with a tote label indicating it contains refrigerated items, but the tote is warm and the cold packs are melted. What should the technician do with the affected items?

- a) Stock them quickly to get them back to the correct temperature
- b) Quarantine them and alert the pharmacist/manager per cold-chain procedure
- c) Place them in the freezer for 30 minutes, then stock
- d) Dispense them first to avoid waste

13) A patient's profile lists anaphylaxis to penicillin. A new eRx arrives for amoxicillin. What should the technician do?

- a) Proceed because allergy documentation is patient-reported
- b) Change to cephalexin and continue
- c) Place an auxiliary label "Stop if rash occurs"
- d) Route to the pharmacist for allergy review before dispensing

14) A prescription reads: "Take 1 tablet PO q6h prn pain. Max 4 tablets/day." Quantity is #24. What is the maximum days' supply?

- a) 4 days
- b) 5 days
- c) 6 days
- d) 8 days

15) A patient requests pseudoephedrine. Which action is required under the Combat Methamphetamine Epidemic Act (CMEA)?

- a) Keep product on open shelves but limit to one per customer
- b) Record purchaser information and verify ID in a logbook/electronic log
- c) Require a prescription for all pseudoephedrine
- d) Allow sale only to patients with a pharmacy profile

16) A prescription is written: "levothyroxine 0.075 mg daily." Which strength should be selected?

- a) 7.5 mcg
- b) 75 mcg
- c) 0.75 mg
- d) 750 mcg

17) While entering a prescription, the technician types "q4h" instead of "qhs" from the prescriber's directions. Which best reduces this error before sending to verification?

- a) Use only free-text directions
- b) Skip the image/hard copy review to save time
- c) Perform a final compare of typed directions against the original order
- d) Ask the cashier to double-check the label at pickup

18) A prescription is for albuterol HFA: "Inhale 2 puffs every 6 hours PRN wheeze." The inhaler contains 200 puffs. What is the days' supply for insurance purposes using the maximum daily use?

- a) 12 days
- b) 20 days
- c) 25 days
- d) 50 days

19) A controlled substance safe count is short by 1 tablet during end-of-shift count. What is the best technician action?

- a) Adjust the on-hand count to match the expected quantity
- b) Recount and immediately report the discrepancy per procedure
- c) Wait until the next delivery arrives and recheck then
- d) Remove the drug from the safe and place it on the shelf

20) A prescription reads: "prednisone 10 mg tabs: Take 4 tabs daily x 3 days, then 2 tabs daily x 3 days, then 1 tab daily x 3 days." How many tablets should be dispensed?

- a) 18
- b) 21
- c) 24

d) 27

21) A claim rejects "refill too soon" for losartan. The patient reports the prescriber increased the dose, but no new prescription is on file. What should the technician do?

- a) Change the directions to match the patient statement and rebill
- b) Decrease the days' supply to force approval
- c) Gather details and route for pharmacist/prescriber follow-up on a new order or override
- d) Tell the patient to pay cash until insurance allows the refill

22) A prescription label prints "Take 1 tablet daily," but the eRx image says "Take 1 tablet twice daily." The medication has not been verified yet. What should the technician do?

- a) Correct the directions in the system and reprint the label
- b) Handwrite "twice daily" on the label
- c) Bag it and let the pharmacist fix it at pickup
- d) Dispense as printed and note the discrepancy in profile notes

23) A patient asks which OTC product is best for seasonal allergies given their other medications. What is the technician's best response?

- a) Recommend loratadine because it is non-drowsy
- b) Recommend diphenhydramine for strongest effect
- c) Ask more medical history and choose based on interactions
- d) Offer to have the pharmacist assist with selection

24) A prescription is written for "Zithromax Z-Pak." Which generic drug is this?

- a) azithromycin
- b) clarithromycin
- c) erythromycin
- d) doxycycline

25) A pharmacy receives a manufacturer notice: "Class I recall" for a specific lot of infant ibuprofen. What should the technician do first?

- a) Pull all ibuprofen products from the shelf
- b) Quarantine the affected lot and start the recall workflow
- c) Continue dispensing until returns are processed
- d) Dispose of the product in regular trash

26) A prescription reads: "Take 2.5 mL PO BID" of a solution labeled 20 mg/5 mL. How many mg per dose?

- a) 5 mg
- b) 10 mg
- c) 20 mg
- d) 40 mg

27) A prescriber e-prescribes "Adderall XR 20 mg daily #30." The patient asks to transfer it to another pharmacy across town before it is filled. What is the correct handling?

- a) Transfer it like any noncontrolled eRx
- b) Cancel at the original pharmacy and have the prescriber send a new prescription to the other pharmacy

- c) Convert it to a fax and send it to the other pharmacy
- d) Split-fill 15 capsules and transfer the remainder

28) A technician selects "hydroxyzine 25 mg" for a prescription written for "hydralazine 25 mg." Which system safeguard most directly prevents this from reaching the patient?

- a) Patient signature at pickup
- b) Hard-copy filing by date
- c) Barcode scan at product selection matched to the label/NDC
- d) Counseling offered at pickup

29) A prescription is for "nitroglycerin 0.4 mg SL tablets." The technician accidentally selects nitroglycerin 0.4 mg/hr patches. What should the technician do?

- a) Dispense the patches and add "place under tongue"
- b) Substitute patches because they are longer acting
- c) Correct the product to sublingual tablets before it goes to verification
- d) Dispense patches if the prescriber indicated "for angina"

30) A prescription reads: "Take 1 tablet PO every other day." Quantity is #30. What is the days' supply?

- a) 15
- b) 30
- c) 45
- d) 60

31) A patient returns an opened bottle of tablets and requests that the pharmacy "put it back on the shelf." What should the technician do?

- a) Return it to stock if the safety seal is intact
- b) Quarantine and process as return/disposal; do not re-dispense returned patient medication
- c) Count remaining tablets and top off with stock
- d) Give store credit and then restock

32) Which medication is an ACE inhibitor?

- a) losartan
- b) amlodipine
- c) lisinopril
- d) metoprolol

33) A technician receives a request from another pharmacy to transfer a clonazepam prescription with refills remaining. What is the correct action?

- a) Transfer it if the patient agrees
- b) Transfer it only if it is less than 30 days old
- c) Refuse because controlled substances can never be transferred
- d) Route to the pharmacist to perform the transfer per applicable rules

34) A prescription is written: "metformin ER 500 mg: take 2 tablets daily #60." What is the days' supply?

- a) 15

- b) 30
- c) 45
- d) 60

35) A DSCSA documentation issue is identified: transaction information is missing for a received bottle of brand-name tablets. What should the technician do with that bottle?

- a) Stock it but mark the shelf tag "pending documents"
- b) Quarantine it until required transaction documentation is obtained/verified
- c) Dispense it only for emergency fills
- d) Return all products in the entire delivery tote

36) A patient's profile shows a severe sulfonamide antibiotic allergy (SJS/TEN). A new prescription arrives for sulfamethoxazole/trimethoprim. What should the technician do?

- a) Fill it and add "take with food"
- b) Fill it because it's commonly used
- c) Route to the pharmacist due to severe allergy risk
- d) Substitute nitrofurantoin and continue

37) A pharmacy receives a DEA Form 222 order that was filled last year. Under federal law, how long must Schedule II purchase records be kept?

- a) 1 year
- b) 2 years
- c) 3 years
- d) 5 years

38) A prescription reads: "buprenorphine/naloxone 8 mg/2 mg films: dissolve 1 film SL daily #30." The technician enters 8 mg/0.2 mg. What should be done?

- a) Keep it; naloxone amount is not clinically relevant
- b) Correct the strength to match the prescription before filling continues
- c) Change to buprenorphine-only to simplify billing
- d) Dispense fewer films and note the correction on the label

39) A patient's warfarin prescription is due today. At pickup, the patient says, "My doctor changed my dose yesterday." What is the best technician action?

- a) Change the directions to match what the patient says
- b) Dispense the prescription and advise the patient to call the doctor
- c) Split tablets to approximate the new dose and dispense
- d) Refer the patient to the pharmacist to determine if a new order is needed

40) A prescription reads: "doxycycline 100 mg capsules BID #14." Which auxiliary instruction is most appropriate?

- a) Avoid sunlight and separate from antacids/iron/dairy
- b) Take with grapefruit juice
- c) Do not drink alcohol for 72 hours after last dose
- d) Take only at bedtime

41) A prescription reads: "Take 0.5 tablet PO daily." The product selected is a modified-release tablet that must not be split. What should the technician do?

- a) Dispense it as written; splitting is always allowed
- b) Change the quantity to half and dispense
- c) Flag it for pharmacist review before filling proceeds
- d) Crush the tablet and provide an oral syringe

42) A patient is prescribed "Trulicity 1.5 mg/0.5 mL: inject 0.5 mL weekly." The carton contains 4 single-dose pens. What days' supply should be billed?

- a) 14
- b) 21
- c) 28
- d) 56

43) A prescription is for "Augmentin 875/125 mg." Which generic matches?

- a) amoxicillin/clavulanate
- b) ampicillin/sulbactam
- c) amoxicillin/sulbactam
- d) cefdinir

44) A technician is asked which identifier is primarily used to verify a prescriber's authority to prescribe controlled substances.

- a) NPI
- b) DEA registration number
- c) State driver's license number
- d) Medicare PTAN

45) A prescription reads: "sertraline 50 mg PO daily." Which brand corresponds to sertraline?

- a) Prozac
- b) Zoloft
- c) Cymbalta
- d) Wellbutrin

46) A prescription reads: "Apply 2 grams topically BID." The jar contains 120 grams. What is the days' supply?

- a) 15
- b) 20
- c) 30
- d) 60

47) During pickup, two patients with similar names are waiting. What is the best practice before handing over any prescription bag?

- a) Ask which medication they are expecting
- b) Confirm address only
- c) Verify two identifiers (e.g., full name and DOB)
- d) Compare the patient's signature to the profile

48) A new order reads: "methotrexate 2.5 mg: take 6 tablets daily" with indication "RA." What should the technician do?

- a) Enter as written; RA dosing varies
- b) Change daily to weekly and continue
- c) Delete the indication and proceed
- d) Route for pharmacist review due to high-risk dosing frequency concern

49) A prescription is for "morphine sulfate ER 15 mg q12h." The technician selects morphine IR 15 mg. What should be done?

- a) Dispense IR with the ER directions
- b) Correct the product selection to ER before verification
- c) Dispense IR and add an auxiliary "extended-release" label
- d) Change the prescription to "q4h" to match IR

50) A nonsterile compounded cream is prepared. Which documentation item is most important for traceability?

- a) Patient insurance BIN
- b) Ingredient lot numbers and beyond-use date
- c) Technician's uniform size
- d) Total gross profit

51) A patient asks for a printed medication list for a specialist appointment. What is the correct technician approach?

- a) Refuse because medication histories cannot be printed
- b) Print after verifying identity and completing any required HIPAA release steps
- c) Print but remove all controlled substances
- d) Provide only if the prescriber calls to request it

52) A prescription reads: "Ciprofloxacin 0.3% ophthalmic solution: instill 1 drop OU q4h while awake." The technician selects ciprofloxacin 500 mg tablets. What should happen?

- a) Dispense tablets; systemic therapy is stronger
- b) Substitute ofloxacin otic solution
- c) Correct to the ophthalmic product before verification
- d) Dispense tablets and add "for eye infection" on label

53) Which medication is a proton pump inhibitor?

- a) famotidine
- b) sucralfate
- c) pantoprazole
- d) dicyclomine

54) A technician is working a rejected claim: "non-covered NDC" for insulin pen needles. What is an appropriate next step within scope?

- a) Select a covered equivalent NDC for the same size/type if permitted and rebill
- b) Change the insulin NDC to match the needles
- c) Override by setting days' supply to 1
- d) Bill the needles under the insulin claim line

55) A controlled substance prescription is missing the prescriber's address. What should the technician do?

- a) Add the address from the pharmacy directory and fill
- b) Ask the patient to write in the address
- c) Fill it because address is optional
- d) Route for pharmacist review to determine if the prescription meets requirements

56) A prescription reads: "cefdinir 250 mg/5 mL: take 6 mL PO BID x 10 days." What volume should be dispensed?

- a) 100 mL
- b) 120 mL
- c) 150 mL
- d) 180 mL

57) A technician is restocking and notices vincristine stored next to vinblastine. What change most directly reduces selection errors?

- a) Store both together alphabetically to keep them "organized"
- b) Remove auxiliary labels to reduce clutter
- c) Separate storage locations and require barcode scanning at selection
- d) Place both in the OTC aisle so they are more visible

58) A prescription is for "fluticasone/salmeterol 250/50 DPI." Which brand is most commonly associated?

- a) Advair Diskus
- b) Symbicort
- c) Spiriva HandiHaler
- d) ProAir HFA

59) A patient has an active prescription for omeprazole and a new eRx for esomeprazole arrives. The system flags duplicate therapy. What should the technician do?

- a) Override because both treat GERD
- b) Inactivate omeprazole and fill esomeprazole
- c) Fill both and let the patient decide
- d) Send to pharmacist review for duplication/therapy change assessment

60) A prescription is for "valacyclovir 1 g BID." The technician selects acyclovir 800 mg. What is the correct action?

- a) Dispense acyclovir; both are antivirals
- b) Correct the drug selection to match the prescription
- c) Change the sig to acyclovir dosing and dispense
- d) Dispense acyclovir and note "therapeutic equivalent"

61) A prescription is for "testosterone cypionate 200 mg/mL." Under federal law, what is its schedule?

- a) C-II
- b) C-III
- c) C-IV
- d) C-V

62) A prescription reads: "tamsulosin 0.4 mg PO daily." What is the most common

indication?

- a) BPH
- b) Hypertension
- c) GERD
- d) Migraine prevention

63) During filling, two different patients' labels are found in the same basket with mixed stock bottles. What should the technician do?

- a) Finish both and separate at the end
- b) Ask the cashier to verify the correct patient at pickup
- c) Stop and separate immediately, re-matching each label to the correct order before continuing
- d) Place both in will-call with a note "verify at register"

64) A patient is prescribed "ondansetron ODT 4 mg." The technician selected ondansetron 4 mg swallow tablets. What is the best action?

- a) Dispense swallow tablets; same drug and strength
- b) Change label to "dissolve on tongue" and dispense swallow tablets
- c) Add an oral syringe and dispense swallow tablets
- d) Correct the product to the ODT formulation before verification

65) A prescription is for "clonidine 0.1 mg patch weekly." The box contains 4 patches. What is the days' supply?

- a) 14
- b) 21
- c) 28
- d) 30

66) A patient's prescription for rosuvastatin is entered. Which brand name corresponds to rosuvastatin?

- a) Lipitor
- b) Crestor
- c) Zocor
- d) Mevacor

67) A patient requests an early refill for zolpidem stating it was lost. It was filled 10 days ago for 30 days. What is the best technician action?

- a) Refill and document "lost"
- b) Offer cash price if insurance rejects
- c) Tell the patient it is impossible to refill early
- d) Collect details and route for pharmacist review per controlled-substance early refill process

68) A pharmacy receives an eRx for "hydrochlorothiazide 25 mg daily" for a patient whose allergy field lists "sulfa—SJS." What should the technician do?

- a) Fill it because it is not an antibiotic
- b) Delete the allergy because it is outdated
- c) Route for pharmacist review due to severe allergy history

d) Substitute furosemide and continue

69) A prescription reads: "Take 1.5 tablets PO daily for 30 days." How many tablets should be dispensed?

- a) 30
- b) 45
- c) 60
- d) 90

70) A technician is asked to calculate a day supply for insulin aspart vial: "Inject 10 units SQ TID with meals." The vial is 10 mL (U-100). What days' supply is appropriate?

- a) 21
- b) 25
- c) 30
- d) 33

71) A prescriber faxes a Schedule II prescription for a patient not in a qualifying exception category. What is the correct handling?

- a) It can be filled if the fax includes the DEA number
- b) It can be filled if the patient pays cash
- c) It can be filled if the pharmacy calls to verify it
- d) A fax is not sufficient in this case; pharmacist must obtain a valid original/eRx per requirements

72) A prescription reads: "Use 2 sprays in each nostril daily." The bottle contains 120 sprays. What is the days' supply?

- a) 15
- b) 20
- c) 30
- d) 60

73) A technician is preparing to report a serious adverse event linked to a medication product. Which FDA program is used for voluntary reporting?

- a) MedWatch
- b) VAERS
- c) REMS
- d) PDMP

74) A prescription is for "Victoza 18 mg/3 mL pen: inject 1.2 mg SQ daily." What volume (mL) is used per day?

- a) 0.06 mL
- b) 0.12 mL
- c) 0.20 mL
- d) 1.20 mL

75) A prescription reads: "Take 1 capsule PO TID." The technician enters "take 3 capsules daily." What is the primary issue?

- a) No issue; total daily amount is the same

- b) Timing/frequency changed and may alter therapy; it must be corrected
- c) It only affects the receipt, not the label
- d) Insurance will automatically correct the directions

76) A Schedule II prescription is partially filled because insufficient stock is available for a typical outpatient. Under federal law, when must the remainder generally be supplied?

- a) Within 24 hours
- b) Within 48 hours
- c) Within 72 hours
- d) Within 7 days

77) A prescription arrives with "DAW" indicated by the prescriber. How should the technician process product selection?

- a) Dispense the brand/DAW product as indicated, unless the prescriber authorizes substitution
- b) Substitute generic automatically to reduce cost
- c) Choose whichever is in stock
- d) Dispense generic and add DAW on the label for billing

78) A hospital medication order shows "KCl 20 mEq IV" without diluent/volume. What should the technician do?

- a) Prepare the most common premix and send
- b) Ask the nurse which concentration is preferred and prepare it
- c) Hold for pharmacist clarification before preparation
- d) Substitute oral potassium to avoid IV risk

79) A prescription is written: "Take 2 capsules PO BID for 10 days." Quantity is #40. What days' supply is this?

- a) 5
- b) 10
- c) 15
- d) 20

80) A technician is asked which medication is commonly monitored by INR.

- a) apixaban
- b) clopidogrel
- c) rivaroxaban
- d) warfarin

81) A prescription reads: "instill 1 gtt OS QID." What is the correct instruction?

- a) Place 1 drop in left eye 4 times daily
- b) Place 1 drop in right eye 4 times daily
- c) Place 1 drop in both eyes 4 times daily
- d) Place 1 drop in left ear 4 times daily

82) A technician is processing a transfer request for a noncontrolled prescription. Which piece of information is required during a transfer?

- a) Patient's insurance BIN

- b) Original date written and date of last fill
- c) Prescriber's personal phone number
- d) Patient's SSN

83) A prescription is for "diltiazem ER 240 mg daily." The technician selects diltiazem IR 60 mg because it can be taken multiple times daily. What should the technician do?

- a) Change to 60 mg QID and proceed
- b) Dispense IR but keep ER directions
- c) Correct the selection to the ER product as prescribed
- d) Split the IR tablets to mimic ER release

84) A technician receives an eRx for "Eliquis 5 mg BID." The will-call bin contains a bag for the same patient filled yesterday for "Xarelto 20 mg daily," not yet picked up. What should the technician do?

- a) Place both in will-call and let the patient decide
- b) Route for pharmacist review due to potential duplicate anticoagulant therapy
- c) Inactivate the older prescription and dispense the newer one
- d) Dispense the older one first and ignore the new order

85) Which medication is a long-acting insulin?

- a) insulin glargine
- b) insulin regular
- c) insulin lispro
- d) insulin aspart

86) A patient presents a prescription and asks the technician to change the dose because "my doctor told me to." What is the technician allowed to do?

- a) Change the dose and document the patient's statement
- b) Change the dose if the total daily amount stays similar
- c) Request the pharmacist to assist because a new order/clarification may be needed
- d) Dispense and instruct the patient to adjust at home

87) A prescription reads: "Take 7.5 mL PO TID x 10 days" of amoxicillin 250 mg/5 mL. What total volume should be dispensed?

- a) 150 mL
- b) 200 mL
- c) 225 mL
- d) 250 mL

88) When must a pharmacy complete an initial inventory of all controlled substances under federal law?

- a) Every month
- b) Every 6 months
- c) Every 2 years
- d) Every 5 years

89) A prescription is written for "Plavix 75 mg daily." Which generic should be selected?

- a) prasugrel

- b) ticagrelor
- c) clopidogrel
- d) dipyridamole

90) A prescription for a nasal spray reads: "Use 1 spray in each nostril BID." The bottle contains 60 metered sprays total. What is the days' supply?

- a) 10
- b) 15
- c) 20
- d) 30

Answer Key

1) b	32) c	63) c
2) c	33) d	64) d
3) b	34) b	65) c
4) b	35) b	66) b
5) b	36) c	67) d
6) c	37) b	68) c
7) c	38) b	69) b
8) c	39) d	70) d
9) c	40) a	71) d
10) b	41) c	72) d
11) b	42) c	73) a
12) b	43) a	74) b
13) d	44) b	75) b
14) c	45) b	76) c
15) b	46) d	77) a
16) b	47) c	78) c
17) c	48) d	79) b
18) c	49) b	80) d
19) b	50) b	81) a
20) b	51) b	82) b
21) c	52) c	83) c
22) a	53) c	84) b
23) d	54) a	85) a
24) a	55) d	86) c
25) b	56) d	87) c
26) b	57) c	88) c
27) b	58) a	89) c
28) c	59) d	90) b
29) c	60) b	
30) d	61) b	
31) b	62) a	

PTCB Sim Test 26

1) A new eRx reads: "Toprol-XL 50 mg: take 1 tablet daily." In data entry, the technician selects metoprolol tartrate 50 mg. What should be done before printing labels?

- a) Keep tartrate and add "ER" to the label
- b) Change directions to twice daily to match tartrate
- c) Switch to metoprolol succinate ER 50 mg to match Toprol-XL
- d) Dispense tartrate and note the change in patient comments

2) A patient requests "Plavix." Which product should be selected?

- a) clopidogrel
- b) prasugrel
- c) ticagrelor
- d) dipyridamole/aspirin

3) A written prescription for warfarin includes "INR in 3 days" in the notes section. Where should that information be placed during processing (if captured)?

- a) In the SIG exactly as written
- b) In a non-label notes field used for clinical communication
- c) Nowhere; delete it to avoid confusion
- d) In the quantity field so it prints for the patient

4) The profile lists "latex—anaphylaxis." The selected insulin pen device is labeled "contains natural rubber latex." What should the technician do next?

- a) Continue because latex allergies only apply to gloves
- b) Remove the allergy since it is not a medication allergy
- c) Hold the fill for pharmacist assessment of an alternative device/product
- d) Dispense and advise the patient to wear gloves when using it

5) Rx: amoxicillin 400 mg/5 mL. Sig: "7.5 mL PO BID x 10 days." How many mL should be dispensed?

- a) 120 mL
- b) 150 mL
- c) 160 mL
- d) 75 mL

6) A person at the counter says they're picking up for a friend and only knows the friend's DOB, not address. What is the best first step to avoid improper disclosure while still helping?

- a) Ask for another identifier and confirm pickup authorization before discussing details
- b) Read the medication names and ask which one sounds correct
- c) Ask them to call the patient on speakerphone and proceed immediately
- d) Refuse all third-party pickups without exception

7) An eRx is for hydroxyzine 25 mg, but the prescriber free-text note says "for muscle spasm." What should the technician do?

- a) Change the medication to cyclobenzaprine based on the note
- b) Remove the note and proceed with hydroxyzine

- c) Enter as written and bypass the note to avoid delays
- d) Send to pharmacist for clarification due to mismatch

8) Rx: Cymbalta 60 mg daily. Which generic should be entered?

- a) duloxetine
- b) venlafaxine ER
- c) desvenlafaxine
- d) bupropion XL

9) A patient presents a hard copy for promethazine with codeine syrup. It is signed and dated, but the prescriber DEA number is not on the paper. What is the best action?

- a) Process as written because signature is present
- b) Ask the patient to provide the prescriber's DEA number
- c) Place on hold for pharmacist review before any filling steps
- d) Add a DEA number from the prescriber directory without contacting anyone

10) A refill is requested for clonazepam with refills remaining. The prescription was written 8 months ago. Under federal law, what is correct?

- a) It may be refilled if refills remain, with no time limit
- b) It cannot be refilled beyond 6 months from the date written
- c) It can be refilled for 12 months if quantity is small
- d) It can be refilled once beyond 6 months if patient is stable

11) Prednisone taper: 10 mg tablets. "4 tabs daily x 3 days, then 3 tabs daily x 3 days, then 2 tabs daily x 3 days, then 1 tab daily x 3 days." How many tablets are needed?

- a) 24
- b) 30
- c) 36
- d) 40

12) A patient asks for "Advair Diskus." Which generic matches?

- a) budesonide/formoterol
- b) fluticasone/salmeterol
- c) fluticasone/vilanterol
- d) albuterol/ipratropium

13) A customer requests pseudoephedrine. What is required at sale under federal Combat Methamphetamine Epidemic Act baseline?

- a) Sell it from the OTC aisle with no documentation
- b) Verify ID and record the sale in a logbook/electronic log
- c) Require a prescription for any amount
- d) Report the purchase to DEA immediately

14) A prescriber calls requesting a transfer of an oxycodone prescription to your store. What is the correct handling expectation?

- a) Accept the transfer if the prescriber requests it
- b) Accept only if both pharmacies are in the same chain
- c) Schedule II prescriptions generally cannot be transferred between pharmacies

d) Accept if no fills have occurred yet

15) Rx: Synthroid 75 mcg daily. Which generic is correct?

- a) levothyroxine
- b) liothyronine
- c) methimazole
- d) propylthiouracil

16) A patient profile lists sertraline 100 mg daily. A new Rx arrives for fluoxetine 40 mg daily. The system flags "therapeutic duplication." What should the technician do?

- a) Override the alert if the patient says they are switching
- b) Discontinue sertraline in the profile and proceed
- c) Continue filling and ignore the alert because SSRIs are common
- d) Route the alert to the pharmacist for review before dispensing

17) Rx: "Apply 1 g topically TID. Dispense 90 g." What is the days' supply?

- a) 15 days
- b) 30 days
- c) 45 days
- d) 90 days

18) During filling, barcode scan shows "NDC mismatch." The selected stock bottle looks similar and the strength appears correct. What should the technician do?

- a) Bypass scanning and continue if strength matches
- b) Replace the label with a new one and re-scan later
- c) Stop and correct the product selection/NDC before proceeding
- d) Dispense and document the mismatch after pickup

19) A controlled substance hard copy has obvious overwriting in the quantity. What should happen next?

- a) Ask the patient what the prescriber intended
- b) Process for the smaller quantity to be safe
- c) Place it in a pharmacist review queue before entry/filling
- d) Void the prescription and tell the patient to return to prescriber

20) A patient asks the technician to "loan a few lisinopril tablets" until the refill is ready. What is the best response within technician scope?

- a) Provide a 3-day supply and document later
- b) Offer to request refill authorization and bring the request to pharmacist for any emergency supply decision
- c) Suggest cutting tablets in half until the new fill is ready
- d) Transfer the prescription automatically and dispense immediately

21) Sig: "instill ii gtt OU qhs." Which label direction is correct?

- a) Instill 2 drops in both eyes at bedtime
- b) Instill 2 drops in the right eye at bedtime
- c) Instill 1 drop in both eyes at bedtime
- d) Instill 2 drops in each eye every morning

22) Which medication is a benzodiazepine?

- a) buspirone
- b) hydroxyzine
- c) zolpidem
- d) alprazolam

23) A vaccine shipment arrives with a temperature indicator showing exposure above range during transit. What should the technician do?

- a) Refrigerate it and monitor for 24 hours
- b) Quarantine and label "DO NOT USE" pending guidance and documentation
- c) Freeze it to preserve potency
- d) Use it first to avoid waste

24) Lantus SoloStar: inject 22 units nightly. Dispense 1 box (5 pens, each 3 mL, U-100). What is the days' supply?

- a) 30 days
- b) 45 days
- c) 68 days
- d) 75 days

25) A patient requests a printout of all prescriptions filled last year for taxes. What is appropriate?

- a) Provide it after verifying identity using required identifiers
- b) Refuse because dispensing history is confidential even to the patient
- c) Provide only if two staff witness the request in writing
- d) Provide it to anyone who knows patient name and DOB

26) Rx: "Zofran ODT 4 mg." The system defaulted to ondansetron tablets (swallow). What should the technician do?

- a) Keep tablets; ODT and tablets are always interchangeable
- b) Change directions to "swallow whole" and continue
- c) Select the ODT dosage form before proceeding
- d) Substitute prochlorperazine due to availability

27) A refill is requested for Adderall XR with refills remaining. Federally, which statement is correct?

- a) Schedule II prescriptions may be refilled up to 5 times in 6 months
- b) Schedule II prescriptions cannot be refilled
- c) Schedule II prescriptions may be refilled once with prescriber approval
- d) Schedule II prescriptions may be refilled if diagnosis code is provided

28) Allergy profile: "penicillin—anaphylaxis." New Rx: amoxicillin/clavulanate. What is the best next step?

- a) Process because clavulanate reduces allergy risk
- b) Change to azithromycin automatically
- c) Hold and route for pharmacist evaluation due to severe documented allergy
- d) Fill if patient says they took it before, no further steps needed

29) Rx: methotrexate 2.5 mg "take 6 tablets weekly." The technician notices the entered SIG says "daily." What should be done?

- a) Correct to weekly and proceed without escalation
- b) Stop processing and route for pharmacist review due to high-alert weekly vs daily risk
- c) Change quantity to 42 and proceed
- d) Leave as entered and let patient clarify at pickup

30) Sig: "Take 2 tsp PO q6h prn cough." Dispense 240 mL. What is the maximum days' supply? (1 tsp = 5 mL)

- a) 6 days
- b) 8 days
- c) 10 days
- d) 12 days

31) A medication requires a Medication Guide each time it is dispensed. What is the technician's responsibility?

- a) Include it only on the first fill
- b) Include it only if the patient requests it
- c) Ensure it is provided with each dispense as required
- d) Replace it with the pharmacy leaflet

32) Which drug is an ACE inhibitor?

- a) losartan
- b) amlodipine
- c) lisinopril
- d) metoprolol

33) A prescriber's office calls asking if an HIV medication is ready. The caller provides only patient first name and DOB. What should the technician do?

- a) Provide status because it's a medical office
- b) Request additional required identifiers/authorization before disclosing any PHI
- c) Provide the medication name but not the status
- d) Provide only the pickup time

34) Azithromycin 200 mg/5 mL: "12.5 mL on day 1, then 6.25 mL daily on days 2–5." Total volume needed?

- a) 31.25 mL
- b) 37.5 mL
- c) 42.5 mL
- d) 50 mL

35) At pickup, a patient says: "My doctor increased it yesterday; I take two now." The Rx on file says 1 tablet daily. What should the technician do?

- a) Update SIG to 2 tablets daily and dispense
- b) Dispense double quantity and keep SIG unchanged
- c) Document the patient statement as prescriber approval and dispense
- d) Place the prescription for pharmacist follow-up before changes are made

36) The prescriber marked DAW 1 for a brand product. The patient requests generic due to cost. What is appropriate?

- a) Substitute generic because patient requested it
- b) Dispense generic but keep DAW 1 so insurance pays
- c) Keep brand on hold and have pharmacist/prescriber handle change request
- d) Split fill: half brand, half generic

37) During restocking, a technician finds nitroglycerin SL tablets in an open stock bottle with no "opened on" date. What should be done?

- a) Keep using until manufacturer expiration date
- b) Transfer tablets to a new bottle to restore stability
- c) Remove from active inventory for proper handling as questionable stock
- d) Refrigerate to extend beyond-use time

38) Which medication is a proton pump inhibitor?

- a) famotidine
- b) sucralfate
- c) omeprazole
- d) calcium carbonate

39) A patient brings an unlabeled vial of tablets and asks the technician to identify them. What is the best action?

- a) Use an online identifier and tell the patient the drug name
- b) Return them to stock if the tablets look intact
- c) Ask the patient to take one to see if it works
- d) Refer to the pharmacist and follow procedure for unknown medications

40) Rx: Eliquis 5 mg. Which generic is correct?

- a) rivaroxaban
- b) apixaban
- c) dabigatran
- d) clopidogrel

41) A pharmacist requests a partial fill of a Schedule II because the pharmacy is short on stock. In a typical retail setting, to dispense the remainder under federal rules, it must be provided within:

- a) 72 hours
- b) 7 days
- c) 30 days
- d) 6 months

42) Symbicort: inhale 2 puffs BID. Inhaler contains 120 actuations. Days' supply?

- a) 15 days
- b) 20 days
- c) 30 days
- d) 60 days

43) Two similar patient profiles exist: "John A. Smith" and "Jon A. Smith," same address, DOB differs by one digit. A new eRx arrives for "Jon." What should the technician do before processing?

- a) Merge profiles immediately
- b) Select the older profile to preserve history
- c) Confirm identifiers and attach the eRx to the correct patient record
- d) Create a third profile to avoid errors

44) A patient brings back unused hydrocodone tablets asking for a refund. What is correct?

- a) Accept return and restock if bottle is sealed
- b) Accept for disposal if permitted, but do not restock or refund as a prescription return
- c) Place them in regular trash after removing the label
- d) Tell the patient to flush them at home

45) Fentanyl 25 mcg/hr patch: apply 1 patch every 72 hours. Dispense #10. Days' supply?

- a) 10 days
- b) 20 days
- c) 30 days
- d) 72 days

46) A prescriber leaves an after-hours voicemail for oxycodone 5 mg. What should happen next day?

- a) Transcribe and fill because prescriber left a message
- b) Fill if the caller ID matches the clinic
- c) Process as a faxed prescription and file it
- d) Route to pharmacist because voicemail is not valid for Schedule II prescribing

47) Child weight 55 lb. Rx: ibuprofen 10 mg/kg per dose. What is the dose in mg? (1 kg = 2.2 lb)

- a) 250 mg
- b) 200 mg
- c) 150 mg
- d) 100 mg

48) Rx: Keflex 500 mg. Which generic should be selected?

- a) cefdinir
- b) cephalexin
- c) cefuroxime
- d) ceftriaxone

49) The prescription says morphine sulfate ER 30 mg. The selected product is morphine sulfate IR 30 mg. What should the technician do?

- a) Continue because the strength matches
- b) Adjust frequency to match ER dosing and proceed
- c) Re-select the correct ER formulation before continuing
- d) Add "do not crush" label and dispense IR

50) During a biennial controlled substance inventory, which task may a technician perform

under appropriate supervision and workflow?

- a) Independently certify/sign the inventory as PIC
- b) Count and record quantities as assigned and provide totals to the pharmacist
- c) Adjust perpetual inventory counts without documentation to match the count
- d) Destroy expired controls during the inventory to reduce discrepancies

51) In data entry, metformin ER 500 mg was prescribed but metformin IR 500 mg was selected. What is the best correction?

- a) Continue; ER and IR are equivalent if taken with food
- b) Change directions to once daily to approximate ER
- c) Dispense IR and counsel patient to split tablets
- d) Correct the dosage form selection before filling

52) Which medication is a beta blocker?

- a) hydrochlorothiazide
- b) carvedilol
- c) losartan
- d) lisinopril

53) A DSCSA notice indicates a specific lot may be illegitimate. What should the technician do?

- a) Dispense it if packaging looks intact
- b) Move it to the front so it sells quickly
- c) Quarantine and follow investigation/verification steps per procedure
- d) Return it to the shelf after checking expiration date

54) Sig: "instill 1 gtt OS BID." Which is correct?

- a) 1 drop in both eyes twice daily
- b) 1 drop in left eye twice daily
- c) 1 drop in right eye twice daily
- d) 2 drops in left eye twice daily

55) A patient wants OTC omeprazole while taking clopidogrel. What is the best technician action?

- a) Recommend omeprazole since it's OTC
- b) Suggest taking clopidogrel twice daily
- c) Tell the patient to stop clopidogrel while on omeprazole
- d) Refer the patient to the pharmacist due to interaction concern

56) Tamsulosin 0.4 mg: "take 1 capsule daily." Quantity #90. Days' supply?

- a) 90 days
- b) 45 days
- c) 60 days
- d) 30 days

57) Printed prescriptions are left on the counter where customers can see names and medications. What is the best corrective action?

- a) Turn them face down and keep them on the same counter

- b) Move them to a secure, non-public area immediately
- c) Shred them even if retention is required
- d) Ask customers not to look at the papers

58) Rx: Xarelto 20 mg daily with evening meal. Which generic is correct?

- a) edoxaban
- b) apixaban
- c) rivaroxaban
- d) dabigatran

59) A patient asks what Nitrostat is for and how to use it. What should the technician do?

- a) Provide full counseling and demonstrate use
- b) Tell the patient it's for daily blood pressure control
- c) Tell the patient to read the package insert and call later
- d) Get the pharmacist to counsel while helping coordinate minimal wait time

60) Eye drops: "1 drop OU QID." Bottle is 10 mL. Assume 20 drops/mL. Days' supply?

- a) 25 days
- b) 10 days
- c) 12 days
- d) 50 days

61) A new Rx arrives for lamotrigine. The allergy list says "Lamictal—Stevens-Johnson syndrome." What should be done?

- a) Proceed because brand allergy doesn't apply to generic
- b) Remove the allergy because it's not a true allergy
- c) Route for pharmacist evaluation before dispensing
- d) Substitute levetiracetam and fill

62) A request is received to transfer a noncontrolled prescription from another pharmacy. Which is the correct expectation?

- a) Technician can independently complete the transfer in all cases
- b) Technician may collect information, but pharmacist must complete/authorize as required
- c) Transfers are allowed only if both stores share the same computer system
- d) Transfers are not allowed for any refills

63) Which medication is a dihydropyridine calcium channel blocker?

- a) diltiazem
- b) verapamil
- c) amlodipine
- d) propranolol

64) Unit-dose packaging is being prepared for a facility. What is the most appropriate quality check before sending?

- a) Confirm label readability, correct drug/strength, and appropriate expiration/BUD for expected use
- b) Check only the first dose packaged for each medication
- c) Skip checks if barcode scanning exists at the facility

d) Check only controlled substances

65) Rx frequency includes “qod.” What should the technician enter on the label to reduce error risk?

- a) Enter “qod” exactly as written
- b) Translate to “every other day” on the label
- c) Change to “once daily” to avoid confusion
- d) Leave frequency blank and print as “use as directed”

66) Insulin aspart U-100: inject 14 units TID with meals. For a 30-day supply, what quantity should be billed in mL? (U-100 = 100 units/mL)

- a) 10 mL
- b) 12.6 mL
- c) 14 mL
- d) 42 mL

67) Allergy profile: “sulfonamide antibiotics—rash.” New Rx: sulfamethoxazole/trimethoprim. What should the technician do?

- a) Fill because it’s a common antibiotic and rash is mild
- b) Fill if the patient is an adult
- c) Change to nitrofurantoin automatically
- d) Route for pharmacist review due to documented sulfonamide antibiotic allergy

68) Metronidazole vaginal gel 0.75%: insert 1 applicatorful (5 g) qhs x 5 days. Package contains 70 g. Days’ supply for one package?

- a) 7 days
- b) 10 days
- c) 14 days
- d) 5 days

69) Refrigerator temperature logs were not recorded for two consecutive days. What should the technician do?

- a) Backfill missing values using today’s temperature
- b) Write “WNL” for the missing days and initial
- c) Document the omission and escalate per QA process for assessment
- d) Discard all refrigerated inventory immediately

70) Cefuroxime suspension: “10 mL PO BID x 7 days.” Total mL to dispense?

- a) 100 mL
- b) 140 mL
- c) 70 mL
- d) 210 mL

71) A patient asks for “Lasix.” Which generic should be selected?

- a) bumetanide
- b) spironolactone
- c) furosemide
- d) hydrochlorothiazide

72) An eRx for a controlled substance has unusual directions and the prescriber address does not match what's on file. What is the best next step?

- a) Fill because it is electronic and therefore verified
- b) Dispense a partial supply while investigating later
- c) Call the patient and ask if they are abusing medication
- d) Send to pharmacist for review due to red flags

73) Rx: ketorolac 10 mg "take 1 tablet QID x 10 days." What should the technician do?

- a) Enter and fill exactly as written
- b) Change to ibuprofen automatically
- c) Route to pharmacist review due to typical duration/safety concerns
- d) Change directions to PRN and fill

74) A technician is applying a manufacturer coupon after insurance has already paid. What is the most appropriate billing workflow?

- a) Add coupon as secondary payer without reversing the paid claim
- b) Reverse and rebill appropriately if allowed by payer/coupon terms
- c) Change the cash price manually while keeping the paid claim
- d) Tell the patient coupons can never be used with insurance

75) Which is a long-acting inhaled anticholinergic used for COPD maintenance?

- a) albuterol
- b) montelukast
- c) tiotropium
- d) fluticasone

76) A recall notice specifies a particular lot of losartan 50 mg. What should the technician do?

- a) Pull all losartan products regardless of lot
- b) Continue dispensing until replacements arrive
- c) Quarantine only the affected lot(s) and prevent dispensing per recall workflow
- d) Return recalled product to the shelf after noting the recall

77) Rx: "Tylenol #3." Which product is correct?

- a) acetaminophen/codeine 300 mg/30 mg
- b) acetaminophen/codeine 300 mg/15 mg
- c) acetaminophen/hydrocodone 325 mg/5 mg
- d) acetaminophen/oxycodone 325 mg/5 mg

78) Another pharmacy calls to transfer a Schedule III prescription. What is the correct approach?

- a) Technician may complete the transfer without pharmacist involvement
- b) Transfers are never allowed for Schedule III prescriptions
- c) Accept only if the patient is present with ID
- d) Gather transfer details and route to pharmacist to complete/authorize per requirements

79) Doxycycline hyclate 100 mg: take 1 capsule BID. Quantity #14. Days' supply?

- a) 5
- b) 7
- c) 10
- d) 14

80) Gabapentin 300 mg: take 1 capsule TID. Quantity #90. Days' supply?

- a) 15
- b) 90
- c) 60
- d) 30

81) An order specifies U-500 insulin, but U-100 regular insulin was selected. What should the technician do?

- a) Continue; units are equivalent regardless of concentration
- b) Adjust the SIG to match U-100 volume and continue
- c) Correct product selection and route for pharmacist review due to high-risk concentration mismatch
- d) Dispense U-100 since it is safer than U-500

82) A shipment includes a hazardous drug with special handling on the outer package. What should the technician do during receiving/stocking?

- a) Stock quickly to minimize handling time
- b) Use required PPE and handling procedures during receiving and stocking
- c) Remove warning labels so staff aren't alarmed
- d) Place it in the waiting area until space is available

83) An eRx comes through: "clindamycin 150 mg/mL; give 5 mL TID." The outpatient products available don't match that concentration. What is the best action?

- a) Select any clindamycin suspension and keep the SIG
- b) Convert to capsules and calculate an equivalent dose
- c) Change 150 mg/mL to 150 mg/5 mL automatically
- d) Route to pharmacist for clarification because the concentration appears incorrect

84) The profile shows active glipizide. A new prescription arrives for glyburide, same day, different prescriber. What should the technician do?

- a) Fill both to improve glucose control
- b) Hold for pharmacist evaluation due to likely therapeutic duplication
- c) Delete the older medication and fill the new one
- d) Dispense both and label one "as needed"

85) A patient says they used to take "Crestor." The new Rx is rosuvastatin 20 mg daily. Which is correct?

- a) Crestor is simvastatin
- b) Crestor is atorvastatin
- c) Crestor is rosuvastatin
- d) Crestor is pravastatin

86) An oral antibiotic requires reconstitution: "Add water to yield 100 mL." The technician

accidentally adds 100 mL of water (instead of adding to final volume). What should be done next?

- a) Dispense; the final concentration is close enough
- b) Add extra powder from another bottle to correct it
- c) Remove some liquid and re-add water to the line
- d) Set aside for pharmacist assessment; product may be incorrect concentration

87) Rx: "Take 1.5 tablets daily." For a 30-day supply, how many tablets are needed?

- a) 30
- b) 40
- c) 45
- d) 60

88) Which medication is an SSRI?

- a) venlafaxine
- b) duloxetine
- c) amitriptyline
- d) sertraline

89) Under DSCSA requirements, what record type must be retrievable upon request for tracing?

- a) OTC sales logs for 2 years
- b) Transaction information/transaction statement for prescription drugs for the required retention period
- c) Daily perpetual inventory logs for all legend drugs
- d) Patient counseling documentation for 10 years

90) A prescriber's office asks the pharmacy to fax a patient's controlled substance fill history for "charting." No patient authorization is on file and the fax number isn't verified. What is the best action?

- a) Fax it because it is for treatment purposes
- b) Fax only last fill date, not medication names
- c) Provide it verbally over the phone instead of faxing
- d) Use the required authorization/verification process before releasing any information

Answer Key

1) c	31) c	61) c
2) a	32) c	62) b
3) b	33) b	63) c
4) c	34) b	64) a
5) b	35) d	65) b
6) a	36) c	66) b
7) d	37) c	67) d
8) a	38) c	68) d
9) c	39) d	69) c
10) b	40) b	70) b
11) c	41) a	71) c
12) b	42) a	72) d
13) b	43) c	73) c
14) c	44) b	74) b
15) a	45) c	75) c
16) d	46) d	76) c
17) b	47) a	77) a
18) c	48) b	78) d
19) c	49) c	79) b
20) b	50) b	80) d
21) a	51) d	81) c
22) d	52) b	82) b
23) b	53) c	83) d
24) c	54) b	84) b
25) a	55) d	85) c
26) c	56) a	86) d
27) b	57) b	87) c
28) c	58) c	88) d
29) b	59) d	89) b
30) b	60) a	90) d

PTCB Sim Test 27

1. A new eRx is received for “metoprolol succinate ER 25 mg: take 1 tablet daily.” While filling, the technician scans a bottle of metoprolol tartrate 25 mg and the system rejects it as a mismatch. What should the technician do next?

- a) Select metoprolol succinate ER 25 mg and rescan the correct product
- b) Proceed with tartrate and add an “ER” auxiliary label
- c) Change directions to “take 1 tablet twice daily” and continue
- d) Bypass the scan since the strength matches

2. A patient picking up says, “I’m here for my wife,” and provides her name and date of birth but is not listed as an authorized pickup person. What is the best technician response?

- a) Ask the spouse to describe the medication and release if correct
- b) Release the prescription because DOB matched
- c) Politely decline and offer to contact the patient to add authorization or have the patient call
- d) Provide a “ready/not ready” status only and release if it is ready

3. A prescription label prints “levothyroxine 0.075 mg daily.” Which strength should be selected?

- a) 7.5 mcg
- b) 0.75 mg
- c) 75 mcg
- d) 0.075 mcg

4. A prescription reads: “Amoxicillin/clavulanate 875/125 mg: take 1 tablet by mouth twice daily for 7 days.” What quantity should be dispensed?

- a) 21
- b) 14
- c) 28
- d) 7

5. A patient is prescribed: “Take 5 mL by mouth three times daily. Dispense 150 mL.” What is the days’ supply?

- a) 10
- b) 15
- c) 20
- d) 30

6. During data entry, a new prescription comes in for methotrexate 2.5 mg tablets with directions typed by the prescriber’s office as “take 6 tablets daily.” What should the technician do?

- a) Enter as written and add “high dose” in the notes
- b) Change the frequency to weekly since that’s typical and process
- c) Put the prescription in a pharmacist review queue due to high-alert dosing concern
- d) Call the patient to confirm the prescriber’s intent and then enter it

7. At the counter, a patient asks, “Can I take ibuprofen with my warfarin?” What should the

technician do?

- a) Recommend naproxen instead of ibuprofen
- b) Suggest taking ibuprofen with food to reduce risk
- c) Tell the patient to separate doses by at least 2 hours
- d) Offer pharmacist consultation for the interaction question

8. A controlled-substance order arrives and the tote seal is broken. The invoice lists several C-II items. What should the technician do first?

- a) Accept the delivery and reconcile counts later
- b) Quarantine the tote and escalate for pharmacist/manager verification before check-in
- c) Stock noncontrols and return the controls to the wholesaler
- d) Refuse the entire delivery immediately without documenting

9. A prescription is written for "diltiazem ER 240 mg daily." Only diltiazem IR 120 mg is on the shelf. What is the best technician action?

- a) Dispense IR 120 mg and label "take 2 daily"
- b) Dispense IR 120 mg with an "ER" auxiliary label
- c) Place on hold until the ER product can be ordered in
- d) Ask the patient if they prefer twice-daily dosing

10. The patient profile shows "penicillin—anaphylaxis." A new prescription for cephalexin is received. What should the technician do?

- a) Process normally because cephalexin is not a penicillin
- b) Remove the allergy since the prescriber ordered the medication
- c) Change the medication to clindamycin and continue processing
- d) Route the prescription for pharmacist assessment of allergy risk

11. Under the Combat Methamphetamine Epidemic Act (CMEA), which is required when selling pseudoephedrine products?

- a) Purchaser must show valid ID and sign the log (paper/electronic)
- b) Purchaser must present a prescription
- c) Only a pharmacist may conduct the sale in all settings
- d) Product must be stored in a refrigerator until sold

12. A label prints for "hydralazine 25 mg." The technician accidentally selects "hydroxyzine 25 mg." Which safeguard most directly prevents this error from reaching the patient?

- a) Counting on a clean tray
- b) Using larger font labels
- c) Barcode scanning the stock bottle NDC against the label
- d) Adding a "may cause drowsiness" auxiliary label

13. An inhaler contains 120 actuations. Directions: "Inhale 2 puffs twice daily." What is the days' supply?

- a) 60
- b) 30
- c) 120
- d) 15

14. A patient requests an early refill for zolpidem claiming the bottle was stolen. The claim rejects as "refill too soon." What should the technician do?

- a) Change the days' supply to force approval
- b) Offer to transfer the prescription to another pharmacy to see if it processes
- c) Tell the patient the pharmacy cannot help with any early refills
- d) Gather details and route the request for pharmacist decision-making/documentation

15. A prednisone taper says: "10 mg tablets: take 4 tablets daily for 3 days, then 3 tablets daily for 3 days, then 2 tablets daily for 3 days, then 1 tablet daily for 3 days." What quantity is needed?

- a) 30
- b) 36
- c) 24
- d) 42

16. DEA Form 222 is used for:

- a) Reporting theft or significant loss of controlled substances
- b) Ordering Schedule II controlled substances
- c) Documenting transfers of Schedule III–V prescriptions
- d) Registering a pharmacy with the DEA

17. A patient returns an opened, labeled vial of tablets and asks the technician to "put them back on the shelf." What is the correct action?

- a) Return to stock if the tablets look intact
- b) Return to stock only if it is not a controlled substance
- c) Decline to return to stock and process through appropriate disposal/return procedure
- d) Repackage into a manufacturer bottle and record the count

18. "Humalog KwikPen: inject 10 units with meals three times daily. Dispense 1 box." The box contains 5 pens, each 3 mL, concentration 100 units/mL. What is the days' supply?

- a) 30
- b) 45
- c) 50
- d) 75

19. A technician prints a patient's profile to resolve an insurance issue. When finished, what should be done with the printout?

- a) Put it in regular trash since it's no longer needed
- b) Place it in the will-call bag for the patient
- c) File it at the workstation for 30 days
- d) Dispose of it using secure HIPAA-compliant shredding/containers

20. A prescription reads: "Take 1 teaspoonful by mouth every 6 hours as needed for cough. Dispense 120 mL." Using maximum frequency, what is the days' supply? (1 tsp = 5 mL)

- a) 6
- b) 4
- c) 5
- d) 10

21. While entering a hard copy, the technician types "morphine ER 30 mg" but the prescription clearly states "morphine IR 30 mg." What should the technician do?
- a) Leave it; final verification will catch it
 - b) Change the directions to q12h to fit ER
 - c) Correct the entry to morphine IR 30 mg before it proceeds
 - d) Keep ER because it reduces dosing frequency
22. A prescriber's office calls asking if a patient picked up an HIV medication. The caller cannot verify any identifiers beyond the patient's name. What should the technician do?
- a) Provide pickup status but not the drug name
 - b) Decline to disclose and request proper verification/authorization
 - c) Provide the medication name only
 - d) Provide last fill date since it's not clinical information
23. A refrigerated biologic arrives with a temperature indicator showing it was out of range for several hours in transit. What should the technician do?
- a) Stock it immediately and mark "use first"
 - b) Place it in the freezer to stabilize it quickly
 - c) Dispense it only if the patient will use it within 48 hours
 - d) Quarantine and route for appropriate disposition before use
24. Which is the generic for Eliquis?
- a) rivaroxaban
 - b) dabigatran
 - c) apixaban
 - d) edoxaban
25. A prescription reads: "Clindamycin oral solution: take 10 mL by mouth three times daily for 7 days." What total volume should be dispensed?
- a) 140 mL
 - b) 210 mL
 - c) 280 mL
 - d) 70 mL
26. A patient's friend asks on the phone, "Is my friend's refill ready?" The friend is not listed as authorized. What is the best technician response?
- a) Provide "ready/not ready" only since no drug names are given
 - b) Provide status if the caller states the patient's DOB
 - c) Provide status if the caller states the prescriber's name
 - d) Decline to disclose and offer steps for the patient to add authorization
27. Under federal rules, which statement about Schedule II prescriptions is correct?
- a) They may be refilled up to 5 times in 6 months
 - b) They may be refilled once with prescriber approval
 - c) They may be transferred one time between pharmacies
 - d) They cannot be refilled; a new prescription is required

28. A prescription is for bupropion SR 150 mg. The technician selected bupropion XL 150 mg. What should be done?

- a) Continue since SR and XL are equivalent at the same strength
- b) Change directions so XL matches SR dosing
- c) Select bupropion SR as written before processing continues
- d) Dispense XL and add "do not crush"

29. A patient reports an allergy to erythromycin. A new prescription arrives for azithromycin. What should the technician do?

- a) Process normally because azithromycin is a different class
- b) Route for pharmacist review due to possible macrolide cross-reactivity
- c) Delete the erythromycin allergy because it's outdated
- d) Substitute doxycycline and continue

30. Eye drops are prescribed: "Instill 1 drop in both eyes four times daily. Dispense 10 mL." If 20 drops = 1 mL, what is the days' supply?

- a) 12
- b) 25
- c) 10
- d) 50

31. Under DSCSA, transaction information is missing for a received shipment of a prescription legend drug. What should the technician do?

- a) Request the missing transaction documentation before the product is distributed/dispensed
- b) Dispense it but store it on a separate shelf
- c) Create transaction data based on the invoice
- d) Ignore it if it is from the usual wholesaler

32. The sig code "qod" means:

- a) once daily
- b) at bedtime
- c) every other day
- d) four times daily

33. A warfarin prescription is written "take as directed." The patient asks, "How many do I take tonight?" What is the best technician response?

- a) Tell them to take 1 tablet nightly
- b) Read the last directions from the previous label and recommend the same
- c) Ask when their next INR is and adjust based on that
- d) Offer pharmacist consultation for dosing clarification

34. A patient presents a paper prescription for a C-II stimulant dated 7 months ago. The technician is unsure about state limits for filling older C-IIs. What should be done?

- a) Fill it because federal law allows 1 year
- b) Refuse because federal law limits to 30 days
- c) Backdate the fill date to make it valid
- d) Route for pharmacist evaluation of validity per state/federal requirements

35. The label directions say: "Take 1.5 tablets daily." Each tablet is 20 mg. How many mg per day is this?

- a) 10 mg
- b) 30 mg
- c) 20 mg
- d) 40 mg

36. A manufacturer recall is issued for one lot of losartan due to an impurity. What is the best technician action?

- a) Pull and quarantine only the affected lot and prevent dispensing from it
- b) Remove all lots of losartan to be safe
- c) Continue dispensing until the replacement shipment arrives
- d) Return the entire stock to patients for exchange

37. A prescription reads: "Take 1 tablet by mouth twice daily. Dispense #120." What is the days' supply?

- a) 120
- b) 30
- c) 60
- d) 90

38. During unit-dose cart fill, two different strengths of the same medication are found in one patient's cassette bin. What should the technician do?

- a) Leave them; nursing will choose the right one
- b) Remove the incorrect strength and correct the bin contents per fill list
- c) Put a note on the bin and continue filling
- d) Move the extra strength to the back of the bin

39. Which pair is a common look-alike/sound-alike risk?

- a) metformin and meloxicam
- b) amlodipine and amoxicillin
- c) pantoprazole and paroxetine
- d) hydralazine and hydroxyzine

40. A prescription is written for Synthroid with "DAW" indicated. What should the technician do during entry?

- a) Select generic levothyroxine and ignore DAW
- b) Select Synthroid and apply the appropriate DAW code/indicator
- c) Select generic and add a note "brand requested"
- d) Change to another brand of levothyroxine

41. A wholesaler delivery includes a bottle with a broken tamper-evident seal. What is the best technician action?

- a) Stock it if the cap is still tightly closed
- b) Put it in will-call to use first
- c) Dispense it only if the patient declines counseling
- d) Quarantine as damaged/suspect product and route for disposition

42. A caller tries to phone in a new prescription for oxycodone/acetaminophen (C-II). What should the technician do?

- a) Take the order and read it back
- b) Ask for the prescriber's NPI and proceed
- c) Transfer the call so the pharmacist can handle it if permitted
- d) Tell the caller to fax it; verbal C-IIs are never allowed

43. A prescription is for furosemide 40 mg daily #90, but only 20 mg tablets are available. What should the technician do?

- a) Dispense #180 of 20 mg and change the sig to "take 2 daily"
- b) Ask the patient if two tablets daily is acceptable, then proceed
- c) Dispense #90 of 20 mg and keep the same directions
- d) Route for pharmacist authorization and/or obtain correct strength before any change

44. A technician almost selects tamoxifen instead of tamsulosin. Which system practice best reduces this risk?

- a) Store all medications strictly alphabetically without separation
- b) Use Tall Man lettering and barcode scanning during product selection
- c) Use brand names only on shelf labels
- d) Remove one of the medications from inventory

45. Tylenol #3 contains:

- a) acetaminophen + oxycodone
- b) ibuprofen + codeine
- c) acetaminophen + codeine
- d) acetaminophen + hydrocodone

46. Azithromycin 250 mg is prescribed: "Take 2 tablets today, then 1 tablet daily for 4 days." What quantity should be dispensed?

- a) 5
- b) 6
- c) 7
- d) 10

47. In nonsterile compounding documentation, the technician realizes the lot number for one ingredient was not recorded. The preparation has not been dispensed yet. What should the technician do?

- a) Dispense it; lot numbers are optional for nonsterile compounds
- b) Estimate the lot number from the invoice date
- c) Record the correct lot number from the ingredient container before release
- d) Record the lot number only if a patient reports a reaction

48. An eRx arrives for lisinopril 10 mg daily. The profile indicates the patient is pregnant. What should the technician do?

- a) Process normally and let counseling address it
- b) Change the medication to labetalol and continue
- c) Ensure the alert is visible and route for pharmacist review before dispensing

d) Delete the pregnancy entry to reduce false alerts

49. While stocking, a technician finds a vaccine that should be refrigerated was placed in the freezer overnight. What should be done?

- a) Thaw it and use if it looks normal
- b) Move it to the refrigerator and use after 24 hours
- c) Use it if it is within the printed expiration date
- d) Quarantine and route for stability guidance and documentation

50. A prescription direction includes "HS." Which interpretation is correct?

- a) at bedtime
- b) every morning
- c) every 4 hours
- d) before meals

51. A patient requests transfer of a gabapentin prescription with refills remaining (assume state law allows). Under federal rules for noncontrolled prescriptions, which is correct?

- a) Transfers are not allowed for any prescription originally sent electronically
- b) Transfer is allowed, and the receiving pharmacy may dispense remaining refills
- c) Transfer requires a DEA Form 222
- d) Transfer is allowed only after all refills are used

52. A prescription is written for promethazine syrup, but promethazine suppositories were selected during product selection. What should the technician do?

- a) Dispense suppositories since they're therapeutically similar
- b) Ask the patient which form they prefer and proceed
- c) Change the route on the label to rectal and continue
- d) Stop and select the correct dosage form before proceeding

53. During the required controlled-substance inventory, which is true under federal rules regarding Schedule II?

- a) Estimated counts are acceptable for Schedule II
- b) Exact counts are required for Schedule II controlled substances
- c) Schedule II can be inventoried as one combined total with Schedule III–V
- d) Only invoices are needed; physical counts are optional

54. A technician finds OTC nicotine gum with a torn package on the shelf. What should be done?

- a) Tape it and return it to the shelf
- b) Mark it down and place it by the register
- c) Remove from sale and process as damaged/return per procedure
- d) Put it behind the counter and sell it only to adults

55. Heparin 25,000 units in 250 mL equals how many units per mL?

- a) 50 units/mL
- b) 75 units/mL
- c) 100 units/mL
- d) 125 units/mL

56. In sterile compounding, an alcohol swab packet is found already opened on the cleanroom counter. What is the correct action?

- a) Use it if it looks clean
- b) Use it only for wiping outer packaging
- c) Reseal it and use it within 1 hour
- d) Discard it and obtain a new intact swab

57. A patient profile shows sertraline as an active medication. A new prescription arrives for paroxetine. What should the technician do?

- a) Process both; SSRIs are commonly combined
- b) Automatically inactivate sertraline
- c) Enter paroxetine as written and route for pharmacist review for duplication/therapy change
- d) Change paroxetine to escitalopram

58. Lantus SoloStar is prescribed: "Inject 18 units subcutaneously nightly. Dispense 2 pens." Each pen is 3 mL of 100 units/mL. What is the days' supply?

- a) 33
- b) 25
- c) 20
- d) 30

59. A subpoena arrives requesting patient prescription records. What should the technician do?

- a) Release the records immediately to avoid being in contempt
- b) Refuse because records can never be released
- c) Provide only controlled-substance records
- d) Route the request for proper handling and authorized release

60. A patient insists they never picked up a medication, but the register shows it was sold yesterday. What should the technician do first?

- a) Reverse the claim immediately
- b) Refill the medication for free and rebill
- c) Review transaction details (time, register, signature/ID documentation) to investigate
- d) Delete the prescription record to avoid an audit

61. What is the generic for Plavix?

- a) prasugrel
- b) ticagrelor
- c) clopidogrel
- d) warfarin

62. A prescription reads: "Take 2 capsules by mouth three times daily for 7 days. Dispense #84." Is the quantity consistent with directions?

- a) No; it should be #42
- b) Yes; #84 matches 7 days
- c) No; it should be #98
- d) Yes; #84 matches 14 days

63. A prescription is written for atorvastatin #30 with 5 refills. The patient requests a 90-day supply, and the plan requires 90 days for maintenance fills. What should the technician do?

- a) Change to #90 and keep refills without authorization
- b) Dispense #90 by changing the directions to "take 3 tablets daily"
- c) Send a request/route per protocol to obtain authorization for a 90-day quantity
- d) Dispense #30 and instruct the patient to cash pay the rest

64. A technician accidentally touched tablets with bare hands while counting. What should be done?

- a) Continue; tablets are coated
- b) Wipe tablets with alcohol and dispense
- c) Place the touched tablets in a separate vial with a warning label
- d) Discard the affected tablets and restart using proper counting technique

65. "Ondansetron ODT 4 mg" — ODT means:

- a) orally disintegrating tablet
- b) oral delayed tablet
- c) ophthalmic dosing tablet
- d) oral dosing tincture

66. A customer asks to purchase syringes without a prescription. Which is the best technician response?

- a) Sell them without restrictions because syringes are never regulated
- b) Require the patient to state their diagnosis and document it
- c) Follow applicable state law and store procedures for nonprescription syringe sales
- d) Refuse because federal law prohibits selling syringes without a prescription

67. DEA Form 106 is used to:

- a) Order Schedule II medications
- b) Report theft or significant loss of controlled substances
- c) Register a new pharmacy location
- d) Document a controlled-substance transfer

68. Eye drops are prescribed: "Instill 1 drop in both eyes twice daily. Dispense 5 mL." If 20 drops = 1 mL, what is the days' supply?

- a) 25
- b) 50
- c) 12
- d) 100

69. A DUR alert flags a major interaction: the patient has sildenafil active and a new prescription for nitroglycerin patches is being processed. What should the technician do?

- a) Override the alert because the prescriber ordered it
- b) Put both medications on hold automatically without review
- c) Tell the patient to stop sildenafil and continue processing
- d) Route the alert for pharmacist evaluation before dispensing

70. Amoxicillin/clavulanate 400 mg/57 mg per 5 mL is prescribed: "Give 7.5 mL by mouth twice daily for 10 days." What quantity should be dispensed?

- a) 100 mL
- b) 150 mL
- c) 200 mL
- d) 75 mL

71. A prescription is for digoxin 0.125 mg daily. The stock bottle is labeled digoxin 125 mcg. Which is correct?

- a) 125 mcg equals 0.0125 mg
- b) 125 mcg equals 0.125 mg
- c) mcg and mg cannot be converted
- d) 125 mcg equals 1.25 mg

72. A Schedule III–V prescription transfer is requested and state law permits it. Under federal rules, which is correct?

- a) It may be transferred unlimited times if refills remain
- b) It may be transferred one time for the purpose of refill dispensing
- c) It may not be transferred if it was originally e-prescribed
- d) It may only be transferred after all refills are used

73. A prescription says "QID," but it was entered as "QD." What should the technician do?

- a) Correct the directions to match the prescription before it proceeds
- b) Leave it and rely on patient counseling
- c) Keep QD to improve adherence
- d) Call the patient to determine what they prefer

74. A new prescription reads "metformin ER 500 mg: take 2 tablets with evening meal." The profile shows metformin IR 500 mg "1 tablet twice daily" filled last week. What should the technician do during processing?

- a) Automatically discontinue metformin IR
- b) Enter metformin IR 500 mg with modified directions since the dose is similar
- c) Enter as written and ensure it is reviewed as a potential therapy change/duplication
- d) Combine directions into "1 tablet AM and 2 tablets PM"

75. A prescription for cephalexin is received for a patient with "penicillin—rash" documented. What should the technician do?

- a) Process without flags because it is not a penicillin
- b) Delete the allergy because rash is not serious
- c) Route for pharmacist review due to allergy history and potential cross-sensitivity
- d) Substitute azithromycin and continue

76. Augmentin is:

- a) amoxicillin
- b) amoxicillin/clavulanate
- c) cefuroxime axetil
- d) ampicillin/sulbactam

77. A faxed prescription for hydrocodone/APAP arrives, but the image is missing the prescriber signature. What should the technician do?

- a) Fill it because the clinic fax header is present
- b) Add the prescriber's name to the signature line and process
- c) Treat it as a verbal order and proceed
- d) Route for resolution because the prescription is incomplete as received

78. During entry, the technician selects lamotrigine 150 mg instead of lamivudine 150 mg. What is the best next step?

- a) Continue; both are commonly used long-term
- b) Dispense both to avoid delay
- c) Correct the drug selection to match the prescription before proceeding
- d) Change to lamotrigine 25 mg to be safer

79. The profile shows an active alprazolam prescription. A new lorazepam prescription triggers a duplicate-therapy alert. What should the technician do?

- a) Override to reduce wait time
- b) Remove alprazolam from the profile to clear the alert
- c) Dispense lorazepam and place alprazolam on hold
- d) Route the alert for pharmacist evaluation before dispensing

80. A liquid antibiotic is prescribed: "Take 7.5 mL by mouth twice daily for 10 days." If the bottle contains 150 mL, how much will remain after the full course?

- a) 0 mL
- b) 15 mL
- c) 30 mL
- d) 75 mL

81. A customer asks for an OTC recommendation for new chest pain that started this morning. What should the technician do?

- a) Recommend an antacid and hydration
- b) Recommend ibuprofen and rest
- c) Recommend acetaminophen and schedule a routine appointment
- d) Promptly involve the pharmacist and advise urgent medical evaluation

82. The pharmacy is out of stock of a C-II medication and a partial fill may be needed. What is the best technician action?

- a) Partially fill now and automatically dispense the remainder when it arrives without further steps
- b) Transfer the prescription to another pharmacy with no documentation
- c) Convert the remainder to refills on the same prescription
- d) Route for pharmacist handling to ensure rules and documentation for partial fills are met

83. A technician must count a hazardous medication (e.g., finasteride) for dispensing. What is the best practice to reduce exposure?

- a) Count by hand to avoid contaminating a tray
- b) Use dedicated equipment and appropriate PPE/containment practices
- c) Crush tablets before counting to reduce errors

d) Open capsules and count the contents

84. What is the generic for Nexium?

- a) omeprazole
- b) pantoprazole
- c) lansoprazole
- d) esomeprazole

85. A new patient eRx prints with an address and phone number that differ from a similar-name patient already in the system. What should the technician do?

- a) Ignore the eRx demographics; they are often wrong
- b) Update the existing profile automatically to match the eRx
- c) Delete the eRx and request it be resent
- d) Verify identifiers and demographics before merging or updating any profile

86. An order is for vancomycin 1 g IV. Available vials are 500 mg each. How many vials are needed for one 1 g dose?

- a) 1
- b) 2
- c) 3
- d) 4

87. Some opioid products require an FDA-required handout to be given each time they are dispensed. What is this handout called?

- a) Vaccine Information Statement (VIS)
- b) Medication Guide
- c) DEA Form 222
- d) Transaction Statement

88. A prescription reads: "Take 1 capsule by mouth twice daily. Dispense #60." How long will it last?

- a) 15 days
- b) 20 days
- c) 30 days
- d) 60 days

89. A customer attempts to buy dextromethorphan-containing cough syrup, and the register prompts for age verification. What should the technician do?

- a) Override the prompt; it applies only to pseudoephedrine
- b) Require a prescription for dextromethorphan
- c) Refuse the sale to anyone under 21 in all states
- d) Follow the prompt and applicable state/store requirements for sale

90. A label prints for "hydralazine 10 mg," and both 10 mg and 25 mg bottles are stored next to each other. Which action most reduces wrong-strength selection?

- a) Pull both bottles and compare tablet appearance
- b) Count in multiples of five to reduce errors
- c) Barcode scan the stock bottle NDC against the label before counting

d) Use a larger vial so the label is easier to read

Answer Key

1) a
2) c
3) c
4) b
5) a
6) c
7) d
8) b
9) c
10) d
11) a
12) c
13) b
14) d
15) b
16) b
17) c
18) c
19) d
20) b
21) c
22) b
23) d
24) c
25) b
26) d
27) d
28) c
29) b
30) b

31) a
32) c
33) d
34) d
35) b
36) a
37) c
38) b
39) d
40) b
41) d
42) c
43) d
44) b
45) c
46) b
47) c
48) c
49) d
50) a
51) b
52) d
53) b
54) c
55) c
56) d
57) c
58) a
59) d
60) c

61) c
62) b
63) c
64) d
65) a
66) c
67) b
68) b
69) d
70) b
71) b
72) b
73) a
74) c
75) c
76) b
77) d
78) c
79) d
80) c
81) d
82) d
83) b
84) d
85) d
86) b
87) b
88) c
89) d
90) c

PTCB Sim Test 28

1) A new eRx is received: "metoprolol succinate ER 50 mg: take 1 tablet twice daily." The patient's profile shows metoprolol tartrate 50 mg "1 tablet twice daily" filled last month. What is the best technician action during data entry?

- a) Enter as written and queue for pharmacist review due to formulation/frequency concern
- b) Substitute tartrate because it matches the prior fill
- c) Change directions to once daily because succinate is ER
- d) Cancel the eRx as a duplicate therapy

2) A patient presents an out-of-state paper prescription for hydrocodone/acetaminophen. The DEA number is present, but the prescriber address is missing. What is the best next step for the technician?

- a) Add the missing address using an online directory and continue
- b) Enter it and annotate "address missing" for later
- c) Hold for pharmacist review because required prescriber information is incomplete
- d) Refuse to accept any out-of-state controlled prescriptions

3) A refill request comes in for sertraline 100 mg. The last fill was 30 tablets with directions "take 1 tablet daily," but the new directions on the prescriber response are "take 1/2 tablet daily." What should the technician do to prevent an incorrect label?

- a) Keep prior directions to avoid confusion
- b) Update the directions exactly as received and route for pharmacist verification
- c) Change the quantity to 15 so it matches half-tablet dosing
- d) Delete the refill and ask the patient to bring a new prescription

4) While filling levothyroxine 75 mcg, the NDC scanned does not match the NDC on the label, although the strength matches. What is the best technician action?

- a) Override the mismatch because strength matches
- b) Reprint the label with the scanned NDC and proceed
- c) Re-select the correct product/NDC per system and restart the fill
- d) Cross out the NDC on the label and initial it

5) A patient requests pseudoephedrine 30 mg tablets. What must occur before sale under federal requirements?

- a) The patient must present a valid photo ID and sign the logbook/electronic log
- b) The patient must have a prescription
- c) The patient must be 21 years old
- d) The patient must provide proof of address

6) A prescription is for amoxicillin/clavulanate suspension 600-42.9 mg/5 mL: "5 mL PO BID x 10 days." How many mL should be dispensed?

- a) 80 mL
- b) 90 mL
- c) 100 mL
- d) 120 mL

7) At pickup, a patient asks how to use their new albuterol HFA inhaler. What is the most appropriate technician response?

- a) Demonstrate technique and counsel on priming and dosing
- b) Offer to have the pharmacist counsel and provide the device instructions
- c) Tell the patient to watch an online video later
- d) Ask the patient if they have asthma or COPD before answering

8) A prescription for clindamycin 150 mg capsules reads: "1 cap PO QID x 7 days." What quantity should be dispensed?

- a) 21
- b) 24
- c) 28
- d) 56

9) A patient's profile lists "anaphylaxis to sulfa." A new prescription arrives for sulfamethoxazole/trimethoprim DS. What is the best technician action?

- a) Fill it because allergy notes are often inaccurate
- b) Ask the patient if they can "tolerate it now" and proceed if yes
- c) Enter the prescription and override the allergy warning
- d) Stop processing and ensure the pharmacist evaluates the allergy conflict

10) During receiving, a tote with insulin pens arrives with a temperature indicator showing it exceeded recommended range for several hours. What should the technician do?

- a) Refrigerate immediately and use first to reduce waste
- b) Quarantine the shipment and escalate for pharmacist decision per procedure
- c) Restock if the boxes are sealed
- d) Dispense but shorten the expiration date on the label

11) A new eRx is for "metformin ER 500 mg: take 2 tablets BID." The technician accidentally selects metformin IR 500 mg. The barcode scan flags a mismatch. What should happen next?

- a) Override because metformin is not high risk
- b) Continue if the patient has taken metformin before
- c) Correct the selection to the ER product and continue the fill
- d) Change directions to once daily to match ER

12) A patient requests a printed copy of their prescription profile to bring to a new provider. What is the best technician action?

- a) Provide it immediately since it's their information
- b) Provide it only after verifying identity and documenting the release per HIPAA workflow
- c) Refuse because profiles can never be printed
- d) Fax it to the new provider without patient authorization

13) The sig reads: "i gtt ou bid." What does this mean?

- a) Instill 1 drop in each ear twice daily
- b) Instill 1 drop in both eyes twice daily
- c) Inject 1 drop in both eyes twice daily
- d) Insert 1 tablet in both eyes twice daily

14) A patient brings in an older prescription for amphetamine/dextroamphetamine (Schedule II). It was written 8 months ago and never filled. What is the best action?

- a) Fill it because it has not been dispensed before
- b) Partially fill it and complete later
- c) Transfer it to a different pharmacy for processing
- d) Do not fill; it is outside the typical Schedule II validity window and needs pharmacist handling/new prescription

15) The pharmacy receives a manufacturer notice of a Class I recall for a specific lot of losartan. That lot is on the shelf. What should the technician do?

- a) Keep dispensing until replacement arrives
- b) Remove from active stock, quarantine it, and alert the pharmacist per recall process
- c) Place a recall sticker and return to shelf
- d) Return all losartan regardless of lot number

16) A heparin infusion order reads: "Heparin 25,000 units in 250 mL D5W. Infuse at 18 units/kg/hr. Weight 70 kg." What rate in mL/hr should be entered?

- a) 10.1 mL/hr
- b) 12.6 mL/hr
- c) 15.0 mL/hr
- d) 18.0 mL/hr

17) A prescriber writes "Zestril 10 mg PO daily." No DAW is indicated. What product should the technician select?

- a) Benazepril 10 mg
- b) Enalapril 10 mg
- c) Lisinopril 10 mg
- d) Ramipril 10 mg

18) A hospital order is entered as "morphine 2 mg q2h PRN pain" with no route. What is the technician's best action?

- a) Assume IV and enter it
- b) Assume PO because it's safest
- c) Enter as IM because it's common in hospitals
- d) Hold for pharmacist review/clarification due to missing route

19) A patient asks which OTC product will treat their child's ear infection. What should the technician do?

- a) Recommend an OTC decongestant and acetaminophen
- b) Suggest leftover antibiotics if available
- c) Refer to the pharmacist because symptoms may require medical evaluation
- d) Recommend hydrogen peroxide drops as first-line therapy

20) The standard azithromycin 250 mg 5-day regimen is prescribed: "2 tablets day 1, then 1 tablet daily days 2–5." What quantity is dispensed?

- a) 5
- b) 6

- c) 7
- d) 10

21) During a controlled substance inventory count, a technician notices the count is short by 6 tablets compared to the perpetual inventory. What is the best action?

- a) Adjust the count to match the system to avoid an audit issue
- b) Recount once, document the discrepancy, and promptly report it to the pharmacist
- c) Ignore it if the shortage is under 10 tablets
- d) Move the bottle to the back and monitor next month

22) A prescription is written for omeprazole 20 mg capsules BID. Only omeprazole 20 mg OTC tablets are available. What is the best technician action?

- a) Substitute OTC tablets automatically
- b) Change to pantoprazole 40 mg daily
- c) Proceed if the patient agrees at pickup
- d) Route for pharmacist review because dosage form substitution may not be equivalent

23) A claim rejects "Refill too soon; plan allows 75% usage." The patient is at day 20 of a 30-day supply. What should the technician do first?

- a) Verify last fill/date and days' supply entered, then proceed appropriately based on findings
- b) Change days' supply to 20 so insurance pays
- c) Automatically reverse and re-bill as lost medication
- d) Tell the patient insurance never allows early refills

24) A new prescription for amlodipine 5 mg daily is received. The patient profile shows nifedipine ER 60 mg daily active. What is the best technician action?

- a) Fill both because they are different drugs
- b) Discontinue nifedipine in the profile before filling
- c) Queue the new prescription for pharmacist review for possible therapeutic duplication/therapy change
- d) Reduce amlodipine to 2.5 mg and fill

25) The prescription is for Latuda 40 mg daily. Which generic should be selected?

- a) Lurasidone
- b) Lamotrigine
- c) Lisdexamfetamine
- d) Levetiracetam

26) A prescription for promethazine-codeine syrup has quantity written as "120" with no unit. What should the technician do?

- a) Assume mL and process
- b) Assume 120 teaspoons and convert to mL
- c) Enter 120 mL and note "unit assumed"
- d) Send to pharmacist for clarification because quantity units are unclear on a controlled medication

27) A label prints "take 1 tablet once daily," but the original prescription image says "take 1 tablet twice daily." What is the technician's first step?

- a) Dispense and verbally correct at pickup
- b) Correct the data entry/label so it matches the prescription before it proceeds
- c) Ask the patient which they prefer
- d) Change the prescription image to match the label

28) A patient's spouse is picking up a prescription. What is the best technician action?

- a) Release it without questions if the last name matches
- b) Verify appropriate identifiers per pickup procedure before release
- c) Refuse; only the patient may pick up
- d) Provide the spouse the patient's full medication list to help adherence

29) Gentamicin 0.3% ophthalmic solution comes in a 5 mL bottle. Assuming 20 drops/mL, how many drops are in one bottle?

- a) 50
- b) 75
- c) 100
- d) 150

30) While pulling a vaccine for an appointment, you find the vial expired yesterday. What should the technician do?

- a) Use it because it's only one day expired
- b) Use it if it has been refrigerated continuously
- c) Quarantine/remove it from use and escalate for replacement
- d) Freeze it to extend stability until replacement arrives

31) A prescription for tramadol has already used all authorized refills. The patient requests another refill. What should the technician do?

- a) Refill because tramadol is not controlled
- b) Provide a small emergency supply without authorization
- c) Transfer the prescription to another pharmacy for additional refills
- d) Initiate a refill authorization request per workflow and route to the pharmacist as required

32) The profile shows simvastatin. A new prescription for clarithromycin is entered and an interaction alert appears. What is the best technician action?

- a) Override because antibiotics are temporary
- b) Ensure the alert is reviewed by the pharmacist before dispensing
- c) Tell the patient to stop simvastatin for a week
- d) Substitute azithromycin without prescriber involvement

33) The sig reads: "Apply AA BID PRN." What does AA mean?

- a) Both arms
- b) Before meals
- c) Affected area
- d) As directed

34) When receiving a controlled substance order from the wholesaler, which action is most appropriate?

- a) Leave the tote in receiving until the end of the day

- b) Verify items against invoice and secure controlled substances promptly
- c) Stock controlled substances with OTC items to reduce attention
- d) Open each bottle to confirm tablet imprint before stocking

35) A patient asks how to store nitroglycerin sublingual tablets. Which response is within technician scope and most appropriate?

- a) Store in the original, tightly closed container away from heat/moisture; pharmacist can provide additional details
- b) Store in a weekly pill organizer to prevent missed doses
- c) Store in the refrigerator to keep potency
- d) Store with cotton in the bottle to reduce moisture

36) Gabapentin 300 mg is prescribed "1 capsule TID" for 90 days. How many capsules should be dispensed?

- a) 180
- b) 240
- c) 270
- d) 300

37) An order is written "MSO4 2 mg IV q2h PRN." What is the best technician action given safety standards?

- a) Enter as written; abbreviations are acceptable if common
- b) Interpret MSO4 as magnesium sulfate and proceed
- c) Change to "morphine" without pharmacist involvement
- d) Route for pharmacist review because it is an error-prone abbreviation that can cause harm

38) A patient is starting a REMS medication and is missing required enrollment documentation. What can a technician appropriately do?

- a) Complete required administrative forms and coordinate documentation while pharmacist completes clinical requirements
- b) Waive REMS since the prescriber ordered it
- c) Provide clinical counseling on major risks to expedite pickup
- d) Substitute to a non-REMS alternative to avoid delays

39) The prescription is for Keflex 500 mg. Which generic should be selected?

- a) Cefdinir
- b) Cephalixin
- c) Ceftriaxone
- d) Cefuroxime

40) At the register, the screen shows the patient's diagnosis code. The technician needs to discuss a cash price. What is the best privacy action?

- a) Discuss the diagnosis to justify the medication
- b) Turn the screen away from public view and discuss only necessary billing details
- c) Announce the drug and diagnosis to confirm the correct patient
- d) Print the patient profile and hand it to the patient for review in line

41) A prescription for alendronate 70 mg weekly was entered as daily. What is the best technician quality step before it reaches the patient?

- a) Compare label and entry to the original prescription and correct the frequency
- b) Dispense and let the patient call if it seems wrong
- c) Change to alendronate 10 mg daily to match "daily"
- d) Delete the prescription to avoid harm

42) A prescription for hydrocodone/acetaminophen reads "1–2 tablets q4–6h PRN pain" with no maximum daily dose noted. What should the technician do?

- a) Enter as written and continue
- b) Add "max 8 tablets/day" based on typical dosing
- c) Send for pharmacist review because directions may be incomplete for safe dispensing
- d) Change to "1 tablet q6h" to be safer

43) A compounded topical preparation appears grainy and separated after mixing. What is the best technician action?

- a) Dispense; appearance does not matter if ingredients are correct
- b) Add additional base to improve consistency and dispense
- c) Remix quickly and skip documentation
- d) Hold the product for pharmacist assessment before dispensing

44) Xarelto 20 mg daily is prescribed. The profile shows warfarin active. What is the best technician action?

- a) Fill Xarelto; the patient will stop warfarin automatically
- b) Delete warfarin from the profile to prevent duplication
- c) Queue for pharmacist review due to possible duplicate anticoagulation/therapy change
- d) Change Xarelto to 10 mg to reduce bleeding risk

45) An antibiotic suspension is reconstituted to a final volume of 100 mL. Directions: 10 mL PO BID for 7 days. How many mL will remain?

- a) 0 mL
- b) 10 mL
- c) 20 mL
- d) 30 mL

46) The patient asks why ProAir was changed to Ventolin. What is the best technician response?

- a) Explain Ventolin is stronger and lasts longer
- b) Explain it was a pharmacy choice and nothing else
- c) Change it back to ProAir without involving the pharmacist
- d) Explain it may be a formulary/coverage change and offer pharmacist counseling for clinical questions

47) A faxed refill request is received for a Schedule III medication with no refills remaining. What is the best technician action?

- a) Add one refill because Schedule III allows it
- b) Request prescriber authorization through the standard process and route appropriately
- c) Transfer the prescription to another pharmacy for refills

d) Convert it to a Schedule IV alternative and refill

48) A caregiver asks for only part of a non-controlled antibiotic today due to cost. What is the best technician action?

- a) Dispense a partial quantity without documentation to help the patient
- b) Change the quantity on the prescription to match what they can afford
- c) Route for pharmacist involvement because partial dispensing and labeling/documentation requirements must be met
- d) Tell them partial fills are illegal for non-controlled medications

49) A prescription is for dorzolamide/timolol ophthalmic solution "1 gtt OU BID." The patient asks if it treats glaucoma. What should the technician do?

- a) Confirm the diagnosis and explain glaucoma treatment goals
- b) Tell them it is an antibiotic for eye infections
- c) Provide non-clinical assistance and offer pharmacist counseling for indication/clinical questions
- d) Recommend stopping if it causes burning

50) A secondary wholesaler offers deeply discounted injectable product but cannot provide transaction history. Under DSCSA, what should the pharmacy do?

- a) Purchase it if the seal is intact
- b) Accept it and create internal documentation to replace missing history
- c) Accept only if expiration is more than 1 year away
- d) Do not accept without required tracing documentation; treat as suspect

51) A prescription is for diltiazem ER 120 mg daily. The technician selects diltiazem IR 120 mg because it is in stock. What is the best action?

- a) Dispense IR because the mg is the same
- b) Route for pharmacist review because ER vs IR substitution is significant
- c) Split the IR dose into smaller doses to mimic ER
- d) Change directions to "take 1 tablet BID" and proceed

52) The plan requires days' supply for a 5 mL eye drop bottle to be entered as 15 days for adjudication. The prescription directions are 1 gtt OU QID. What should the technician do?

- a) Enter 30 days because that is typical for a bottle
- b) Change directions to BID to make 30 days
- c) Enter 15 days as required by the plan rule while keeping directions accurate
- d) Bypass insurance and bill cash automatically

53) A patient picking up clopidogrel says they also take aspirin 81 mg OTC and asks if it's safe to take both. What should the technician do?

- a) Tell them to stop aspirin immediately
- b) Tell them it is always safe because aspirin is OTC
- c) Recommend taking aspirin only on alternate days
- d) Offer pharmacist counseling for the safety question

54) Valacyclovir 1 g is prescribed: "Take 2 tablets PO BID for 1 day." What quantity should be dispensed?

- a) 2
- b) 4
- c) 6
- d) 8

55) A patient asks the pharmacy to mail a controlled medication to a different address than on file. What is the best technician action?

- a) Ship to any requested address if the patient can state DOB
- b) Update the address immediately to avoid delay
- c) Verify identity and escalate for pharmacist/manager review due to diversion risk and shipping requirements
- d) Refuse all mailing of controlled substances in every circumstance

56) Ciprodex otic is dispensed as a 7.5 mL bottle. Assuming 20 drops/mL, how many drops are in the bottle?

- a) 150
- b) 175
- c) 200
- d) 225

57) A renal dosing alert appears during entry for metformin on a patient flagged with chronic kidney disease. What should the technician do?

- a) Override because metformin is common
- b) Reduce the dose and proceed
- c) Ask the patient for lab values and decide
- d) Ensure the pharmacist reviews the alert before dispensing

58) While restocking, you find multiple stock bottles with torn safety seals. What is the best action?

- a) Quarantine the bottles and escalate for integrity review
- b) Tape the seals and return to stock
- c) Use them first to reduce waste
- d) Dispense only for inpatients and not outpatients

59) A prescription says: "Take 2.5 mL PO q8h." The concentration is 250 mg/5 mL. How many mg per dose?

- a) 62.5 mg
- b) 100 mg
- c) 125 mg
- d) 250 mg

60) A Schedule II stimulant was partially filled due to insufficient stock. Under federal rules for a typical outpatient (not LTC/hospice), within what time must the remainder generally be filled?

- a) 24 hours
- b) 48 hours
- c) 72 hours
- d) 7 days

61) During final assembly, the tablet imprint in the vial does not match the product image for the selected NDC. What should the technician do first?

- a) Dispense; images are often outdated
- b) Remove the odd tablet and replace it from the same stock bottle
- c) Stop and re-verify the stock bottle, NDC, and selection before continuing
- d) Ask the patient at pickup if it looks right

62) Eliquis 5 mg BID is prescribed. What is the generic name?

- a) Apixaban
- b) Dabigatran
- c) Edoxaban
- d) Rivaroxaban

63) The profile lists both losartan and lisinopril as active, and refills are requested for both. What is the best technician action?

- a) Process both refills to avoid therapy interruption
- b) Discontinue one medication in the system
- c) Reduce one dose by half and fill both
- d) Route for pharmacist review due to possible duplicate RAAS therapy

64) "Take 1 tsp PO QID." How many mL per dose?

- a) 2.5 mL
- b) 5 mL
- c) 10 mL
- d) 15 mL

65) A prescriber calls in a verbal prescription for a non-controlled legend drug. In many settings, what may a technician do?

- a) Take and transcribe the verbal order if allowed by state law and pharmacy policy
- b) Take verbal orders only for Schedule II drugs
- c) Refuse; verbal prescriptions are illegal
- d) Take the order but sign as the prescriber to speed processing

66) An eRx for methotrexate 2.5 mg says "take 1 tablet daily." The profile notes rheumatoid arthritis. What is the best technician action?

- a) Fill as written; daily is typical for RA
- b) Change to weekly dosing and proceed
- c) Immediately route for pharmacist review due to high-alert weekly-vs-daily error risk
- d) Delete the prescription as an obvious error

67) A patient asks whether grapefruit juice interacts with atorvastatin. What should the technician do?

- a) Say there is no interaction with any statin
- b) Say grapefruit is always life-threatening with statins
- c) Suggest the patient stop atorvastatin if they drink grapefruit
- d) Offer pharmacist counseling for interaction guidance

68) Tamsulosin 0.4 mg is prescribed, but the system is set to an NDC for tamsulosin 0.8 mg. What is the best technician action?

- a) Correct the product/NDC to match 0.4 mg before dispensing
- b) Dispense 0.8 mg because it is a common dose
- c) Fill 0.4 mg but keep the 0.8 mg NDC for inventory balance
- d) Open capsules and remove half the contents

69) A prescription specifies U-500 insulin, but the technician selected U-100 in the system. What is the best action?

- a) Keep U-100 because it is most common
- b) Switch to U-500 and continue without pharmacist involvement
- c) Continue with U-100 and advise the patient to take one-fifth the dose
- d) Stop processing and route for pharmacist verification due to high-alert concentration risk

70) A patient asks you to run a discount card instead of insurance for a controlled medication. What is the best technician action?

- a) Run the discount card without questions since it's patient preference
- b) Confirm the patient request and process appropriately while escalating any diversion/red flags per workflow
- c) Refuse because discount cards are illegal for controlled substances
- d) Bill both insurance and the discount card simultaneously

71) Carvedilol 12.5 mg "1 tablet BID" is prescribed with quantity "#30." What should the technician do?

- a) Dispense #30 as written and move on
- b) Change quantity to #60 without contacting anyone
- c) Change directions to once daily so #30 matches
- d) Route for pharmacist clarification because quantity conflicts with directions

72) A patient reports a serious adverse effect from a new medication and asks how to report it. What is the best technician response?

- a) Provide information on FDA MedWatch reporting and offer pharmacist involvement for clinical follow-up
- b) Tell them to call their insurance company
- c) Tell them to post a review online
- d) Document it only in the patient profile; no reporting system exists

73) Fluconazole 150 mg "take 1 tablet once" is prescribed. The profile shows warfarin. What should the technician do?

- a) Fill quickly; it's only one dose
- b) Ensure the pharmacist reviews the interaction alert before dispensing
- c) Tell the patient to skip warfarin for 3 days
- d) Substitute nystatin tablets

74) A prescription for a topical cream has quantity "QS" with directions "apply thin layer to affected area BID x 14 days," and no grams are specified. What is the best technician action?

- a) Dispense the smallest tube available

- b) Select a standard tube size and document without further review
- c) Route for pharmacist determination of appropriate quantity/package size
- d) Change directions to once daily to make it last longer

75) In a hospital MAR, potassium chloride ER tablets are active, and an oral solution order is discontinued. A technician is asked to prepare the solution anyway. What is the best action?

- a) Prepare the solution because both are potassium
- b) Prepare both to avoid missing a dose
- c) Substitute IV potassium to avoid oral confusion
- d) Stop and clarify because the requested product conflicts with order status

76) A patient brings unused fentanyl patches and asks how to dispose of them. What should the technician do?

- a) Tell them to flush them in all cases
- b) Recommend mixing with coffee grounds and placing in household trash
- c) Provide take-back options information and offer pharmacist guidance for safe disposal specifics
- d) Accept them at the counter and throw them in regular trash

77) A patient is started on lisinopril. Which common adverse effect should be flagged for counseling by the pharmacist if the patient reports it?

- a) Dry cough
- b) Purple urine
- c) Tinnitus
- d) Hair loss

78) A prescription is for "Synthroid 125 mcg." Which generic is correct for selection?

- a) Liothyronine
- b) Levothyroxine
- c) Levofloxacin
- d) Levocetirizine

79) A patient asks for a refill of clonazepam and says they lost the bottle. The refill is 12 days early. What is the best technician action?

- a) Process a refill immediately and note "lost"
- b) Refuse because early refills are never allowed for controlled substances
- c) Offer a small emergency supply without authorization
- d) Gather relevant details and route for pharmacist decision on an early-fill request

80) A prescription label says "Take 1 tablet PO QD" but the eRx says "Take 1 tablet PO BID." Which is the best technician action?

- a) Dispense and counsel the patient verbally to take BID
- b) Correct the entry and reprint the label to match the eRx before dispensing
- c) Change the eRx directions to match the label
- d) Leave it; the pharmacist will catch it later

81) A patient's allergy profile lists penicillin rash. A prescription for amoxicillin is received. What should the technician do?

- a) Delete the allergy since it was only a rash
- b) Enter and override the allergy warning to avoid delays
- c) Stop and ensure pharmacist evaluates the allergy before filling
- d) Substitute azithromycin automatically

82) The technician is entering a prescription that reads "Take 0.5 mg PO HS." The product is available as 1 mg tablets. What is the correct quantity to dispense for 30 days?

- a) 15 tablets
- b) 30 tablets
- c) 45 tablets
- d) 60 tablets

83) A long-term care facility sends a request to transfer a non-controlled prescription between two pharmacies that share a real-time, common electronic database. Which is permitted under federal law for refill transfers in this setup?

- a) Transfers are prohibited because refills can never be transferred
- b) Transfers may occur up to the number of refills authorized, if the system meets federal requirements
- c) Only one transfer is permitted total
- d) Transfers are permitted only for Schedule II prescriptions

84) A technician is processing a return-to-stock of a refrigerated biologic that was picked up and returned unopened the next day. What is the safest action?

- a) Return to stock if it was in a cooler bag
- b) Return to stock if the patient says it stayed cold
- c) Do not return to stock; quarantine and follow destruction/return procedures
- d) Return to stock after relabeling with a shorter expiration

85) A patient presents a Medicaid card and asks to pay cash instead "to keep it off insurance." What should the technician do?

- a) Bill cash to respect patient privacy with no further steps
- b) Explain that billing choices may be restricted and route appropriately for pharmacist/manager handling
- c) Refuse service
- d) Bill Medicaid anyway without telling the patient

86) A prescription is written for prednisone 10 mg tablets: "4 tablets daily x 3 days, then 3 daily x 3 days, then 2 daily x 3 days, then 1 daily x 3 days." How many tablets are needed?

- a) 24
- b) 30
- c) 36
- d) 42

87) A prescription is written for insulin lispro U-100: "Inject 12 units before meals TID." One 10 mL vial is dispensed (100 units/mL). What is the days' supply?

- a) 21 days
- b) 27 days
- c) 30 days

d) 45 days

88) A patient is prescribed escitalopram. Which brand name matches?

- a) Celexa
- b) Lexapro
- c) Zoloft
- d) Cymbalta

89) During data entry, a prescriber's name is similar to another provider in the system. The NPI entered auto-populates a different address and phone than the prescription shows. What is the best technician action?

- a) Keep the auto-populated provider to save time
- b) Edit the prescription to match the system provider address even if different
- c) Verify and select the correct prescriber profile to match the prescription details before processing
- d) Remove all prescriber information and leave it blank

90) A DEA Form 222 is being used for ordering Schedule II medications. Which is a technician-appropriate role in this process?

- a) Signing the form on behalf of the registrant to speed ordering
- b) Completing ordering details for pharmacist review and ensuring records are filed as required
- c) Faxing a copy to any supplier as an open order authorization
- d) Using it to order Schedule III–V medications only

Answer Key

1) a	31) d	61) c
2) c	32) b	62) a
3) b	33) c	63) d
4) c	34) b	64) b
5) a	35) a	65) a
6) c	36) c	66) c
7) b	37) d	67) d
8) c	38) a	68) a
9) d	39) b	69) d
10) b	40) b	70) b
11) c	41) a	71) d
12) b	42) c	72) a
13) b	43) d	73) b
14) d	44) c	74) c
15) b	45) c	75) d
16) b	46) d	76) c
17) c	47) b	77) a
18) d	48) c	78) b
19) c	49) c	79) d
20) b	50) d	80) b
21) b	51) b	81) c
22) d	52) c	82) a
23) a	53) d	83) b
24) c	54) b	84) c
25) a	55) c	85) b
26) d	56) a	86) b
27) b	57) d	87) b
28) b	58) a	88) b
29) c	59) c	89) c
30) c	60) c	90) b

PTCB Sim Test 29

- 1) A new eRx reads: "rosuvastatin 20 mg tablet, take 1 tablet PO nightly, #90." While selecting the product, the technician notices the shelf has rosuvastatin 10 mg and 20 mg in similar bottles. What is the safest next step before printing the fill label?
- a) Select the 10 mg and plan to counsel the patient to take 2 tablets
 - b) Select the 20 mg by matching drug name, strength, dosage form, and NDC to the selected item
 - c) Select either strength and let the pharmacist change it at verification
 - d) Choose the bottle with the soonest expiration date
- 2) A prescription says: "amoxicillin 400 mg/5 mL; give 6 mL PO BID x 10 days." How many mL should be dispensed?
- a) 100 mL
 - b) 110 mL
 - c) 120 mL
 - d) 130 mL
- 3) A patient's profile shows an anaphylactic reaction to penicillin. A new prescription arrives for dicloxacillin. What should the technician do next?
- a) Continue processing because dicloxacillin is narrow-spectrum
 - b) Remove the allergy because it may be outdated
 - c) Place the order on hold and route it for pharmacist review of the allergy conflict
 - d) Change dicloxacillin to cephalexin and continue
- 4) At drop-off, a patient asks for an early refill of clonazepam because they "left it at a hotel." The last fill was 8 days ago for a 30-day supply. What is the best technician action?
- a) Process as cash to bypass insurance limits
 - b) Collect details (date lost, remaining quantity, police report if any) and route the request for pharmacist decision before promising a refill
 - c) Tell the patient the pharmacy is required to replace lost medication once per year
 - d) Transfer the prescription to another store to fill sooner
- 5) A prescriber writes: "metoprolol succinate ER 50 mg daily." During entry, the system suggests metoprolol tartrate 50 mg as the top match. What should the technician do?
- a) Select tartrate because it's the same drug
 - b) Select succinate ER and verify the dosage form matches the order
 - c) Select tartrate and change directions to BID
 - d) Enter succinate but change it to immediate-release for faster onset
- 6) A technician is asked to sell pseudoephedrine. Which step is required before completing the sale under federal requirements?
- a) Verify a government-issued photo ID and record required purchaser information in the logbook/electronic system
 - b) Require a prescriber authorization on a pharmacy letterhead
 - c) Obtain a signed HIPAA release from the purchaser
 - d) Dispense only if the patient is at least 21 years old

7) A label direction reads: "Instill 1 gtt OU qid." How many drops per day is that?

- a) 4
- b) 6
- c) 8
- d) 10

8) An eRx for "Keflex 500 mg" is received. Which generic should be selected?

- a) cefdinir
- b) cephalexin
- c) cefuroxime
- d) ceftriaxone

9) A patient is at pickup for a new warfarin prescription and says they started apixaban yesterday from a different clinic. What should the technician do?

- a) Complete the sale because both prevent clots
- b) Remove apixaban from the patient profile to avoid confusion
- c) Flag the profile and route for pharmacist review before dispensing due to duplicate anticoagulants
- d) Tell the patient to take warfarin only and stop apixaban

10) A carton fill in the hospital reads: "heparin 25,000 units in 250 mL D5W." What is the concentration?

- a) 50 units/mL
- b) 80 units/mL
- c) 100 units/mL
- d) 125 units/mL

11) A prescription is written for "azithromycin 200 mg/5 mL: 12.5 mL day 1, then 6.25 mL daily days 2–5." How many mL should be dispensed?

- a) 31.25 mL
- b) 35 mL
- c) 37.5 mL
- d) 42.5 mL

12) A patient requests their medication profile be emailed to their personal email. What is the best technician response?

- a) Email it if the patient confirms their date of birth
- b) Offer to print it or use an approved secure release method after verifying identity and authorization requirements
- c) Refuse because patients are not allowed access to pharmacy records
- d) Read the list aloud over the phone instead

13) A new prescription arrives for "Toprol-XL 25 mg daily." Which generic product matches?

- a) metoprolol tartrate
- b) metoprolol succinate ER
- c) propranolol ER
- d) atenolol

14) A prescription reads: "prednisone 10 mg: take 4 tabs daily x 3 days, then 3 tabs daily x 3 days, then 2 tabs daily x 3 days, then 1 tab daily x 3 days." How many tablets should be dispensed?

- a) 24
- b) 27
- c) 30
- d) 36

15) The technician finds two will-call bags stapled together; each contains a different patient leaflet with PHI visible through the bag window. What is the best immediate action?

- a) Separate, correct the bag contents, and document/report the privacy incident per the pharmacy's QA process
- b) Leave them stapled to prevent mix-ups
- c) Remove both leaflets and throw them away to eliminate PHI exposure
- d) Wait until the end of the shift to fix it

16) A prescriber calls in a prescription by voicemail for alprazolam. The message includes patient name, drug, strength, directions, quantity, refills, and prescriber name—but no callback number. What should the technician do?

- a) Process it; a callback number is optional for Schedule IV
- b) Hold for pharmacist and obtain the missing required information before it can be dispensed
- c) Ask the patient to provide the prescriber's phone number and proceed
- d) Convert to lorazepam and process as equivalent

17) Which drug is a proton pump inhibitor?

- a) famotidine
- b) pantoprazole
- c) sucralfate
- d) dicyclomine

18) A prescription says: "Ventolin HFA: inhale 2 puffs q4h prn SOB." The inhaler contains 200 puffs. If used at the maximum frequency, what is the days' supply?

- a) 10 days
- b) 14 days
- c) 16 days
- d) 25 days

19) A refrigerated shipment arrives with a temperature indicator showing out-of-range during transit. What should the technician do?

- a) Stock it immediately and mark "use first"
- b) Place it in will-call since it's already assigned to patients
- c) Quarantine the shipment and route to pharmacist/manager for disposition per procedure
- d) Put it in the refrigerator and ignore the indicator

20) A patient is prescribed "Lantus SoloStar inject 18 units SQ at bedtime; dispense 1 box." One box contains five 3 mL pens (100 units/mL). What is the days' supply?

- a) 50 days
- b) 60 days
- c) 75 days
- d) 83 days

21) An eRx reads: "Lamictal 200 mg daily." Which generic should be selected?

- a) lamotrigine
- b) levothyroxine
- c) levetiracetam
- d) lithium carbonate

22) A new technician is preparing to compound a nonsterile oral suspension but notices the balance calibration sticker expired last month. What should be done next?

- a) Proceed; small errors are acceptable for oral preparations
- b) Use a measuring cylinder instead of weighing and continue
- c) Stop and obtain a calibrated balance/device before compounding
- d) Estimate the weight from the label and document it

23) A patient returns a bottle of tablets they say were dispensed in error and asks the pharmacy to "put them back on the shelf." What should the technician do?

- a) Restock if the seal looks intact
- b) Restock if the tablets match the imprint for the intended drug
- c) Place the bottle in a designated returns/destruction area per procedure; do not restock for dispensing
- d) Give the tablets to another patient if the label is removed

24) A prescription is written: "Apply 2 g topically BID." The product is a 120 g jar. What is the days' supply?

- a) 15 days
- b) 25 days
- c) 30 days
- d) 60 days

25) A DEA audit question: which controlled substances require a separate, exact inventory count (not "estimated") at inventory time?

- a) Schedule II
- b) Schedule III
- c) Schedule IV
- d) Schedule V

26) During entry, the technician sees an interaction alert: simvastatin and clarithromycin. What is the most appropriate next step?

- a) Override because interaction alerts are for counseling only
- b) Continue and add an auxiliary label to take at different times
- c) Route for pharmacist review before filling due to clinically significant interaction risk
- d) Delete simvastatin from the profile to resolve the alert

27) A hospital label prints "morphine ER 30 mg," but the tech accidentally selected morphine

IR 30 mg in the cart-fill system. What is the best technician action when the mismatch is noticed?

- a) Continue; the nurse can adjust administration timing
- b) Correct the selection to ER and reprint/reprocess per workflow so the correct product is dispensed
- c) Dispense IR and add "do not crush" label
- d) Change the dose to 15 mg IR to reduce risk

28) Which medication commonly requires a Medication Guide to be provided at dispensing?

- a) isotretinoin
- b) amoxicillin
- c) metformin
- d) albuterol HFA

29) A prescription is written for "clopidogrel 75 mg daily." Which brand name matches?

- a) Plavix
- b) Coumadin
- c) Eliquis
- d) Brilinta

30) A paper prescription for a Schedule II medication is dated 45 days ago and has no "do not fill until" date. What should the technician do?

- a) Fill it because federal law has no expiration for Schedule II
- b) Refuse and return it without review
- c) Bring it to the pharmacist to evaluate validity under state law, prescriber intent, and pharmacy policy before processing
- d) Change the written date in the system to today and process

31) A prescription says: "Take 1 tablet PO every 6 hours x 7 days." How many tablets are needed?

- a) 21
- b) 24
- c) 28
- d) 42

32) A patient asks what "qhs" means on their label. What is the best technician response?

- a) "Every morning"
- b) "At bedtime"
- c) "As needed"
- d) "Four times daily"

33) A prescription transfer request is received for a patient's tramadol with refills remaining. What is the best technician action?

- a) Complete the transfer verbally technician-to-technician for any controlled substance
- b) Decline because controlled prescriptions cannot be transferred
- c) Route to the pharmacist to complete the transfer if permitted by federal/state rules
- d) Transfer only the remaining refills and void the original automatically

34) A pediatric prescription is entered: "acetaminophen 160 mg/5 mL; give 20 mL q4h prn fever." What should the technician do?

- a) Enter as written because acetaminophen is OTC
- b) Change to 10 mL automatically to reduce overdose risk
- c) Flag and route for pharmacist review due to potentially excessive dose
- d) Ask the parent to decide the dose at pickup

35) A prescription is for "Bactroban ointment." What product should be dispensed?

- a) bacitracin
- b) mupirocin
- c) metronidazole
- d) triamcinolone

36) A patient is picking up "glimepiride 2 mg daily" and asks if they can skip breakfast. What is within technician scope?

- a) Tell them it's fine as long as they drink juice
- b) Tell them to take it at bedtime instead
- c) Ask them to reduce the dose if skipping meals
- d) Refer to the pharmacist because meal timing affects hypoglycemia risk

37) A prescription reads: "gabapentin 300 mg: take 1 cap TID; dispense #270." What is the days' supply?

- a) 60 days
- b) 75 days
- c) 90 days
- d) 120 days

38) A DSCSA alert indicates a specific NDC/lot may be illegitimate. What should the technician do first?

- a) Keep dispensing until the manufacturer confirms
- b) Quarantine the affected product and alert the pharmacist/manager for investigation steps
- c) Return all products from that manufacturer to the wholesaler
- d) Re-label the product with an internal barcode and continue use

39) A technician is entering "benzonatate 200 mg TID PRN cough," but the system defaults to 100 mg. What is the correct action?

- a) Accept the default and proceed
- b) Change directions to "take 2 capsules TID" and keep 100 mg
- c) Correct the strength to 200 mg and confirm quantity matches the prescribed directions
- d) Substitute dextromethorphan because it's OTC

40) A patient's spouse arrives to pick up the patient's oxycodone prescription. What is the best technician action at pickup?

- a) Refuse; only the patient can pick up controlled substances
- b) Release it if the spouse knows the patient's address
- c) Verify identity/authorization per pharmacy procedure and document required information before release
- d) Release it if the spouse can state the drug name and strength

41) A prescription is received for "Xarelto 20 mg daily." Which generic name matches?

- a) rivaroxaban
- b) apixaban
- c) dabigatran
- d) clopidogrel

42) A technician notices a vaccine refrigerator temperature log has two missing entries from yesterday. What should be done?

- a) Leave blanks; missing entries are acceptable
- b) Estimate temperatures based on today's range and fill them in
- c) Document the omission and notify the appropriate supervisor per quality process for follow-up
- d) Discard all vaccines because the log is incomplete

43) A prescription is written: "cephalexin 500 mg: take 1 cap PO QID x 10 days." What quantity should be dispensed?

- a) 20
- b) 30
- c) 40
- d) 60

44) A technician is asked to restock returned medications from an inpatient unit. Which item should NOT be returned to inventory under typical rules?

- a) An unopened manufacturer-sealed bottle of tablets
- b) A sealed manufacturer carton of topical cream
- c) A unit-dose blister pack that remained unopened and properly stored
- d) A compounded sterile syringe returned from the floor

45) A prescription is for "nitroglycerin 0.4 mg SL tablets." The patient asks where to store it. Which response is appropriate for technician scope?

- a) Store in the bathroom medicine cabinet for quick access
- b) Keep in the original tightly closed container away from heat and moisture; the pharmacist can provide additional counseling
- c) Transfer into a weekly pillbox to prevent missing doses
- d) Refrigerate after opening to maintain potency

46) A technician receives a prescriber call authorizing refills for lisinopril. What is most important to document?

- a) The patient's weight
- b) The caller's name, role, and callback number plus the exact authorization details
- c) The patient's diagnosis
- d) The patient's insurance group number

47) A label prints "Do not crush or chew." Which medication is most likely to carry this warning due to dosage form?

- a) bupropion XL
- b) amoxicillin

- c) amlodipine
- d) doxycycline hyclate

48) A prescription says: "Take 2 tablets in the morning and 1 tablet in the evening." The quantity is 180 tablets. What is the days' supply?

- a) 45 days
- b) 60 days
- c) 75 days
- d) 90 days

49) A patient drops off a prescription that appears altered (quantity overwritten). What should the technician do?

- a) Fill it as written because the prescriber signed it
- b) Ask the patient to initial the change and proceed
- c) Refuse and shred it immediately
- d) Hold the prescription and route to the pharmacist for verification/clarification before processing

50) A prescription is written for "Augmentin 875/125 mg: 1 tab PO BID x 7 days." What quantity should be dispensed?

- a) 10
- b) 14
- c) 20
- d) 28

51) A patient profile has documented pregnancy. A new prescription arrives for lisinopril. What should the technician do next?

- a) Process normally; warnings are for counseling only
- b) Change to labetalol and continue
- c) Route for pharmacist review before dispensing due to contraindication concerns
- d) Ask the patient to sign a waiver at pickup

52) Which medication is a beta-blocker?

- a) losartan
- b) hydrochlorothiazide
- c) metoprolol
- d) amlodipine

53) A prescription is received for "methylprednisolone dose pack: use as directed." During entry, what is most appropriate?

- a) Enter the standard manufacturer taper directions associated with that product
- b) Replace with prednisone equivalent dosing automatically
- c) Enter "take 1 tablet daily" to simplify
- d) Reject because "use as directed" is always invalid

54) A controlled-substance prescription is partially filled due to insufficient stock. What is the best technician action regarding the partial fill?

- a) Complete the partial fill without documentation; the label is enough

- b) Ensure the pharmacist is involved so partial-fill documentation and timelines are handled correctly
- c) Owe the balance indefinitely until the patient returns
- d) Convert the remaining quantity to a new prescription number without note

55) A prescription reads: "fluoxetine oral solution 20 mg/5 mL: take 5 mL daily; dispense 120 mL." What is the days' supply?

- a) 12 days
- b) 18 days
- c) 24 days
- d) 30 days

56) A patient asks for "the generic for Norvasc." Which is correct?

- a) amlodipine
- b) amiodarone
- c) atenolol
- d) amoxicillin

57) While typing, the technician accidentally scans a prescription image into the wrong patient profile with a similar name. What should be done first?

- a) Continue; verification will catch it
- b) Correct the patient selection immediately and document per workflow/QA requirements
- c) Delete the image and ask the patient to return later
- d) Fill it and confirm at pickup

58) A prescription reads: "Take 1 tablet PO every 8 hours; dispense #21." What is the days' supply?

- a) 5 days
- b) 7 days
- c) 10 days
- d) 14 days

59) A technician is stocking insulin. Which product is typically stored refrigerated prior to dispensing when unopened?

- a) insulin aspart
- b) levothyroxine
- c) sertraline
- d) atorvastatin

60) An eRx is received for "Klonopin 0.5 mg." Which generic matches?

- a) clonidine
- b) clotrimazole
- c) clonazepam
- d) clopidogrel

61) A manufacturer recall notice is received for a specific lot of valsartan. What should the technician do first?

- a) Remove all valsartan regardless of lot number

- b) Quarantine the affected lot(s) immediately to prevent dispensing
- c) Dispense remaining bottles quickly before the return window closes
- d) File the notice and wait for the next inventory count

62) A prescription is written for "tamsulosin 0.4 mg daily." The technician selects "tamoxifen 20 mg" due to name similarity. Which practice best prevents this error?

- a) Store both medications next to each other for faster selection
- b) Use barcode scanning and verify the full drug name/strength before counting
- c) Choose based on indication because both are commonly used long-term
- d) Select the most commonly dispensed item in the list

63) A technician receives a phone call from someone claiming to be a prescriber's office requesting a patient's full medication list, but they cannot verify identity beyond the clinic name. What should the technician do?

- a) Provide the list because it's for treatment
- b) Provide only noncontrolled medications
- c) Refuse all requests from prescribers without a written release
- d) Obtain a verified callback number and route to the pharmacist/authorized staff per privacy procedure before releasing PHI

64) A prescription is for "digoxin 0.125 mg daily." Two stock bottles are available: 0.125 mg and 0.25 mg. What is the best selection approach?

- a) Choose the bottle with the newest manufacturer
- b) Choose the bottle closest to expiration to reduce waste
- c) Match drug name, strength, dosage form, and NDC to the label/order before counting
- d) Use 0.25 mg and instruct the patient to split tablets

65) A prescription reads: "enoxaparin 40 mg/0.4 mL: inject 0.4 mL SQ daily x 10 days." How many syringes should be dispensed?

- a) 5
- b) 8
- c) 10
- d) 20

66) A prescription for a Schedule III medication has refills remaining but the last fill was 7 months ago. What is the best technician action?

- a) Fill it because refills remain
- b) Route to the pharmacist because federal law limits dispensing of Schedule III/IV refills to 6 months from issue date
- c) Transfer it to another pharmacy to restart the refill window
- d) Convert it to a noncontrolled alternative automatically

67) A prescription is written for "valacyclovir 1 g: take 2 tablets PO BID for 1 day." What quantity should be dispensed?

- a) 2
- b) 4
- c) 6
- d) 8

68) In the IV room, the technician observes another staff member with exposed wrists and no hair cover actively compounding. What is the best immediate action?

- a) Ignore it if they washed hands
- b) Wait until the batch is finished to avoid waste
- c) Stop the activity and escalate per sterile compounding garbing requirements so it can be corrected promptly
- d) Move to a different hood and continue compounding

69) A prescription is written for "omeprazole 20 mg daily," but the technician selects esomeprazole 20 mg because both are PPIs. What should the technician do?

- a) Dispense esomeprazole because it's equivalent
- b) Change the prescription to esomeprazole and document interchange
- c) Change directions to every other day to reduce risk
- d) Correct the selection to omeprazole as prescribed before proceeding

70) A prescription says: "ciprofloxacin 0.3% otic solution." The pharmacy has only ciprofloxacin 0.3% ophthalmic solution in stock. What should the technician do?

- a) Substitute ophthalmic automatically because eye drops can be used in ears
- b) Change the prescription to ophthalmic and process
- c) Route to the pharmacist to determine if substitution is appropriate and whether prescriber contact is needed
- d) Tell the patient to use the ophthalmic drops in the eye instead

71) A technician is preparing amoxicillin suspension and accidentally adds too much water. What should be done next?

- a) Add extra powder to compensate
- b) Dispense and tell the patient to take more mL per dose
- c) Shake longer to correct the concentration
- d) Stop and route to the pharmacist; the product may need to be remade

72) A patient requests an OTC recommendation for severe chest pain "so I don't have to go in." What should the technician do?

- a) Recommend naproxen and a heating pad
- b) Recommend aspirin and rest
- c) Recommend acetaminophen 1000 mg
- d) Get the pharmacist immediately and advise urgent medical evaluation based on symptoms

73) A patient presents for isotretinoin pickup. What step is commonly required before dispensing?

- a) Confirm required REMS authorization/requirements are met before release
- b) Verify the patient's A1c is controlled
- c) Ensure the prescription includes at least 3 refills
- d) Dispense only if the patient signs a HIPAA notice

74) A prescription reads: "Apply 1 patch daily; #30." The product comes in boxes of 28 patches. What is the best action?

- a) Dispense 28 and change the quantity in the system to match the box size
- b) Dispense 56 so the patient has at least 30
- c) Dispense 28 and owe the remaining 2 without documenting
- d) Verify packaging constraints and adjust quantity/dispense plan appropriately (e.g., dispense 28 with pharmacist-approved change or obtain clarification) before filling

75) A technician is asked which drug is an SSRI. Which is correct?

- a) sertraline
- b) bupropion
- c) duloxetine
- d) amitriptyline

76) A technician is entering an inpatient order for "pantoprazole 40 mg IV daily," but the system defaults route to PO. What should be done?

- a) Leave PO; the same drug will be sent
- b) Correct the route to IV and ensure the selected dosage form matches the order
- c) Change to omeprazole capsules for convenience
- d) Keep PO and add "NPO" in the notes

77) A patient asks to pay cash for a controlled prescription because insurance rejects "too soon." What should the technician do?

- a) Run as cash immediately to help the patient
- b) Tell the patient to come back when insurance allows it
- c) Split the quantity into smaller fills to bypass the reject
- d) Route to the pharmacist for evaluation of early refill, corresponding responsibility, and appropriate next steps

78) A patient's profile shows amlodipine 5 mg daily filled last week, and a new prescription arrives for amlodipine 10 mg daily. What should the technician do?

- a) Fill both strengths so the patient can choose
- b) Discontinue the 5 mg automatically
- c) Flag for pharmacist review as a likely dose change/duplication before dispensing
- d) Change the 10 mg directions to "take 1/2 tablet daily"

79) Which medication is an ACE inhibitor?

- a) valsartan
- b) lisinopril
- c) metoprolol
- d) hydrochlorothiazide

80) A prescription reads: "Take 5 mL PO TID x 7 days." What total volume is needed?

- a) 90 mL
- b) 100 mL
- c) 105 mL
- d) 120 mL

81) A technician discovers a stock bottle on the shelf with no label and unknown tablets inside. What should be done?

- a) Identify by imprint and relabel for use
- b) Mix with the most similar-looking stock bottle
- c) Keep it aside to identify later and continue filling
- d) Quarantine it and alert the pharmacist; do not dispense

82) Which medication is a thiazide diuretic?

- a) furosemide
- b) spironolactone
- c) hydrochlorothiazide
- d) bumetanide

83) A prescription is written for "Eliquis 5 mg." Which generic matches?

- a) edoxaban
- b) apixaban
- c) rivaroxaban
- d) dabigatran

84) A patient is prescribed "cefdinir 300 mg: take 1 capsule PO BID x 7 days." What quantity should be dispensed?

- a) 10
- b) 12
- c) 14
- d) 20

85) A technician is asked to print a hard copy of an e-prescription for filing. What item is most important for audit/traceability?

- a) The patient's email address
- b) The eRx unique identifier and date/time received
- c) The wholesaler invoice number for the medication
- d) The prescriber's office hours

86) A patient asks, "Is lisinopril the same thing as losartan?" What is the best technician response?

- a) Yes, they are the same medication
- b) No, but both lower blood pressure; the pharmacist can explain differences and why one was chosen
- c) Yes, both are beta-blockers
- d) No, losartan is a diuretic and lisinopril is a statin

87) A prescription is written for "clindamycin 150 mg: take 1 capsule QID x 10 days." What quantity should be dispensed?

- a) 30
- b) 40
- c) 60
- d) 80

88) Which action best supports HIPAA privacy at the pickup window when a caregiver requests speakerphone discussion of the patient's meds?

- a) Proceed because the caregiver asked
- b) Speak louder so the caregiver can hear clearly
- c) Move to a more private area and limit PHI to what's necessary, or involve the pharmacist as appropriate
- d) Provide the full medication list quickly to reduce time at the counter

89) A prescription reads: "Take 1 capsule PO BID; dispense #20." What is the days' supply?

- a) 5
- b) 7
- c) 10
- d) 20

90) A technician is entering an order for "U-500 insulin," but the prescription lacks clear device/syringe directions and dosing details are ambiguous. What is the best next action?

- a) Convert to U-100 insulin and keep the same unit dose
- b) Select any regular insulin vial and dispense U-100 syringes
- c) Put the prescription on hold until the patient confirms what they used last time
- d) Route to the pharmacist for clarification due to high-alert concentration and dosing safety

Answer Key

1) b	31) c	61) b
2) c	32) b	62) b
3) c	33) c	63) d
4) b	34) c	64) c
5) b	35) b	65) c
6) a	36) d	66) b
7) c	37) c	67) b
8) b	38) b	68) c
9) c	39) c	69) d
10) c	40) c	70) c
11) a	41) a	71) d
12) b	42) c	72) d
13) b	43) c	73) a
14) c	44) d	74) d
15) a	45) b	75) a
16) b	46) b	76) b
17) b	47) a	77) d
18) c	48) b	78) c
19) c	49) d	79) b
20) b	50) b	80) c
21) a	51) c	81) d
22) c	52) c	82) c
23) c	53) a	83) b
24) a	54) b	84) c
25) a	55) d	85) b
26) c	56) a	86) b
27) b	57) b	87) b
28) a	58) b	88) c
29) a	59) a	89) c
30) c	60) c	90) d

PTCB Sim Test 30

1) During data entry, an eRx for “Wellbutrin XL 150 mg QAM” is selected as bupropion SR 150 mg BID because both show “bupropion 150.” What should the technician do before continuing?

- a) Keep SR and change the sig to QAM
- b) Switch to the XL product and keep the sig as written
- c) Keep SR and reduce quantity so it lasts 30 days
- d) Dispense SR and add an auxiliary label “do not crush”

2) A patient asks for the “generic for Norvasc” while at the counter. Which product should the technician select?

- a) Amlodipine
- b) Atenolol
- c) Lisinopril
- d) Losartan

3) An order reads: “Azithromycin 250 mg: take 2 tablets once, then 1 tablet daily for 4 days.” What quantity is needed?

- a) 5
- b) 6
- c) 7
- d) 10

4) A controlled-substance invoice is filed, but the pharmacy needs to be able to retrieve it quickly during an audit. Under federal rules, what is the minimum retention period for controlled-substance records?

- a) 1 year
- b) 2 years
- c) 3 years
- d) 5 years

5) A patient is picking up and says, “My doctor changed my dose yesterday.” The will-call bag contains the previous dose filled from an older eRx with refills. What is the best technician step before completing the sale?

- a) Sell the medication and tell the patient to call the prescriber if it's wrong
- b) Ask what the new dose is and change the label to match
- c) Check the profile for a newer prescription or message and confirm which order is intended for pickup
- d) Return it to stock and tell the patient to come back tomorrow

6) A new eRx arrives for a patient with an allergy on file: “penicillin—anaphylaxis.” The prescription is for amoxicillin 500 mg. What should the technician do?

- a) Enter the prescription and select “override allergy”
- b) Stop processing and route the order for pharmacist review due to the allergy conflict
- c) Substitute doxycycline and continue
- d) Remove the allergy because it may be outdated

7) A patient asks, "What's the generic for Zoloft?"

- a) Fluoxetine
- b) Paroxetine
- c) Sertraline
- d) Escitalopram

8) The refrigerator log shows 12°C at opening (acceptable range 2–8°C). What is the most appropriate technician action?

- a) Continue using products if they still feel cold
- b) Quarantine potentially affected items and alert the pharmacist for excursion assessment and documentation
- c) Put all items in the freezer for 30 minutes
- d) Discard all refrigerated medications immediately

9) A prescription is for albuterol HFA: "Inhale 2 puffs q4h PRN. Max 12 puffs/day." One inhaler contains 200 puffs. What day supply should be billed using the maximum?

- a) 8 days
- b) 10 days
- c) 16 days
- d) 30 days

10) At pickup, two patients with the same last name are waiting. Which verification is best before handing over the bag?

- a) Ask, "Are you here for antibiotics?"
- b) Ask the patient to state date of birth and confirm it matches the label/profile
- c) Ask the patient to confirm the prescriber name only
- d) Ask for the last name and address, then hand over the top bag

11) A prescription reads: "Prednisone 10 mg: 4 tabs daily x 3 days, then 3 tabs daily x 3 days, then 2 tabs daily x 3 days, then 1 tab daily x 3 days." Quantity to dispense?

- a) 24
- b) 30
- c) 36
- d) 40

12) The pharmacy receives a manufacturer notice that a specific lot number of metformin 500 mg tablets is recalled. What is the technician's best first action?

- a) Pull only the affected lot from inventory and quarantine it per workflow
- b) Pull all metformin strengths from the shelf to be safe
- c) Continue dispensing until the wholesaler issues a replacement
- d) Delete metformin from all patient profiles

13) A patient asks for the generic of Crestor.

- a) Atorvastatin
- b) Rosuvastatin
- c) Simvastatin
- d) Pravastatin

14) While processing a new profile, the DOB on the eRx differs by one digit from the existing profile. What should the technician do?

- a) Ignore it if the address matches
- b) Change the profile DOB to match the eRx
- c) Create a second profile to avoid altering data
- d) Verify DOB using a reliable source before proceeding to avoid wrong-patient errors

15) A prescription is written for clindamycin 75 mg/5 mL: "Take 10 mL PO TID x 7 days." How many mL should be dispensed?

- a) 140 mL
- b) 180 mL
- c) 200 mL
- d) 210 mL

16) A patient asks which medication is the generic for Eliquis.

- a) Apixaban
- b) Rivaroxaban
- c) Dabigatran
- d) Clopidogrel

17) A patient calls asking if their spouse's prescription is ready. The caller is not documented as an authorized representative. What can the technician disclose?

- a) The medication name and whether it is ready
- b) Only that you cannot confirm prescription status without authorization
- c) The prescriber name and directions but not the drug name
- d) Everything if the caller verifies address and DOB

18) A prescription for a Schedule III medication was written 7 months ago with refills remaining. Under federal law, what happens?

- a) It may be refilled if refills remain, up to 12 months
- b) It may be refilled only if the patient pays cash
- c) It cannot be refilled because it is beyond the 6-month limit from the date written
- d) It can be refilled only after obtaining a new DEA number

19) A label prints for "metformin ER 500 mg," but the stock bottle pulled is "metformin 500 mg" (immediate-release). What should the technician do?

- a) Fill with IR because the strength is the same
- b) Change the directions to BID to match IR
- c) Dispense IR and add "do not crush"
- d) Return the IR product and pull/select the ER product that matches the label and order

20) The system flags a major interaction: warfarin and metronidazole. The prescription is ready to fill. What should the technician do?

- a) Override the alert since the prescriber ordered it
- b) Remove warfarin from the profile as inactive
- c) Route the order/alert for pharmacist review before dispensing
- d) Add a "take with food" auxiliary label and proceed

21) A patient asks for the generic of Flomax.

- a) Tamsulosin
- b) Terazosin
- c) Finasteride
- d) Tadalafil

22) A hardcopy prescription for promethazine-codeine is presented without the prescriber's signature. What is the appropriate next step?

- a) Fill it if the DEA number is present
- b) Hold the prescription as invalid until a valid, signed prescription is obtained before dispensing
- c) Have the patient sign in the prescriber signature area
- d) Convert it to an eRx and proceed

23) You find a paper with patient name, DOB, and medication list left on the counter. What should you do immediately?

- a) Place it in regular trash
- b) Put it into the pharmacy's shred bin/secure destruction container
- c) Staple it to the register log for later
- d) Leave it on the counter until the pharmacist returns

24) A patient requests pseudoephedrine tablets. What is required at the point of sale under federal law (CMEA)?

- a) A prescription for all pseudoephedrine products
- b) ID verification, sales log entry, and compliance with daily/monthly limits
- c) Mandatory insurance billing
- d) Limiting sales to unit-dose packaging only

25) An eRx indicates "DAW 1" for levothyroxine. What should the technician do?

- a) Dispense any AB-rated generic levothyroxine
- b) Dispense the brand as written with no substitution
- c) Substitute liothyronine
- d) Change manufacturers monthly to reduce cost

26) A patient profile shows "ACE inhibitors—angioedema." A new eRx arrives for lisinopril. What is the correct technician action?

- a) Process it because it came from the prescriber
- b) Substitute losartan automatically and continue
- c) Route the order for pharmacist review due to documented angioedema with ACE inhibitors
- d) Ask the patient if they want to try it and document consent

27) A prescription reads: "Gabapentin 300 mg: take 1 cap PO TID. Dispense #90." What is the day supply?

- a) 15
- b) 30
- c) 45

d) 90

28) A patient brings back an unopened vial of tablets dispensed yesterday and asks you to "put it back on the shelf." What is typical community pharmacy practice?

- a) Return to stock if the seal is intact
- b) Return to stock only if the patient paid cash
- c) Return to stock only if it is noncontrolled
- d) Do not return to stock once it has left the pharmacy

29) Which brand/generic pair is correct?

- a) Toprol-XL / metoprolol tartrate
- b) Lopressor / metoprolol succinate ER
- c) Toprol-XL / metoprolol succinate ER
- d) Inderal LA / metoprolol succinate ER

30) A technician notices hydroxyzine and hydralazine stored side-by-side in similar bottles. What is the best error-prevention step?

- a) Keep them adjacent so staff compare labels more often
- b) Separate storage locations and add visual cues to reduce look-alike/sound-alike errors
- c) Remove one from inventory to prevent confusion
- d) Store both in will-call bins

31) A prescription is for timolol 0.5% eye drops: "Instill 1 drop OU BID." Bottle size is 5 mL. Use 20 drops/mL. What day supply should be billed for 1 bottle?

- a) 12 days
- b) 25 days
- c) 50 days
- d) 100 days

32) A patient asks for the generic of Xarelto.

- a) Apixaban
- b) Rivaroxaban
- c) Dabigatran
- d) Edoxaban

33) A new eRx for "ondansetron ODT 4 mg" is entered, but the product selected is ondansetron 4 mg standard tablets. What should the technician do?

- a) Keep tablets; same strength is equivalent
- b) Select the ODT dosage form to match the prescription before continuing
- c) Change the directions to "dissolve on tongue" and keep tablets
- d) Change the route to IM to match an available product

34) A faxed Schedule II prescription is received for a routine outpatient (not hospice/LTC/emergency exception). What is the best handling?

- a) Treat the fax as the original and fill it
- b) Accept it only if the prescriber is in the same state
- c) Route for pharmacist review because the fax is generally not the valid original for routine outpatient C-II dispensing

d) Fill it only if the patient pays cash

35) A prescription says: "Take 1 tablet PO q6h PRN pain. Max 4 tablets/day." Quantity is 60. What day supply should be billed using the maximum?

- a) 10
- b) 12
- c) 15
- d) 30

36) During receiving, a sealed stock bottle's tablets do not match the expected imprint for that NDC. What should the technician do?

- a) Stock it because the bottle is sealed from the wholesaler
- b) Quarantine the product as suspect and escalate per pharmacy procedure
- c) Re-label the bottle with a different NDC that matches the tablets
- d) Dispense it only to established patients

37) A prescription is for fentanyl 25 mcg/hr patches: "Apply 1 patch every 72 hours. Dispense 10 patches." What day supply is this?

- a) 10 days
- b) 20 days
- c) 30 days
- d) 72 days

38) A patient asks, "Is Plavix for cholesterol?" What is Plavix primarily used for?

- a) Antiplatelet therapy to reduce clot risk
- b) Statin to lower LDL cholesterol
- c) Beta-blocker for heart rate control
- d) Diuretic for swelling

39) A new prescription for insulin aspart vial reads: "Inject 8 units before meals TID." Dispense 1 vial (10 mL, 100 units/mL). What day supply is correct?

- a) 30
- b) 41
- c) 45
- d) 60

40) An eRx arrives for methotrexate 2.5 mg with directions "Take 6 tablets daily for rheumatoid arthritis." What should the technician do?

- a) Enter as written and process
- b) Change "daily" to "weekly" and continue
- c) Route for pharmacist review due to high-alert dosing frequency concern
- d) Dispense with an auxiliary label "avoid alcohol" and continue

41) Which medication is a proton pump inhibitor (PPI)?

- a) Famotidine
- b) Sucralfate
- c) Omeprazole
- d) Dicyclomine

42) The pharmacy is asked to transfer a clonazepam prescription (Schedule IV) to another pharmacy, and state law allows it. Under federal rules, what is generally true?

- a) It may be transferred once for the purpose of refill dispensing between pharmacies
- b) Controlled substances cannot be transferred
- c) It may be transferred unlimited times
- d) Only Schedule II prescriptions can be transferred

43) A patient asks for the generic of Zyrtec.

- a) Cetirizine
- b) Levocetirizine
- c) Loratadine
- d) Fexofenadine

44) A heparin infusion is labeled 25,000 units in 250 mL D5W. The order is to infuse 1,000 units/hour. What is the rate in mL/hour?

- a) 8 mL/hr
- b) 10 mL/hr
- c) 12 mL/hr
- d) 25 mL/hr

45) A technician receives an early refill request for alprazolam; insurance rejects "refill too soon." What is the best technician action?

- a) Change days' supply to force the claim through
- b) Tell the patient to take half-doses until the refill date
- c) Collect relevant details (e.g., travel date) and route through the pharmacy's early-refill documentation/approval process
- d) Dispense now and bill later

46) A controlled-substance hardcopy is missing the patient's address. What should the technician do?

- a) Fill it; patient address is never required
- b) Add an address found online and proceed
- c) Have the patient write an address on the prescription
- d) Route for pharmacist review to determine whether it can be resolved appropriately before dispensing

47) Which medication is an SSRI?

- a) Amitriptyline
- b) Duloxetine
- c) Bupropion
- d) Sertraline

48) A prescription is for valacyclovir 1 g: "Take 1 tablet TID x 7 days." What quantity should be dispensed?

- a) 14
- b) 21
- c) 28

d) 30

49) While filling, several tablets fall onto the floor. What should the technician do?

- a) Wipe them and return to stock
- b) Place them in a separate vial for later use
- c) Dispose of them per procedure and replace with tablets from stock
- d) Dispense them if unbroken and placed in a clean vial

50) A patient profile lists "sulfa antibiotics—rash." An eRx arrives for sulfamethoxazole/trimethoprim. What should the technician do?

- a) Process it because it is a combination product
- b) Enter and override the allergy alert
- c) Substitute nitrofurantoin and continue
- d) Route for pharmacist review due to the documented sulfa allergy

51) Which controlled substance schedule is lisdexamfetamine (Vyvanse)?

- a) Schedule II
- b) Schedule III
- c) Schedule IV
- d) Schedule V

52) A new warfarin prescription for a patient says "Use as directed" with no dosing calendar or specified dose. What is the best technician step?

- a) Enter "1 tablet daily" so the label prints
- b) Enter "UAD" and proceed without clarification
- c) Hold the order for clarification so specific directions are available prior to dispensing
- d) Ask the patient what dose they think they are taking and enter that

53) During data entry, the eRx drug field says "digoxin 0.25 mg," but the directions say "take 0.125 mg daily." What should the technician do?

- a) Dispense 0.25 mg because the drug field controls
- b) Dispense 0.125 mg because the sig is more specific
- c) Dispense 0.25 mg tablets and label "take 1/2 tablet daily"
- d) Route for pharmacist clarification due to conflicting strength information

54) Which brand/generic pair is correct?

- a) Januvia / canagliflozin
- b) Farxiga / sitagliptin
- c) Jardiance / empagliflozin
- d) Invokana / empagliflozin

55) A prescription for Humalog KwikPen reads: "Inject 12 units TID with meals." Dispense 1 box (5 pens x 3 mL; 100 units/mL). What day supply is correct?

- a) 41
- b) 30
- c) 34
- d) 21

56) A patient asks a technician to decide whether they qualify for a vaccine today based on medical conditions and timing. This task is:

- a) Within technician scope if the patient signs consent
- b) Within technician scope if a standing order exists
- c) Not within typical technician scope; eligibility assessment requires authorized clinician/pharmacist involvement
- d) Required for technicians before printing a VIS

57) A prescription is for fluconazole 150 mg: "Take 1 tablet now and repeat in 3 days." Quantity?

- a) 1
- b) 2
- c) 3
- d) 4

58) A frozen vaccine product was left at room temperature for 2 hours. What should the technician do?

- a) Put it back in the freezer and use it today
- b) Refrigerate it and extend the beyond-use date
- c) Shake it well and proceed
- d) Quarantine it and route for pharmacist guidance per excursion procedure

59) Which medication is a beta-blocker?

- a) Losartan
- b) Amlodipine
- c) Metoprolol
- d) Hydrochlorothiazide

60) A liquid medication is labeled 10 mg/5 mL. The directions are 2.5 mL PO daily. How many mg per day is the dose?

- a) 2.5 mg
- b) 5 mg
- c) 10 mg
- d) 20 mg

61) A patient says they have a penicillin allergy, but the allergy section in the profile is blank. What should the technician do?

- a) Ignore it because allergies are optional
- b) Document the allergy in the patient profile using the system's allergy entry process
- c) Add "take with food" to reduce risk
- d) Tell the patient to contact the prescriber; do not document it

62) A prescription is written for "ciprofloxacin otic." The pharmacy only has ciprofloxacin ophthalmic solution. What is the best technician step?

- a) Dispense ophthalmic automatically because it is sterile
- b) Change the prescription to ophthalmic and proceed
- c) Route for pharmacist review because otic/ophthalmic interchangeability requires pharmacist decision and policy compliance

d) Dispense ophthalmic and handwrite "for ear" on the label

63) Which medication commonly requires REMS procedures before dispensing in outpatient settings?

- a) Metformin
- b) Isotretinoin
- c) Amoxicillin
- d) Amlodipine

64) A prescription reads: "Take 2 tablets PO q12h." Each tablet is 500 mg. How many mg per day is ordered?

- a) 500 mg/day
- b) 1,000 mg/day
- c) 2,000 mg/day
- d) 4,000 mg/day

65) At pickup, the barcode scan shows the POS patient does not match the will-call bag label. What should the technician do first?

- a) Override and finish the sale because barcodes fail sometimes
- b) Stop the transaction and re-verify the correct patient and correct bag before proceeding
- c) Change the POS patient to match the bag
- d) Complete the sale and fix the mismatch later

66) A patient has an active pantoprazole 40 mg daily on the profile. A new eRx arrives for omeprazole 20 mg daily with no note. What should the technician do?

- a) Process both; two PPIs are common
- b) Inactivate pantoprazole automatically
- c) Route for pharmacist review for potential duplication/therapy change
- d) Replace both with famotidine during entry

67) A prescription is for benzonatate 200 mg: "Take 1 capsule TID PRN cough." Which auxiliary instruction is most appropriate?

- a) Avoid grapefruit juice
- b) Swallow whole; do not chew or dissolve
- c) May cause photosensitivity
- d) Take with dairy products

68) A prescription is for ceftriaxone 1 g IM once. The pharmacy stocks 500 mg vials. How many vials are needed for a 1 g dose?

- a) 1
- b) 2
- c) 3
- d) 4

69) A patient asks the pharmacy to "take my old meds back so someone else can use them." What is the most appropriate response?

- a) Accept and restock if unopened
- b) Accept and restock if noncontrolled

- c) Decline restocking and provide disposal options (e.g., take-back/disposal guidance) if available
- d) Accept and restock if expiration is more than 6 months away

70) A prescription is for insulin glargine U-100 pens: "Inject 22 units nightly." Dispense 1 box (5 pens x 3 mL; 100 units/mL). What day supply is correct?

- a) 68
- b) 75
- c) 34
- d) 30

71) Which antibiotic is the generic for Keflex?

- a) Cefdinir
- b) Cephalexin
- c) Cefuroxime
- d) Cefoxitin

72) Two different strengths of glimepiride are mixed in the same shelf slot. What should the technician do?

- a) Leave them; the pharmacist will catch it at final verification
- b) Combine open bottles to reduce clutter
- c) Separate by strength/NDC and correct shelf placement immediately
- d) Move both strengths to will-call to prevent selection errors

73) A patient asks, "Is Xanax the same as clonazepam?"

- a) Yes, Xanax is clonazepam
- b) Yes, clonazepam is the brand for Xanax
- c) No, Xanax is alprazolam; clonazepam is a different benzodiazepine
- d) No, Xanax is diazepam

74) A third party calls to pay with the patient's card on file and asks, "What meds am I paying for?" The caller is not authorized. What is the best technician approach?

- a) Read the medication names to confirm accuracy
- b) Refuse all third-party payments as always illegal
- c) Accept payment while minimizing PHI disclosure and avoid discussing medication details with an unauthorized caller
- d) Provide full counseling to the cardholder

75) A prescription is for polyethylene glycol 3350 powder: "17 g PO daily." Container size is 510 g. What is the approximate day supply?

- a) 15 days
- b) 20 days
- c) 30 days
- d) 60 days

76) During entry, the technician types amlodipine 50 mg instead of 5 mg. The prescription clearly shows 5 mg. What should the technician do?

- a) Leave it for the pharmacist to catch later

- b) Correct the entry to match the prescription and continue processing for verification
- c) Dispense 5 mg tablets but keep the label at 50 mg
- d) Cancel the prescription and tell the patient to call the prescriber

77) A patient profile shows “morphine—anaphylaxis.” A new eRx arrives for hydromorphone 2 mg. What should the technician do?

- a) Process as written; all opioid allergies are intolerances
- b) Substitute oxycodone/acetaminophen and continue
- c) Remove the allergy from the profile
- d) Route for pharmacist review due to severe opioid allergy history

78) A label is printed for losartan 50 mg, but lisinopril 20 mg was counted into the vial. What is the correct technician action?

- a) Change the label to lisinopril 20 mg and proceed
- b) Remove the incorrect tablets, correct the fill using the right medication, and continue through verification steps
- c) Leave it; the pharmacist will catch it at final check
- d) Mix both and label as “blood pressure medication”

79) Another pharmacy calls requesting a transfer of tramadol with refills remaining. What should the technician do?

- a) Provide the transfer verbally if the caller confirms the DEA number
- b) Refuse because tramadol can never be transferred
- c) Route the request to the pharmacist because controlled-substance transfers require pharmacist involvement and documentation
- d) Transfer only directions and omit refill information to reduce errors

80) Barcode verification flags “wrong NDC” after a look-alike product was selected. What should the technician do next?

- a) Override because the drug name is similar
- b) Reprint the label to match the product that was pulled
- c) Stop and correct the product selection to the exact prescribed product/NDC before proceeding
- d) Continue; counseling will catch it

81) A prescription is for Lantus SoloStar. Insulin glargine is best classified as:

- a) Rapid-acting insulin
- b) Short-acting insulin
- c) Intermediate-acting insulin
- d) Long-acting insulin

82) An eRx for Adderall XR 20 mg is missing the patient address field. What should the technician do?

- a) Add the address from the patient profile and proceed without review
- b) Change to Adderall IR 10 mg BID and proceed
- c) Delete the eRx and tell the patient to contact the prescriber
- d) Route for pharmacist review because controlled-substance validation may require confirmation of required elements

83) A prescription is for 1.5 liters of normal saline to infuse over 12 hours. What is the infusion rate in mL/hour?

- a) 100 mL/hr
- b) 125 mL/hr
- c) 150 mL/hr
- d) 175 mL/hr

84) A DSCSA shipment arrives with transaction information and a transaction statement. What is the primary purpose of these documents?

- a) To establish therapeutic equivalence for substitution
- b) To trace products through the supply chain and support verification of product legitimacy
- c) To set the usual and customary price
- d) To provide patient counseling instructions

85) A patient asks, "What is the generic for Lasix?"

- a) Spironolactone
- b) Torsemide
- c) Bumetanide
- d) Furosemide

86) A prescription is for hydrocortisone 2.5% cream: "Apply thin layer to affected area BID." The prescriber writes "Dispense 30 g." Which entry is most appropriate for quantity?

- a) 30 mL
- b) 30 g
- c) 2 tubes
- d) 60 g

87) A patient says they need "my water pill refilled," and the profile shows furosemide 20 mg daily and hydrochlorothiazide 25 mg daily from different prescribers. The patient doesn't know which one. What should the technician do first?

- a) Refill the most recently filled diuretic
- b) Refill both and let the patient decide at pickup
- c) Ask the patient to confirm by reading the last label or describing the tablet/prescriber, then match to the profile
- d) Choose based on what condition the patient reports

88) A new prescription for a controlled substance is phoned in to the pharmacy. Who may take the verbal order (federal standard)?

- a) A technician, if the prescriber provides a DEA number
- b) A technician, if the patient is established
- c) Any staff member as long as it is documented
- d) A pharmacist (or appropriately authorized individual per law); the technician should route per workflow

89) A prescription is written for "Ventolin HFA DAW 1." The insurance prefers generic albuterol HFA. What should the technician do?

- a) Dispense generic because it's therapeutically equivalent

- b) Dispense brand because DAW 1 indicates no substitution
- c) Change DAW to 0 to get coverage
- d) Dispense generic and annotate "brand unavailable"

90) A prescription is for sertraline 100 mg tablets with directions: "Take 1/2 tablet daily for 7 days, then 1 tablet daily." The prescriber wants a 30-day supply. How many tablets should be dispensed?

- a) 26 tablets
- b) 27 tablets
- c) 30 tablets
- d) 33 tablets

Answer Key

1) b	31) b	61) b
2) a	32) b	62) c
3) b	33) b	63) b
4) b	34) c	64) c
5) c	35) c	65) b
6) b	36) b	66) c
7) c	37) c	67) b
8) b	38) a	68) b
9) a	39) b	69) c
10) b	40) c	70) a
11) c	41) c	71) b
12) a	42) a	72) c
13) b	43) a	73) c
14) d	44) b	74) c
15) d	45) c	75) d
16) a	46) d	76) b
17) b	47) d	77) d
18) c	48) b	78) b
19) d	49) c	79) c
20) c	50) d	80) c
21) a	51) a	81) d
22) b	52) c	82) d
23) b	53) d	83) b
24) b	54) c	84) b
25) b	55) c	85) d
26) c	56) c	86) b
27) b	57) b	87) c
28) d	58) d	88) d
29) c	59) c	89) b
30) b	60) b	90) a

PTCB Sim Test 31

1. A new eRx arrives for duloxetine 60 mg daily. The patient profile shows duloxetine 30 mg daily filled 5 days ago, and a note says "titrating." The patient says the prescriber increased it yesterday. What is the best technician action?

- a) Enter the new prescription as written and route for pharmacist verification as a dose increase
- b) Discontinue the 30 mg prescription in the system without review
- c) Change the new prescription to 30 mg to match the most recent fill
- d) Tell the patient to finish all 30 mg capsules before starting 60 mg

2. A customer asks to buy pseudoephedrine 30 mg tablets, quantity 96. Which step is required before completing the sale?

- a) Document the sale only if the customer pays cash
- b) Verify identity and complete the required electronic or logbook record
- c) Require a prescription because 96 tablets exceeds the federal daily limit
- d) Refuse the sale unless the customer is at least 21 years old

3. During filling, the barcode scan shows the product in your hand is hydroxyzine 25 mg, but the label is for hydralazine 25 mg. What should the technician do next?

- a) Override the scan because both start with "hydra-"
- b) Reprint the label to match the scanned product
- c) Stop and select the correct product/NDC before continuing
- d) Continue filling and rely on final verification to catch it

4. A prescription is written for "metoprolol succinate ER 50 mg: take 1 tablet twice daily." The patient's prior directions were "take 1 tablet daily." What is the best technician action?

- a) Enter as written and route for pharmacist review due to ER formulation/direction mismatch
- b) Change directions to once daily to match the previous label
- c) Substitute metoprolol tartrate 50 mg twice daily because it matches dosing
- d) Enter as written and proceed to fill because it is not a controlled substance

5. A spouse tries to pick up a prescription and asks, "What is this medication for?" There is no documented authorization on file. What should the technician do?

- a) Provide the indication if the spouse can confirm the address
- b) Provide the drug name only, but not the directions
- c) Explain you can't discuss medication details without authorization and offer to contact the patient
- d) Provide the full counseling points because pickup implies permission

6. Amoxicillin/clavulanate suspension is prescribed: "Take 7.5 mL by mouth twice daily for 10 days." How many mL should be dispensed?

- a) 150 mL
- b) 100 mL
- c) 120 mL
- d) 180 mL

7. A patient returns an unopened box of insulin pens stating it was left in a hot car all day. What is the appropriate technician action?

- a) Place it back into inventory since it is unopened
- b) Quarantine it and route to the pharmacist/manager for disposition decision
- c) Exchange it immediately and document "temperature excursion"
- d) Tell the patient to use it if the solution is still clear

8. You are processing isotretinoin for a new patient. What must be confirmed before dispensing?

- a) Only that a Medication Guide prints with the label
- b) That the prescriber is enrolled and the patient authorization is current in the REMS
- c) That the patient signs a waiver acknowledging pregnancy risk
- d) That the patient is at least 18 years old

9. The hardcopy directions state "take 1 tablet twice daily," but the label prints "take 1 tablet daily." What should the technician do?

- a) Cross out the label direction and handwrite the correct one
- b) Call the prescriber directly to verify the intended dose
- c) Stop processing and route for correction/verification before dispensing
- d) Dispense as printed and have the patient confirm at pickup

10. During entry for lisinopril, the profile lists "ACE inhibitors—angioedema." What is the best technician action?

- a) Proceed because lisinopril is standard therapy for hypertension
- b) Remove the allergy because the reaction is not specific to lisinopril
- c) Change the medication to losartan without a new prescription
- d) Enter and route to the pharmacist because it's a high-severity allergy flag

11. A prescription is written for diltiazem ER 240 mg daily, but only diltiazem IR 60 mg is in stock. What should the technician do?

- a) Substitute IR 60 mg and change directions to four times daily
- b) Hold and route because changing formulation requires pharmacist decision/prescriber authorization
- c) Dispense IR 60 mg and note "ER out of stock" on the label
- d) Call the prescriber and request a new prescription without involving the pharmacist

12. A recall notice identifies a specific lot number of losartan currently on your shelf. What should the technician do first?

- a) Pull and quarantine the affected lot per recall procedure
- b) Continue dispensing until replacement stock arrives
- c) Remove only opened bottles; sealed bottles can remain
- d) Call every patient who has ever received losartan to stop therapy

13. A paper prescription for alprazolam is presented with no prescriber DEA number on the hardcopy. What is the best technician action?

- a) Look up the prescriber's DEA online and add it to the prescription
- b) Fill a partial supply and obtain the DEA later
- c) Treat it as a noncontrolled prescription because the prescriber is known

d) Hold and route for pharmacist review because required controlled-substance elements are missing

14. While preparing a nonsterile compound, you notice the balance calibration sticker is expired. What should you do?

- a) Proceed if another technician double-checks the weights
- b) Estimate weights using volume and document the estimate
- c) Remove the balance from use and use a properly calibrated/verified balance
- d) Proceed as long as the final mixture looks uniform

15. A new prescription is for metformin ER 500 mg twice daily. The system defaulted to metformin IR 500 mg. What should the technician do?

- a) Keep IR because the dose is the same
- b) Change to ER selection to match the prescription before processing
- c) Keep IR and add "do not crush" to make it equivalent
- d) Change to ER only if the patient requests it

16. Another pharmacy requests a transfer for clonazepam with refills remaining. What is the correct technician action?

- a) Complete the transfer yourself because Schedule IV transfers are allowed
- b) Decline because controlled substances can never be transferred
- c) Route to the pharmacist to complete the controlled-substance transfer
- d) Provide the information by fax and document it under your initials

17. A patient picking up warfarin asks what foods they should avoid. What should the technician do?

- a) Tell them to avoid all green vegetables completely
- b) Recommend taking warfarin with a high-fat meal for consistency
- c) Tell them to increase vitamin K foods to "cancel out" warfarin
- d) Offer pharmacist counseling for diet consistency and monitoring guidance

18. Gentamicin ophthalmic solution: "Instill 2 drops in both eyes every 4 hours for 7 days." If 1 mL = 20 drops, how many mL are needed for the full course (assume around-the-clock use)?

- a) 5.6 mL
- b) 8.4 mL
- c) 7.0 mL
- d) 14.0 mL

19. A refrigerated vaccine shipment arrives and the temperature monitor indicates out-of-range exposure during transit. What should the technician do?

- a) Put it in the refrigerator and use it first
- b) Quarantine it and route to the pharmacist/manager for guidance before use
- c) Discard immediately to prevent accidental use
- d) Freeze it briefly to restore potency

20. A new prescription for naproxen is entered and an alert shows an active ibuprofen prescription. What should the technician do?

- a) Override because patients often take multiple NSAIDs
- b) Delete the naproxen prescription because ibuprofen is already active
- c) Route for pharmacist review of duplication and safety risk
- d) Substitute acetaminophen automatically to avoid duplication

21. The sig reads: "1 tab PO qhs PRN insomnia." What does this mean?

- a) Take 1 tablet by mouth at bedtime as needed for insomnia
- b) Take 1 tablet by mouth every hour as needed for insomnia
- c) Take 1 tablet by mouth every morning as needed for insomnia
- d) Take 1 tablet by mouth four times daily as needed for insomnia

22. Heparin infusion is 25,000 units in 250 mL. The order is 1,200 units/hour. What pump rate (mL/hour) is needed?

- a) 10 mL/hr
- b) 11 mL/hr
- c) 12 mL/hr
- d) 15 mL/hr

23. During a controlled-substance count, an open bottle of hydromorphone tablets is short and there is no documentation. What is the best technician action?

- a) Adjust the inventory count to match what is physically present
- b) Report the discrepancy immediately per controlled-substance procedures
- c) Finish counting all other controlled substances before mentioning it
- d) Combine tablets from another bottle to correct the count

24. A patient taking apixaban asks which OTC pain reliever is safest. What should the technician do?

- a) Recommend ibuprofen because it's effective for pain
- b) Recommend naproxen because it lasts longer
- c) Recommend aspirin because it's cardioprotective
- d) Refer to the pharmacist for OTC selection and bleeding-risk counseling

25. Azithromycin 250 mg tablets: "Take 2 tablets today, then 1 tablet daily for 4 days." What quantity should be dispensed?

- a) 5 tablets
- b) 6 tablets
- c) 7 tablets
- d) 8 tablets

26. An eRx is written for "Lamictal 100 mg daily." The patient previously received lamotrigine 100 mg. What is the best technician action in a state that allows generic substitution unless DAW is indicated?

- a) Contact the prescriber to rewrite it as lamotrigine
- b) Dispense brand only unless the patient signs a waiver
- c) Process per substitution rules/system settings and continue workflow
- d) Hold because brand-name prescriptions cannot be substituted

27. A patient requests a printed medication profile and presents a photo ID. What is the best

technician action?

- a) Refuse because profiles can only be sent directly to prescribers
- b) Provide the profile after verifying identity per pharmacy procedure
- c) Provide it but omit any controlled substances automatically
- d) Provide it without verification because the patient is in person

28. Your pharmacy has a policy requiring at least 6 months to expiration when dispensing nitroglycerin SL tablets. Only stock expiring in 3 months is available. What should the technician do?

- a) Dispense anyway because it is not expired
- b) Repackage into an amber vial to extend dating
- c) Obtain acceptable stock or route for appropriate next steps
- d) Reduce quantity to a smaller amount so it's used sooner

29. The profile shows anaphylaxis to penicillin. A new prescription is for cephalexin. What should the technician do?

- a) Fill because cephalexin is not a penicillin
- b) Route for pharmacist review due to severity and potential cross-reactivity
- c) Substitute clindamycin because it's commonly used
- d) Ask the patient if they have taken cephalexin before and decide based on that

30. Insurance rejects a refill as "too soon." The patient says the prescriber increased the dose yesterday. What should the technician do?

- a) Change the days' supply to a smaller number so it pays
- b) Submit an override without new directions since it's a dose change
- c) Obtain the updated prescription/directions and process appropriately through verification/insurance steps
- d) Tell the patient insurance never covers early refills

31. An eRx arrives for zolpidem 10 mg, #30 with 5 refills. What is the best technician action?

- a) Process exactly as written because it was transmitted electronically
- b) Change refills to 0 because zolpidem is Schedule II
- c) Delete because controlled substances cannot be e-prescribed
- d) Route for pharmacist review because it is Schedule IV and refill/appropriateness must be checked

32. A technician accidentally selected prednisolone instead of prednisone during entry. Which workflow step most directly helps prevent this error from reaching the patient?

- a) Compare tablet color during counting
- b) Use barcode/NDC verification during filling and do not bypass alerts
- c) Store similar names next to each other to reduce search time
- d) Rely on the patient to catch the mistake at pickup

33. A new prescription for sertraline is being processed. The patient says they started linezolid 2 days ago from urgent care. What should the technician do?

- a) Continue because linezolid is short-term therapy
- b) Change sertraline to citalopram to avoid the interaction
- c) Route for pharmacist review due to a high-risk interaction concern

d) Tell the patient to stop linezolid immediately

34. Albuterol HFA: inhale 2 puffs every 4 hours as needed. Inhaler contains 200 actuations. If used at the maximum frequency around the clock, what is the days' supply?

a) 25 days

b) 12 days

c) 14 days

d) 16 days

35. A prescriber's office calls asking you to "approve" a tramadol refill. What should the technician do?

a) Approve it because tramadol is not Schedule II

b) Add refills based on the patient's previous history

c) Route the request through the normal workflow for pharmacist authorization

d) Transfer the prescription to another pharmacy to avoid delays

36. In a hospital cart-fill area, oral syringes are stored next to hypodermic needles. What action best reduces wrong-route risk?

a) Leave them together because both are "syringes"

b) Separate oral syringes from injectable supplies

c) Relabel oral syringes "for injection" to avoid confusion

d) Discard oral syringes because they are not sterile

37. A prescription is written for "metoprolol 25 mg" without specifying formulation. The patient history shows metoprolol succinate ER once daily. What should the technician do?

a) Select metoprolol tartrate because it is most common

b) Select succinate ER to match the profile without asking

c) Split the quantity between tartrate and succinate

d) Route for pharmacist review because formulation cannot be assumed

38. A patient asks if they can drink alcohol while taking metronidazole. What is the best technician response?

a) Advise spacing alcohol and doses by 2 hours

b) Recommend moderate alcohol is acceptable

c) Offer pharmacist counseling for alcohol-avoidance guidance

d) Suggest alcohol improves absorption

39. A stock bottle of tablets without a label is found in the return-to-stock bin. What should the technician do?

a) Identify by imprint and relabel it for use

b) Return to stock based on tablet color and shape

c) Place tablets in a vial labeled "unknown" to use later

d) Quarantine and route per procedure for proper disposal/handling

40. Cyanocobalamin 1,000 mcg/mL injection: "Inject 0.6 mL IM monthly." How many mcg is each dose?

a) 1,000 mcg

b) 600 mcg

- c) 1,600 mcg
- d) 6,000 mcg

41. A patient presents a paper prescription for hydrocodone/acetaminophen dated 7 months ago that has never been filled. What should the technician do?

- a) Fill because controlled prescriptions never expire
- b) Fill a partial supply and treat the rest as refills
- c) Route for pharmacist review because controlled-substance validity depends on time limits and state law
- d) Convert to acetaminophen-only and dispense

42. DSCSA documentation is missing transaction information for a delivered product. What should the technician do?

- a) Stock it and resolve paperwork later
- b) Quarantine and route for investigation before dispensing
- c) Dispense it only for cash patients
- d) Return it to the wholesaler without documenting anything

43. Furosemide 40 mg: "Take 1 tablet by mouth daily." Quantity 90. What is the days' supply?

- a) 45 days
- b) 60 days
- c) 30 days
- d) 90 days

44. Another pharmacy requests a transfer for a Schedule III prescription with refills remaining. What should the technician do?

- a) Take the transfer by phone and document it yourself
- b) Decline because Schedule III prescriptions are not transferable
- c) Route to the pharmacist to complete the controlled-substance transfer
- d) Accept the transfer by fax and process it immediately

45. A profile shows simvastatin active. A new prescription arrives for atorvastatin and the system flags duplicate therapy. What should the technician do?

- a) Fill both because some patients take two statins
- b) Automatically inactivate simvastatin and fill atorvastatin
- c) Ask the patient which one they want and dispense that one
- d) Route for pharmacist review to confirm switch vs duplication

46. A prescription is written for Humalog. During entry, "Humalog Mix 75/25" was selected. What is the best technician action?

- a) Proceed because both contain insulin lispro
- b) Dispense the mix and add "use as directed"
- c) Correct the selection and route for verification because mix insulin is not interchangeable
- d) Change to NovoLog because it is also rapid-acting insulin

47. In will-call, two prescriptions are for "Maria L." and "Marie L." with the same last name. What is the best technician step to reduce mix-ups?

- a) Bag them together so the family can pick up at once
- b) Remove DOB from labels to protect privacy
- c) Separate bags and use at least two identifiers at pickup (e.g., name and DOB)
- d) Place both in the front to speed checkout

48. A label says: "Take 1.5 tablets daily." Quantity dispensed is 45 tablets. What is the days' supply?

- a) 25 days
- b) 30 days
- c) 20 days
- d) 45 days

49. A caller asks the status of a prescription and provides the patient name and DOB but cannot confirm address. What should the technician do?

- a) Provide full details since DOB matches
- b) Provide only store hours and refuse any prescription information
- c) Ask for additional verification before disclosing protected information
- d) Provide the drug name but not the directions

50. You enter clopidogrel and see omeprazole OTC listed as active. What should the technician do?

- a) Ignore OTC products because they are not prescriptions
- b) Remove omeprazole from the profile since it's OTC
- c) Route for pharmacist review due to a potential interaction concern
- d) Tell the patient to stop clopidogrel immediately

51. Azithromycin suspension 200 mg/5 mL: "12.5 mL on day 1, then 6.25 mL daily on days 2–5." What total volume should be dispensed?

- a) 37.5 mL
- b) 31.25 mL
- c) 43.75 mL
- d) 50 mL

52. A prescription for methylphenidate IR 10 mg is written "1 tablet three times daily," #90, with 2 refills. What is the best technician action?

- a) Process as written because it is maintenance therapy
- b) Convert to extended-release and keep refills
- c) Dispense #30 and treat the remainder as refills
- d) Route for pharmacist review because Schedule II prescriptions cannot have refills

53. A medication requires REMS documentation prior to dispensing, but the authorization cannot be confirmed. What should the technician do?

- a) Dispense and document "REMS pending"
- b) Hold and route for completion of required REMS steps before dispensing
- c) Substitute a similar medication not in REMS
- d) Have the patient sign a waiver and dispense

54. Morphine sulfate oral solution 20 mg/mL: "Take 0.25 mL every 4 hours as needed." How

many mg per dose?

- a) 10 mg
- b) 2.5 mg
- c) 5 mg
- d) 20 mg

55. The profile lists an allergy to Norvasc. A new prescription is for amlodipine. What should the technician do?

- a) Fill because brand-name allergies don't apply to generics
- b) Delete the allergy because it's likely intolerance
- c) Substitute nifedipine ER because it's in the same class
- d) Route for pharmacist review because Norvasc is amlodipine

56. When reconstituting an antibiotic, the bottle says "Add 62 mL water," but the computer instruction says "Add 64 mL." What should the technician do?

- a) Add 63 mL to split the difference
- b) Follow the computer instruction because it prints from the system
- c) Verify using product labeling and correct the workflow before mixing
- d) Add 64 mL and shake longer to compensate

57. A patient is picking up buprenorphine/naloxone films. Another customer asks what medication it is. What should the technician do?

- a) State the medication name because it is commonly used
- b) Explain the indication so the customer understands
- c) Confirm only if the customer also uses the pharmacy
- d) Decline to disclose and keep the transaction discreet

58. A prescription is entered for glipizide XL 10 mg, but glipizide IR 10 mg was selected. What should the technician do?

- a) Dispense IR because the ingredient is the same
- b) Change directions to twice daily and dispense IR
- c) Correct to XL selection (or route) because IR and XL are not interchangeable
- d) Dispense IR and add "do not crush"

59. A technician is handling multiple hazardous tablets for returns processing. Which PPE is most appropriate?

- a) Cloth gloves to improve grip
- b) Chemotherapy-rated hazardous-drug gloves
- c) Safety glasses only
- d) No PPE is needed if bottles are sealed

60. Directions read: "Take 2 tsp by mouth every 6 hours as needed." How many mL is each dose?

- a) 7.5 mL
- b) 5 mL
- c) 10 mL
- d) 15 mL

61. A patient asks whether gabapentin is controlled and whether it can be refilled early. What is the best technician response?

- a) Confirm it is Schedule V federally and early refills are always allowed
- b) State it is never controlled anywhere
- c) Explain scheduling rules vary by state and advise review through the pharmacy's process for early refills
- d) State early refills are illegal for gabapentin in all states

62. Potassium chloride ER 20 mEq: "Take 2 tablets daily," #60. The profile shows potassium chloride ER 10 mEq daily still active. What should the technician do?

- a) Inactivate the 10 mEq automatically and fill the new prescription
- b) Fill both strengths so the patient can choose
- c) Change the new prescription to 10 mEq to match the profile
- d) Route for pharmacist review of potential duplication and dose change

63. A 17-year-old requests Zyrtec-D (pseudoephedrine-containing) but has no ID. What should the technician do?

- a) Sell it because it is OTC
- b) Sell it and complete the log later
- c) Require a prescription because the buyer is under 18
- d) Do not complete the sale without required ID/log steps; offer non-pseudoephedrine alternatives

64. A sealed stock bottle shows visible moisture/condensation inside. What should the technician do?

- a) Use the tablets because the seal indicates safety
- b) Pour tablets into a dry vial and use them first
- c) Quarantine the bottle and route for return/quality disposition
- d) Dispense only the dry tablets and discard damp ones

65. Budesonide/formoterol 160/4.5: inhale 2 puffs twice daily. Inhaler contains 120 inhalations. What is the days' supply?

- a) 60 days
- b) 20 days
- c) 30 days
- d) 15 days

66. An eRx for a Schedule II stimulant arrives missing the patient address field. What should the technician do?

- a) Add the address from the profile and proceed
- b) Dispense a small emergency supply and request corrected eRx later
- c) Ask the patient to write the address on the printed copy
- d) Route for pharmacist review because required elements may need correction before dispensing

67. A patient requests "Prograf," but the prescription is written for tacrolimus 1 mg capsules. What should the technician do?

- a) Refuse substitution because brand and generic are never interchangeable

- b) Automatically dispense brand Prograf regardless of the prescription
- c) Process tacrolimus as prescribed; handle brand requests through appropriate substitution/billing steps
- d) Change the prescription in the system to brand and rebill without review

68. Clonidine patch 0.2 mg/day: apply 1 patch weekly. A box contains 4 patches. What is the days' supply for one box?

- a) 30 days
- b) 21 days
- c) 28 days
- d) 14 days

69. The prescription is for metronidazole vaginal gel, but metronidazole topical skin gel was selected. What should the technician do?

- a) Dispense topical gel because it is the same drug
- b) Change directions to apply externally to match the selected product
- c) Dispense both and let the patient choose
- d) Correct the product selection because route/dosage form must match

70. You are taking a transfer-in for a noncontrolled prescription. Which information is most important to capture accurately?

- a) Wholesaler invoice number
- b) Original prescription number and refills remaining
- c) Patient's preferred pickup time
- d) Insurance BIN and PCN

71. A wrong strength (Eliquis 2.5 mg instead of 5 mg) was selected because barcode scanning was skipped. What is the best prevention step?

- a) Store all strengths together so they are easy to find
- b) Remove NDC data to force manual checks
- c) Stop technicians from stocking shelves
- d) Require and enforce barcode/NDC scanning during the fill process

72. Cephalexin 500 mg: "Take 1 capsule by mouth four times daily." Quantity 28. What is the days' supply?

- a) 5 days
- b) 14 days
- c) 7 days
- d) 10 days

73. You discover two profiles for the same patient with different DOBs. The patient is at the counter. What should the technician do?

- a) Merge immediately because the names match
- b) Verify identifiers with the patient and correct/merge per workflow to prevent errors
- c) Ignore it if the address matches
- d) Delete the older profile and keep the newest

74. Augmentin 875/125: "Take 1 tablet by mouth twice daily for 7 days." What quantity

should be dispensed?

- a) 21 tablets
- b) 10 tablets
- c) 7 tablets
- d) 14 tablets

75. The pharmacy has only part of a Schedule II medication in stock and the patient requests a partial fill. What should the technician do?

- a) Decide a partial quantity, dispense it, and document in the notes
- b) Decline because partial fills are never allowed for Schedule II
- c) Route for pharmacist handling because Schedule II partial fills have specific rules and documentation
- d) Convert to a Schedule III alternative so refills are allowed

76. A prescriber's office calls: "Change escitalopram from 10 mg to 20 mg; the doctor approved." What should the technician do?

- a) Change the strength based on the call and document the caller's name
- b) Change directions to "take 2 tablets daily" and keep 10 mg
- c) Dispense 10 mg now and change next month
- d) Process the change through the required authorization pathway and route appropriately before updating

77. A prescription is written for glyburide. The profile has multiple notes of hypoglycemia episodes. What should the technician do?

- a) Substitute glipizide automatically because it is safer
- b) Dispense and add "take with food"
- c) Change to metformin because it is first-line
- d) Route for pharmacist review before dispensing

78. The profile lists a latex allergy. Vaccine supplies available indicate latex in the syringe stopper. What should the technician do?

- a) Proceed because latex allergy is unrelated to vaccines
- b) Ask if the allergy is "still active," then proceed
- c) Proceed and shorten observation time
- d) Route to ensure latex-free supplies or an appropriate plan is used

79. A noncontrolled hardcopy for doxycycline has a prescriber signature but no date written. What should the technician do?

- a) Fill because it is not a controlled substance
- b) Add today's date to complete the prescription
- c) Route for pharmacist review to determine whether clarification is needed before dispensing
- d) Tell the patient to write in a date at the counter

80. Insulin glargine U-100 vial: inject 18 units at bedtime. A 10 mL vial contains 1,000 units. What is the days' supply based on units only?

- a) 60 days
- b) 45 days

- c) 55 days
- d) 50 days

81. A prescription is for "Lantus SoloStar," but "Lantus vial" was selected in the system. What should the technician do?

- a) Dispense the vial because it contains the same insulin
- b) Dispense the vial and include pen needles
- c) Correct to the pen product because pen vs vial is a dosage-form selection issue
- d) Change the prescription to vial to match stock

82. A patient asks the pharmacy to dispose of unused fentanyl patches. What is the best technician action?

- a) Accept them and discard in regular trash behind the counter
- b) Provide appropriate disposal options (take-back/collection site) and process via the pharmacy's approved method
- c) Tell the patient to flush them immediately at home
- d) Place them into the pharmacy's regular returns bin

83. Levetiracetam ER is on backorder. The prescription says "do not substitute." What should the technician do?

- a) Dispense levetiracetam IR and change directions to twice daily
- b) Tell the patient to skip doses until ER is available
- c) Transfer the prescription without telling the patient
- d) Route for pharmacist contact with prescriber/patient for an appropriate plan

84. A topical medication directions are "Apply 2 grams daily." Tube size is 60 grams. What is the days' supply?

- a) 60 days
- b) 15 days
- c) 20 days
- d) 30 days

85. A filled prescription has the correct drug and directions but the wrong patient name on the label. It has not left the pharmacy. What should the technician do?

- a) Reprint the label and continue without documenting since it was caught early
- b) Cross out the name and handwrite the correct one
- c) Dispense if the correct patient confirms DOB at pickup
- d) Correct the labeling error and route through the quality process required for documentation before dispensing

86. Valacyclovir 1 g: "Take 2 tablets twice daily for 1 day." Quantity is 4 tablets. The note indicates cold sore treatment. What should the technician do?

- a) Increase quantity to 8 because cold sores require 2 days of therapy
- b) Process as written because quantity and directions match the intended 1-day regimen
- c) Change to 1 tablet daily because that is typical antiviral dosing
- d) Reject because valacyclovir is not used for cold sores

87. An order is for vancomycin 1 g IV every 12 hours. Vials available are 500 mg. How many

vials are needed per dose?

- a) 3 vials
- b) 2 vials
- c) 1 vial
- d) 4 vials

88. A patient calls for a lorazepam refill. The profile shows 0 refills remaining and the prescription is expired. What should the technician do?

- a) Provide a courtesy 3-day supply and request authorization later
- b) Transfer it to another pharmacy to obtain refills
- c) Tell the patient to take half doses until the next appointment
- d) Send a refill request/new prescription request through the workflow for appropriate authorization

89. A new prescription is for rosuvastatin. The profile shows atorvastatin active with supply on hand. The patient says, "My cardiologist switched it." What should the technician do?

- a) Cancel rosuvastatin because two statins should never appear
- b) Tell the patient to finish atorvastatin first
- c) Enter/process the new prescription and ensure it is reviewed as a switch vs duplication before dispensing
- d) Inactivate atorvastatin immediately and fill rosuvastatin without review

90. A patient presents a paper prescription for promethazine-codeine syrup. It has a prescriber signature and DEA, but the patient address is missing. What should the technician do?

- a) Enter and dispense because you can pull the address from the profile
- b) Ask the patient to write their address on the prescription
- c) Convert it to promethazine without codeine and dispense
- d) Hold and route for pharmacist review because controlled-substance validity elements must be confirmed

Answer Key

1) a	31) d	61) c
2) b	32) b	62) d
3) c	33) c	63) d
4) a	34) b	64) c
5) c	35) c	65) c
6) a	36) b	66) d
7) b	37) d	67) c
8) b	38) c	68) c
9) c	39) d	69) d
10) d	40) b	70) b
11) b	41) c	71) d
12) a	42) b	72) c
13) d	43) d	73) b
14) c	44) c	74) d
15) b	45) d	75) c
16) c	46) c	76) d
17) d	47) c	77) d
18) b	48) a	78) d
19) b	49) c	79) c
20) c	50) c	80) d
21) a	51) a	81) c
22) c	52) d	82) b
23) b	53) b	83) d
24) d	54) c	84) d
25) b	55) d	85) d
26) c	56) c	86) b
27) b	57) d	87) b
28) c	58) c	88) d
29) b	59) b	89) c
30) c	60) c	90) d

PTCB Sim Test 32

1) At drop-off, a paper prescription reads "levothyroxine 0.05 mg PO daily." The system is prompting for strength in mcg. Which entry matches the written dose?

- a) 5 mcg
- b) 50 mcg
- c) 0.5 mcg
- d) 500 mcg

2) A new eRx arrives: "metformin ER 500 mg: take 2 tabs daily with evening meal." Which quantity should be entered for a 90-day supply?

- a) 90
- b) 180
- c) 270
- d) 360

3) A patient asks to buy two boxes of pseudoephedrine 30 mg tablets, each box containing 48 tablets. How many grams of pseudoephedrine is this total?

- a) 1.44 g
- b) 2.88 g
- c) 3.60 g
- d) 4.80 g

4) A pharmacist asks the technician to take a new verbal prescription for amoxicillin from a prescriber. What is the correct technician action?

- a) Take the verbal and process it if the prescriber provides an NPI
- b) Take the verbal only if it is non-controlled and state law allows technician receipt
- c) Refuse all verbals; only eRx is legal
- d) Take the verbal only if the prescriber is in the same health system

5) During filling, the label is for "sertraline 25 mg," but the scanned stock bottle is "sertraline 50 mg." What should the technician do next?

- a) Continue and counsel the patient to take half a tablet
- b) Change the strength on the label to 50 mg to match the bottle
- c) Stop, obtain the correct 25 mg product, and rescan before continuing
- d) Fill 50 mg and reduce the quantity by half

6) The refrigerator temperature log shows 12°C (53.6°F) for the past 3 hours. Insulin and vaccines were stored there. What should the technician do first?

- a) Move items to the freezer to stabilize them
- b) Continue use since 12°C is close to refrigerated range
- c) Quarantine the affected products and escalate per excursion process
- d) Discard everything immediately without documentation

7) A prescription reads: "amoxicillin/clavulanate 400 mg/57 mg per 5 mL; give 7.5 mL PO BID x 10 days." What total volume should be dispensed?

- a) 120 mL

- b) 150 mL
- c) 165 mL
- d) 180 mL

8) An eRx is written for “Toprol-XL 50 mg daily.” Which product should be selected?

- a) Metoprolol tartrate 50 mg
- b) Metoprolol succinate ER 50 mg
- c) Atenolol 50 mg
- d) Propranolol ER 60 mg

9) A patient profile shows allergy: “sulfonamides—anaphylaxis.” A new eRx is received for sulfamethoxazole/trimethoprim. What is the technician’s best next step?

- a) Process it because it is not a penicillin
- b) Change to nitrofurantoin without contacting the prescriber
- c) Enter the prescription and ensure it is flagged for pharmacist clinical review before dispensing
- d) Remove the allergy because it is nonspecific

10) A controlled substance invoice is being filed. Under federal law, controlled substance records must generally be retained for at least:

- a) 1 year
- b) 2 years
- c) 3 years
- d) 5 years

11) A prescriber’s office calls to correct a non-controlled eRx quantity (should be #90, not #30). What should the technician do?

- a) Change it immediately and write “per office”
- b) Ask the office to resend and delete the original without documenting
- c) Take the details and route to the pharmacist for authorization/acceptance and documentation per requirements
- d) Change it only if the patient confirms they take it daily

12) An eRx arrives for “Xarelto 20 mg daily.” Which generic is Xarelto?

- a) Apixaban
- b) Dabigatran
- c) Rivaroxaban
- d) Edoxaban

13) A paper prescription is written: “Apply 2 grams to affected area BID.” The tube size is 60 grams. What is the days’ supply?

- a) 10 days
- b) 12 days
- c) 15 days
- d) 30 days

14) A technician is entering a new prescription for diltiazem “240 mg ER” with no additional descriptor. The system lists ER, CD, and LA products. What should the technician do?

- a) Select any 240 mg extended-release option; they are equivalent
- b) Select the cheapest NDC to reduce copay
- c) Enter diltiazem IR 60 mg QID because it matches total daily dose
- d) Set aside for pharmacist review because ER formulations may not be interchangeable

15) The pharmacy receives a manufacturer recall notice for one lot of losartan 50 mg tablets. What is the correct technician response?

- a) Remove only opened bottles; keep sealed bottles for later
- b) Pull the affected lot from stock and quarantine per recall process
- c) Keep dispensing until replacement arrives to prevent therapy gaps
- d) Call every patient and instruct them to stop immediately

16) A technician is asked which schedule has no refills allowed under federal law. Which is correct?

- a) C-II
- b) C-III
- c) C-IV
- d) C-V

17) A prescription reads: "ii gtt OU QID x 7 days." The drops are 20 gtt/mL. How many mL will be used?

- a) 2.8 mL
- b) 4.2 mL
- c) 5.6 mL
- d) 7.0 mL

18) The patient says, "What is lisinopril for?" What is the technician's best response?

- a) "It's for blood pressure; it can cause cough."
- b) "It's a water pill; expect increased urination."
- c) "It's for blood pressure; take only when your BP is high."
- d) "The pharmacist can discuss what it's for and how to take it."

19) A prescription reads: "Take 1 tablet PO q6h PRN pain. Max 8 tablets/day. Qty 80." What is the days' supply using the maximum directions?

- a) 5
- b) 8
- c) 10
- d) 20

20) A tech is preparing a pseudoephedrine sale. Which information is required by federal law for the sales log?

- a) Prescriber NPI and diagnosis
- b) Purchaser ID verification and amount sold
- c) Patient allergy list and DOB
- d) Insurance BIN/PCN and copay

21) A technician sees two patient profiles with the same name and similar DOBs. One has a severe latex allergy. Before entering the prescription, what should the technician do?

- a) Choose the profile with the most history to reduce duplicates
- b) Merge the profiles immediately
- c) Verify additional identifiers to select the correct patient profile
- d) Ignore allergies since latex is not a drug

22) A barcode scan fails because the NDC on the stock bottle doesn't match the NDC tied to the label. The drug name and strength appear identical on screen. What should the technician do?

- a) Bypass scanning and complete the fill
- b) Reprint the label and try again; it will usually pass
- c) Change the NDC in the system to match the bottle and continue
- d) Stop and resolve the NDC mismatch using the correct product/NDC before proceeding

23) A prescription is written for "Augmentin ES-600." The pharmacy only has Augmentin 400/57 per 5 mL. What should the technician do?

- a) Dispense 400/57 and increase the volume to compensate
- b) Dispense amoxicillin alone
- c) Set aside for pharmacist review because ES-600 is a specific formulation and not an automatic substitution
- d) Change the concentration in the system and keep the same directions

24) A patient returns an opened, partially used bottle of doxycycline capsules and requests a refund. What should the technician do with the medication?

- a) Return it to stock if the expiration date is valid
- b) Return it to stock if it was opened within 14 days
- c) Repackage and dispense to a different patient
- d) Do not return to stock; treat as nonreturnable and handle per disposal policy

25) A patient requests an early refill of clonazepam 1 mg. Last fill was 12 days ago for #60 with directions "1 tablet BID." Which next step is most appropriate?

- a) Process it if the patient states it was lost
- b) Deny it because controlled substances can never be early
- c) Send the request for pharmacist review based on controlled status and fill history
- d) Offer an emergency loan of 10 tablets and bill cash

26) Which brand/generic pair is correct?

- a) Lexapro — escitalopram
- b) Cymbalta — venlafaxine
- c) Zoloft — fluoxetine
- d) Prozac — paroxetine

27) A prescriber faxes a prescription for alprazolam that is missing the prescriber signature. What should the technician do?

- a) Fill it if the fax header shows the clinic number
- b) Hold it for pharmacist review because it is not a valid signed prescription
- c) Ask the patient to sign for the prescriber at pickup
- d) Convert it by entering it as a verbal and filling

28) A prescription reads: "Take 2 capsules PO TID. Qty 180." What is the days' supply?

- a) 15
- b) 30
- c) 45
- d) 90

29) A patient asks the technician to transfer an alprazolam prescription to another pharmacy. What should the technician do?

- a) Complete the transfer immediately to avoid delays
- b) Tell the patient controlled substances cannot be transferred
- c) Print the Rx and hand it to the patient to take over
- d) Gather destination information and route to the pharmacist to evaluate eligibility and perform the transfer

30) A medication is labeled "For external use only." Which prescription most appropriately requires this auxiliary label?

- a) Triamcinolone 0.1% cream
- b) Amoxicillin 500 mg capsules
- c) Metformin 500 mg tablets
- d) Warfarin 5 mg tablets

31) A new eRx is received for isotretinoin. The system requires REMS steps before it can be dispensed. What should the technician do?

- a) Override the REMS requirement and proceed
- b) Cancel the prescription and tell the patient to contact the prescriber
- c) Complete technician-allowed data entry steps and route for pharmacist completion of required verification before dispensing
- d) Dispense a reduced quantity to bypass the restriction

32) A patient brings a paper C-II prescription dated today that also says "Do not fill until 05/10." The patient wants it filled on 05/01. What should the technician do?

- a) Fill it because it is dated today
- b) Partially fill now and owe the rest on 05/10
- c) Refuse permanently and return it to the patient
- d) Set the prescription to be filled on/after 05/10 per the instruction

33) A technician enters "Wellbutrin XL 150 mg." Which is the correct generic selection?

- a) Bupropion IR 150 mg
- b) Bupropion SR 150 mg
- c) Bupropion XL 150 mg
- d) Bupropion 150 mg (any formulation)

34) A prescription reads "qod" in the directions. How should this be interpreted?

- a) Every other day
- b) Four times daily
- c) Every morning
- d) At bedtime

35) A tech is asked to calculate: stock solution is 50 mg/mL. A compounded preparation needs 2,500 mg of the drug. How many mL are needed?

- a) 25 mL
- b) 40 mL
- c) 50 mL
- d) 60 mL

36) A patient's insurance rejects omeprazole with "non-preferred; use esomeprazole." What should the technician do?

- a) Substitute esomeprazole automatically and dispense
- b) Submit a prior authorization without involving the pharmacist
- c) Route to the pharmacist for therapeutic interchange decision and any needed prescriber authorization
- d) Tell the patient to buy OTC and close the prescription

37) A prescription is for "Symbicort." Which ingredients are in Symbicort?

- a) Fluticasone/salmeterol
- b) Budesonide/formoterol
- c) Mometasone/formoterol
- d) Fluticasone/vilanterol

38) A prescription reads: "Inject 0.6 mL." The concentration is 150 mg/mL. How many mg is the dose?

- a) 60 mg
- b) 75 mg
- c) 90 mg
- d) 150 mg

39) Which medication is a proton pump inhibitor (PPI)?

- a) Famotidine
- b) Omeprazole
- c) Sucralfate
- d) Dicyclomine

40) An eRx is received for "insulin glargine-yfgn." On the shelf are Lantus and Semglee. What is the best technician action?

- a) Dispense Lantus because all glargine products are interchangeable
- b) Dispense Semglee and label it as Lantus for familiarity
- c) Confirm the exact product selected/available in the system and route to pharmacist if substitution/interchangeability is unclear
- d) Dispense whichever is cheaper and document "same ingredient"

41) A technician drops one tablet onto a counter near a tray previously used for amoxicillin. What should be done with the tablet?

- a) Put it back into the stock bottle
- b) Use it if the patient has no penicillin allergy
- c) Wipe with alcohol and use it
- d) Discard it to prevent contamination risk

42) Which schedule is correct for tramadol under federal law?

- a) C-II
- b) C-III
- c) C-IV
- d) C-V

43) A prescription label prints "take 1 tab daily," but the hardcopy indicates "take 1 tab BID." What should the technician do?

- a) Dispense as daily to reduce side effects
- b) Correct the sig in the system to match the prescription before proceeding
- c) Ask the patient at pickup which they prefer
- d) Leave the label and write "BID" by hand

44) A patient requests their spouse's medication list over the phone and can verify the spouse's DOB, but is not listed as an authorized representative. What should the technician do?

- a) Provide the list since DOB was verified
- b) Provide only the medication names
- c) Decline and request the patient or authorized person provide permission per HIPAA requirements
- d) Fax the list to the address on file

45) A prescription is written for "Crestor 20 mg." Which generic is correct?

- a) Atorvastatin
- b) Rosuvastatin
- c) Pravastatin
- d) Simvastatin

46) A prescription is written for "glipizide 10 mg, take 1 tab TID with meals." The technician selected glipizide ER 10 mg. What is the best next step?

- a) Dispense ER and keep TID directions
- b) Dispense ER and change to daily without review
- c) Change to IR automatically and proceed
- d) Set aside for pharmacist review because IR vs ER selection changes dosing

47) A patient brings a DSCSA return to the pharmacy from a wholesaler tote. The seal is broken and the 2D barcode is not readable. What should the technician do?

- a) Return it to stock if the expiration date is acceptable
- b) Relabel with a new barcode and place into inventory
- c) Quarantine as a suspect product and escalate per procedure
- d) Dispense it first to avoid waste

48) A prescription reads: "Take 1.5 tsp PO q8h." How many mL is each dose?

- a) 5 mL
- b) 7.5 mL
- c) 10 mL
- d) 15 mL

49) A technician is asked to restock an automated dispensing cabinet with concentrated potassium chloride injection. What is the safest practice?

- a) Stock it in any open pocket to finish quickly
- b) Place it next to normal saline since both are clear liquids
- c) Remove warning labels so it is less alarming
- d) Use required segregated storage and double-check pocket selection before completing the restock

50) A prescription is for "nitroglycerin 0.4 mg SL tablets," but the filling tech selected nitroglycerin transdermal patches. When is this most likely caught if barcode workflow is followed?

- a) At will-call filing
- b) At product scan during filling
- c) When the patient signs at pickup
- d) After the first dose is taken

51) A controlled substance is discovered missing during end-of-shift inventory reconciliation. What is the best initial technician action?

- a) Adjust the on-hand count to match the computer
- b) Wait 48 hours to see if it turns up
- c) Document the discrepancy and immediately alert the pharmacist/manager per diversion procedure
- d) Call the DEA directly before notifying anyone internally

52) Which abbreviation means "left ear"?

- a) OS
- b) AS
- c) AD
- d) AU

53) A prescription is for "albuterol HFA 90 mcg/actuation." Which product matches?

- a) Levalbuterol HFA 45 mcg/actuation
- b) Albuterol nebulizer solution 0.083%
- c) Fluticasone HFA inhaler
- d) Albuterol HFA 90 mcg/actuation inhaler

54) A patient requests a partial fill of a C-II e-prescription for methylphenidate due to cost. What should the technician do?

- a) Partial fill and owe the remainder any time within 6 months
- b) Decline; partial fills are never allowed for C-II
- c) Route to pharmacist because partial-fill rules depend on circumstances and timing for remainder
- d) Dispense the full quantity but bill insurance for a smaller amount

55) A prescription reads: "Take 2 puffs QID for 30 days." The inhaler contains 120 actuations. How many inhalers are needed?

- a) 1

- b) 2
- c) 3
- d) 4

56) A technician notices a coworker bypassing barcode scans repeatedly to save time. What is the best technician action?

- a) Ignore it if no errors have been reported
- b) Disable scanning so everyone works the same way
- c) Confront the coworker loudly in front of patients
- d) Report the behavior through the appropriate supervisor/quality channel

57) A new prescription arrives for "losartan 50 mg daily." The profile has lisinopril 20 mg daily active, and the patient says, "My doctor switched me." What should the technician do?

- a) Inactivate lisinopril and fill losartan immediately
- b) Process losartan and flag for pharmacist review of therapy change/duplication
- c) Fill both and tell the patient to take whichever they want
- d) Refuse to fill losartan due to duplication

58) A prescription is written "DAW 1" for brand Adderall XR. What does DAW indicate?

- a) Dispense as written; do not substitute
- b) Dispense any AB-rated generic automatically
- c) Dispense immediate-release in place of XR
- d) Dispense a different strength if insurance prefers it

59) A prescription is for "Keflex 500 mg." The profile shows "penicillin—anaphylaxis." What should the technician do?

- a) Process normally because cephalexin is unrelated to penicillin allergy
- b) Enter the prescription and ensure it is flagged for pharmacist review given the severe allergy history
- c) Change to azithromycin automatically
- d) Delete the penicillin allergy to avoid alerts

60) A prescription states: "Take 1 tab q4–6h PRN pain, not to exceed 6 tabs/day. Qty 90." Using maximum dosing, what is the days' supply?

- a) 12
- b) 15
- c) 18
- d) 22

61) A prescription is written for "Percocet." What is the correct generic combination?

- a) Hydrocodone/acetaminophen
- b) Oxycodone/acetaminophen
- c) Oxycodone/ibuprofen
- d) Codeine/acetaminophen

62) The pharmacy receives a shipment of refrigerated GLP-1 pens. The tote arrives warm and has no temperature indicator. What should the technician do?

- a) Refrigerate immediately and place into active stock

- b) Freeze for 2 hours, then refrigerate
- c) Quarantine and escalate so cold-chain integrity can be assessed before use
- d) Place at room temperature to “acclimate,” then dispense

63) A prescription is for “vancomycin 125 mg capsules.” Only vancomycin 1 g vials for injection are in stock. What should the technician do?

- a) Dispense the injectable vials; it’s the same drug
- b) Change the dosage form to injection and process
- c) Substitute metronidazole because it treats the same condition
- d) Set aside for pharmacist review because dosage forms/products are not equivalent

64) Which medication commonly requires a Medication Guide to be dispensed with each fill?

- a) Isotretinoin
- b) Amoxicillin
- c) Metformin
- d) Hydrochlorothiazide

65) A technician receives a request to disclose whether a prescription is ready to a caregiver who is listed on the patient’s HIPAA authorization. What is appropriate?

- a) Provide nothing; only the patient can receive any information
- b) Provide full medication history and prescriber notes
- c) Provide readiness status and pickup logistics as allowed by the authorization
- d) Provide diagnosis and lab results if present

66) A prescription is written for “EpiPen 0.3 mg.” Which description is correct?

- a) Epinephrine auto-injector 0.3 mg
- b) Norepinephrine auto-injector 0.3 mg
- c) Epinephrine prefilled syringe 3 mg
- d) Albuterol auto-injector 0.3 mg

67) During order entry, an alert appears for methotrexate directions entered as “Take 6 tablets daily.” What should the technician do?

- a) Override the alert because it was e-prescribed
- b) Change “daily” to “weekly” because that is typical
- c) Continue processing and let the patient clarify at pickup
- d) Stop and route for pharmacist review due to high-alert dosing risk

68) Which situation best describes a “near miss”?

- a) A patient experiences hives after taking a dose
- b) A wrong strength is detected and corrected before the medication reaches the patient
- c) A manufacturer issues a nationwide recall
- d) A patient refuses counseling at pickup

69) A prescription is for “warfarin 5 mg daily.” Which drug class is warfarin?

- a) Antiplatelet
- b) Anticoagulant
- c) ACE inhibitor
- d) Beta blocker

70) A technician is asked to accept a voicemail left after hours that contains a new order for a C-II opioid. What is the correct action when the pharmacy opens?

- a) Transcribe it and fill it if caller ID matches the clinic
- b) Fill it as an emergency prescription without further steps
- c) Route it to the pharmacist because C-II prescriptions require proper legal format and emergency rules are strict
- d) Delete the message because controlled substances are never allowed

71) A technician is entering "hydroxyzine 25 mg: take 1 tab TID PRN anxiety." Which auxiliary label is most appropriate?

- a) "Finish all medication"
- b) "May cause drowsiness"
- c) "Take on an empty stomach"
- d) "Do not crush or chew"

72) A technician is asked to interpret "Instill 1 gtt OU BID." A 5 mL bottle is dispensed and drops are 20 gtt/mL. What is the days' supply?

- a) 25 days
- b) 50 days
- c) 75 days
- d) 100 days

73) A prescription is written for "clopidogrel 75 mg daily." Which class is clopidogrel?

- a) Antiplatelet
- b) Anticoagulant
- c) Statin
- d) Beta blocker

74) A technician sees simvastatin active on the profile. A new prescription arrives for clarithromycin. What should the technician do?

- a) Process normally because interactions are not relevant
- b) Tell the patient to stop simvastatin during antibiotics
- c) Change clarithromycin to azithromycin without authorization
- d) Enter and flag for pharmacist review due to significant interaction risk

75) A technician is asked which action best reduces expired medication dispensing risk. What is best practice?

- a) Check expirations only during annual inventory
- b) Rotate stock and check expiration during stocking and dispensing activities
- c) Remove expiration dates from bottles after opening
- d) Dispense the oldest stock even if it expires next week

76) A prescription reads: "azithromycin 200 mg/5 mL; give 12.5 mL day 1, then 6.25 mL daily days 2–5." What total volume should be dispensed?

- a) 25 mL
- b) 30 mL
- c) 37.5 mL

d) 50 mL

77) A technician is asked which action is within technician scope when a patient requests counseling. What is appropriate?

- a) Provide a full side-effect discussion and recommend therapy changes
- b) Interpret lab results to explain monitoring
- c) Refer the patient to the pharmacist for counseling
- d) Recommend stopping the medication if dizziness occurs

78) A prescription is written for "triamcinolone 0.1% cream: apply thin layer BID." Which auxiliary label is most appropriate?

- a) "May cause drowsiness"
- b) "Take with food"
- c) "For external use only"
- d) "Finish all medication"

79) A patient asks the pharmacy to deliver a prescription to a neighbor's address and says the neighbor can sign. What should the technician do?

- a) Change the address and send it out
- b) Deliver to the neighbor if they can confirm the patient's DOB
- c) Remove the label to protect privacy and deliver anyway
- d) Follow delivery verification requirements; do not redirect to an unverified address/recipient

80) A prescription reads: "latanoprost 0.005% 2.5 mL; instill 1 gtt OU qhs." Using 20 gtt/mL, what is the days' supply?

- a) 12.5
- b) 25
- c) 50
- d) 75

81) A technician is selecting a product for "morphine sulfate ER 30 mg," but both ER and IR 30 mg bottles are on the counter. What is the best step to prevent an error?

- a) Choose IR because it is safer
- b) Use either; both are morphine 30 mg
- c) Dispense both and let the patient decide
- d) Confirm the ER formulation matches the label/prescription and verify via barcode/NDC before filling

82) A prescription is written for "furosemide 40 mg daily," but the technician selects torsemide 20 mg. This is most consistent with what error type?

- a) Look-alike/sound-alike selection error
- b) REMS dispensing error
- c) DEA ordering limit error
- d) Therapeutic equivalence substitution

83) A patient wants to purchase pseudoephedrine but refuses to show photo ID. What is the appropriate technician response?

- a) Complete the sale using a stored patient profile

- b) Complete the sale if the patient pays cash
- c) Decline the sale because ID verification is required for the log
- d) Sell only one box without logging it

84) A prescription is for "gabapentin 300 mg TID." The profile shows an active pregabalin prescription. What should the technician do?

- a) Inactivate pregabalin automatically
- b) Change gabapentin to pregabalin because it is stronger
- c) Process both without any review
- d) Enter and flag for pharmacist review due to possible therapeutic duplication

85) Which brand/generic pair is correct?

- a) Lamictal — lamotrigine
- b) Keppra — carbamazepine
- c) Norvasc — losartan
- d) Synthroid — liothyronine

86) A prescription for a non-controlled medication is transferred from another pharmacy. What is the best technician action?

- a) Refuse; transfers must come only from prescribers
- b) Take the transfer information allowed by law/policy and route per workflow for pharmacist verification/acceptance
- c) Accept but omit original refill information to avoid confusion
- d) Require the other pharmacy to resend as an eRx because transfers are illegal

87) A technician is disposing of a returned oral chemotherapy tablet. What is the best action?

- a) Place it in regular trash after removing the label
- b) Flush it to prevent diversion
- c) Return it to stock if the bottle is sealed
- d) Place it into the hazardous drug waste stream per procedure/vendor

88) Which medication is an SSRI antidepressant?

- a) Duloxetine
- b) Amitriptyline
- c) Sertraline
- d) Bupropion

89) A prescription is written for "phentermine 37.5 mg." Under federal law, phentermine is a:

- a) C-II
- b) C-III
- c) C-IV
- d) Non-controlled legend drug

90) A prescription for "prednisone 10 mg tablets" is entered, but the technician selects prednisolone oral solution. What best describes the error?

- a) Therapeutic interchange
- b) Dosage form/drug selection error

- c) Quantity not sufficient error
- d) Billing/administrative error

Answer Key

1) b	31) c	61) b
2) b	32) d	62) c
3) b	33) c	63) d
4) b	34) a	64) a
5) c	35) c	65) c
6) c	36) c	66) a
7) c	37) b	67) d
8) b	38) c	68) b
9) c	39) b	69) b
10) b	40) c	70) c
11) c	41) d	71) b
12) c	42) c	72) b
13) c	43) b	73) a
14) d	44) c	74) d
15) b	45) b	75) b
16) a	46) d	76) b
17) c	47) c	77) c
18) d	48) b	78) c
19) c	49) d	79) d
20) b	50) b	80) b
21) c	51) c	81) d
22) d	52) b	82) a
23) c	53) d	83) c
24) d	54) c	84) d
25) c	55) b	85) a
26) a	56) d	86) b
27) b	57) b	87) d
28) b	58) a	88) c
29) d	59) b	89) c
30) a	60) b	90) b

PTCB Sim Test 33

1) While entering an eRx, you see: "levothyroxine 0.075 mg tab, take 1 tab PO daily." Which strength should you select?

- a) 7.5 mcg
- b) 75 mcg
- c) 0.75 mg
- d) 750 mcg

2) A patient asks to buy a box of pseudoephedrine 30 mg tablets (96 count). What is required under CMEA at the time of sale?

- a) Record purchaser info and obtain a signature in the required log system
- b) Dispense only with a written prescription
- c) Call the prescriber to verify medical necessity
- d) Package must be placed in a child-resistant container

3) A hardcopy prescription for hydrocodone/acetaminophen is missing the prescriber's address. What should you do?

- a) Add the address from a previous prescription and continue
- b) Ask the patient to write the prescriber's address on the prescription
- c) Set it aside for pharmacist review before processing
- d) Process it as long as the DEA number is present

4) A patient asks for a printed copy of their spouse's full medication profile. No authorization is on file. What should you do?

- a) Provide it if the spouse shares the same address
- b) Provide only the last 6 months to limit PHI exposure
- c) Provide it if the requester knows the spouse's DOB
- d) Decline and request appropriate authorization before releasing PHI

5) You are filling hydroxyzine 25 mg, but the bottle in your hand is hydralazine 25 mg. What is the best next step?

- a) Continue and rely on pharmacist final check
- b) Stop, correct the product selection, and rescan/verify the NDC before proceeding
- c) Cross out the label and handwrite the correct medication name
- d) Dispense because the strength matches

6) A new prescription is for metoprolol succinate ER 50 mg with directions "take 1 tablet twice daily." What should you do?

- a) Change to metoprolol tartrate 50 mg and continue
- b) Change the directions to once daily and continue
- c) Enter as written and print the label
- d) Route for pharmacist review/clarification before dispensing

7) A prescription for Augmentin ES-600 reads: "7.5 mL PO BID x 10 days." How many mL should be dispensed?

- a) 150 mL

- b) 75 mL
- c) 100 mL
- d) 200 mL

8) A prescription reads: "timolol 0.5% ophthalmic solution: instill 1 gtt OU BID." What does OU mean?

- a) Left eye
- b) Both eyes
- c) Right eye
- d) Both ears

9) An eRx for lisdexamfetamine (Vyvanse) 30 mg daily arrives with "Refills: 2." What should happen next?

- a) Change refills to 0 and proceed without notifying anyone
- b) Dispense 30 capsules and put remaining refills on hold
- c) Route for pharmacist review because Schedule II prescriptions cannot have refills
- d) Reject the prescription and delete it from the queue

10) A vaccine shipment arrives with a temperature indicator showing an excursion. What is the correct technician action?

- a) Stock it immediately to prevent further warming
- b) Place it in the refrigerator and use first
- c) Leave it at room temperature until the pharmacist is free
- d) Quarantine the shipment and alert the pharmacist/manager for guidance

11) Prednisone taper: "10 mg tabs: 4 tabs daily x 3 days, then 3 tabs daily x 3 days, then 2 tabs daily x 3 days, then 1 tab daily x 3 days." How many tablets total?

- a) 24
- b) 40
- c) 30
- d) 36

12) A patient says their levetiracetam tablets look different this month. What is the best technician response?

- a) "That means it's counterfeit—don't take it."
- b) "Different manufacturers can change appearance; the pharmacist can discuss any concerns with you."
- c) "Generics work differently; expect different effects."
- d) "Take two tablets today to make up for the change."

13) A Medication Guide is required for a prescription at will-call. What must occur at dispensing?

- a) Provide the Medication Guide to the patient with the prescription
- b) Provide it only if the patient requests it
- c) Place it in the patient profile and do not print
- d) Provide it only for controlled substances

14) Diclofenac 1% gel: "Apply 4 g to knee QID." You dispense a 100 g tube. What days'

supply should be billed?

- a) 6 days
- b) 12 days
- c) 25 days
- d) 3 days

15) A new prescription for cephalexin is entered; the profile lists "penicillin—hives." What is the best next step?

- a) Continue because cephalexin is not penicillin
- b) Remove the allergy because it's unrelated
- c) Route for pharmacist evaluation before dispensing
- d) Change cephalexin to azithromycin and process

16) DSCSA sends a notice to investigate a suspect product for a specific NDC/lot of carvedilol. Matching stock is on your shelf. What should you do?

- a) Keep dispensing until the wholesaler confirms
- b) Quarantine the matching stock and alert the pharmacist/manager to follow DSCSA steps
- c) Pull all beta-blockers from the shelf
- d) Only remove expired bottles

17) A prescription reads: "Apply 2 g topically to affected area QID. Dispense 120 g." What is the days' supply?

- a) 15 days
- b) 10 days
- c) 30 days
- d) 12 days

18) A patient requests insulin syringes. In your state, syringes require proof of a current prescription for sale. What should you do?

- a) Sell them because syringes are OTC everywhere
- b) Ask for proof of a current prescription/documented authorization before selling
- c) Suggest reusing old syringes
- d) Use a family member's prescription on file to complete the sale

19) At pickup, a patient requests counseling. The pharmacist is on a phone call. What should you do?

- a) Provide counseling yourself to avoid delays
- b) Complete the transaction; counseling can occur after the sale
- c) Pause the sale and ensure the pharmacist provides counseling before the medication is handed over
- d) Tell the patient counseling isn't available today

20) A customer asks for an OTC recommendation for sudden shortness of breath and chest tightness. What is the best action?

- a) Recommend an OTC bronchodilator and send them home
- b) Ask what flavor they prefer in a cough syrup
- c) Suggest ibuprofen and rest
- d) Involve the pharmacist promptly and advise urgent medical evaluation

21) Gabapentin 300 mg: "Take 1 capsule TID. Qty 270." What is the days' supply?

- a) 60
- b) 30
- c) 90
- d) 270

22) Albuterol HFA: "Inhale 2 puffs q4h PRN." Inhaler contains 200 puffs. If used at the maximum frequency, what days' supply should be billed?

- a) 8 days
- b) 10 days
- c) 16 days
- d) 25 days

23) Insulin aspart pens: "Inject 12 units SQ TID with meals. Dispense 1 box (5 pens)." Each pen contains 300 units. What is the days' supply?

- a) 35 days
- b) 25 days
- c) 30 days
- d) 42 days

24) Latanoprost 0.005%: "Instill 1 gtt OU qHS." Bottle size 2.5 mL. Use 20 gtt = 1 mL. What is the days' supply?

- a) 12 days
- b) 50 days
- c) 25 days
- d) 100 days

25) Nystatin oral suspension: "Swish and swallow 5 mL QID x 14 days." How many mL should be dispensed?

- a) 200 mL
- b) 240 mL
- c) 140 mL
- d) 280 mL

26) A standard azithromycin "Z-Pak" (250 mg) is typically dispensed as which quantity?

- a) 5 tablets
- b) 10 tablets
- c) 6 tablets
- d) 12 tablets

27) Augmentin 875/125 mg: "Take 1 tab PO BID x 10 days." What quantity should be dispensed?

- a) 14
- b) 28
- c) 10
- d) 20

28) A patient's label says "Take 1/2 tab PO BID." The tablet strength is 20 mg. How many mg per dose?

- a) 20 mg
- b) 10 mg
- c) 5 mg
- d) 40 mg

29) Morphine oral solution 20 mg/mL: "Take 0.5 mL PO q4h PRN pain." How many mg per dose?

- a) 5 mg
- b) 20 mg
- c) 10 mg
- d) 40 mg

30) A prescription states: "Take 2 tablets now, then 1 tablet in 8 hours." Tablets are 250 mg. What total dose will be taken?

- a) 750 mg
- b) 500 mg
- c) 1,000 mg
- d) 1,250 mg

31) Breo Ellipta 100/25: eRx includes "inhale 1 puff once daily" and "disp: 1 inhaler (30 doses)." What days' supply should be submitted?

- a) 15
- b) 60
- c) 30
- d) 45

32) Fluticasone nasal spray: "2 sprays in each nostril daily." Bottle contains 120 sprays. What days' supply?

- a) 60
- b) 30
- c) 45
- d) 15

33) While restocking an ADC, you see potassium chloride injection concentrate. Why does it require extra caution?

- a) It is a hazardous antineoplastic
- b) It is a high-alert medication with significant harm potential if misused
- c) It is a vaccine requiring freezing
- d) It is a LASA antibiotic

34) A compounded oral liquid is returned from a unit without a patient label and with an unclear beyond-use date. What should be done?

- a) Return it to inventory if it looks intact
- b) Relabel it with today's date and restock
- c) Dispense it first to reduce waste
- d) Quarantine for pharmacist review because identity/integrity cannot be verified

35) You are preparing a compounded suspension and realize the ingredient lot number was not recorded. Product has not been dispensed. What should you do?

- a) Estimate lot number from the invoice date
- b) Stop and record the actual lot number from the ingredient container before continuing
- c) Dispense it; lot numbers are only required for controls
- d) Discard the preparation every time this occurs

36) During unit-dose packaging, the barcode scan fails on a look-alike vial. What is the best next step?

- a) Bypass scanning because the label is readable
- b) Manually type any NDC matching the strength
- c) Stop and obtain a scannable product or follow the required secondary verification process
- d) Move a barcode label from a different vial onto this one

37) A patient's label prints with an old address and phone number. The prescription is bagged but not sold. What should you do?

- a) Dispense anyway; demographics do not affect safety
- b) Correct the demographics and reprint required labels/bag tags before dispensing
- c) Cross out the address and proceed
- d) Update the profile after pickup

38) A customer describes sudden facial droop and slurred speech while asking for an OTC item. What should you do?

- a) Recommend aspirin and have them rest
- b) Suggest an antihistamine for swelling
- c) Offer a blood pressure check later
- d) Involve the pharmacist immediately and advise emergency care

39) A pharmacy may keep controlled-substance prescriptions using which federally permitted filing method?

- a) Three separate files: C-II, C-III–V, and noncontrols
- b) Controls must always be filed with noncontrols in a single file
- c) C-II must be mixed with noncontrols; C-III–V must be separate
- d) C-II hardcopies cannot be filed at all

40) A supplier is ordering Schedule II medications from a wholesaler. Which DEA form is used?

- a) DEA Form 106
- b) DEA Form 41
- c) DEA Form 222 (or electronic equivalent)
- d) DEA Form 363

41) Which is required information on a completed DEA Form 222 record?

- a) Patient date of birth
- b) Diagnosis code
- c) Prescriber NPI
- d) Supplier name and quantity ordered for each item

42) A recall notice lists a specific lot number of valsartan. That lot is on your shelf. What should you do?

- a) Keep dispensing until the wholesaler pickup arrives
- b) Remove and quarantine the affected lot per recall instructions
- c) Return all valsartan regardless of lot number
- d) Relabel the lot with a new expiration date

43) A patient is picking up a Schedule II prescription and the electronic signature pad is down. What is the best next step?

- a) Dispense without a signature because payment is recorded
- b) Tell the patient controls cannot be dispensed today
- c) Use an approved alternate signature/log method per workflow and complete required documentation
- d) Ask the patient to text a photo of their signature

44) A prescriber's office calls and asks to "change Adderall XR 20 mg to 30 mg." What should you do as a technician?

- a) Update the dose if the caller provides the prescriber NPI
- b) Change it and document "per office"
- c) Change it only if the patient agrees
- d) Route to the pharmacist because controlled-substance changes require proper authorization/handling

45) A Schedule III prescription was written 7 months ago and has refills remaining. Under federal law, can it be refilled?

- a) Yes, for 1 year if refills remain
- b) Yes, if the prescriber allows it verbally
- c) No, it is limited to 6 months from the date written
- d) No, Schedule III cannot be refilled

46) A pharmacy is disposing of expired controlled substances through a reverse distributor. Which DEA form is commonly used to document destruction of controlled substances?

- a) DEA Form 41
- b) DEA Form 222
- c) DEA Form 106
- d) DEA Form 363

47) An eRx for isotretinoin is received and the patient asks why it takes longer than usual. What is the best technician response?

- a) It is a Schedule II medication and requires extra ID steps
- b) It is limited to hospital-only dispensing
- c) It requires REMS steps that must be completed before dispensing
- d) It cannot be dispensed to patients of childbearing potential

48) Which storage statement is correct?

- a) Shingrix is stored frozen
- b) Shingrix is stored refrigerated and protected from freezing

- c) Zostavax is stored at room temperature
- d) Shingrix can be stored at room temperature for 30 days after delivery

49) A patient asks the pharmacy to fax their complete medication profile to a new physician. No authorization is on file. What should you do?

- a) Fax it because it is for treatment and always allowed without documentation
- b) Fax only the controlled-substance list
- c) Tell the patient to photograph the screen and send it
- d) Obtain appropriate authorization/documentation per workflow before sending

50) At the register, another customer is close enough to hear. A parent is authorized to pick up their child's prescription. What should you do?

- a) Speak louder so the parent hears clearly
- b) Move to a more private area or lower your voice to reduce PHI exposure
- c) Provide the child's full medication history since the parent is authorized
- d) Refuse to release prescriptions to caregivers

51) A claim is rejected "refill too soon" for a noncontrolled medication; the patient is traveling. What is an appropriate technician step?

- a) Submit a vacation override if permitted by the plan and document appropriately
- b) Change the previous fill date to make it go through
- c) Reduce quantity to a 7-day supply automatically
- d) Tell the patient early refills are illegal

52) A patient brings back an unopened bottle of lisinopril dispensed yesterday and asks for a refund. What is the best action?

- a) Return it to stock if the seal appears intact
- b) Re-dispense it to another patient if packaging is unopened
- c) Decline restocking and route to pharmacist/manager for return/refund handling per workflow
- d) Accept only if the patient signs a waiver

53) You are processing "Toprol-XL 100 mg daily" and need the generic. What is the correct selection?

- a) Metoprolol succinate ER
- b) Metoprolol tartrate
- c) Propranolol ER
- d) Carvedilol

54) A prescription is for bupropion XL 150 mg daily. Which substitution is most likely to cause a problem if dispensed without clarification?

- a) Bupropion XL 150 mg from a different manufacturer
- b) Bupropion SR 150 mg
- c) Bupropion XL 150 mg with a different tablet color
- d) Bupropion XL 150 mg with a different bottle size

55) The prescription is for immediate-release glipizide 10 mg, but the label prints "glipizide ER 10 mg." What should you do?

- a) Dispense ER because it lasts longer
- b) Dispense IR but keep ER directions
- c) Stop, correct the dosage form, and reprocess for pharmacist verification
- d) Split the fill between IR and ER

56) A prescription is written for "Klonopin 1 mg." Which generic should be selected?

- a) Clonidine
- b) Cyclobenzaprine
- c) Clonazepam
- d) Clopidogrel

57) "Plavix 75 mg daily" is prescribed. What is the correct generic?

- a) Dipyridamole
- b) Prasugrel
- c) Clopidogrel
- d) Cilostazol

58) The profile lists phenelzine. A new prescription for sertraline is received. What should you do?

- a) Process it; counseling can occur at pickup
- b) Route for pharmacist review due to a potentially serious interaction
- c) Change sertraline to venlafaxine and proceed
- d) Remove phenelzine since it's older than 30 days

59) The profile shows losartan active. A new prescription arrives for lisinopril. What should you do?

- a) Process it because both treat blood pressure
- b) Delete losartan because lisinopril is newer
- c) Change lisinopril to amlodipine
- d) Route for pharmacist review due to possible duplication/therapy change concerns

60) A patient's active meds include atorvastatin 40 mg daily. A new prescription arrives for simvastatin 40 mg daily. What should you do?

- a) Process both; two statins are common
- b) Route for pharmacist review for therapeutic duplication
- c) Discontinue atorvastatin automatically
- d) Change simvastatin to rosuvastatin and proceed

61) A prescription for "Eliquis 5 mg BID" is dropped off. A profile note states "mechanical heart valve." What should you do?

- a) Process normally; diagnosis notes are not used in dispensing
- b) Substitute warfarin automatically
- c) Route for pharmacist review because therapy may be inappropriate for that condition
- d) Tell the patient to stop anticoagulants

62) Methotrexate 2.5 mg for rheumatoid arthritis: "Take 6 tablets weekly." What is the most critical label detail to reduce serious error?

- a) "Take with food"

- b) "May cause drowsiness"
- c) "Refrigerate after opening"
- d) Clear weekly dosing with a warning such as "Do not take daily"

63) A prescription is for amphetamine-dextroamphetamine ER 10 mg capsules, but the product pulled is IR 10 mg tablets. What should you do?

- a) Dispense IR because mg strength matches
- b) Dispense IR and change directions to BID
- c) Stop and select the correct dosage form; route for pharmacist help if substitution is needed
- d) Substitute methylphenidate ER 10 mg

64) A prescription for vancomycin 125 mg PO QID x 10 days is entered, but the system default selects IV vancomycin vials. What should you do?

- a) Dispense IV vials and instruct patient to drink it
- b) Change directions to BID to match typical antibiotic dosing
- c) Ensure an appropriate oral product/formulation is selected; route if compounding is needed
- d) Substitute metronidazole automatically

65) A patient asks if they can crush oxybutynin ER. What should you do?

- a) Tell them yes so it works faster
- b) Tell them to split it in half instead
- c) Recommend they stop taking it
- d) Refer to the pharmacist for counseling on dosage form modification

66) A DUR alert appears: "Duplicate therapy—two opioids." The profile shows an active hydrocodone/APAP prescription, and a new oxycodone prescription is ready. What should you do?

- a) Override the alert because both were prescribed
- b) Ask the patient which one they prefer and dispense only that
- c) Hold and route to the pharmacist for DUR review before dispensing
- d) Dispense both and tell the patient to alternate doses

67) While receiving stock, you find bottles with altered/covered lot numbers and the invoice count does not match the delivered quantity. What should you do?

- a) Stock them; lot numbers are rarely used
- b) Quarantine as suspect product and alert the pharmacist/manager
- c) Remove the covering and place on shelf if tablets look normal
- d) Assign an internal lot number and dispense as usual

68) A prescription is for pantoprazole 40 mg daily, but insurance prefers omeprazole. What can you do as a technician?

- a) Substitute omeprazole because they're both PPIs
- b) Bill pantoprazole as omeprazole to get paid
- c) Route to the pharmacist because changing the prescribed drug requires pharmacist/prescriber involvement
- d) Change to omeprazole only if the strengths match

69) A patient requests "my water pill" and mentions a "sulfa allergy." The ready prescription is furosemide 20 mg. What is the best next step?

- a) Cancel and refuse to dispense
- b) Ask the pharmacist to speak with the patient before completing the sale
- c) Tell the patient it's safe because it's not an antibiotic
- d) Recommend switching to hydrochlorothiazide at pickup

70) An eRx says verapamil ER 240 mg, and your pharmacy stocks multiple ER formulations. The prescriber selected "SR" in the dropdown. What should you do?

- a) Select any verapamil ER 240 mg because ER products are interchangeable
- b) Choose the cheapest NDC and proceed
- c) Substitute immediate-release verapamil 80 mg TID
- d) Route for pharmacist confirmation of the intended formulation before dispensing

71) Nitroglycerin SL 0.4 mg directions include "max 3 doses." The patient asks what to do if pain persists after 3 tablets. What should you do?

- a) Tell them to take 3 more tablets
- b) Tell them to chew aspirin and lie down
- c) Refer to the pharmacist for counseling
- d) Tell them to stop nitroglycerin permanently

72) Doxycycline 100 mg is being picked up. The patient asks about taking it with calcium supplements. What should you do?

- a) Tell them it doesn't matter
- b) Refer to the pharmacist for counseling on administration and interactions
- c) Tell them to stop calcium permanently
- d) Tell them to take both together with milk to prevent nausea

73) A new prescription is received for metformin 500 mg BID. The profile shows a recently documented eGFR of 22 mL/min/1.73 m². What should you do?

- a) Fill immediately; dosing is prescriber responsibility
- b) Change directions to once daily and proceed
- c) Enter and route for pharmacist clinical review before dispensing
- d) Place on hold until patient asks

74) A prescriber's office asks you to change a noncontrolled prescription to a different drug in the same class due to cost. What should you do?

- a) Make the interchange to save the patient money
- b) Route to the pharmacist because therapy changes require pharmacist involvement and often prescriber authorization
- c) Ask the patient to choose and then change it
- d) Change it and document "therapeutic interchange"

75) A controlled-substance inventory is being performed. Under federal requirements, when must a pharmacy take an initial controlled-substance inventory?

- a) Every 30 days
- b) Only when DEA requests it

- c) Only when the pharmacist-in-charge changes
- d) On the date the pharmacy first engages in controlled-substance activities

76) A pharmacy reports a significant theft of controlled substances. Which DEA form is used to report it?

- a) DEA Form 106
- b) DEA Form 41
- c) DEA Form 222
- d) DEA Form 363

77) A prescriber calls in a verbal prescription for a Schedule II medication to a retail pharmacy for routine pain control. Under federal law, what is generally required for dispensing?

- a) The verbal order is sufficient if documented
- b) A written/electronic Schedule II prescription is generally required (except true emergency rules)
- c) The patient can sign a waiver in place of the hardcopy
- d) The pharmacy can dispense if quantity is under 7 days

78) A patient wants to return an opened bottle of tablets for credit. What is the safest correct outcome in most community pharmacy workflows?

- a) Return to stock if more than half remains
- b) Accept and re-dispense if the lot number matches
- c) Do not return to stock; handle refund/return per workflow without re-dispensing
- d) Pour into a counting tray to verify quantity and restock

79) During order entry, you notice two patient profiles with the same name and DOB, but different addresses. A new eRx arrives. What should you do first?

- a) Pick the profile with the most recent fill and proceed
- b) Merge the profiles yourself immediately
- c) Confirm identifiers (address/phone) and ensure the prescription is placed on the correct profile before processing
- d) Delete one profile to prevent duplication

80) A prescriber sends an eRx with directions "UAD" for an ointment. What is the best action for accurate processing and billing?

- a) Enter "use as directed" and bill a standard 30-day supply
- b) Hold until the patient picks up and then ask how they use it
- c) Enter "as directed" and select any days' supply; it will be corrected at pickup
- d) Obtain more specific directions/quantity-use information through appropriate workflow before billing

81) A prescription is written for "Lantus Solostar: inject 18 units SQ nightly. Dispense 1 box (5 pens)." Each pen contains 3 mL of U-100 insulin. What days' supply is 1 box?

- a) 83 days
- b) 28 days
- c) 50 days
- d) 16 days

82) A prescription is for warfarin 2 mg daily. The pharmacy has 1 mg and 5 mg tablets in stock but no 2 mg. What is the correct technician action?

- a) Substitute two 1 mg tablets without contacting anyone
- b) Route to the pharmacist to determine if a change/new prescription is needed before dispensing
- c) Dispense 5 mg and instruct patient to take half
- d) Dispense 1 mg and change directions to "take 2 tablets daily"

83) A prescription is for "amoxicillin 400 mg/5 mL: take 7.5 mL PO BID x 10 days." How many mL should be dispensed?

- a) 75 mL
- b) 100 mL
- c) 150 mL
- d) 200 mL

84) A patient is prescribed "clindamycin 150 mg: take 2 capsules PO QID x 7 days." How many capsules should be dispensed?

- a) 28
- b) 56
- c) 42
- d) 84

85) A prescription is for "apixaban (Eliquis) 5 mg: take 1 tablet PO BID. Qty 60." What is the days' supply?

- a) 60
- b) 15
- c) 45
- d) 30

86) A patient's third-party claim rejects "refill too soon" for a controlled medication; the patient reports it was stolen. What is the best technician next step?

- a) Change the days' supply on the claim until it pays
- b) Apply a lost medication override without documentation
- c) Tell the patient to pay cash to avoid delays
- d) Route to the pharmacist for evaluation and next steps per requirements (documentation/plan rules/prescriber contact)

87) You receive an eRx with "UD" in the directions field only. How should this be interpreted for order entry?

- a) As directed
- b) Unit dose
- c) Use diluted
- d) Once daily

88) While typing a prescription for insulin, the prescriber wrote "Humalog U-200 KwikPen." Your shelf has Humalog U-100 pens. What is the correct action?

- a) Substitute U-100; the patient can inject double the volume

- b) Substitute U-100 because it's the same insulin
- c) Hold and route for pharmacist review because concentration/product selection must match exactly
- d) Change to Novolog U-100 because it's interchangeable

89) A patient asks whether they should avoid spinach while taking warfarin. What should you do?

- a) Tell them to stop all leafy greens
- b) Tell them spinach has no interaction with warfarin
- c) Refer to the pharmacist for counseling
- d) Recommend switching to apixaban without contacting anyone

90) At pickup for sertraline 50 mg daily, the patient mentions taking St. John's wort every day. What should you do?

- a) Tell them to stop sertraline immediately
- b) Tell them it's safe because it's herbal
- c) Recommend doubling sertraline
- d) Refer to the pharmacist for interaction counseling

Answer Key

1) b	31) c	61) c
2) a	32) b	62) d
3) c	33) b	63) c
4) d	34) d	64) b
5) b	35) b	65) d
6) d	36) c	66) c
7) a	37) b	67) b
8) b	38) d	68) c
9) c	39) a	69) b
10) d	40) c	70) d
11) c	41) d	71) c
12) b	42) b	72) b
13) a	43) c	73) c
14) b	44) d	74) b
15) c	45) c	75) d
16) b	46) a	76) a
17) a	47) c	77) b
18) b	48) b	78) c
19) c	49) d	79) c
20) d	50) b	80) d
21) c	51) a	81) a
22) a	52) c	82) b
23) a	53) a	83) c
24) b	54) b	84) b
25) b	55) c	85) d
26) c	56) c	86) d
27) d	57) c	87) a
28) b	58) b	88) c
29) c	59) d	89) c
30) a	60) b	90) d

PTCB Sim Test 34

1. An eRx comes in for “Xarelto 20 mg: Take 1 tablet by mouth daily with evening meal.” In data entry, the technician accidentally selects Eliquis 5 mg because the patient previously used it. What is the best technician action before it goes to verification?

- a) Keep Eliquis and add a note that both are anticoagulants
- b) Correct the drug selection to rivaroxaban 20 mg and recheck the strength and formulation
- c) Change directions to match Eliquis dosing and proceed
- d) Cancel the prescription and ask the patient to request a new one

2. A written prescription states: “Amoxicillin 500 mg: Take 1 capsule by mouth three times daily for 10 days.” What quantity should be dispensed?

- a) 30
- b) 20
- c) 10
- d) 60

3. A patient asks to buy pseudoephedrine. The technician completes the electronic log but does not verify the purchaser’s photo ID. What requirement was missed?

- a) Confirming the patient has insurance on file
- b) Recording the prescriber’s DEA number
- c) Verifying photo identification and purchaser eligibility under limits
- d) Packaging the product in child-resistant packaging

4. A prescription is entered as “metoprolol succinate ER 50 mg daily,” but the product pulled is metoprolol tartrate 50 mg tablets. What is the correct next step for the technician?

- a) Replace with metoprolol succinate ER 50 mg before continuing
- b) Change the sig to BID so tartrate matches typical use
- c) Dispense tartrate and let the patient know it’s similar
- d) Split the tablets to reduce risk and proceed

5. An eRx for “TMP-SMX DS: 1 tablet by mouth twice daily for 5 days” arrives. The patient profile shows “sulfa—hives.” What should the technician do?

- a) Proceed because hives is not a true allergy
- b) Change to nitrofurantoin per UTI protocols
- c) Enter the prescription and bypass interaction screens
- d) Route the order for pharmacist review before dispensing

6. Insulin glargine U-100 pens: “Inject 18 units subcutaneously nightly.” Box contains five 3 mL pens. Using 100 units/mL, what days’ supply should be billed?

- a) 30
- b) 83
- c) 45
- d) 75

7. A refrigerated vaccine shipment arrives with a temperature indicator showing out-of-range exposure. What should the technician do with the shipment?

- a) Place it in the freezer to stabilize it
- b) Stock it in the fridge and mark "use first"
- c) Quarantine it and escalate through the cold-chain process for disposition guidance
- d) Dispense it only to cash-paying patients to avoid rejects

8. A customer requests a printed medication list for their spouse who is not present and not on the phone. No authorization is documented. What is the appropriate technician response?

- a) Provide the list because spouses are automatically authorized
- b) Provide only non-controlled medications
- c) Decline and offer to obtain patient authorization or speak directly with the patient
- d) Print it and place it in a sealed envelope labeled "confidential"

9. An eRx for Adderall XR 20 mg #30 is on file at your pharmacy. The patient asks to transfer it to another pharmacy. What is the correct handling?

- a) Transfers are permitted for Schedule II if both pharmacies agree
- b) A technician may transfer it if the patient requests it verbally
- c) Copy the prescription details and send them to the other pharmacy
- d) Refer for pharmacist handling; Schedule II transfers are generally not permitted under federal rules, and state rules may vary

10. A new eRx is for levothyroxine 75 mcg daily. The patient filled levothyroxine 88 mcg last month from the same clinic. What is the best technician action?

- a) Enter and fill 75 mcg without concern; thyroid doses change often
- b) Inactivate the 88 mcg prescription automatically
- c) Ensure the pharmacist reviews for possible dose change/duplicate therapy before dispensing
- d) Tell the patient to alternate strengths until the next appointment

11. A manufacturer recall notice lists one specific lot of amlodipine 10 mg. What should the technician do first?

- a) Pull all amlodipine strengths from the shelf
- b) Quarantine the affected lot(s) and start recall documentation and inventory reconciliation
- c) Relabel the product with a new beyond-use date
- d) Continue dispensing until the replacement shipment arrives

12. A label prints for "hydroxyzine 25 mg," but the technician accidentally pulled "hydralazine 25 mg." Which step best prevents this error during filling?

- a) Compare tablet shape and color only
- b) Rely on shelf position and bin label
- c) Scan the stock bottle barcode and match drug/strength/dosage form to the label
- d) Choose the bottle with the closest NDC number

13. A Schedule II purchase invoice is filed with routine wholesaler invoices without being separated or marked. What requirement is not being met?

- a) Schedule II records must be retained for 6 months only
- b) Schedule II records must be kept separately or be readily retrievable
- c) Schedule II invoices must be sent to the DEA weekly

d) Schedule II invoices must be stored in the will-call area for audit access

14. A claim rejects "Refill too soon." The patient says they are leaving town tomorrow and want the full quantity now. What should the technician do first?

- a) Use a vacation override code automatically to force coverage
- b) Ask the patient to pay cash and rebill later without documentation
- c) Reduce quantity to a 7-day supply to bypass the reject
- d) Collect travel dates and needed supply details and route through the appropriate override/authorization workflow

15. A pediatric prescription reads: "Amoxicillin 400 mg/5 mL: Give 6 mL by mouth twice daily for 10 days." How many mL should be dispensed?

- a) 60 mL
- b) 120 mL
- c) 100 mL
- d) 150 mL

16. A patient requests counseling on new nitroglycerin sublingual tablets at pickup. What is the technician's best response?

- a) Provide full counseling to reduce the pharmacist's workload
- b) Tell the patient to read the package insert at home
- c) Offer pharmacist counseling and pause the sale until counseling is offered
- d) Advise taking one tablet daily to prevent chest pain

17. The prescriber indicates "DAW 1" on a brand-name prescription. What does the technician do during processing?

- a) Select the generic equivalent to reduce cost
- b) Dispense the brand/product specified and prevent substitution
- c) Substitute any therapeutically similar drug in the same class
- d) Dispense whichever NDC is currently on sale

18. Prednisone taper: "Take 4 tablets daily x 3 days, then 3 tablets daily x 3 days, then 2 tablets daily x 3 days, then 1 tablet daily x 3 days." Tablets are 10 mg. What quantity is needed?

- a) 24
- b) 30
- c) 36
- d) 40

19. A new prescription for cephalexin is entered. The patient profile lists "penicillin— anaphylaxis." What should the technician do?

- a) Fill because cephalexin is not a penicillin
- b) Document "OK" because the prescriber ordered it
- c) Route to the pharmacist for allergy assessment before dispensing
- d) Suggest the patient try one dose and watch for symptoms

20. A patient presents a paper prescription for oxycodone IR. The prescriber signature is present, but the prescriber address is missing. What is the appropriate technician action?

- a) Add the address from an older prescription and process
- b) Enter and dispense; address is optional if DEA number is present
- c) Return it to the patient and refuse to discuss further
- d) Route for pharmacist review to resolve missing required elements prior to dispensing

21. A nasal spray is billed for days' supply. Directions: "2 sprays in each nostril once daily." Bottle contains 120 sprays. What days' supply should be submitted?

- a) 15
- b) 30
- c) 60
- d) 120

22. While counting tablets, the technician finds two different imprints mixed in a stock bottle labeled "losartan 50 mg." What should the technician do?

- a) Use only tablets that match the most common imprint
- b) Separate them by imprint and keep using the bottle
- c) Quarantine the bottle and remove it from dispensing stock for investigation
- d) Discard the tablets in the regular trash immediately

23. An eRx is written for "Crestor 10 mg daily." Which product should be selected?

- a) Rosuvastatin 10 mg
- b) Atorvastatin 10 mg
- c) Pravastatin 10 mg
- d) Simvastatin 10 mg

24. Law enforcement requests a patient's prescription records without a subpoena, court order, or written patient authorization. What should the technician do?

- a) Provide only controlled substance records
- b) Provide records immediately to avoid obstruction
- c) Refuse and shred the request
- d) Refer to the pharmacist/manager for proper legal review and HIPAA handling

25. A prescription is written for "Flonase: 2 sprays each nostril once daily." Which generic is correct?

- a) Mometasone
- b) Fluticasone
- c) Azelastine
- d) Ipratropium

26. A cephalexin suspension label indicates "Discard 14 days after reconstitution." Which auxiliary label is most appropriate?

- a) "Shake well"
- b) "Avoid sunlight"
- c) "Do not take with dairy"
- d) "For external use only"

27. A Schedule III prescription was written 7 months ago and has refills remaining. The patient requests a refill today. What is the correct response under federal rules?

- a) Refill because refills remain on the label
- b) Refill once, then require a new prescription next month
- c) Do not refill; it is beyond the 6-month refill limit
- d) Convert to Schedule IV and refill

28. A new prescription for warfarin is received. The profile shows apixaban was filled 5 days ago by a different prescriber. What should the technician do?

- a) Fill warfarin and discontinue apixaban in the profile
- b) Route to the pharmacist due to potential duplicate anticoagulation/therapy change
- c) Reduce warfarin strength during data entry to be safer
- d) Tell the patient to stop apixaban immediately

29. Azithromycin directions: "Take 2 tablets on day 1, then 1 tablet daily on days 2–5." Tablets are 250 mg. What quantity is needed?

- a) 5
- b) 6
- c) 7
- d) 10

30. A prescriber's office calls in a verbal prescription for clonazepam. What should the technician do?

- a) Take the verbal and enter it under the pharmacist's credentials
- b) Tell the caller verbal prescriptions are never allowed
- c) Transfer the call to the pharmacist to receive the verbal order
- d) Take the verbal if it's under a 30-day supply

31. A patient asks why their metronidazole label says "take with food." What is the technician's best response?

- a) Explain the mechanism and side effect profile in detail
- b) Remove the auxiliary label to avoid confusion
- c) Tell them it is optional and doesn't matter
- d) Offer pharmacist counseling for the question

32. The sig reads: "Instill 1 gtt OU BID." What does "OU" mean?

- a) Both ears
- b) Right eye
- c) Both eyes
- d) Left eye

33. Eye drops: "1 drop in left eye QID." Assume 20 drops/mL. A 5 mL bottle is dispensed. What days' supply should be billed?

- a) 10
- b) 20
- c) 25
- d) 50

34. A patient returns an unopened prescription bottle that left the pharmacy yesterday and requests it be restocked. What is correct in most community pharmacy settings?

- a) Restock if the seal is intact and the bottle is unopened
- b) Restock only nonrefrigerated medications
- c) Restock if it is not a controlled substance
- d) Do not restock medication that has left the pharmacy; handle refund/disposal per policy

35. During DSCSA receiving, a sealed case appears re-taped with a broken manufacturer seal. What should the technician do?

- a) Accept it if the NDC matches the invoice
- b) Stock it in a separate section and use last
- c) Quarantine as suspect product and escalate for investigation
- d) Remove outer packaging and stock the inner bottles

36. The prescription is for "Zoloft 50 mg daily." Which generic is correct?

- a) Fluoxetine
- b) Sertraline
- c) Citalopram
- d) Paroxetine

37. A patient profile lists a latex allergy. A vial product has a latex-containing stopper per manufacturer information. What is the best technician action?

- a) Proceed; stopper latex is only an issue for injections
- b) Remove the stopper and dispense the vial
- c) Add a warning sticker and continue
- d) Route to the pharmacist for assessment and product selection guidance

38. A transfer-in is being documented for a noncontrolled prescription. Which additional item is essential to document for a valid transfer?

- a) The patient's email address
- b) The transferring and receiving pharmacist/authorized individual identification (name/initials)
- c) The prescriber's social security number
- d) The wholesaler invoice number for the product

39. "Keflex 500 mg PO QID x 7 days" is entered. Keflex is:

- a) Cefdinir
- b) Ceftriaxone
- c) Cephalixin
- d) Cefazolin

40. The hard copy says "Take 1 tablet PO qAM," and the label prints "Take 1 tablet by mouth daily" for furosemide. What is being checked?

- a) Controlled substance schedule accuracy
- b) Sig translation/entry consistency
- c) NPI validity
- d) REMS enrollment requirement

41. Diclofenac 1% gel: "Apply 2 g to affected area QID." Tube contains 100 g. What days' supply should be billed?

- a) 12
- b) 25
- c) 10
- d) 5

42. At pickup, the customer's name matches the will-call bag, but the DOB does not. What should the technician do?

- a) Complete the sale because the last name matches
- b) Ask which medications they take and match the bag
- c) Stop the sale and locate the correct prescription using two identifiers
- d) Remove DOB from the receipt and proceed

43. A prescription is for "Wellbutrin XL 150 mg daily." The technician pulled bupropion SR 150 mg because it's available. What is the correct action?

- a) Dispense SR; same ingredient and strength
- b) Change directions to once daily to match XL
- c) Split SR tablets and take once daily
- d) Replace with the correct XL formulation before proceeding

44. During controlled inventory, which federal statement is correct?

- a) Schedule II inventory may be estimated even if containers are opened
- b) Schedule II inventory records must be kept separate from other schedules
- c) Schedule II counts are required only every 3 years
- d) Schedule II records can be discarded after 1 year

45. The directions say "Take 1.5 tablets daily." Each tablet is 8 mg. How many mg per day is this dose?

- a) 10 mg
- b) 12 mg
- c) 16 mg
- d) 24 mg

46. A patient requests an early refill for hydrocodone/acetaminophen stating the medication was stolen. What should the technician do?

- a) Process the refill and document "stolen" in notes
- b) Tell the patient it is always illegal to refill early
- c) Ask the patient to bring a police report and then refill automatically
- d) Collect the details and route the request for pharmacist review and decision

47. Albuterol HFA: "Inhale 2 puffs every 4 hours as needed." Inhaler contains 200 puffs. If used around-the-clock at the maximum frequency, what is the days' supply?

- a) 8
- b) 10
- c) 12
- d) 25

48. A technician enters cyanocobalamin 1000 mcg/mL as "1000 mg/mL." What type of error is this?

- a) Wrong patient error
- b) Duplicate therapy error
- c) Magnitude/unit conversion error
- d) Wrong route error

49. A prescription is for Latuda 60 mg daily. What is the generic name?

- a) Lurasidone
- b) Lamotrigine
- c) Liraglutide
- d) Levetiracetam

50. An insurance claim rejects: "PA required." The patient wants the medication today. Which technician action is appropriate?

- a) Switch to a different drug in the class and bill it
- b) Force an override code without authorization
- c) Split the prescription into smaller quantities until it pays
- d) Initiate the standard PA process steps allowed in your setting and route appropriately for completion

51. A controlled prescription includes "Do not fill until 05/15." Today is 05/10. What is the correct handling?

- a) Fill now and post-date the label to 05/15
- b) Hold the prescription so it is processed/dispensed on or after the authorized date
- c) Change the "do not fill" date to today since the patient presented it
- d) Reject it because future-dated controlled prescriptions are invalid

52. The sig says: "Take ii caps PO qhs." What does "qhs" mean?

- a) Every morning
- b) Every 6 hours
- c) Twice daily
- d) At bedtime

53. Unit-dose blister cards arrive with an unreadable lot number, but the expiration date is readable. What should the technician do before restocking?

- a) Restock anyway; expiration date is all that matters
- b) Assign a new lot number based on another card in the box
- c) Quarantine/remove from inventory because traceability is compromised
- d) Send it to the floor stock area and document later

54. Suboxone 8 mg/2 mg film contains which active ingredients?

- a) Buprenorphine/naloxone
- b) Buprenorphine/methadone
- c) Naloxone/naltrexone
- d) Morphine/naloxone

55. A patient has "pantoprazole—hives" listed as an allergy. A new prescription is for Protonix 40 mg daily. What should the technician do?

- a) Fill Protonix because it's a different name

- b) Route for pharmacist review because Protonix is pantoprazole
- c) Substitute omeprazole without authorization
- d) Delete the allergy because PPIs are interchangeable

56. A patient asks whether they can crush diltiazem ER capsules. What is the technician's best response?

- a) Advise crushing only half the capsule contents
- b) Advise opening and sprinkling on applesauce
- c) Recommend switching to immediate-release without a new prescription
- d) Offer pharmacist counseling for the question

57. Valacyclovir 1 g: "Take 1 tablet by mouth twice daily for 7 days." What quantity should be dispensed?

- a) 7
- b) 10
- c) 14
- d) 28

58. A prescription is for omeprazole 20 mg, but only Prilosec OTC is in stock. What is the best next step for the technician?

- a) Dispense OTC without a prescription label because it's nonprescription
- b) Confirm the OTC product/NDC and package size are appropriate for prescription dispensing and billing before filling
- c) Substitute esomeprazole without prescriber authorization
- d) Change the directions to "PRN heartburn" to match OTC labeling

59. Which record/form is used to order Schedule II medications from a wholesaler?

- a) DEA Form 106
- b) DEA Form 41
- c) DEA Form 224
- d) DEA Form 222 or an approved electronic CSOS record

60. The prescription is for Eliquis 5 mg twice daily. Eliquis is:

- a) Rivaroxaban
- b) Dabigatran
- c) Apixaban
- d) Edoxaban

61. Under federal requirements, how long must prescription records (controlled and noncontrolled) generally be retained?

- a) 1 year
- b) 2 years
- c) 5 years
- d) 10 years

62. The label prints "gabapentin: Take 1 capsule BID," but the eRx directions were TID. What should the technician do?

- a) Handwrite "TID" on the label and dispense

- b) Correct the directions in the system, reprint the label, and continue normal verification workflow
- c) Tell the patient verbally to take TID
- d) Leave BID because it is safer

63. A prescription is for lisinopril/HCTZ 20/12.5 mg daily. The technician selected lisinopril 20 mg. What should be done?

- a) Dispense lisinopril alone; HCTZ is optional
- b) Add HCTZ as an OTC recommendation and proceed
- c) Select the correct combination product and strength before filling
- d) Change the prescription to lisinopril/HCTZ 10/12.5 mg for safety

64. A new prescription for sertraline 50 mg daily is received. The profile shows phenelzine filled 12 days ago. What should the technician do?

- a) Fill sertraline and add "stop phenelzine" to the label
- b) Inactivate phenelzine and proceed
- c) Reduce sertraline to 25 mg and fill
- d) Route for pharmacist review due to serious interaction/washout concerns

65. A will-call bag is found with the correct patient name but the wrong address on the leaflet. What is the best technician action before handing it out?

- a) Cross out the address and initial it
- b) Verify two identifiers with the patient and confirm the correct profile and prescription match before sale
- c) Dispense anyway since the name matches
- d) Remove the leaflet and provide only the vial

66. At fill, barcode scanning shows the bottle is finasteride 5 mg, but the label is for tamsulosin 0.4 mg. What should the technician do?

- a) Override the scan because the shelf location is correct
- b) Stop and resolve the mismatch; do not dispense until correct product is selected
- c) Dispense finasteride since both treat BPH
- d) Reprint the label for finasteride to match the scan

67. Singulair 10 mg nightly is prescribed. Which generic is correct?

- a) Montelukast
- b) Loratadine
- c) Fluticasone
- d) Albuterol

68. Which DEA form is associated with documenting destruction of controlled substances when required?

- a) DEA Form 41
- b) DEA Form 106
- c) DEA Form 222
- d) DEA Form 224a

69. A prescription says carvedilol 12.5 mg: "Take 1 tablet twice daily." The technician enters

once daily. Which best prevents this type of error at entry?

- a) Skip sig review if quantity seems reasonable
- b) Default all beta-blockers to once daily
- c) Compare the typed sig to the original prescription before finalizing
- d) Avoid entering frequencies; type "as directed" instead

70. Clopidogrel 75 mg daily is prescribed. Which brand name matches?

- a) Plaquenil
- b) Paxil
- c) Pravachol
- d) Plavix

71. Liquid medication: "Take 5 mL by mouth TID for 10 days." How many mL should be dispensed?

- a) 100 mL
- b) 120 mL
- c) 150 mL
- d) 180 mL

72. A patient asks what "take on an empty stomach" means. What should the technician do?

- a) Explain it fully because it is nonclinical
- b) Tell them it means no food for the entire day
- c) Suggest ignoring the label to avoid nausea
- d) Offer pharmacist counseling for clarification

73. Spiriva HandiHaler: "Inhale contents of 1 capsule daily." Which generic is correct?

- a) Ipratropium
- b) Tiotropium
- c) Salmeterol
- d) Budesonide

74. During data entry, the eRx DOB differs from the profile by one digit. What is the best technician action?

- a) Update the profile to match the eRx automatically
- b) Ignore it; eRx demographics are often wrong
- c) Create a new profile immediately
- d) Verify identity using additional identifiers and correct the record appropriately before proceeding

75. Doxycycline 100 mg: "Take 1 capsule BID for 10 days." What quantity should be dispensed?

- a) 10
- b) 14
- c) 20
- d) 30

76. While restocking an automated dispensing cabinet, a pocket labeled "morphine ER 30 mg" contains morphine IR 30 mg. What should the technician do?

- a) Leave it because the strength matches
- b) Add an "IR" sticker and keep it in the pocket
- c) Fill the pocket with both ER and IR to reduce stockouts
- d) Remove the incorrect medication, quarantine per process, and correct the pocket contents

77. Pen needles: "Use 1 pen needle with each insulin injection QID." Box contains 100 pen needles. What days' supply should be billed?

- a) 20
- b) 25
- c) 30
- d) 10

78. Humalog KwikPen is which insulin?

- a) Insulin aspart
- b) Insulin glargine
- c) Insulin lispro
- d) Insulin detemir

79. A label with PHI prints and is left on the counter visible to customers. What is the best immediate action?

- a) Throw it into regular trash
- b) Put it back into the printer tray
- c) Place it into the secure shred/PHI destruction container
- d) Tape it to the hard copy so it isn't lost

80. A prescriber sends a refill authorization for a Schedule II medication. Which statement is correct?

- a) Schedule II refills are allowed if the prescriber approves
- b) Schedule II refills are allowed within 6 months only
- c) Schedule II refills are not permitted; each fill requires a new prescription
- d) Schedule II refills are allowed if the patient signs a controlled log

81. Nitrostat 0.4 mg SL is:

- a) Isosorbide mononitrate ER
- b) Nitroglycerin sublingual tablets
- c) Propranolol
- d) Ranolazine

82. Potassium chloride oral solution: "KCl 20 mEq/15 mL: Take 15 mL daily." How many mEq per dose?

- a) 10 mEq
- b) 15 mEq
- c) 30 mEq
- d) 20 mEq

83. A patient reports a serious adverse reaction and asks how to report it. What is the best technician response?

- a) Direct them to report it to the DEA
- b) Tell them only the prescriber can report adverse events
- c) Refer them to the pharmacist and mention FDA MedWatch as the reporting program
- d) Tell them to submit it to the insurance company

84. A Schedule II medication is partially filled due to insufficient stock. Which action is appropriate within technician workflow?

- a) Add refills to cover the remainder automatically
- b) Dispense the remainder later without additional documentation
- c) Convert the remainder to a new prescription under the same Rx number
- d) Ensure the partial fill is documented and routed for pharmacist completion per legal requirements

85. Metformin ER 500 mg: "Take 2 tablets daily with dinner." The technician has both IR and ER in stock. What should the technician do?

- a) Use IR since the strength matches
- b) Dispense ER and ensure the NDC and formulation match the ER product on the label
- c) Use either and just write "ER" on the label
- d) Dispense IR and change directions to BID

86. Entresto 49/51 mg BID contains:

- a) Losartan/hydrochlorothiazide
- b) Sacubitril/valsartan
- c) Amlodipine/benazepril
- d) Lisinopril/amlodipine

87. A technician is asked to help validate a DEA number entered from a hard copy. Which method is used to verify the check digit format?

- a) Add all digits and divide by 7
- b) Use the DEA check-digit algorithm (odd/even position sums and final digit check)
- c) Match the DEA number to the patient's insurance ID
- d) Check digits cannot be validated; only the DEA can confirm

88. Which brand name matches lansoprazole?

- a) Prevacid
- b) Nexium
- c) Protonix
- d) Prilosec

89. Before a compounded oral rinse is released to the counter, what best supports quality assurance at the technician level?

- a) Use the manufacturer expiration date of the main ingredient only
- b) Remove auxiliary labels to reduce confusion
- c) Provide it without labeling since it is compounded
- d) Confirm the container has required labeling including directions, ingredients/strength as applicable, beyond-use date, and appropriate auxiliary labels

90. A prescription is for rosuvastatin 40 mg daily. The technician accidentally counted 20 mg

tablets but labeled them as 40 mg. Which technician safeguard best prevents this at filling?

- a) Ask the patient which strength they usually take
- b) Keep both strengths in the same bin to reduce walking time
- c) Verify the NDC/strength at fill using barcode scanning and match it to the printed label
- d) Count by tablet appearance and proceed if it looks right

Answer Key

1) b	31) d	61) b
2) a	32) c	62) b
3) c	33) c	63) c
4) a	34) d	64) d
5) d	35) c	65) b
6) b	36) b	66) b
7) c	37) d	67) a
8) c	38) b	68) a
9) d	39) c	69) c
10) c	40) b	70) d
11) b	41) b	71) c
12) c	42) c	72) d
13) b	43) d	73) b
14) d	44) b	74) d
15) b	45) b	75) c
16) c	46) d	76) d
17) b	47) a	77) b
18) c	48) c	78) c
19) c	49) a	79) c
20) d	50) d	80) c
21) c	51) b	81) b
22) c	52) d	82) d
23) a	53) c	83) c
24) d	54) a	84) d
25) b	55) b	85) b
26) a	56) d	86) b
27) c	57) c	87) b
28) b	58) b	88) a
29) b	59) d	89) d
30) c	60) c	90) c