Antibiotic	Dose for Normal Renal Function	Dosage Adjustment for Renal Failure According to Estimated Creatinine Clearance			Doses for Dialysis		
		>50 mL/min	10-50 mL/min	<10 mL/min	HD	PD	CRRT
Aminoglycosides							
Amikacin IV	**Single Daily Dose (SDD):** 15 mg/kg q24H (max 1.5 gm/day) **Use adjusted body weight if actual body weight > 30% ideal body weight**	**>60 mL/min:** 15 mg/kg q24H	**20-39 mL/min:** 15 mg/kg q48H	15 mg/kg single dose then monitor serum levels	5-12.5 mg/kg three times a week after dialysis on dialysis days[1]	5-12.5 mg/kg q48-72H[1]	7.5 mg/kg q24H
		40-59 mL/min: 15 mg/kg q36H	**≤20 mL/min:** 15 mg/kg single dose then monitor serum levels				
	Multiple Daily Dose (MDD): 15 mg/kg q24H or 7.5 mg/kg q12H **Use adjusted body weight if actual body weight > 30% ideal body weight**	7.5 mg/kg q12H	7.5 mg/kg q24H	7.5 mg/kg q48H			
Gentamicin IV	**High-dose, extended-interval dosing (Gram-negative infections):** 5 mg/kg q24H (in septic shock, up to 7 mg/kg q24H)[1] **Use adjusted body weight if actual body weight > 30% ideal body weight**	**>60 mL/min:** 5-7 mg/kg q24H **40-59 mL/min:** 5-7 mg/kg q36H	**40-59 mL/min:** 5-7 mg/kg q36H[1]	5-7 mg/kg single dose and monitor serum levels[1]	2-3 mg/kg loading dose, then 2-3 mg/kg three times a week after dialysis on dialysis days[1]	1-3 mg/kg q48-72H[1]	1.7-2 mg/kg q24H
			20-39 mL/min: 5-7 mg/kg q48H[1]				
			≤20 mL/min: 5-7 mg/kg single dose and monitor serum levels[1]				
	Synergy dosing (Gram-positive infections): 1 mg/kg q8H **Use adjusted body weight if actual body weight > 30% ideal body weight**	Normal	**40-59 mL/min:** 1 mg/kg q12H **20-39 mL/min:** 1 mg/kg q24H **<20 mL/min:** 1 mg/kg single dose and monitor serum levels	1 mg/kg single dose and monitor serum levels	1 mg/kg three times a week after dialysis on dialysis days		
Streptomycin IV/IM	15 mg/kg (max 1 gm) q24H	Normal	**>30 mL/min:** Normal	15 mg/kg two to three times a week	15 mg/kg two to three times a week after dialysis on dialysis days[1]	15 mg/kg (max 1 gm) q72-96H + 20-40 mg lost per litre dialysate per day[1]	15 mg/kg q24-72H

Antibiotic	Dose for Normal Renal Function	Dosage Adjustment for Renal Failure According to Estimated Creatinine Clearance			Doses for Dialysis		
		>50 mL/min	10-50 mL/min	<10 mL/min	HD	PD	CRRT
			<30 mL/min: 15 mg/kg two to three times a week			dialysate per day	
Carbapenems							
Ertapenem IV	1 gm q24H	Normal	**>30 mL/min:** 1 gm q24H **<30 mL/min:** 0.5 gm q24H	0.5 gm q24H	0.5-1 gm three times a week after dialysis on dialysis days[1]	0.5 gm q24H	0.5-1 gm q24H
Meropenem IV	1 gm q8H	Normal	**26-50 mL/min:** 1 gm q12H **10-25 mL/min:** 500 mg q12H	500 mg q24H	500 mg q24H, dose after dialysis on dialysis days	500 mg q24H	1 gm q8-12H or 0.5-1 gm q8H[1]
	Meningitis and CRE: 2 gm q8H **Note:** For treatment of CRE, to administer as extended infusion over 3 hours.	Normal	**26-50 mL/min:** 2 gm q12H **10-25 mL/min:** 1 gm q12H	1 gm q24H	1 gm q24H, dose after dialysis on dialysis days	1 gm q24H	2 gm q8-12H
Cephalosporins							
Cefazolin IV	2 gm q8H (max 12 gm/day)	Normal	**>30 mL/min:** 2 gm q8-12H[1] **11-29 mL/min:** 1 gm q12H (up to 2 gm q12H in severe infection)[1]	1 gm q24H	2 gm after dialysis if next dialysis expected in 48 hours, and 3 gm after dialysis is expected in 72 hours or 1 gm q24H or 2 gm q48-72H, dose after dialysis only on dialysis days[1]	0.5 gm q12H	2 gm single dose, then 2 gm q12H[1]
Cefalexin PO	250-1000 mg q6H	Normal	**15-29 mL/min:** 250-500 mg q8-12H[1]	250-500 mg q12-24H[1]	250-500 mg q12-24H, dose after dialysis on	250-500 mg q12-24H	250-500 mg q8-12H[3]

Antibiotic	Dose for Normal Renal Function	Dosage Adjustment for Renal Failure According to Estimated Creatinine Clearance			Doses for Dialysis		
		>50 mL/min	10-50 mL/min	<10 mL/min	HD	PD	CRRT
Cefalexin PO	(max 4 gm/day)	Normal	**<15 mL/min:** 250-500 mg q12-24H[1]	250-500 mg q12-24H[1]	dose after dialysis on dialysis days	250-500 mg q12-24H	250-500 mg q8-12H[1]
Cefuroxime IV	1500 mg q6-8H	Normal	**10-29 mL/min:** 1500 mg q12H[1]	1500 mg q24H	1500 mg q24H, dose after dialysis on dialysis days	1500 mg q24H	1500 mg q8-12H
Cefuroxime PO	250-500 mg q12H	Normal[3]	**10-29 mL/min:** 250 mg q12H[1] or 250-500 mg q24H[1] or Normal[3]	250 mg q24H[1] or 250-500 mg q48H[1] or Normal[3]	250 mg q24H, dose after dialysis on dialysis days[1] or 250-500 mg q48H, dose after dialysis on dialysis days[1] or Normal[3]	250 mg q24H[1] or 250-500 mg q48H[1]	Normal[3]
Ceftriaxone IV	1-2 gm q12-24H	Normal	Normal	Normal	Normal No supplement dose required after dialysis	Normal	Normal
Ceftazidime IV	1-2 gm q8H	Normal[1]	**31-50mL/min:** 1-2 gm q12H[1] **16-30mL/min:** 1-2 gm q24H[1] **<15 mL/min:** 0.5-1 gm q24H[1]	0.5-1 gm q24H[1]	0.5-1 gm q24H or 2 gm q48-72H, dose after dialysis only on dialysis days	1 gm q24H[1]	2 gm q8-12H[1]
Cefoperazone IV	1-2 gm q12H	Normal	Normal	Normal	Normal No supplement dose required after dialysis	Normal	Normal
Cefepime IV	2 gm q8H	Normal	**30-60 mL/min:** 2 gm q12H **11-29 mL/min:** 2 gm q24H	1 gm q24H	1 gm q24H, dose after dialysis on dialysis days[1] or 2 gm q48-72H, dose after dialysis only on dialysis days[1]	2 gm q48H or 1 gm q24H[1]	2 gm q8-12H[1]
			30-50 mL/min: 1.25 gm q8H[1]		0.94 gm q24-48H, dose after dialysis on dialysis	0.94 gm q24-48H, dose after dialysis on dialysis	

Antibiotic	Dose for Normal Renal Function	Dosage Adjustment for Renal Failure According to Estimated Creatinine Clearance			Doses for Dialysis		
		>50 mL/min	10-50 mL/min	<10 mL/min	HD	PD	CRRT
Ceftazidime/Avibactam IV	2.5 gm q8H[1] **Note:** Administer as infusion over 2 hours.	Normal[1]	**15-30 mL/min:** 0.94 gm q12H[1] **5-15 mL/min:** 0.94 gm q24H[1]	**<5 mL/min:** 0.94 gm q48H[1]	after dialysis on dialysis days (Administer q48H in patients with minimal residual kidney function and less severe infections)[1]	after dialysis on dialysis days (Administer q48H in patients with minimal residual kidney function and less severe infections)[1]	1.25 gm q8H[1]
Fluoroquinolones							
Ciprofloxacin IV	200-400 mg q8-12H[1] **Deep-seated infections, *Pseudomonas spp.* infections, meningitis:** 400 mg q8H[1]	Normal	**> 30 mL/min:** Normal **5-29 mL/min:** 200-400 mg q12-24H[1]	200-400 mg q12-24H[1]	200-400 mg q24H, dose after dialysis on dialysis days	200 mg q12H[3]	200-400 mg q8-12H[1]
Ciprofloxacin PO	500-750 mg q12H	Normal	**30-49 mL/min:** 250-500 mg q12H **5-29 mL/min:** 500-750 mg q24H	500-750 mg q24H	250-500 mg q24H, dose after dialysis on dialysis days	250-500 mg q24H[1] or 250mg q8-12H[3]	250-750 mg q12H[1]
Levofloxacin IV / PO	500-750 mg q24H[1]	Normal	**20-49 mL/min:** 500-750 mg loading dose, then 250 mg q24H or 750 mg q48H[1] **10-19 mL/min:** 500-750 mg loading dose, then 250-500 mg q48H[1]	500-750 mg loading dose, then 250-500 mg q48H[1]	500-750 mg loading dose, then 250-500 mg q48H[1]	500-750 mg loading dose, then 250-500 mg q48H[1]	500-750 mg loading dose, then 250-500 mg q24H[1]
	Tuberculosis: 500-1000 mg q24H[1]	Normal[1]	**>30 mL/min:** Normal[1] **<30 mL/min:** 500 mg or 1 gm three times per week[1]	500 mg or 1 gm three times per week[1]	500 mg or 1 gm three times per week, dose after dialysis on dialysis days[1]	500 mg or 1 gm three times per week[1]	No data[1]
Moxifloxacin PO	400 mg q24H	Normal	Normal	Normal	Normal No supplemental dose necessary after dialysis	Normal	Normal
Macrolides							
Azithromycin IV / PO	500 mg q24H	Normal	Normal	Normal	Normal No supplemental dose necessary after dialysis	Normal	Normal

Antibiotic	Dose for Normal Renal Function	Dosage Adjustment for Renal Failure According to Estimated Creatinine Clearance			Doses for Dialysis		
		>50 mL/min	10-50 mL/min	<10 mL/min	HD	PD	CRRT
Erythromycin PO	400-800 mg q6-12H	Normal	Normal	Normal	Normal No supplemental dose necessary after dialysis	Normal	Normal
Clarithromycin PO	250-500 mg q12H	Normal	**>30 mL/min:** Normal[1] **<30 mL/min:** 250-500 mg q24H[1]	250-500 mg q24H[1]	250-500 mg q24H, dose after dialysis on dialysis days[1]	250-500 mg q24H[1]	250-500 mg q12-24H
Penicillins							
Amoxicillin PO	250-1000 mg q8H	Normal	**>30 mL/min:** Normal **10-30 mL/min:** 250-1000 mg q12H[1]	250-500 mg q12-24H[1]	250-500 mg q12-24H, dose after dialysis on dialysis days[1]	250-500 mg q12H	250-500 mg q8-12H
Ampicillin IV	1-2 gm q4-6H	Normal	**30-50 mL/min:** 1-2 gm q6-8H **10-29 mL/min:** 1-2 gm q8-12H	1-2 gm q12H	1-2 gm q12H, dose after dialysis on dialysis days	500-1000 mg q12H	1-2 gm q8-12H
Ampicillin/Sulbactam IV	1.5-3 gm q6H	Normal	**>30 mL/min:** Normal **15-29 mL/min:** 1.5-3 gm q12H[1]	1.5-3 gm q24H[1]	1.5-3 gm q12-24H, dose after dialysis[1]	1.5-3 gm q12-24H[1]	1.5-3 gm q8-12H
	Carbapenem-resistant *Acinetobacter* spp.: 9 gm q8H over 4 hours[1] Dilute in 200 mL NS or WFI **Alternative regimen for intolerance or toxicity from high-dose** 3 gm q4H[1]	**>90 mL/min:** Normal	**30-50 mL/min:** 3 gm q6H[1]	3 gm q12H[1]	3 gm q12H, dose after dialysis on dialysis days[1]	3 gm q12H[1]	3 gm q6H[1]
		60-90 mL/min: 6 gm q8H[1]	**15-29 mL/min:** 3 gm q8H[1]				
		51-59 mL/min: 3 gm q6H[1]	**10-15 mL/min:** 3 gm q12H[1]				

Antibiotic	Dose for Normal Renal Function	Dosage Adjustment for Renal Failure According to Estimated Creatinine Clearance			Doses for Dialysis		
		>50 mL/min	10-50 mL/min	<10 mL/min	HD	PD	CRRT
Ampicillin/Sulbactam PO (Sultamicillin)	375-750 mg q12H	Normal	Normal	Normal	Normal	Normal	Normal
Benzylpenicillin IV (Penicillin G)	1-4 MU q4-6H[1]	Normal	75% of dose q4-6H[1]	20-50% of dose q4-6H[1]	25-50% of dose q4-6H, dose after dialysis[1]	25-50% of dose q4-6H[1]	2-4 MU q4-6H[1]
Cloxacillin IV / PO	1-2 gm q4-6H	Normal	Normal	Normal	Normal	Normal	Normal
Co-amoxiclav PO (Amoxicillin 500 mg/ Clavulanic Acid 125 mg)	625 mg q8-12H	Normal	**>30 mL/min:** Normal **10-30 mL/min:** 625 mg q12-24H	625 mg q24H	625 mg q24H, dose after dialysis on dialysis days	625 mg q12H[1]	No data
Co-amoxiclav PO (Amoxicillin 875 mg/ Clavulanic Acid 125 mg)	**Deep-seated infection:** 1 gm q8-12H	Normal	**> 30 mL/min:** Normal **<30 mL/min:** Avoid use[1]	Avoid use[1]	Avoid use[1]	No data	No data
Co-amoxiclav IV	1.2 gm q8H **Severe infection:** 1.2 gm q6H	Normal	**>30 mL/min:** Normal **10-30 mL/min:** 1.2 g q12H[3] or 1.2 gm STAT then 600 mg q12H	1.2 gm q12H[3] or 1.2 gm STAT then 600 mg q24H	1.2 gm q12H[3] or 1.2 gm STAT then 600 mg q24H, dose after dialysis on dialysis days	1.2 gm q12H[3]	1.2 gm q12H[3]
Phenoxymethylpenicillin PO (Penicillin V)	250-1000 mg q6-12H[1]	Normal	Normal. Use with caution; excretion is prolonged in patients with kidney impairment.[1]	Normal. Use with caution; excretion is prolonged in patients with kidney impairment.[1]	Normal. Use with caution; excretion is prolonged in patients with kidney impairment.[1]	Normal. Use with caution; excretion is prolonged in patients with kidney impairment.[1]	Normal. Use with caution; excretion is prolonged in patients with kidney impairment.[1]
Piperacillin/Tazobactam IV	4.5 gm q6H	Normal	**20-40 mL/min** 4.5 gm q8H[1]	2.25 gm q6H	2.25 gm q8H[1]	2.25 gm q8H	4.5 gm q8H[1]

Antibiotic	Dose for Normal Renal Function	Dosage Adjustment for Renal Failure According to Estimated Creatinine Clearance			Doses for Dialysis		
		>50 mL/min	10-50 mL/min	<10 mL/min	HD	PD	CRRT
Piperacillin/ Tazobactam IV	4.5 gm q6H	Normal	**10-20 mL/min** 2.25 gm q6H	2.25 gm q6H	2.25 gm q8H[1]	2.25 gm q8H	4.5 gm q8H[1]
Tetracyclines							
Doxycycline PO	100 mg q12H	Normal	Normal	Normal	Normal	Normal	Normal
Tigecycline IV	100-200 mg loading dose, then 50-100 mg q12H[1]	Normal	Normal	Normal	Normal	Normal	Normal
Miscellaneous							
Aztreonam IV	2 gm q8H **Note:** Use q6H for meningitis	Normal	**>30 mL/min:** Normal **10-29 mL/min:** 2 gm q12H	2 gm q24H	2 gm q24H, dose after dialysis on dialysis days	2 gm q24H	2 gm q12H
Clindamycin IV / PO	600 mg q6-8H or 900 mg q8H	Normal	Normal	Normal	Normal	Normal	Normal
Colistin IV	**Loading dose:** 9 MU STAT[1] **Maintenance dose (to start 12 hours after LD):** 4.5-5.4 MU q12H[1] ***Note:** Refer to 'Colistin Dosing' tab for detailed renal dose adjustment	4.5 MU q12H[1] Patients with good kidney function (CrCl ≥80 mL/min) may require doses of CBA >300 mg per day.	**30-50 mL/min:** 3 MU q12H[1] **10-30 mL/min:** 2.5 MU q12H[1]	1.75 MU q12H[1]	**Loading dose:** 9 MU STAT[1] **Maintenance dose (to start 12 hours after LD):** 2 MU q12H, dose on non-dialysis days[1] + 2.7 MU q12H, dose after dialysis on dialysis days[1]	**Loading dose:** 9 MU STAT[1] **Maintenance dose (to start 12 hours after LD):** 4.5-6 MU q24H[1]	**Loading dose:** 9 MU STAT[1] **Maintenance dose (to start 12 hours after LD):** 6.6 MU q12H[1]
Daptomycin IV	4-12 mg/kg q24H (according to indication) **Use adjusted body weight for obese patients**	Normal	**30-50 mL/min:** Normal **<30 mL/min:** 4-6 mg/kg q48H	4-6 mg/kg q48H	4-6 mg/kg q48H, administer after dialysis on dialysis days	4-6 mg/kg q48H	6 mg/kg q24H

Antibiotic	Dose for Normal Renal Function	Dosage Adjustment for Renal Failure According to Estimated Creatinine Clearance			Doses for Dialysis		
		>50 mL/min	10-50 mL/min	<10 mL/min	HD	PD	CRRT
Linezolid IV / PO	600 mg q12H	Normal	Normal	Normal	Normal	Normal	Normal
Metronidazole IV	500 mg q6-12H or 7.5 mg/kg q6H (max 4 gm/day)	Normal	Normal	500 mg q8-12H or 7.5 mg/kg q12H	500 mg q8-12H, dose after dialysis or 7.5 mg/kg q12H, dose after dialysis on dialysis days	500 mg q8-12H or 7.5 mg/kg q12H	500 mg q6-12H or 7.5 mg/kg q6H
Metronidazole PO	200-800 mg q8-12H	Normal	Normal	Normal	Normal	Normal	Normal
Nitrofurantoin PO	50-100 mg q6H	Normal	**>30 mL/min:** Normal **<30 mL/min:** Avoid use	Avoid use	Avoid use	Avoid use	Avoid use
Polymyxin B IV	**Loading dose:** 2.5 mg/kg STAT over 2 hours **Maintenance dose (to start 12 hours after LD):** 1.5 mg/kg q12H, each dose over 1 hour **Use adjusted body weight for obese patients**	Normal	Normal	Normal	Normal	Normal	Normal
Trimethoprim/ Sulfamethoxazole IV/PO	**Treatment:** Trimethoprim component 8-20 mg/kg/day divided to q6 - 12H **Use adjusted body weight for obese patient**[9-10]	Normal	**30-50 mL/min:** Normal **10-30 mL/min:** Trimethoprim component 2.5-5 mg/kg q12H	Use not recommended, but if required, Trimethoprim component 5-10 mg/kg q24H	Use not recommended, but if required, Trimethoprim component 5-10 mg/kg q24H	Use not recommended, but if required, Trimethoprim component 5-10 mg/kg q24H	Trimethoprim component 5-10 mg/kg q12H
	Prophylaxis: 1-2 tabs PO q24H	Normal	Normal	Normal	Normal	Normal	Normal

Antibiotic	Dose for Normal Renal Function	Dosage Adjustment for Renal Failure According to Estimated Creatinine Clearance			Doses for Dialysis		
		>50 mL/min	10-50 mL/min	<10 mL/min	HD	PD	CRRT
Vancomycin IV	**Loading Dose[1]:** **>90 mL/min:** 25-30 mg/kg single dose **<90 mL/min:** 20-25 mg/kg single dose (maximum 3 gm per dose) **Maintenance Dose[1]:** 15-20 mg/kg q8-12H (suggest maximum 1 gm per dose to avoid toxicity) **Max infusion rate[1]:** 1 gm/hour **Max dilution concentration[1]:** 1 gm/100mL **Obese patient requires 20-25 mg/kg of loading dose (maximum 3 gm)[1]**	**>90 mL/min:** 15-20 mg/kg q8-12H[1] **50-90 mL/min:** 15-20 mg/kg q12H[1]	**15-50 mL/min:** 10-15 mg/kg q24H[1]	**<15 mL/min:** 20-25 mg/kg single dose, then send a random TDM sample next morning. If TDM is not available the next day, give 7.5-10 mg/kg single dose 24 hours after the initial loading dose. Send a random TDM sample the next working day.	20-25 mg/kg single dose, then send a random TDM sample next morning. If TDM is not available the next day, give 7.5-10 mg/kg single dose 24 hours after the initial loading dose. Send a random TDM sample the next working day.	20-25 mg/kg single dose, then send a random TDM sample next morning.	20-25 mg/kg single dose, then 7.5-10 mg/kg q12-24H
		Refer to pharmacy when in doubt.					
Antifungals							
Amphotericin B Deoxycholate IV	0.3-1 mg/kg q24H	Normal	Normal	Normal	Normal	Normal	Normal
Liposomal Amphotericin B IV	3-5 mg/kg q24H	Normal	Normal	Normal	Normal	Normal	Normal
Fluconazole IV / PO	100-400 mg q24H	Normal	50-200 mg q24H	50-200 mg q24H	50-200 mg q24H, dose after dialysis on dialysis days or 100-400 mg q48-72H, dose after dialysis	50-200 mg q24H	200-400 mg q24H
Itraconazole PO	100-200 mg q12H (max 600 mg/day)	Normal	Normal	No data	No data	100 mg q12-24H	100-200 mg q12H

Antibiotic	Dose for Normal Renal Function	Dosage Adjustment for Renal Failure According to Estimated Creatinine Clearance			Doses for Dialysis		
		>50 mL/min	10-50 mL/min	<10 mL/min	HD	PD	CRRT
Voriconazole IV	6 mg/kg q12H for 2 doses then 4 mg/kg q12H	Normal	Not recommended due to accumulation of IV vehicle (cyclodextrin), oral route preferred	Not recommended due to accumulation of IV vehicle (cyclodextrin), oral route preferred	No data. Oral route preferred	No data. Oral route preferred	No data. Oral route preferred
Voriconazole PO	**> 40 kg:** 400 mg q12H for 2 doses, then 200 mg q12H **< 40 kg:** 200 mg q12H for 2 doses, then 100 mg q12H	Normal	Normal	Normal	Normal	Normal	Normal
Micafungin IV	100 mg q24H	Normal	Normal	Normal	Normal	Normal	150-200 mg q24H
Flucytosine PO	25 mg/kg q6H	25 mg/kg q6H	25 mg/kg q12H	25 mg/kg q24H	25 mg/kg q24H	0.5-1 gm q24H	25 mg/kg q12H
Antiparasitic							
Pentamidine IV	4 mg/kg q24H	Normal	Normal	4 mg/kg q24-36H	4 mg/kg q48H, dose after dialysis	4 mg/kg q24-36H	4 mg/kg q24H
Antituberculosis							
Rifampicin PO	10 mg/kg q24H	Normal	Normal	Normal	Normal	Normal	Normal
Isoniazid PO	5 mg/kg q24H	Normal	Normal	Normal	Normal	Normal	Normal
Ethambutol PO	15-25 mg/kg q24H	Normal	**>30 mL/min:** 15-25 mg/kg q24H **10-30 mL/min:** 20-25 mg/kg three times a week	20-25 mg/kg three times a week	20-25 mg/kg three times a week, after dialysis on dialysis days	20-25 mg/kg three times a week	15-25 mg/kg q24H
Pyrazinamide PO	20-25 mg/kg q24H (max 2.5 gm/day)	Normal	**>30 mL/min:** 20-25 mg/kg q24H **<30 mL/min:** 25-35 mg/kg three times a week	25-35 mg/kg three times a week	25-35 mg/kg three times a week, after dialysis on dialysis days	20-25 mg/kg q24H	20-25 mg/kg q24H

Antibiotic	Dose for Normal Renal Function	Dosage Adjustment for Renal Failure According to Estimated Creatinine Clearance			Doses for Dialysis		
		>50 mL/min	10-50 mL/min	<10 mL/min	HD	PD	CRRT
Antivirals							
Acyclovir IV	5-10 mg/kg q8H[1] **Use IBW in obese patient**	Normal	**25-50 mL/min:** 5-10 mg/kg q12H[1] **10-25 mL/min:** 5-10 mg/kg q24H[1]	2.5-5 mg/kg q24H[1]	2.5-5 mg/kg q24H, dose after dialysis[1]	2.5-5 mg/kg q24H[1]	5-10 mg/kg q12-24H[1]
Acyclovir PO	200-800 mg q8H or 5 times daily	Normal	**>25mL/min:** Normal **10-25 mL/min:** 200-800 mg q8H[1]	200-800 mg q12H[1]	200-800 mg q12H, dose after dialysis on dialysis days	200-800 mg q12H	No data
Entecavir PO	0.5 mg q24H	Normal	0.15-0.25 mg q24H	0.05 mg q24H	0.05 mg q24H, dose after dialysis on dialysis days or 0.5 mg once weekly, dose after dialysis on dialysis days[1]	0.05 mg q24H	No data
Ganciclovir IV	**Induction Dose:** 5 mg/kg q12H	**> 70 mL/min:** Normal **50-69 mL/min:** 2.5 mg/kg q12H	**25-49 mL/min:** 2.5 mg/kg q24H **10-24 mL/min:** 1.25 mg/kg q24H	1.25 mg/kg three times a week	1.25 mg/kg three times a week, dose after dialysis on dialysis days	1.25 mg/kg three times a week	2.5 mg/kg q24H[1]
	Maintenance Dose: 5 mg/kg q24H	**> 70 mL/min:** Normal **50-69 mL/min:** 2.5 mg/kg q24H	**25-49 mL/min:** 1.25 mg/kg q24H **10-24 mL/min:** 0.625 mg/kg q24H	0.625 mg/kg three times a week	0.625 mg/kg three times a week, dose after dialysis on dialysis days	0.625 mg/kg three times a week	1.25 mg/kg q24H[1]
Lamivudine PO	**Hepatitis B:** 100 mg q24H[1]	Normal	**30-49 mL/min:** 100 mg STAT then 50 mg q24H[1] **15-29 mL/min:** 100 mg STAT then 25 mg q24H[1] **5-14 mL/min:** 35 mg STAT then 15 mg q24H[1]	**<5 mL/min:** 35 mg STAT then 10 mg q24H[1]	35 mg STAT then 10 mg q24H, dose after dialysis on dialysis days[1]	35 mg STAT then 10 mg q24H[1]	100 mg STAT

Antibiotic	Dose for Normal Renal Function	Dosage Adjustment for Renal Failure According to Estimated Creatinine Clearance			Doses for Dialysis		
		>50 mL/min	10-50 mL/min	<10 mL/min	HD	PD	CRRT
Lamivudine PO	**HIV:** 150 mg q12H[1] or 300 mg q24H[1]	Normal	**30-49 mL/min:** 150 mg q24H[1] **15-29 mL/min:** 150 mg STAT then 100 mg q24H[1] **5-14 mL/min:** 150 mg STAT then 50 mg q24H[1]	**<5 mL/min:** 50 mg STAT then 25 mg q24H[1]	50 mg STAT then 25 mg q24H, dose after dialysis on dialysis days[1]	50 mg STAT then 25 mg q24H[1]	then 50 mg q24H[1]
Oseltamivir PO	75 mg q12H	Normal	**30-60 mL/min:** 30 mg q12H **10-29 mL/min:** 30 mg q24H	75 mg single dose[3]	30 mg STAT then 30 mg after each dialysis day, no drug on non-dialysis days. Treatment duration not to exceed 5 days. Assumes 3 HD sessions in 5 days period.[3]	**CAPD:** 30 mg single dose **APD:** 75 mg single dose	75 mg q24H[3] or 30 mg q12H[3]
Nirmatrelvir / Ritonavir PO	Nirmatrelvir 300 mg q12H + Ritonavir 100 mg q12H	**>60 mL/min:** Normal	**30-60 mL/min:** Nirmatrelvir 150 mg q12H + Ritonavir 100 mg q12H **< 30mL/min:** Not recommended (no data)	Not recommended (no data)	Not recommended (no data)	Not recommended (no data)	Not recommended (no data)
Telbivudine PO	600 mg q24H	Normal	**30-49 mL/min:** 600 mg q48H **10-29 mL/min:** 600 mg q72H	600 mg q96H	600 mg q96H, dose after dialysis on dialysis days	No data	No data
Tenofovir Disoproxil Fumarate PO	300 mg q24H	Normal	**30-49 mL/min:** 300 mg q48H **10-29 mL/min:** 300 mg q72-96H	No data	300 mg after every 3rd dialysis or every 7 days if no dialysis	No data	No data
Tenofovir Disoproxil Fumarate /	300 / 200 mg (1 tab) q24H[1]	Normal	**30-49 mL/min:** 1 tab q48H[1]	Not recommended[1]	Not recommended[1]	Not recommended[1]	Not recommended[1]

Antibiotic	Dose for Normal Renal Function	Dosage Adjustment for Renal Failure According to Estimated Creatinine Clearance			Doses for Dialysis		
		>50 mL/min	10-50 mL/min	<10 mL/min	HD	PD	CRRT
Emtricitabine PO	300 / 200 mg (1 tab) q24H[1]	Normal	**< 30 mL/min:** Not recommended[1]	Not recommended[1]	Not recommended[1]	Not recommended[1]	Not recommended[1]
Zidovudine PO	300 mg q12H	Normal	**>15 mL/min:** 300 mg q12H	**< 15 mL/min:** 100 mg q8H or 300 mg q24H	100 mg q8H, dose after dialysis on dialysis days	100 mg q8H	300 mg q12H