

# Cell decomposition in $P$ -minimal structures

Saskia Chambille

Joint work with Pablo Cubides Kovacsics & Eva Leenknegt


Departement of Mathematics  
KU Leuven

Konstanz, July 22, 2016


## Tree pictures of the $p$ -adic numbers




# Tree pictures of the $p$ -adic numbers




# Tree pictures of the $p$ -adic numbers





# Tree pictures of the $p$ -adic numbers





# Tree pictures of the $p$ -adic numbers



# Tree pictures of the $p$ -adic numbers



# Tree pictures of the $p$ -adic numbers





# Definitions

Let  $p$  be a prime number.  $\mathbb{Q}_p$  is the field of the *p-adic numbers*.

$$\mathbb{Q}_p^\times = \{\sum_{i=k}^{\infty} a_i p^i \mid k \in \mathbb{Z}, a_i \in \{0, 1, \dots, p-1\}, a_k \neq 0\}.$$

# Definitions

Let  $p$  be a prime number.  $\mathbb{Q}_p$  is the field of the  $p$ -adic numbers.

$$\mathbb{Q}_p^\times = \{\sum_{i=k}^{\infty} a_i p^i \mid k \in \mathbb{Z}, a_i \in \{0, 1, \dots, p-1\}, a_k \neq 0\}.$$

$$\text{ord} : \mathbb{Q}_p \rightarrow \mathbb{Z} \cup \{\infty\} : a \mapsto \begin{cases} k & \text{if } a = \sum_{i=k}^{\infty} a_i p^i; \\ \infty & \text{if } a = 0. \end{cases}$$

# Definitions

Let  $p$  be a prime number.  $\mathbb{Q}_p$  is the field of the  $p$ -adic numbers.

$$\mathbb{Q}_p^\times = \{\sum_{i=k}^{\infty} a_i p^i \mid k \in \mathbb{Z}, a_i \in \{0, 1, \dots, p-1\}, a_k \neq 0\}.$$

$$\text{ord} : \mathbb{Q}_p \rightarrow \mathbb{Z} \cup \{\infty\} : a \mapsto \begin{cases} k & \text{if } a = \sum_{i=k}^{\infty} a_i p^i; \\ \infty & \text{if } a = 0. \end{cases}$$

For each  $m \in \mathbb{N}_0$ ,

$$\text{ac}_m : \mathbb{Q}_p \rightarrow \mathbb{Z}_p / p^m \mathbb{Z}_p : a \mapsto \begin{cases} \sum_{i=0}^{m-1} a_{i+k} p^i & \text{if } a = \sum_{i=k}^{\infty} a_i p^i; \\ 0 & \text{if } a = 0. \end{cases}$$

# Definitions

Let  $p$  be a prime number.  $\mathbb{Q}_p$  is the field of the  $p$ -adic numbers.

$$\mathbb{Q}_p^\times = \{\sum_{i=k}^{\infty} a_i p^i \mid k \in \mathbb{Z}, a_i \in \{0, 1, \dots, p-1\}, a_k \neq 0\}.$$

$$\text{ord} : \mathbb{Q}_p \rightarrow \mathbb{Z} \cup \{\infty\} : a \mapsto \begin{cases} k & \text{if } a = \sum_{i=k}^{\infty} a_i p^i; \\ \infty & \text{if } a = 0. \end{cases}$$

For each  $m \in \mathbb{N}_0$ ,

$$\text{ac}_m : \mathbb{Q}_p \rightarrow \mathbb{Z}_p / p^m \mathbb{Z}_p : a \mapsto \begin{cases} \sum_{i=0}^{m-1} a_{i+k} p^i & \text{if } a = \sum_{i=k}^{\infty} a_i p^i; \\ 0 & \text{if } a = 0. \end{cases}$$

Clopen balls  $B_\gamma(a) := \{x \in \mathbb{Q}_p \mid \text{ord}(x - a) \geq \gamma\}$ ,  $\gamma \in \mathbb{Z}$ .

# Semialgebraic sets

Let  $K$  be a field and  $\mathcal{L}_{\text{ring}} = \{+, -, \cdot, 0, 1\}$ .

## Definition

The *semialgebraic sets* in  $K$  are the  $\mathcal{L}_{\text{ring}}$ -definable subsets of  $K$ .

# Semialgebraic sets

Let  $K$  be a field and  $\mathcal{L}_{\text{ring}} = \{+, -, \cdot, 0, 1\}$ .

## Definition

The *semialgebraic sets* in  $K$  are the  $\mathcal{L}_{\text{ring}}$ -definable subsets of  $K$ .

## *o-minimality*

- $K$  is a *real closed field* if it is  $\mathcal{L}_{\text{ring}}$ -elementary equivalent to  $\mathbb{R}$ .
- $x < y \Leftrightarrow \exists z(x + z^2 = y)$ .
- $\mathcal{L}_{\text{or}} := \mathcal{L}_{\text{ring}} \cup \{<\}$ .
- $(K, \mathcal{L}_{\text{or}})$  has QE.

# Semialgebraic sets

Let  $K$  be a field and  $\mathcal{L}_{\text{ring}} = \{+, -, \cdot, 0, 1\}$ .

## Definition

The *semialgebraic sets* in  $K$  are the  $\mathcal{L}_{\text{ring}}$ -definable subsets of  $K$ .

### *o-minimality*

- $K$  is a *real closed field* if it is  $\mathcal{L}_{\text{ring}}$ -elementary equivalent to  $\mathbb{R}$ .
- $x < y \Leftrightarrow \exists z(x + z^2 = y)$ .
- $\mathcal{L}_{\text{or}} := \mathcal{L}_{\text{ring}} \cup \{<\}$ .
- $(K, \mathcal{L}_{\text{or}})$  has QE.

### *P-minimality*

- $K$  is a  *$p$ -adically closed field* if it is  $\mathcal{L}_{\text{ring}}$ -elementary equivalent to a finite field extension of  $\mathbb{Q}_p$ .
- $x \in P_k \Leftrightarrow \exists y \neq 0(y^k = x)$ .
- $\mathcal{L}_{\text{Mac}} := \mathcal{L}_{\text{ring}} \cup \{P_k\}_{k>0}$ .
- $(K, \mathcal{L}_{\text{Mac}})$  has QE. (Prestel & Roquette, '84)

# Examples of semialgebraic sets

$X \subseteq \mathbb{R}$  semialgebraic  $\Rightarrow X$  is a Boolean combination of sets of the form

$$\{x \in \mathbb{R} \mid f(x) = 0\} \quad \{x \in \mathbb{R} \mid g(x) > 0\}.$$



# Examples of semialgebraic sets

$X \subseteq \mathbb{R}$  semialgebraic  $\Rightarrow X$  is a Boolean combination of sets of the form

$$\{x \in \mathbb{R} \mid f(x) = 0\} \quad \{x \in \mathbb{R} \mid g(x) > 0\}.$$

$X \subseteq \mathbb{Q}_p$  semialgebraic  $\Rightarrow X$  is a Boolean combination of sets of the form

$$\{x \in \mathbb{Q}_p \mid f(x) = 0\} \quad \{x \in \mathbb{Q}_p \mid g(x) \in P_k\}.$$

# Examples of semialgebraic sets

$X \subseteq \mathbb{R}$  semialgebraic  $\Rightarrow X$  is a Boolean combination of sets of the form


$$\{x \in \mathbb{R} \mid f(x) = 0\} \quad \{x \in \mathbb{R} \mid g(x) > 0\}.$$

$X \subseteq \mathbb{Q}_p$  semialgebraic  $\Rightarrow X$  is a Boolean combination of sets of the form

$$\{x \in \mathbb{Q}_p \mid f(x) = 0\} \quad \{x \in \mathbb{Q}_p \mid g(x) \in P_k\}.$$

Instead of  $(P_k)_{k>0}$  we prefer  $(Q_{m,n})_{m,n>0}$ , where

$$Q_{m,n} := \{x \in \mathbb{Q}_p^\times \mid \text{ord}(x) \equiv 0 \pmod n, \text{ac}_m(x) = 1\}.$$



# $\mathcal{o}$ -minimality and $\mathcal{P}$ -minimality

## Definition

Let  $\mathcal{L} \supseteq \mathcal{L}_{\text{ring}}$  and let  $K$  be a real closed field. The structure  $(K, \mathcal{L})$  is  *$\mathcal{o}$ -minimal* if every  $\mathcal{L}$ -definable subset  $X \subseteq K$  is also  $\mathcal{L}_{\text{ring}}$ -definable.

# $\mathcal{o}$ -minimality and $P$ -minimality

## Definition

Let  $\mathcal{L} \supseteq \mathcal{L}_{\text{ring}}$  and let  $K$  be a real closed field. The structure  $(K, \mathcal{L})$  is  *$\mathcal{o}$ -minimal* if every  $\mathcal{L}$ -definable subset  $X \subseteq K$  is also  $\mathcal{L}_{\text{ring}}$ -definable.

## Definition

Let  $\mathcal{L} \supseteq \mathcal{L}_{\text{ring}}$  and let  $K$  be a  $p$ -adically closed field. The structure  $(K, \mathcal{L})$  is  *$P$ -minimal* if for every  $(K', \mathcal{L}) \equiv (K, \mathcal{L})$ , every  $\mathcal{L}$ -definable subset  $X \subseteq K'$  is also  $\mathcal{L}_{\text{ring}}$ -definable.

# Classical cells


# Classical cells



# Classical cells




(0)-cell.




(1)-cell.

# Classical cells






# Classical cells



# Classical cells



# Classical cells

$$C = \left\{ (s, x) \in S \times K \mid \begin{array}{l} \text{ord}(\alpha(s)) \square_1 \text{ord}(x - \sigma(s)) \square_2 \text{ord}(\beta(s)), \\ x - \sigma(s) \in \lambda Q_{m,n} \end{array} \right\}$$

$S$  is a definable set,  $\alpha, \beta : S \rightarrow K^\times$  and  $\sigma : S \rightarrow K$  are definable functions,  $\square_i$  can be either  $<$  or 'no condition',  $n, m \in \mathbb{N}_0$  and  $\lambda \in K$ .

We call  $\sigma$  the *center* of the cell.

# Classical cells

$$C = \left\{ (s, x) \in S \times K \mid \begin{array}{l} \text{ord}(\alpha(s)) \square_1 \text{ord}(x - \sigma(s)) \square_2 \text{ord}(\beta(s)), \\ x - \sigma(s) \in \lambda Q_{m,n} \end{array} \right\}$$


$S$  is a definable set,  $\alpha, \beta : S \rightarrow K^\times$  and  $\sigma : S \rightarrow K$  are definable functions,  $\square_i$  can be either  $<$  or 'no condition',  $n, m \in \mathbb{N}_0$  and  $\lambda \in K$ .

We call  $\sigma$  the *center* of the cell.

$\sigma(s)$



$\lambda = 0.$



$\lambda \neq 0$  and  $\square_1 = \square_2 = <.$

# Definable Skolem functions

## Definition

A structure  $(K, \mathcal{L})$  has *definable Skolem functions* if for every definable  $X \subseteq K^{r+1}$  there exists a definable section  $\sigma : \pi(X) \rightarrow K$ . This means that  $(x, \sigma(x)) \in X$  for all  $x \in \pi(X)$ . By  $\pi : K^{r+1} \rightarrow K^r$  we mean the projection onto the first  $r$  coordinates.

## Theorem (Mourgues, '09)

A  $P$ -minimal structure  $(K, \mathcal{L})$  has cell decomposition (with classical cells) if and only if it has definable Skolem functions.

# Definable Skolem functions

## Definition

A structure  $(K, \mathcal{L})$  has *definable Skolem functions* if for every definable  $X \subseteq K^{r+1}$  there exists a definable section  $\sigma : \pi(X) \rightarrow K$ . This means that  $(x, \sigma(x)) \in X$  for all  $x \in \pi(X)$ . By  $\pi : K^{r+1} \rightarrow K^r$  we mean the projection onto the first  $r$  coordinates.

## Theorem (Mourgues, '09)

A  $P$ -minimal structure  $(K, \mathcal{L})$  has cell decomposition (with classical cells) if and only if it has definable Skolem functions.

## Remarks:


- All  $\mathcal{o}$ -minimal structures  $(K, \mathcal{L})$  have definable Skolem functions (and cell decomposition).
- Cubides and Nguyen have found a  $P$ -minimal structure that does not have definable Skolem functions!

# Centers of cells

Cell condition:  $C(s, y, x) =$   
 $\text{ord}(\alpha(s)) \square_1 \text{ord}(x - y) \square_2 \text{ord}(\beta(s))$   
 $\wedge (x - y) \in \lambda Q_{m,n} \wedge s \in S.$

# Centers of cells

Cell condition:  $C(s, y, x) = \text{ord}(\alpha(s)) \sqcap_1 \text{ord}(x - y) \sqcap_2 \text{ord}(\beta(s))$   
 $\wedge (x - y) \in \lambda Q_{m,n} \wedge s \in S.$




For any function  $\sigma : S \rightarrow K$ ,  
 we define

$$C^\sigma := \{(s, x) \in S \times K \mid C(s, \sigma(s), x)\}.$$



# Centers of cells

Cell condition:  $C(s, y, x) = \text{ord}(\alpha(s)) \square_1 \text{ord}(x - y) \square_2 \text{ord}(\beta(s))$   
 $\wedge (x - y) \in \lambda Q_{m,n} \wedge s \in S.$




For any function  $\sigma : S \rightarrow K$ ,  
 we define

$$C^\sigma := \{(s, x) \in S \times K \mid C(s, \sigma(s), x)\}.$$

We have  $C^\sigma = C^{\sigma'}$ .

# Centers of cells

Cell condition:  $C(s, y, x) = \text{ord}(\alpha(s)) \square_1 \text{ord}(x - y) \square_2 \text{ord}(\beta(s))$   
 $\wedge (x - y) \in \lambda Q_{m,n} \wedge s \in S.$



For any function  $\sigma : S \rightarrow K$ ,  
 we define

$$C^\sigma := \{(s, x) \in S \times K \mid C(s, \sigma(s), x)\}.$$


We have  $C^\sigma = C^{\sigma'}$ .

The possible centers are the  
 sections of

$$\Sigma := \bigcup_{s \in S} \{s\} \times B_{\rho_{\max}(s) + m(\sigma(s))} \subseteq S \times K.$$


Then  $C^\sigma = \{(s, x) \mid \exists \sigma' \in \Sigma_s C(s, \sigma', x)\}.$

# Inseparably clustered cells



$$X = C^{\sigma_1} \sqcup \dots \sqcup C^{\sigma_k}.$$

# Inseparably clustered cells



$$X = C^{\sigma_1} \sqcup \dots \sqcup C^{\sigma_k}.$$

There exists a definable  $\Sigma \subseteq S \times K$  for which each  $\Sigma_s$  is a union of  $k$  balls of the same size and such that

$$X = \{(s, x) \mid \exists \sigma' \in \Sigma_s C(s, \sigma', x)\}.$$