

Nash groups

Anand Pillay

Konstanz, July 2016

Nash groups

Anand Pillay

Konstanz, July 2016

Background

- ▶ The original motivation is model theoretic: the description/classification of definable groups in the structure $(\mathbb{R}, +, \times)$, up to definable isomorphism.

Background

- ▶ The original motivation is model theoretic: the description/classification of definable groups in the structure $(\mathbb{R}, +, \times)$, up to definable isomorphism.
- ▶ Definable equals semialgebraic, so this is the classification of (real) semialgebraic groups up to semialgebraic isomorphism.

Background

- ▶ The original motivation is model theoretic: the description/classification of definable groups in the structure $(\mathbb{R}, +, \times)$, up to definable isomorphism.
- ▶ Definable equals semialgebraic, so this is the classification of (real) semialgebraic groups up to semialgebraic isomorphism.
- ▶ We are not a priori imposing any topological structure on G , or making any assumptions about continuity of the group operation (but see later).

Background

- ▶ The original motivation is model theoretic: the description/classification of definable groups in the structure $(\mathbb{R}, +, \times)$, up to definable isomorphism.
- ▶ Definable equals semialgebraic, so this is the classification of (real) semialgebraic groups up to semialgebraic isomorphism.
- ▶ We are not a priori imposing any topological structure on G , or making any assumptions about continuity of the group operation (but see later).
- ▶ But we usually assume G to be semialgebraically connected in the sense of having no proper semialgebraic subgroup of finite index.

Background

- ▶ The original motivation is model theoretic: the description/classification of definable groups in the structure $(\mathbb{R}, +, \times)$, up to definable isomorphism.
- ▶ Definable equals semialgebraic, so this is the classification of (real) semialgebraic groups up to semialgebraic isomorphism.
- ▶ We are not a priori imposing any topological structure on G , or making any assumptions about continuity of the group operation (but see later).
- ▶ But we usually assume G to be semialgebraically connected in the sense of having no proper semialgebraic subgroup of finite index.
- ▶ A typical example is the interval $[0, 1)$ equipped with addition modulo 1, which is a semialgebraic group in the above sense, but not a topological group in the naive sense.

Real algebraic groups

- By a real algebraic group I will mean the (topological) connected component of the group $H(\mathbb{R})$ of real points of a (connected) algebraic group H defined over \mathbb{R} .

Real algebraic groups

- ▶ By a real algebraic group I will mean the (topological) connected component of the group $H(\mathbb{R})$ of real points of a (connected) algebraic group H defined over \mathbb{R} .
- ▶ An example is $GL_n(\mathbb{R})^+$, the group of nonsingular $n \times n$ matrices over \mathbb{R} with positive determinant, which is obviously a semialgebraic group in the “naive” sense above.

Real algebraic groups

- ▶ By a real algebraic group I will mean the (topological) connected component of the group $H(\mathbb{R})$ of real points of a (connected) algebraic group H defined over \mathbb{R} .
- ▶ An example is $GL_n(\mathbb{R})^+$, the group of nonsingular $n \times n$ matrices over \mathbb{R} with positive determinant, which is obviously a semialgebraic group in the “naive” sense above.
- ▶ In fact any real algebraic group (such as $A(\mathbb{R})^0$, A an abelian variety over \mathbb{R}) “is” a semialgebraically connected semialgebraic group in the sense above.

Real algebraic groups

- ▶ By a real algebraic group I will mean the (topological) connected component of the group $H(\mathbb{R})$ of real points of a (connected) algebraic group H defined over \mathbb{R} .
- ▶ An example is $GL_n(\mathbb{R})^+$, the group of nonsingular $n \times n$ matrices over \mathbb{R} with positive determinant, which is obviously a semialgebraic group in the “naive” sense above.
- ▶ In fact any real algebraic group (such as $A(\mathbb{R})^0$, A an abelian variety over \mathbb{R}) “is” a semialgebraically connected semialgebraic group in the sense above.
- ▶ This can be seen model theoretically via elimination of imaginaries in $(\mathbb{R}, +, \times)$. But we will see later on a more interesting reason.

Real algebraic groups

- ▶ By a real algebraic group I will mean the (topological) connected component of the group $H(\mathbb{R})$ of real points of a (connected) algebraic group H defined over \mathbb{R} .
- ▶ An example is $GL_n(\mathbb{R})^+$, the group of nonsingular $n \times n$ matrices over \mathbb{R} with positive determinant, which is obviously a semialgebraic group in the “naive” sense above.
- ▶ In fact any real algebraic group (such as $A(\mathbb{R})^0$, A an abelian variety over \mathbb{R}) “is” a semialgebraically connected semialgebraic group in the sense above.
- ▶ This can be seen model theoretically via elimination of imaginaries in $(\mathbb{R}, +, \times)$. But we will see later on a more interesting reason.
- ▶ In any case we want to classify semialgebraic groups in terms of real algebraic groups and various constructions.

Nash manifolds I

- ▶ Nash manifolds originated in work of John Nash (1952) where he introduced “real algebraic manifolds”, which in current parlance are “compact affine Nash manifolds”. (The complex analogues are the Moishezon compact complex manifolds.)

Nash manifolds I

- ▶ Nash manifolds originated in work of John Nash (1952) where he introduced “real algebraic manifolds”, which in current parlance are “compact affine Nash manifolds”. (The complex analogues are the Moishezon compact complex manifolds.)
- ▶ Nash manifolds appear again in work of Artin and Mazur (1965) in connection with dynamical systems, and were systematically studied in books by Shiota and Coste-Roy in the 80's.

Nash manifolds I

- ▶ Nash manifolds originated in work of John Nash (1952) where he introduced “real algebraic manifolds”, which in current parlance are “compact affine Nash manifolds”. (The complex analogues are the Moishezon compact complex manifolds.)
- ▶ Nash manifolds appear again in work of Artin and Mazur (1965) in connection with dynamical systems, and were systematically studied in books by Shiota and Coste-Roy in the 80's.
- ▶ Firstly if U is an open semialgebraic subset of \mathbb{R}^n , then a function $f : U \rightarrow \mathbb{R}$ is said to be Nash if f is both analytic and semialgebraic, which essentially means that there is a polynomial $P(\bar{x}, y)$ such that $P(\bar{x}, f(\bar{x})) = 0$ for all $\bar{x} \in U$. (e.g. $f(x) = +\sqrt{x}$ on $(0, \infty)$).

Nash manifolds I

- ▶ Nash manifolds originated in work of John Nash (1952) where he introduced “real algebraic manifolds”, which in current parlance are “compact affine Nash manifolds”. (The complex analogues are the Moishezon compact complex manifolds.)
- ▶ Nash manifolds appear again in work of Artin and Mazur (1965) in connection with dynamical systems, and were systematically studied in books by Shiota and Coste-Roy in the 80's.
- ▶ Firstly if U is an open semialgebraic subset of \mathbb{R}^n , then a function $f : U \rightarrow \mathbb{R}$ is said to be Nash if f is both analytic and semialgebraic, which essentially means that there is a polynomial $P(\bar{x}, y)$ such that $P(\bar{x}, f(\bar{x})) = 0$ for all $\bar{x} \in U$. (e.g. $f(x) = +\sqrt{x}$ on $(0, \infty)$).
- ▶ Semialgebraic functions are piecewise Nash.

Nash manifolds II

- ▶ An (n -dimensional) Nash manifold is a Hausdorff topological space X with a covering by *finitely many* open sets V_i , each homeomorphic via some f_i to an open semialgebraic subset U_i of \mathbb{R}^n such that the transition maps $f_i \circ f_j^{-1}$ between the open semialgebraic sets $f_j(V_i \cap V_j)$ and $f_i(V_i \cap V_j)$ are Nash (i.e each coordinate is Nash).

Nash manifolds II

- ▶ An (n -dimensional) Nash manifold is a Hausdorff topological space X with a covering by *finitely many* open sets V_i , each homeomorphic via some f_i to an open semialgebraic subset U_i of \mathbb{R}^n such that the transition maps $f_i \circ f_j^{-1}$ between the open semialgebraic sets $f_j(V_i \cap V_j)$ and $f_i(V_i \cap V_j)$ are Nash (i.e each coordinate is Nash).
- ▶ Intuitively a Nash manifold is a *definable* or *semialgebraic* real analytic manifold.

Nash manifolds II

- ▶ An (n -dimensional) Nash manifold is a Hausdorff topological space X with a covering by *finitely many* open sets V_i , each homeomorphic via some f_i to an open semialgebraic subset U_i of \mathbb{R}^n such that the transition maps $f_i \circ f_j^{-1}$ between the open semialgebraic sets $f_j(V_i \cap V_j)$ and $f_i(V_i \cap V_j)$ are Nash (i.e each coordinate is Nash).
- ▶ Intuitively a Nash manifold is a *definable* or *semialgebraic* real analytic manifold.
- ▶ By a *locally Nash manifold* we mean as above, except that we allow the possibility of countably many V_i . And again this is intuitively a “locally definable” or “ind-definable” real analytic manifold.

Nash manifolds II

- ▶ An (n -dimensional) Nash manifold is a Hausdorff topological space X with a covering by *finitely many* open sets V_i , each homeomorphic via some f_i to an open semialgebraic subset U_i of \mathbb{R}^n such that the transition maps $f_i \circ f_j^{-1}$ between the open semialgebraic sets $f_j(V_i \cap V_j)$ and $f_i(V_i \cap V_j)$ are Nash (i.e. each coordinate is Nash).
- ▶ Intuitively a Nash manifold is a *definable* or *semialgebraic* real analytic manifold.
- ▶ By a *locally Nash manifold* we mean as above, except that we allow the possibility of countably many V_i . And again this is intuitively a “locally definable” or “ind-definable” real analytic manifold.
- ▶ There is a natural notion of a Nash map between Nash manifolds, and a Nash group is a Nash manifold X with group operation given by a Nash map $X \times X \rightarrow X$ (i.e. a group object in the category of Nash manifolds).

Nash manifolds III

- ▶ Likewise for local versions (locally Nash group).

Nash manifolds III

- ▶ Likewise for local versions (locally Nash group).
- ▶ A Nash manifold is said to be affine if it has a Nash embedding in some \mathbb{R}^n .

Nash manifolds III

- ▶ Likewise for local versions (locally Nash group).
- ▶ A Nash manifold is said to be affine if it has a Nash embedding in some \mathbb{R}^n .
- ▶ Somewhat paradoxically real projective space is affine as a Nash manifold, via stereographic projection, whereby any real nonsingular quasiprojective variety (such as $H(\mathbb{R})$ for H any algebraic group over \mathbb{R}) is affine as a Nash manifold.

Nash manifolds III

- ▶ Likewise for local versions (locally Nash group).
- ▶ A Nash manifold is said to be affine if it has a Nash embedding in some \mathbb{R}^n .
- ▶ Somewhat paradoxically real projective space is affine as a Nash manifold, via stereographic projection, whereby any real nonsingular quasiprojective variety (such as $H(\mathbb{R})$ for H any algebraic group over \mathbb{R}) is affine as a Nash manifold.
- ▶ Conversely it was proved (Tognoli ?) that any affine Nash manifold is Nash isomorphic to a nonsingular real algebraic variety.

Nash manifolds III

- ▶ Likewise for local versions (locally Nash group).
- ▶ A Nash manifold is said to be affine if it has a Nash embedding in some \mathbb{R}^n .
- ▶ Somewhat paradoxically real projective space is affine as a Nash manifold, via stereographic projection, whereby any real nonsingular quasiprojective variety (such as $H(\mathbb{R})$ for H any algebraic group over \mathbb{R}) is affine as a Nash manifold.
- ▶ Conversely it was proved (Tognoli ?) that any affine Nash manifold is Nash isomorphic to a nonsingular real algebraic variety.
- ▶ \mathbb{R}/\mathbb{Z} has naturally the structure of a Nash manifold (in fact group): $[0, 1]$ with $0, 1$ identified, and addition modulo 1 but it is non affine.

Nash manifolds III

- ▶ Likewise for local versions (locally Nash group).
- ▶ A Nash manifold is said to be affine if it has a Nash embedding in some \mathbb{R}^n .
- ▶ Somewhat paradoxically real projective space is affine as a Nash manifold, via stereographic projection, whereby any real nonsingular quasiprojective variety (such as $H(\mathbb{R})$ for H any algebraic group over \mathbb{R}) is affine as a Nash manifold.
- ▶ Conversely it was proved (Tognoli ?) that any affine Nash manifold is Nash isomorphic to a nonsingular real algebraic variety.
- ▶ \mathbb{R}/\mathbb{Z} has naturally the structure of a Nash manifold (in fact group): $[0, 1]$ with $0, 1$ identified, and addition modulo 1 but it is non affine.
- ▶ In general little is known about arbitrary (not necessarily affine) Nash manifolds but there is a chance to classify the Nash groups.

Nash groups I

We mention some early results:

Theorem 0.1

(P 1986) Any semialgebraic group G can be semialgebraically equipped with the structure of a Nash group (in particular a Lie group) unique up to Nash isomorphism. The category of semialgebraic groups coincides with that of Nash groups.

Nash groups I

We mention some early results:

Theorem 0.1

(P 1986) Any semialgebraic group G can be semialgebraically equipped with the structure of a Nash group (in particular a Lie group) unique up to Nash isomorphism. The category of semialgebraic groups coincides with that of Nash groups.

We have five categories: real algebraic groups \subset affine Nash groups \subset Nash groups \subset locally Nash groups \subset Lie groups.

Nash groups I

We mention some early results:

Theorem 0.1

(P 1986) Any semialgebraic group G can be semialgebraically equipped with the structure of a Nash group (in particular a Lie group) unique up to Nash isomorphism. The category of semialgebraic groups coincides with that of Nash groups.

We have five categories: real algebraic groups \subset affine Nash groups \subset Nash groups \subset locally Nash groups \subset Lie groups.

Theorem 0.2

(Hrushovski-P, 1994, with model theory proof) Any (even locally) Nash group G is Nash, locally isomorphic, to a real algebraic group H in the sense that there is Nash isomorphism between open semialgebraic neighbourhoods of the identity of G and of H .

Nash groups II

- ▶ Earlier, in the late 80's Madden-Stanton had noted the one-dimensional case of Theorem 0.2 and used it to classify one-dimensional Nash groups (included in our main result, to be stated later). In particular there are many distinct incarnations of the 1-dimensional compact Lie group S^1 as Nash groups.

Nash groups II

- ▶ Earlier, in the late 80's Madden-Stanton had noted the one-dimensional case of Theorem 0.2 and used it to classify one-dimensional Nash groups (included in our main result, to be stated later). In particular there are many distinct incarnations of the 1-dimensional compact Lie group S^1 as Nash groups.
- ▶ We used Theorem 0.2 to prove the following (Hrushovski-P, 1994, with corrected proof 2012):

Theorem 0.3

The affine Nash groups are precisely the finite covers of real algebraic groups.

Nash groups II

- ▶ Earlier, in the late 80's Madden-Stanton had noted the one-dimensional case of Theorem 0.2 and used it to classify one-dimensional Nash groups (included in our main result, to be stated later). In particular there are many distinct incarnations of the 1-dimensional compact Lie group S^1 as Nash groups.
- ▶ We used Theorem 0.2 to prove the following (Hrushovski-P, 1994, with corrected proof 2012):

Theorem 0.3

The affine Nash groups are precisely the finite covers of real algebraic groups.

Note this already takes us out of the category of algebraic groups: the proper finite covers of $SL(2, \mathbb{R})$ are neither real algebraic nor linear.

Nash groups III

We have the following easy fact about locally Nash groups:

Theorem 0.4

The class of locally Nash groups is closed under taking universal covers and quotients by discrete subgroups.

Nash groups III

We have the following easy fact about locally Nash groups:

Theorem 0.4

The class of locally Nash groups is closed under taking universal covers and quotients by discrete subgroups.

We conclude from Theorem 0.2 a classification of *locally Nash groups*.

Nash groups III

We have the following easy fact about locally Nash groups:

Theorem 0.4

The class of locally Nash groups is closed under taking universal covers and quotients by discrete subgroups.

We conclude from Theorem 0.2 a classification of *locally Nash groups*.

Corollary 0.5

The locally Nash groups are, up to local Nash isomorphism precisely the quotients of universal covers of real algebraic groups by discrete subgroups.

Nash groups IV

The following easy result of Shiota is useful and yields some quick conclusions:

Theorem 0.6

Let X and Y be Nash manifolds with Y affine and let $f : X \rightarrow Y$ be a locally Nash map. Then f is a Nash map.

Corollary 0.7

- ▶ *(i) Any simply connected Nash group is a finite cover of a real algebraic group.*
- ▶ *(ii) Any torsion-free Nash group is a real algebraic group.*

Proof. ??

Problem

- ▶ To answer the original problem of classifying Nash groups up to Nash isomorphism, by Theorem 0.5, we must carry out the following:

Problem

- ▶ To answer the original problem of classifying Nash groups up to Nash isomorphism, by Theorem 0.5, we must carry out the following:
- ▶ (i) For any real algebraic group G , describe those discrete subgroups Γ of the universal cover \tilde{G} such that the locally Nash group \tilde{G}/Γ has a compatible structure of Nash group. (For example the universal cover of $SL(2, \mathbb{R})$ does not itself have the structure of a Nash group.)

Problem

- ▶ To answer the original problem of classifying Nash groups up to Nash isomorphism, by Theorem 0.5, we must carry out the following:
- ▶ (i) For any real algebraic group G , describe those discrete subgroups Γ of the universal cover \tilde{G} such that the locally Nash group \tilde{G}/Γ has a compatible structure of Nash group. (For example the universal cover of $SL(2, \mathbb{R})$ does not itself have the structure of a Nash group.)
- ▶ (ii) Classify the resulting Nash groups up to Nash isomorphism.

Problem

- ▶ To answer the original problem of classifying Nash groups up to Nash isomorphism, by Theorem 0.5, we must carry out the following:
- ▶ (i) For any real algebraic group G , describe those discrete subgroups Γ of the universal cover \tilde{G} such that the locally Nash group \tilde{G}/Γ has a compatible structure of Nash group. (For example the universal cover of $SL(2, \mathbb{R})$ does not itself have the structure of a Nash group.)
- ▶ (ii) Classify the resulting Nash groups up to Nash isomorphism.
- ▶ We will answer question (i) in the commutative case, and give some examples showing some subtleties around (ii) compared to the one-dimensional case.

Commutative real algebraic groups

- ▶ We need to know something of the structure of commutative real algebraic groups G .

Commutative real algebraic groups

- ▶ We need to know something of the structure of commutative real algebraic groups G .
- ▶ A basic fact is that there is an exact sequence (of real algebraic groups) $0 \rightarrow L \rightarrow G \rightarrow C \rightarrow 0$, where

Commutative real algebraic groups

- ▶ We need to know something of the structure of commutative real algebraic groups G .
- ▶ A basic fact is that there is an exact sequence (of real algebraic groups) $0 \rightarrow L \rightarrow G \rightarrow C \rightarrow 0$, where
- ▶ L is a finite product of copies of $(\mathbb{R}, +)$ and $(\mathbb{R}_{>0}, \times)$ and C is compact.

Commutative real algebraic groups

- ▶ We need to know something of the structure of commutative real algebraic groups G .
- ▶ A basic fact is that there is an exact sequence (of real algebraic groups) $0 \rightarrow L \rightarrow G \rightarrow C \rightarrow 0$, where
- ▶ L is a finite product of copies of $(\mathbb{R}, +)$ and $(\mathbb{R}_{>0}, \times)$ and C is compact.
- ▶ C itself is understood by an exact sequence $0 \rightarrow C_1 \rightarrow C \rightarrow C_2 \rightarrow 0$ where C_1 is a product of copies of $SO(2, \mathbb{R})$ and C_2 a real abelian variety (i.e. its topological connected component).

Commutative real algebraic groups

- ▶ We need to know something of the structure of commutative real algebraic groups G .
- ▶ A basic fact is that there is an exact sequence (of real algebraic groups) $0 \rightarrow L \rightarrow G \rightarrow C \rightarrow 0$, where
- ▶ L is a finite product of copies of $(\mathbb{R}, +)$ and $(\mathbb{R}_{>0}, \times)$ and C is compact.
- ▶ C itself is understood by an exact sequence $0 \rightarrow C_1 \rightarrow C \rightarrow C_2 \rightarrow 0$ where C_1 is a product of copies of $SO(2, \mathbb{R})$ and C_2 a real abelian variety (i.e. its topological connected component).
- ▶ This passes to universal covers: in particular we have $0 \rightarrow L \rightarrow \tilde{G} \rightarrow_{\pi} \tilde{C} \rightarrow 0$.

Result

We have, with above notation the solution to (i) in the commutative case:

Theorem 0.8

Let G be a commutative real algebraic group, and let Γ be a discrete subgroup of \tilde{G} (maybe trivial). Then \tilde{G}/Γ has a (compatible) structure of Nash group just if $\tilde{C}/\pi(\Gamma)$ is compact.

Result

We have, with above notation the solution to (i) in the commutative case:

Theorem 0.8

Let G be a commutative real algebraic group, and let Γ be a discrete subgroup of \tilde{G} (maybe trivial). Then \tilde{G}/Γ has a (compatible) structure of Nash group just if $\tilde{C}/\pi(\Gamma)$ is compact.

- Some ingredients in the proof are:

Result

We have, with above notation the solution to (i) in the commutative case:

Theorem 0.8

Let G be a commutative real algebraic group, and let Γ be a discrete subgroup of \tilde{G} (maybe trivial). Then \tilde{G}/Γ has a (compatible) structure of Nash group just if $\tilde{C}/\pi(\Gamma)$ is compact.

- ▶ Some ingredients in the proof are:
- ▶ The (obvious) fact that a compact locally Nash group is Nash.

Result

We have, with above notation the solution to (i) in the commutative case:

Theorem 0.8

Let G be a commutative real algebraic group, and let Γ be a discrete subgroup of \tilde{G} (maybe trivial). Then \tilde{G}/Γ has a (compatible) structure of Nash group just if $\tilde{C}/\pi(\Gamma)$ is compact.

- ▶ Some ingredients in the proof are:
- ▶ The (obvious) fact that a compact locally Nash group is Nash.
- ▶ The universal cover of either $SO(2, \mathbb{R})$ or a real abelian variety does not have the compatible structure of a Nash group (a point that seems to have been overlooked by Madden-Stanton).

We have, with above notation the solution to (i) in the commutative case:

Theorem 0.8

Let G be a commutative real algebraic group, and let Γ be a discrete subgroup of \tilde{G} (maybe trivial). Then \tilde{G}/Γ has a (compatible) structure of Nash group just if $\tilde{C}/\pi(\Gamma)$ is compact.

- ▶ Some ingredients in the proof are:
- ▶ The (obvious) fact that a compact locally Nash group is Nash.
- ▶ The universal cover of either $SO(2, \mathbb{R})$ or a real abelian variety does not have the compatible structure of a Nash group (a point that seems to have been overlooked by Madden-Stanton).
- ▶ \tilde{G}/Γ has structure of Nash (definable) group if and only if \tilde{G} has a “definable fundamental region” for Γ .

Fundamental region I

- ▶ A fundamental region Ω of \tilde{G} for Γ is simply a neighbourhood Ω of 0 in \tilde{G} such that every coset of Γ in \tilde{G} meets Ω and both Ω and $\Omega + \Omega$ contain only finitely many elements of Γ .

Fundamental region I

- ▶ A fundamental region Ω of \tilde{G} for Γ is simply a neighbourhood Ω of 0 in \tilde{G} such that every coset of Γ in \tilde{G} meets Ω and both Ω and $\Omega + \Omega$ contain only finitely many elements of Γ .
- ▶ Such a fundamental region is *definable* if Ω is a definable subset of the locally definable \tilde{G} and the restriction of $+$ to Ω is definable.

Fundamental region I

- ▶ A fundamental region Ω of \tilde{G} for Γ is simply a neighbourhood Ω of 0 in \tilde{G} such that every coset of Γ in \tilde{G} meets Ω and both Ω and $\Omega + \Omega$ contain only finitely many elements of Γ .
- ▶ Such a fundamental region is *definable* if Ω is a definable subset of the locally definable \tilde{G} and the restriction of $+$ to Ω is definable.
- ▶ The choice of a definable fundamental region in \tilde{G} for Γ determines a Nash or definable group structure on \tilde{G}/Γ (in the obvious way; explain).

Fundamental region I

- ▶ A fundamental region Ω of \tilde{G} for Γ is simply a neighbourhood Ω of 0 in \tilde{G} such that every coset of Γ in \tilde{G} meets Ω and both Ω and $\Omega + \Omega$ contain only finitely many elements of Γ .
- ▶ Such a fundamental region is *definable* if Ω is a definable subset of the locally definable \tilde{G} and the restriction of $+$ to Ω is definable.
- ▶ The choice of a definable fundamental region in \tilde{G} for Γ determines a Nash or definable group structure on \tilde{G}/Γ (in the obvious way; explain).
- ▶ The interesting new phenomena, compared with the one-dimensional case, is that different choices of definable fundamental region may give non isomorphic Nash group structures on the same locally Nash group \tilde{G}/Γ .

Fundamental region II

- ▶ An example is $\tilde{G} = G = (\mathbb{R}, +) \times (\mathbb{R}_{>0}, \times)$.

Fundamental region II

- ▶ An example is $\tilde{G} = G = (\mathbb{R}, +) \times (\mathbb{R}_{>0}, \times)$.
- ▶ By Theorem 0.8 any quotient of G by a discrete subgroup has a Nash group structure.

Fundamental region II

- ▶ An example is $\tilde{G} = G = (\mathbb{R}, +) \times (\mathbb{R}_{>0}, \times)$.
- ▶ By Theorem 0.8 any quotient of G by a discrete subgroup has a Nash group structure.
- ▶ Let Γ be the cyclic subgroup of G generated by $(1, 2)$.

Fundamental region II

- ▶ An example is $\tilde{G} = G = (\mathbb{R}, +) \times (\mathbb{R}_{>0}, \times)$.
- ▶ By Theorem 0.8 any quotient of G by a discrete subgroup has a Nash group structure.
- ▶ Let Γ be the cyclic subgroup of G generated by $(1, 2)$.
- ▶ Then we have at least two definable fundamental regions in L for Γ :

Fundamental region II

- ▶ An example is $\tilde{G} = G = (\mathbb{R}, +) \times (\mathbb{R}_{>0}, \times)$.
- ▶ By Theorem 0.8 any quotient of G by a discrete subgroup has a Nash group structure.
- ▶ Let Γ be the cyclic subgroup of G generated by $(1, 2)$.
- ▶ Then we have at least two definable fundamental regions in L for Γ :
- ▶ $\Omega_1 = \{(x, y) : x \in \mathbb{R}, 1 \leq y \leq 2\}$.

Fundamental region II

- ▶ An example is $\tilde{G} = G = (\mathbb{R}, +) \times (\mathbb{R}_{>0}, \times)$.
- ▶ By Theorem 0.8 any quotient of G by a discrete subgroup has a Nash group structure.
- ▶ Let Γ be the cyclic subgroup of G generated by $(1, 2)$.
- ▶ Then we have at least two definable fundamental regions in L for Γ :
- ▶ $\Omega_1 = \{(x, y) : x \in \mathbb{R}, 1 \leq y \leq 2\}$.
- ▶ $\Omega_2 = \{(x, y) : 0 \leq x \leq 1, y \in \mathbb{R}_{>0}\}$.

Fundamental region II

- ▶ An example is $\tilde{G} = G = (\mathbb{R}, +) \times (\mathbb{R}_{>0}, \times)$.
- ▶ By Theorem 0.8 any quotient of G by a discrete subgroup has a Nash group structure.
- ▶ Let Γ be the cyclic subgroup of G generated by $(1, 2)$.
- ▶ Then we have at least two definable fundamental regions in L for Γ :
- ▶ $\Omega_1 = \{(x, y) : x \in \mathbb{R}, 1 \leq y \leq 2\}$.
- ▶ $\Omega_2 = \{(x, y) : 0 \leq x \leq 1, y \in \mathbb{R}_{>0}\}$.
- ▶ For the first choice G/Γ contains a definable copy of $(\mathbb{R}, +)$ but not of $(\mathbb{R}_{>0}, \times)$.

Fundamental region II

- ▶ An example is $\tilde{G} = G = (\mathbb{R}, +) \times (\mathbb{R}_{>0}, \times)$.
- ▶ By Theorem 0.8 any quotient of G by a discrete subgroup has a Nash group structure.
- ▶ Let Γ be the cyclic subgroup of G generated by $(1, 2)$.
- ▶ Then we have at least two definable fundamental regions in L for Γ :
- ▶ $\Omega_1 = \{(x, y) : x \in \mathbb{R}, 1 \leq y \leq 2\}$.
- ▶ $\Omega_2 = \{(x, y) : 0 \leq x \leq 1, y \in \mathbb{R}_{>0}\}$.
- ▶ For the first choice G/Γ contains a definable copy of $(\mathbb{R}, +)$ but not of $(\mathbb{R}_{>0}, \times)$.
- ▶ For the second choice it is the other way round, so they cannot be Nash isomorphic.