

# Definable Extension Theorems in O-minimal Structures

Matthias Aschenbrenner  
University of California, Los Angeles

**UCLA**

- 1 Nonconstructive nature of extension theorems
- 2 Definable extension theorems
- 3 Open questions

# Nonconstructive nature of extension theorems

## Question

Let  $f: A \rightarrow \mathbb{R}^n$  ( $A \subseteq \mathbb{R}^m$ ) be  $L$ -LIPSCHITZ, i.e.,

$$\|f(x) - f(y)\| \leq L \cdot \|x - y\| \quad \text{for all } x, y \in A.$$

Can one extend  $f$  to an  $L$ -LIPSCHITZ map  $\mathbb{R}^m \rightarrow \mathbb{R}^n$ ?

# Nonconstructive nature of extension theorems

## Question

Let  $f: A \rightarrow \mathbb{R}^n$  ( $A \subseteq \mathbb{R}^m$ ) be  $L$ -LIPSCHITZ, i.e.,

$$\|f(x) - f(y)\| \leq L \cdot \|x - y\| \quad \text{for all } x, y \in A.$$

Can one extend  $f$  to an  $L$ -LIPSCHITZ map  $\mathbb{R}^m \rightarrow \mathbb{R}^n$ ?

**MACSHANE & WHITNEY (1934):** yes if  $n = 1$

$$f_a(x) := f(a) + L \cdot \|x - a\| \quad (a \in A)$$

is  $L$ -LIPSCHITZ on  $\mathbb{R}^m$ , hence so is  $F(x) := \inf\{f_a(x) : a \in A\}$ ,  
and clearly  $F \upharpoonright A = f$ .

# Nonconstructive nature of extension theorems

## Question

Let  $f: A \rightarrow \mathbb{R}^n$  ( $A \subseteq \mathbb{R}^m$ ) be  $L$ -LIPSCHITZ, i.e.,

$$\|f(x) - f(y)\| \leq L \cdot \|x - y\| \quad \text{for all } x, y \in A.$$

Can one extend  $f$  to an  $L$ -LIPSCHITZ map  $\mathbb{R}^m \rightarrow \mathbb{R}^n$ ?

**MACSHANE & WHITNEY (1934):** yes if  $n = 1$

$$f_a(x) := f(a) + L \cdot \|x - a\| \quad (a \in A)$$

is  $L$ -LIPSCHITZ on  $\mathbb{R}^m$ , hence so is  $F(x) := \inf\{f_a(x) : a \in A\}$ ,  
and clearly  $F \upharpoonright A = f$ .

Applying this to the coordinate functions of  $f$  yields an  
 $L\sqrt{n}$ -LIPSCHITZ extension.

# Nonconstructive nature of extension theorems

## Question

Let  $f: A \rightarrow \mathbb{R}^n$  ( $A \subseteq \mathbb{R}^m$ ) be  $L$ -LIPSCHITZ, i.e.,

$$\|f(x) - f(y)\| \leq L \cdot \|x - y\| \quad \text{for all } x, y \in A.$$

Can one extend  $f$  to an  $L$ -LIPSCHITZ map  $\mathbb{R}^m \rightarrow \mathbb{R}^n$ ?

**MACSHANE & WHITNEY (1934):** yes if  $n = 1$

$$f_a(x) := f(a) + L \cdot \|x - a\| \quad (a \in A)$$

is  $L$ -LIPSCHITZ on  $\mathbb{R}^m$ , hence so is  $F(x) := \inf\{f_a(x) : a \in A\}$ ,  
and clearly  $F \upharpoonright A = f$ .

Applying this to the coordinate functions of  $f$  yields an  $L\sqrt{n}$ -LIPSCHITZ extension. But one can do better!

# Nonconstructive nature of extension theorems

KIRSZBRAUN (1934): *yes* for all  $n$

There always exists an  $L$ -LIPSCHITZ extension  $\mathbb{R}^m \rightarrow \mathbb{R}^n$  of  $f$ .

# Nonconstructive nature of extension theorems

KIRSZBRAUN (1934): *yes* for all  $n$

There always exists an  $L$ -LIPSCHITZ extension  $\mathbb{R}^m \rightarrow \mathbb{R}^n$  of  $f$ .

The usual proofs all use some sort of transfinite induction and:



# Nonconstructive nature of extension theorems

KIRSZBRAUN (1934): *yes* for all  $n$

There always exists an  $L$ -LIPSCHITZ extension  $\mathbb{R}^m \rightarrow \mathbb{R}^n$  of  $f$ .

The usual proofs all use some sort of transfinite induction and:

## Basic Extension Lemma

Let  $f: A \rightarrow \mathbb{R}^n$ ,  $A \subseteq \mathbb{R}^m$ , be 1-LIPSCHITZ, and let  $x \in \mathbb{R}^m \setminus A$ .

# Nonconstructive nature of extension theorems

KIRSZBRAUN (1934): *yes* for all  $n$

There always exists an  $L$ -LIPSCHITZ extension  $\mathbb{R}^m \rightarrow \mathbb{R}^n$  of  $f$ .

The usual proofs all use some sort of transfinite induction and:

## Basic Extension Lemma

Let  $f: A \rightarrow \mathbb{R}^n$ ,  $A \subseteq \mathbb{R}^m$ , be 1-LIPSCHITZ, and let  $x \in \mathbb{R}^m \setminus A$ .  
Then  $f$  extends to a 1-LIPSCHITZ map  $A \cup \{x\} \rightarrow \mathbb{R}^n$ .

# Nonconstructive nature of extension theorems

KIRSZBRAUN (1934): *yes* for all  $n$

There always exists an  $L$ -LIPSCHITZ extension  $\mathbb{R}^m \rightarrow \mathbb{R}^n$  of  $f$ .

The usual proofs all use some sort of transfinite induction and:

## Basic Extension Lemma

Let  $f: A \rightarrow \mathbb{R}^n$ ,  $A \subseteq \mathbb{R}^m$ , be 1-LIPSCHITZ, and let  $x \in \mathbb{R}^m \setminus A$ . Then  $f$  extends to a 1-LIPSCHITZ map  $A \cup \{x\} \rightarrow \mathbb{R}^n$ . I.e.,

$$\bigcap_{a \in A} \underbrace{\{y \in \mathbb{R}^n : \|y - f(a)\| \leq \|x - a\|\}}_{\overline{B}_{\|x-a\|}(f(a))} \quad \text{is non-empty.}$$

# Nonconstructive nature of extension theorems

KIRSZBRAUN (1934): *yes* for all  $n$

There always exists an  $L$ -LIPSCHITZ extension  $\mathbb{R}^m \rightarrow \mathbb{R}^n$  of  $f$ .

The usual proofs all use some sort of transfinite induction and:

## Basic Extension Lemma

Let  $f: A \rightarrow \mathbb{R}^n$ ,  $A \subseteq \mathbb{R}^m$ , be 1-LIPSCHITZ, and let  $x \in \mathbb{R}^m \setminus A$ . Then  $f$  extends to a 1-LIPSCHITZ map  $A \cup \{x\} \rightarrow \mathbb{R}^n$ . I.e.,

$$\bigcap_{a \in A} \underbrace{\{y \in \mathbb{R}^n : \|y - f(a)\| \leq \|x - a\|\}}_{\overline{B}_{\|x-a\|}(f(a))} \quad \text{is non-empty.}$$

By compactness one first reduces to the case of a *finite* set  $A$ .

# Nonconstructive nature of extension theorems

## Basic Extension Lemma, finite version

# Nonconstructive nature of extension theorems

## Basic Extension Lemma, finite version

Let  $x_1, \dots, x_k \in \mathbb{R}^m$ ,  $y_1, \dots, y_k \in \mathbb{R}^n$ , such that

$$\|y_i - y_j\| \leq \|x_i - x_j\| \quad (1 \leq i, j \leq k).$$

# Nonconstructive nature of extension theorems

## Basic Extension Lemma, finite version

Let  $x_1, \dots, x_k \in \mathbb{R}^m$ ,  $y_1, \dots, y_k \in \mathbb{R}^n$ , such that

$$\|y_i - y_j\| \leq \|x_i - x_j\| \quad (1 \leq i, j \leq k).$$

Let  $r_1, \dots, r_k \in \mathbb{R}^{>0}$ .

# Nonconstructive nature of extension theorems

## Basic Extension Lemma, finite version

Let  $x_1, \dots, x_k \in \mathbb{R}^m$ ,  $y_1, \dots, y_k \in \mathbb{R}^n$ , such that

$$\|y_i - y_j\| \leq \|x_i - x_j\| \quad (1 \leq i, j \leq k).$$

Let  $r_1, \dots, r_k \in \mathbb{R}^{>0}$ . Then

$$\overline{B}_{r_1}(x_1) \cap \dots \cap \overline{B}_{r_k}(x_k) \neq \emptyset \quad \Rightarrow \quad \overline{B}_{r_1}(y_1) \cap \dots \cap \overline{B}_{r_k}(y_k) \neq \emptyset.$$



# Nonconstructive nature of extension theorems

## Basic Extension Lemma, finite version

Let  $x_1, \dots, x_k \in \mathbb{R}^m$ ,  $y_1, \dots, y_k \in \mathbb{R}^n$ , such that

$$\|y_i - y_j\| \leq \|x_i - x_j\| \quad (1 \leq i, j \leq k).$$

Let  $r_1, \dots, r_k \in \mathbb{R}^{>0}$ . Then

$$\overline{B}_{r_1}(x_1) \cap \dots \cap \overline{B}_{r_k}(x_k) \neq \emptyset \quad \Rightarrow \quad \overline{B}_{r_1}(y_1) \cap \dots \cap \overline{B}_{r_k}(y_k) \neq \emptyset.$$

Idea of the proof: the function

$$\mathbb{R}^m \rightarrow \mathbb{R}: y \mapsto G(y) := \max \{ \|y - y_i\|/r_i : i = 1, \dots, k \}$$

is LIPSCHITZ with  $\lim_{\|y\| \rightarrow \infty} G(y) \rightarrow \infty$ .

# Nonconstructive nature of extension theorems

## Basic Extension Lemma, finite version

Let  $x_1, \dots, x_k \in \mathbb{R}^m$ ,  $y_1, \dots, y_k \in \mathbb{R}^n$ , such that

$$\|y_i - y_j\| \leq \|x_i - x_j\| \quad (1 \leq i, j \leq k).$$

Let  $r_1, \dots, r_k \in \mathbb{R}^{>0}$ . Then

$$\overline{B}_{r_1}(x_1) \cap \dots \cap \overline{B}_{r_k}(x_k) \neq \emptyset \quad \Rightarrow \quad \overline{B}_{r_1}(y_1) \cap \dots \cap \overline{B}_{r_k}(y_k) \neq \emptyset.$$

Idea of the proof: the function

$$\mathbb{R}^m \rightarrow \mathbb{R}: y \mapsto G(y) := \max \{ \|y - y_i\|/r_i : i = 1, \dots, k \}$$

is LIPSCHITZ with  $\lim_{\|y\| \rightarrow \infty} G(y) \rightarrow \infty$ . Thus  $G$  achieves a minimum at some  $w \in \mathbb{R}^m$ ; show  $G(w) \leq 1$ .

# Nonconstructive nature of extension theorems

By an **extension problem** we will mean a situation of the following kind:

Let  $\mathcal{C}$  be a class of maps  $\mathbb{R}^m \rightarrow \mathbb{R}^n$ . Find a necessary and sufficient condition for some given map  $X \rightarrow \mathbb{R}^n$  ( $X \subseteq \mathbb{R}^m$ ), possibly equipped with additional data, to have an extension to a map  $\mathbb{R}^m \rightarrow \mathbb{R}^n$  from  $\mathcal{C}$ .

# Nonconstructive nature of extension theorems

By an **extension problem** we will mean a situation of the following kind:

Let  $\mathcal{C}$  be a class of maps  $\mathbb{R}^m \rightarrow \mathbb{R}^n$ . Find a necessary and sufficient condition for some given map  $X \rightarrow \mathbb{R}^n$  ( $X \subseteq \mathbb{R}^m$ ), possibly equipped with additional data, to have an extension to a map  $\mathbb{R}^m \rightarrow \mathbb{R}^n$  from  $\mathcal{C}$ .

KIRSZBRAUN: for

$\mathcal{C}$  = class of  $L$ -LIPSCHITZ maps  $\mathbb{R}^m \rightarrow \mathbb{R}^n$ ,

no condition is necessary.

# Nonconstructive nature of extension theorems

By an **extension problem** we will mean a situation of the following kind:

Let  $\mathcal{C}$  be a class of maps  $\mathbb{R}^m \rightarrow \mathbb{R}^n$ . Find a necessary and sufficient condition for some given map  $X \rightarrow \mathbb{R}^n$  ( $X \subseteq \mathbb{R}^m$ ), possibly equipped with additional data, to have an extension to a map  $\mathbb{R}^m \rightarrow \mathbb{R}^n$  from  $\mathcal{C}$ .

KIRSZBRAUN: for

$\mathcal{C}$  = class of  $L$ -LIPSCHITZ maps  $\mathbb{R}^m \rightarrow \mathbb{R}^n$ ,

no condition is necessary.

We'll look at some other examples, associated with WHITNEY.

# Nonconstructive nature of extension theorems

From now on,  $X \subseteq \mathbb{R}^n$  is closed, and

$$\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n, \quad D^\alpha = \frac{\partial^{\alpha_1}}{\partial x_1^{\alpha_1}} \cdots \frac{\partial^{\alpha_n}}{\partial x_n^{\alpha_n}}, \quad |\alpha| = \alpha_1 + \cdots + \alpha_n.$$

## Definition

A **jet of order  $m$  on  $X$**  is a family  $F = (F^\alpha)_{|\alpha| \leq m}$  of continuous functions  $F^\alpha: X \rightarrow \mathbb{R}$ .

# Nonconstructive nature of extension theorems

From now on,  $X \subseteq \mathbb{R}^n$  is closed, and

$$\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n, \quad D^\alpha = \frac{\partial^{\alpha_1}}{\partial x_1^{\alpha_1}} \cdots \frac{\partial^{\alpha_n}}{\partial x_n^{\alpha_n}}, \quad |\alpha| = \alpha_1 + \cdots + \alpha_n.$$

## Definition

A **jet of order  $m$  on  $X$**  is a family  $F = (F^\alpha)_{|\alpha| \leq m}$  of continuous functions  $F^\alpha: X \rightarrow \mathbb{R}$ .

For  $f \in C^m(\mathbb{R}^n)$ , we obtain a jet  $J_X^m(f) := (D^\alpha f \upharpoonright X)_{|\alpha| \leq m}$  of order  $m$  on  $X$ .

# Nonconstructive nature of extension theorems

From now on,  $X \subseteq \mathbb{R}^n$  is closed, and

$$\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n, \quad D^\alpha = \frac{\partial^{\alpha_1}}{\partial x_1^{\alpha_1}} \cdots \frac{\partial^{\alpha_n}}{\partial x_n^{\alpha_n}}, \quad |\alpha| = \alpha_1 + \cdots + \alpha_n.$$

## Definition

A **jet of order  $m$  on  $X$**  is a family  $F = (F^\alpha)_{|\alpha| \leq m}$  of continuous functions  $F^\alpha: X \rightarrow \mathbb{R}$ .

For  $f \in C^m(\mathbb{R}^n)$ , we obtain a jet  $J_X^m(f) := (D^\alpha f \upharpoonright X)_{|\alpha| \leq m}$  of order  $m$  on  $X$ .

## Question

Let  $F$  be a jet of order  $m$  on  $X$ . *What is a necessary and sufficient condition to guarantee the existence of a  $C^m$ -function  $f: \mathbb{R}^n \rightarrow \mathbb{R}$  such that  $J_X^m(f) = F$ ?*



# Nonconstructive nature of extension theorems

Let  $F = (F^\alpha)_{|\alpha| \leq m}$  be a jet of order  $m$  on  $X$  and  $a \in X$ . Put

$$T_a^m F(x) := \sum_{|\alpha| \leq m} \frac{F^\alpha(a)}{\alpha!} (x - a)^\alpha, \quad R_a^m F := F - J_X^m(T_a^m F).$$

## Definition

A jet  $F$  of order  $m$  is a  $C^m$ -**WHITNEY field** ( $F \in \mathcal{E}^m(X)$ ) if for  $x_0 \in X$  and  $|\alpha| \leq m$ ,

$$(R_x^m F)^\alpha(y) = o(|x - y|^{m-|\alpha|}) \quad \text{as } X \ni x, y \rightarrow x_0.$$

By TAYLOR's Formula, for each  $f \in C^m(\mathbb{R}^n)$ ,

$$J_X^m(f) := (D^\alpha f \upharpoonright X)_{|\alpha| \leq m} \in \mathcal{E}^m(X).$$

# Nonconstructive nature of extension theorems

## WHITNEY Extension Theorem (H. WHITNEY, 1934)

For every  $F \in \mathcal{C}^m(X)$ , there is an  $f \in C^m(\mathbb{R}^n)$  with  $J_X^m(f) = F$ .

### Proof outline

- Decompose  $\mathbb{R}^n \setminus X$  into countably many cubes with disjoint interior satisfying some inequality regarding their diameter and distance from  $X$ . (“WHITNEY decomposition”)
- Use this to get a “special” partition of unity  $(\phi_i)$  on  $\mathbb{R}^n \setminus X$ .
- Pick  $x_i \in X$  such that  $d(x_i, \text{supp}(\phi_i)) = d(X, \text{supp}(\phi_i))$ .
- $$f(x) = \begin{cases} F^0(x), & \text{if } x \in X; \\ \sum_i \phi_i(x) T_{x_i}^m F(x), & \text{if } x \notin X. \end{cases}$$

# Nonconstructive nature of extension theorems

## WHITNEY Extension Theorem (H. WHITNEY, 1934)

For every  $F \in \mathcal{C}^m(X)$ , there is an  $f \in C^m(\mathbb{R}^n)$  with  $J_X^m(f) = F$ .

### Proof outline

- Decompose  $\mathbb{R}^n \setminus X$  into countably many cubes with disjoint interior satisfying some inequality regarding their diameter and distance from  $X$ . (“WHITNEY decomposition”)
- Use this to get a “special” partition of unity  $(\phi_i)$  on  $\mathbb{R}^n \setminus X$ .
- Pick  $x_i \in X$  such that  $d(x_i, \text{supp}(\phi_i)) = d(X, \text{supp}(\phi_i))$ .
- $$f(x) = \begin{cases} F^0(x), & \text{if } x \in X; \\ \sum_i \phi_i(x) T_{x_i}^m F(x), & \text{if } x \notin X. \end{cases}$$

Again, this is very non-explicit!

# Nonconstructive nature of extension theorems

WHITNEY actually asked a somewhat different question (and answered it for  $n = 1$ ):

## WHITNEY's Extension Problem

How can we determine whether a function  $X \rightarrow \mathbb{R}$  is the restriction of a  $C^m$ -function  $\mathbb{R}^n \rightarrow \mathbb{R}$ ?

A complete answer was only given by C. FEFFERMAN in the early 2000s.

An answer in the case  $m = 1$  was found earlier by G. GLAESER in 1958, and simplified by B. KLARTAG and N. ZOBIN (2007).

# Nonconstructive nature of extension theorems

WHITNEY actually asked a somewhat different question (and answered it for  $n = 1$ ):

## WHITNEY's Extension Problem

How can we determine whether a function  $X \rightarrow \mathbb{R}$  is the restriction of a  $C^m$ -function  $\mathbb{R}^n \rightarrow \mathbb{R}$ ?

A complete answer was only given by C. FEFFERMAN in the early 2000s.

An answer in the case  $m = 1$  was found earlier by G. GLAESER in 1958, and simplified by B. KLARTAG and N. ZOBIN (2007).

More about that later.

It is interesting to ask whether one can solve a given extension problem “constructively”, e.g.:

It is interesting to ask whether one can solve a given extension problem “constructively”, e.g.:

An “explicit” version of KIRSZBRAUN? (CHRIS MILLER)

It is interesting to ask whether one can solve a given extension problem “constructively”, e.g.:

An “explicit” version of KIRSZBRAUN? (CHRIS MILLER)

Let  $f: A \rightarrow \mathbb{R}^n$  ( $A \subseteq \mathbb{R}^m$ ) be  $L$ -LIPSCHITZ and *semialgebraic*. Is there a *semialgebraic*  $L$ -LIPSCHITZ map on  $\mathbb{R}^m$  extending  $f$ ?



It is interesting to ask whether one can solve a given extension problem “constructively”, e.g.:

## An “explicit” version of KIRSZBRAUN? (CHRIS MILLER)

Let  $f: A \rightarrow \mathbb{R}^n$  ( $A \subseteq \mathbb{R}^m$ ) be  $L$ -LIPSCHITZ and *semialgebraic*. Is there a *semialgebraic*  $L$ -LIPSCHITZ map on  $\mathbb{R}^m$  extending  $f$ ?

To answer this question, one has to forgo all the “nonconstructive” tricks in KIRSZBRAUN’s proof. But we gain the nice (topological, geometric) properties of semialgebraic sets.

It is interesting to ask whether one can solve a given extension problem “constructively”, e.g.:

## An “explicit” version of KIRSZBRAUN? (CHRIS MILLER)

Let  $f: A \rightarrow \mathbb{R}^n$  ( $A \subseteq \mathbb{R}^m$ ) be  $L$ -LIPSCHITZ and *semialgebraic*. Is there a *semialgebraic*  $L$ -LIPSCHITZ map on  $\mathbb{R}^m$  extending  $f$ ?

To answer this question, one has to forgo all the “nonconstructive” tricks in KIRSZBRAUN’s proof. But we gain the nice (topological, geometric) properties of semialgebraic sets.

Of course, one can also ask this question for maps definable in an o-minimal expansion of a real closed ordered field.

Assuming o-minimality turned out to be unnecessarily strong:

## Theorem (A.-FISCHER, 2011)

*Let  $R$  be any definably complete expansion of a real closed ordered field. Every definable  $L$ -LIPSCHITZ map  $A \rightarrow R^n$  ( $A \subseteq R^m$ ) has a definable  $L$ -LIPSCHITZ extension  $R^m \rightarrow R^n$ .*

Assuming o-minimality turned out to be unnecessarily strong:

## Theorem (A.-FISCHER, 2011)

*Let  $R$  be any definably complete expansion of a real closed ordered field. Every definable  $L$ -LIPSCHITZ map  $A \rightarrow R^n$  ( $A \subseteq R^m$ ) has a definable  $L$ -LIPSCHITZ extension  $R^m \rightarrow R^n$ .*

The heavy lifting for the proof of this theorem was already done in convex analysis, based on a relationship between LIPSCHITZ maps and (certain) set-valued maps (MINTY; more recently, BAUSCHKE & WANG).

Assuming o-minimality turned out to be unnecessarily strong:

## Theorem (A.-FISCHER, 2011)

*Let  $R$  be any definably complete expansion of a real closed ordered field. Every definable  $L$ -LIPSCHITZ map  $A \rightarrow R^n$  ( $A \subseteq R^m$ ) has a definable  $L$ -LIPSCHITZ extension  $R^m \rightarrow R^n$ .*

The heavy lifting for the proof of this theorem was already done in convex analysis, based on a relationship between LIPSCHITZ maps and (certain) set-valued maps (MINTY; more recently, BAUSCHKE & WANG).

A version for  $p$ -adic semialgebraic (or  $p$ -adic subanalytic maps) has recently been shown by CLUCKERS-MARTIN (2015).

Assuming o-minimality turned out to be unnecessarily strong:

## Theorem (A.-FISCHER, 2011)

*Let  $R$  be any definably complete expansion of a real closed ordered field. Every definable  $L$ -LIPSCHITZ map  $A \rightarrow R^n$  ( $A \subseteq R^m$ ) has a definable  $L$ -LIPSCHITZ extension  $R^m \rightarrow R^n$ .*

The heavy lifting for the proof of this theorem was already done in convex analysis, based on a relationship between LIPSCHITZ maps and (certain) set-valued maps (MINTY; more recently, BAUSCHKE & WANG).

A version for  $p$ -adic semialgebraic (or  $p$ -adic subanalytic maps) has recently been shown by CLUCKERS-MARTIN (2015).

What about a definable WHITNEY Extension Theorem?

Let  $X \subseteq \mathbb{R}^n$  be closed.

## Theorem (KURDYKA & PAWŁUCKI, 1997)

Let  $F \in \mathcal{C}^m(X)$  be subanalytic. Then there is a subanalytic  $C^m$ -function  $f: \mathbb{R}^n \rightarrow \mathbb{R}$  such that  $J_X^m(f) = F$ .

Their proof used tools very specific to the subanalytic context (e.g., reduction to the case  $X$  compact; quasiconvexity).

# Definable extension theorems

Let  $X \subseteq \mathbb{R}^n$  be closed.

## Theorem (KURDYKA & PAWŁUCKI, 1997)

Let  $F \in \mathcal{E}^m(X)$  be subanalytic. Then there is a subanalytic  $C^m$ -function  $f: \mathbb{R}^n \rightarrow \mathbb{R}$  such that  $J_X^m(f) = F$ .

Their proof used tools very specific to the subanalytic context (e.g., reduction to the case  $X$  compact; quasiconvexity).

## Theorem (PAWŁUCKI, 2008)

Suppose  $X$  is definable in an o-minimal expansion  $R$  of the ordered field of reals. There is a linear extension operator

$$\mathcal{E}_{\text{def}}^m(X) \rightarrow C^m(\mathbb{R}^n)$$

which is a finite composition of operators each of which preserves definability, or is integration w.r.t. a parameter.



# Definable extension theorems

Let now  $\mathcal{R}$  be an o-minimal expansion of a real closed field.

Theorem (THAMRONGTHANYALAK, 2012;  
KURDYKA-PAWŁUCKI, 2014)

*Let  $F \in \mathcal{C}^m(X)$  be definable. Then there is a definable  $C^m$ -function  $f: R^n \rightarrow R$  such that  $J_X^m(f) = F$ .*

# Definable extension theorems

Let now  $\mathcal{R}$  be an o-minimal expansion of a real closed field.

Theorem (THAMRONGTHANYALAK, 2012;  
KURDYKA-PAWŁUCKI, 2014)

*Let  $F \in \mathcal{C}^m(X)$  be definable. Then there is a definable  $C^m$ -function  $f: \mathcal{R}^n \rightarrow \mathcal{R}$  such that  $J_X^m(f) = F$ .*

The construction is uniform (for definable families of WHITNEY fields).

# Definable extension theorems

Let now  $R$  be an o-minimal expansion of a real closed field.

Theorem (THAMRONGTHANYALAK, 2012;  
KURDYKA-PAWŁUCKI, 2014)

*Let  $F \in \mathcal{C}^m(X)$  be definable. Then there is a definable  $C^m$ -function  $f: R^n \rightarrow R$  such that  $J_X^m(f) = F$ .*

The construction is uniform (for definable families of WHITNEY fields).

THAMRONGTHANYALAK's proof follows the outline of the construction of PAWŁUCKI, combining it with results on  $\Lambda^m$ -stratifications by FISCHER, which help to gain control on the growth of derivatives.

Let  $X \subseteq \mathbb{R}^d$  be open,  $d \geq 1$ , and  $\partial\Omega = \text{cl}(X) \setminus X$ .

Let  $X \subseteq \mathbb{R}^d$  be open,  $d \geq 1$ , and  $\partial\Omega = \text{cl}(X) \setminus X$ .

## Definition

Let  $f: X \rightarrow \mathbb{R}^n$  ( $n \geq 1$ ) be definable and  $C^m$ . One says that  $f$  is  $\Lambda^m$ -**regular** if there exists  $L > 0$  such that

$$\|D^\alpha f(x)\| \leq \frac{L}{d(x, \partial X)^{|\alpha|-1}} \quad \text{for all } x \in X, \alpha \in \mathbb{N}^d, 1 \leq |\alpha| \leq m.$$

Let  $X \subseteq R^d$  be open,  $d \geq 1$ , and  $\partial\Omega = \text{cl}(X) \setminus X$ .

## Definition

Let  $f: X \rightarrow R^n$  ( $n \geq 1$ ) be definable and  $C^m$ . One says that  $f$  is  $\Lambda^m$ -**regular** if there exists  $L > 0$  such that

$$\|D^\alpha f(x)\| \leq \frac{L}{d(x, \partial X)^{|\alpha|-1}} \quad \text{for all } x \in X, \alpha \in \mathbb{N}^d, 1 \leq |\alpha| \leq m.$$

Also declare each map  $R^0 \rightarrow R^n$  and the constant functions  $\pm\infty$  to be  $\Lambda^m$ -regular.

Let  $X \subseteq R^d$  be open,  $d \geq 1$ , and  $\partial\Omega = \text{cl}(X) \setminus X$ .

## Definition

Let  $f: X \rightarrow R^n$  ( $n \geq 1$ ) be definable and  $C^m$ . One says that  $f$  is  $\Lambda^m$ -**regular** if there exists  $L > 0$  such that

$$\|D^\alpha f(x)\| \leq \frac{L}{d(x, \partial X)^{|\alpha|-1}} \quad \text{for all } x \in X, \alpha \in \mathbb{N}^d, 1 \leq |\alpha| \leq m.$$

Also declare each map  $R^0 \rightarrow R^n$  and the constant functions  $\pm\infty$  to be  $\Lambda^m$ -regular.

For example,  $f(x) = \frac{1}{x}$  on  $X = (0, +\infty)$  is *not*  $\Lambda^1$ -regular.

**Standard open  $\Lambda^m$ -regular cells** in  $R^n$  are defined inductively just like open cells, requiring the continuous definable functions arising in the definition to be  $\Lambda^m$ -regular.



**Standard open  $\Lambda^m$ -regular cells** in  $R^n$  are defined inductively just like open cells, requiring the continuous definable functions arising in the definition to be  $\Lambda^m$ -regular.

A **standard  $\Lambda^m$ -regular cell in  $R^n$**  is either

- 1 a standard open  $\Lambda^m$ -regular cell in  $R^n$ ; or
- 2 the graph of a definable  $\Lambda^m$ -regular map  $D \rightarrow R^{n-d}$ , where  $D$  is a standard open  $\Lambda^m$ -regular cell in  $R^d$ , and  $0 \leq d < n$ .

**Standard open  $\Lambda^m$ -regular cells** in  $R^n$  are defined inductively just like open cells, requiring the continuous definable functions arising in the definition to be  $\Lambda^m$ -regular.

A **standard  $\Lambda^m$ -regular cell in  $R^n$**  is either

- 1 a standard open  $\Lambda^m$ -regular cell in  $R^n$ ; or
- 2 the graph of a definable  $\Lambda^m$ -regular map  $D \rightarrow R^{n-d}$ , where  $D$  is a standard open  $\Lambda^m$ -regular cell in  $R^d$ , and  $0 \leq d < n$ .

The image of a standard  $\Lambda^m$ -regular cell in  $R^n$  under an  $R$ -linear orthogonal isomorphism of  $R^n$  is called a  **$\Lambda^m$ -regular cell in  $R^n$** .

## Theorem (FISCHER, 2007)

*Let  $S_1, \dots, S_k$  be definable subsets of  $R^n$ . Then there exists a finite partition  $\mathcal{D}$  of  $R^n$  into  $\Lambda^m$ -regular cells such that*

- each  $\partial D$  ( $D \in \mathcal{D}$ ) and*
- each  $S_i$*

*is a union of sets from  $\mathcal{D}$ .*

## Theorem (FISCHER, 2007)

*Let  $S_1, \dots, S_k$  be definable subsets of  $R^n$ . Then there exists a finite partition  $\mathcal{D}$  of  $R^n$  into  $\Lambda^m$ -regular cells such that*

- each  $\partial D$  ( $D \in \mathcal{D}$ ) and*
- each  $S_i$*

*is a union of sets from  $\mathcal{D}$ .*

Moreover, one has quite some fine control over the cells in  $\mathcal{D}$ ; e.g., they can additionally chosen to be LIPSCHITZ (with rational LIPSCHITZ constant depending only on  $n$ ).

## Theorem (FISCHER, 2007)

*Let  $S_1, \dots, S_k$  be definable subsets of  $R^n$ . Then there exists a finite partition  $\mathcal{D}$  of  $R^n$  into  $\Lambda^m$ -regular cells such that*

- each  $\partial D$  ( $D \in \mathcal{D}$ ) and*
- each  $S_i$*

*is a union of sets from  $\mathcal{D}$ .*

Moreover, one has quite some fine control over the cells in  $\mathcal{D}$ ; e.g., they can additionally chosen to be LIPSCHITZ (with rational LIPSCHITZ constant depending only on  $n$ ).

At the root of this (and also of the PILA-WILKIE parametrization theorem) is a calculus lemma due to GROMOV.

## Lemma

*Let  $h: I \rightarrow R$  be a definable  $C^2$ -function on an interval  $I$  in  $R$  such that  $h, h''$  are semidefinite. Let  $t \in I$  and  $r > 0$  with  $[t - r, t + r] \subseteq I$ . Then*

$$|h'(t)| \leq \frac{1}{r} \sup \{ |h(\xi)| : \xi \in [t - r, t + r] \}.$$

# Definable extension theorems

Now on to the definable WHITNEY extension problem for  $m = 1$ .

Now on to the definable WHITNEY extension problem for  $m = 1$ .

## Theorem (A. & THAMRONGTHANYALAK)

*Let  $(f_a)_{a \in R^N}$  be a definable family of functions  $f_a: X_a \rightarrow R$ , where  $X_a \subseteq R^n$  is closed. Then*

$$A_* := \{a \in R^N : f_a \text{ extends to a definable } C^1\text{-function } R^n \rightarrow R\}$$

*is definable. Moreover, there is a definable family  $(\tilde{f}_a)_{a \in A_*}$  of  $C^1$ -functions  $R^n \rightarrow R$  such that  $\tilde{f}_a \upharpoonright X_a = f_a$  for each  $a \in A_*$ .*



# Definable extension theorems

Now on to the definable WHITNEY extension problem for  $m = 1$ .

## Theorem (A. & THAMRONGTHANYALAK)

*Let  $(f_a)_{a \in R^N}$  be a definable family of functions  $f_a: X_a \rightarrow R$ , where  $X_a \subseteq R^n$  is closed. Then*

*$A_* := \{a \in R^N : f_a \text{ extends to a definable } C^1\text{-function } R^n \rightarrow R\}$*

*is definable. Moreover, there is a definable family  $(\tilde{f}_a)_{a \in A_*}$  of  $C^1$ -functions  $R^n \rightarrow R$  such that  $\tilde{f}_a \upharpoonright X_a = f_a$  for each  $a \in A_*$ .*

## Corollary ( $C^1$ case of a question of VAN DEN DRIES, 1997)

*Let  $f: X \rightarrow R$  be a definable function where  $X \subseteq R^n$  is closed. Suppose that  $f$  locally extends to a definable  $C^1$ -function. Then  $f$  extends to a definable  $C^1$ -function on  $R^n$ .*

# Definable extension theorems

The proof of this theorem (following KLARTAG-ZOBIN) uses the method of *affine bundles*:

## Definition

A definable set-valued map  $H: X \rightrightarrows R^m$  is an **affine bundle** if for every  $x \in X$ ,  $H(x)$  is an affine subspace of  $R^m$  or  $H(x) = \emptyset$ .

# Definable extension theorems

The proof of this theorem (following KLARTAG-ZOBIN) uses the method of *affine bundles*:

## Definition

A definable set-valued map  $H: X \rightrightarrows R^m$  is an **affine bundle** if for every  $x \in X$ ,  $H(x)$  is an affine subspace of  $R^m$  or  $H(x) = \emptyset$ .

The goal is to successively cut down  $H$  in order to find a continuous **selection** (= SKOLEM function) of  $H$ .

# Definable extension theorems

The proof of this theorem (following KLARTAG-ZOBIN) uses the method of *affine bundles*:

## Definition

A definable set-valued map  $H: X \rightrightarrows R^m$  is an **affine bundle** if for every  $x \in X$ ,  $H(x)$  is an affine subspace of  $R^m$  or  $H(x) = \emptyset$ .

The goal is to successively cut down  $H$  in order to find a continuous **selection** (= SKOLEM function) of  $H$ .

This method is actually easier to explain when applied to solving a different problem:

# Definable extension theorems

The proof of this theorem (following KLARTAG-ZOBIN) uses the method of *affine bundles*:

## Definition

A definable set-valued map  $H: X \rightrightarrows R^m$  is an **affine bundle** if for every  $x \in X$ ,  $H(x)$  is an affine subspace of  $R^m$  or  $H(x) = \emptyset$ .

The goal is to successively cut down  $H$  in order to find a continuous **selection** (= SKOLEM function) of  $H$ .

This method is actually easier to explain when applied to solving a different problem:

Let  $f, g_1, \dots, g_m: X \rightarrow R$  be definable. Are there *definable* continuous functions  $y_1, \dots, y_m: X \rightarrow R$  such that

$$f = g_1 y_1 + \dots + g_m y_m? \quad (*)$$

Now  $(*)$  has a solution iff the affine bundle  $H^{(0)}$  on  $R^m$  given by

$$H^{(0)}(x) := \{(y_1, \dots, y_m) \in R^m : f(x) = g_1(x)y_1 + \dots + g_m(x)y_m\}.$$

has a definable continuous selection.

Now  $(*)$  has a solution iff the affine bundle  $H^{(0)}$  on  $R^m$  given by

$$H^{(0)}(x) := \{(y_1, \dots, y_m) \in R^m : f(x) = g_1(x)y_1 + \dots + g_m(x)y_m\}.$$

has a definable continuous selection.

Inductively define  $H^{(l+1)} := (H^{(l)})'$ .

Here,  $H'$  is the GLAESER **refinement** of  $H$ :

$$H'(x_0) := \{y_0 \in H(x_0) : d(y_0, H(x)) \rightarrow 0 \text{ as } x \rightarrow x_0 \text{ in } X\}.$$

Now  $(*)$  has a solution iff the affine bundle  $H^{(0)}$  on  $R^m$  given by

$$H^{(0)}(x) := \{(y_1, \dots, y_m) \in R^m : f(x) = g_1(x)y_1 + \dots + g_m(x)y_m\}.$$

has a definable continuous selection.

Inductively define  $H^{(l+1)} := (H^{(l)})'$ .

Here,  $H'$  is the GLAESER **refinement** of  $H$ :

$$H'(x_0) := \{y_0 \in H(x_0) : d(y_0, H(x)) \rightarrow 0 \text{ as } x \rightarrow x_0 \text{ in } X\}.$$

Easy to check:

- $H' \subseteq H$ , so every selection of  $H'$  is a selection of  $H$ ; conversely, each *continuous* selection of  $H$  is also a selection of  $H'$ ;
- $H$  is an affine bundle  $\Rightarrow H'$  is an affine bundle.



## Lemma

*Let  $x_0 \in X$ . Then  $\dim H'(x_0) \leq \liminf_{X \ni x \rightarrow x_0} \dim H(x)$ .*

Indeed, let  $p_0, \dots, p_d \in H'(x_0)$  be such that  $p_1 - p_0, \dots, p_d - p_0$  are linearly independent,  $d = \dim H'(x_0)$ , and  $\varepsilon > 0$ .

By definition of  $H'$ , there is a  $\delta > 0$  such that for all  $x \in B_\delta(x_0)$  we obtain  $q_0, \dots, q_d \in H(x)$  with  $\|p_i - q_i\| < \varepsilon$ . For small  $\varepsilon$ ,  $q_1 - q_0, \dots, q_d - q_0$  are linearly independent:  $\dim H(x) \geq d$ .  $\square$

## Lemma

*Let  $x_0 \in X$ . Then  $\dim H'(x_0) \leq \liminf_{X \ni x \rightarrow x_0} \dim H(x)$ .*

Indeed, let  $p_0, \dots, p_d \in H'(x_0)$  be such that  $p_1 - p_0, \dots, p_d - p_0$  are linearly independent,  $d = \dim H'(x_0)$ , and  $\varepsilon > 0$ .

By definition of  $H'$ , there is a  $\delta > 0$  such that for all  $x \in B_\delta(x_0)$  we obtain  $q_0, \dots, q_d \in H(x)$  with  $\|p_i - q_i\| < \varepsilon$ . For small  $\varepsilon$ ,  $q_1 - q_0, \dots, q_d - q_0$  are linearly independent:  $\dim H(x) \geq d$ .  $\square$

Another little extra argument based on this lemma yields

$$H^{(*)} := H^{(2m+1)} = H^{(2m+2)} = \dots$$

So we have:

$H$  has a continuous selection



$H^{(*)}(x) \neq \emptyset$  for all  $x \in X$ .

Can we reverse this implication?

So we have:

$H$  has a continuous selection



$H^{(*)}(x) \neq \emptyset$  for all  $x \in X$ .

Can we reverse this implication? To answer this we note that


$H^{(*)}$  is *lower semi-continuous*  
in the sense of the following definition.

# Definable extension theorems


Let  $T: X \rightrightarrows \mathbb{R}^m$  be a set-valued map.

## Definition

One says that  $T$  is **lower semi-continuous (l.s.c.)** if for every  $x \in X$ ,  $y \in T(x)$ , and neighborhood  $V$  of  $y$ , there is a neighborhood  $U$  of  $x$  such that  $T(x') \cap V \neq \emptyset$  for all  $x' \in U \cap X$ .



l.s.c.



not l.s.c.

## Theorem (Definable MICHAEL's Selection Theorem)

*Suppose  $X$  is closed and  $T: X \rightrightarrows \mathbb{R}^m$  is definable and l.s.c. such that for every  $x \in X$ ,  $T(x)$  is nonempty, closed, and convex. Then  $T$  has a continuous definable selection.*

## Theorem (Definable MICHAEL's Selection Theorem)

*Suppose  $X$  is closed and  $T: X \rightrightarrows R^m$  is definable and l.s.c. such that for every  $x \in X$ ,  $T(x)$  is nonempty, closed, and convex. Then  $T$  has a continuous definable selection.*

Classically, this theorem is shown by a nonconstructive iterative procedure. Our proof relies on Cell Decomposition.

## Theorem (Definable MICHAEL's Selection Theorem)

*Suppose  $X$  is closed and  $T: X \rightrightarrows R^m$  is definable and l.s.c. such that for every  $x \in X$ ,  $T(x)$  is nonempty, closed, and convex. Then  $T$  has a continuous definable selection.*

Classically, this theorem is shown by a nonconstructive iterative procedure. Our proof relies on Cell Decomposition.

It does also hold for *bounded*  $X$  in the category of semilinear sets and maps (using a different proof).

[A. & THAMRONGTHANYALAK; simplified by CZAPLA & PAWŁUCKI]



## Corollary

*Let  $f, g_1, \dots, g_m$  be definable maps  $X \rightarrow R$ . If there are continuous functions  $y_1, \dots, y_m: X \rightarrow R$  such that*

$$f = g_1 y_1 + \dots + g_m y_m, \quad (*)$$

*then there are also definable continuous function  $y_i$  solving  $(*)$ .*

For polynomials  $f, g_1, \dots, g_m$ , this was shown by FEFFERMAN and KOLLÁR by other means. (KOLLÁR-NOWAK: in this case one cannot always take the  $y_i$  to be rational.)

## Corollary

*Let  $f, g_1, \dots, g_m$  be definable maps  $X \rightarrow R$ . If there are continuous functions  $y_1, \dots, y_m: X \rightarrow R$  such that*

$$f = g_1 y_1 + \dots + g_m y_m, \quad (*)$$

*then there are also definable continuous function  $y_i$  solving  $(*)$ .*

For polynomials  $f, g_1, \dots, g_m$ , this was shown by FEFFERMAN and KOLLÁR by other means. (KOLLÁR-NOWAK: in this case one cannot always take the  $y_i$  to be rational.)

## Question (GABRIELOV)

If a definable set-valued map  $X \rightrightarrows \mathbb{R}^m$  has a continuous selection, does it always have one that is definable?

Back to the  $C^1$ -WHITNEY Extension Problem.

Back to the  $C^1$ -WHITNEY Extension Problem.

## Definition

Let  $f: X \rightarrow R$ . A **holding space** for  $f$  is an affine bundle

$$H: X \rightrightarrows (R \times R^n)$$

such that whenever  $F \in C^1(R^n)$  is definable,  $F \upharpoonright X = f$ , then

$$\left( F(x), \frac{\partial F}{\partial x_1}(x), \dots, \frac{\partial F}{\partial x_n}(x) \right) \in H(x) \quad \text{for all } x \in X.$$

Back to the  $C^1$ -WHITNEY Extension Problem.

## Definition

Let  $f: X \rightarrow R$ . A **holding space** for  $f$  is an affine bundle

$$H: X \rightrightarrows (R \times R^n)$$

such that whenever  $F \in C^1(R^n)$  is definable,  $F \upharpoonright X = f$ , then

$$\left( F(x), \frac{\partial F}{\partial x_1}(x), \dots, \frac{\partial F}{\partial x_n}(x) \right) \in H(x) \quad \text{for all } x \in X.$$

Identify  $R \times R^n$  with the space  $\mathcal{P}_n$  of linear polynomials in  $n$  indeterminates. Think of a holding space for  $f$  as a collection of potential TAYLOR polynomials of definable  $C^1$ -extensions of  $f$ .

## Definition (the $C^1$ -GLAESER **refinement** $\tilde{H}$ of $H$ )

$p_0 \in \tilde{H}(x_0) :\iff p_0 \in H(x_0)$  and

$(\forall \varepsilon > 0) (\exists \delta > 0) (\forall x_1, x_2 \in X \cap B_\delta(x_0)) (\exists p_1 \in H(x_1), p_2 \in H(x_2))$

$|D^\alpha(p_i - p_j)(x_i)| \leq \varepsilon \|x_i - x_j\|^{1-|\alpha|}$  for  $i, j = 0, 1, 2$  and  $|\alpha| \leq 1$ .

## Definition (the $C^1$ -GLAESER **refinement** $\widetilde{H}$ of $H$ )

$p_0 \in \widetilde{H}(x_0) :\iff p_0 \in H(x_0)$  and

$(\forall \varepsilon > 0) (\exists \delta > 0) (\forall x_1, x_2 \in X \cap B_\delta(x_0)) (\exists p_1 \in H(x_1), p_2 \in H(x_2))$   
 $|D^\alpha(p_i - p_j)(x_i)| \leq \varepsilon \|x_i - x_j\|^{1-|\alpha|}$  for  $i, j = 0, 1, 2$  and  $|\alpha| \leq 1$ .

As before, the sequence  $(H_l)$  where

$$H_0 = \text{trivial holding space for } f, \quad H_{l+1} := \widetilde{H}_l$$

eventually stabilizes. With  $H_*$  its eventual value, by definable WHITNEY extension,

$f$  extends to a definable  $C^1$ -function on  $R^n$   $\iff$   $H_*$  has a *continuous* definable selection.

Let  $R$  be an o-minimal structure on  $\mathbb{R}$ .

- 1 Suppose  $f: A \rightarrow \mathbb{R}^n$  ( $A \subseteq \mathbb{R}^m$ ) is  $L$ -LIPSCHITZ and *locally definable* in  $R$ . Does  $f$  extend to a locally definable  $L$ -LIPSCHITZ function  $\mathbb{R}^m \rightarrow \mathbb{R}^n$ ? (Yes if  $f$  is bounded, or if in the conclusion  $L$  is replaced by  $L + \varepsilon$ .)



Let  $R$  be an o-minimal structure on  $\mathbb{R}$ .

- 1 Suppose  $f: A \rightarrow \mathbb{R}^n$  ( $A \subseteq \mathbb{R}^m$ ) is  $L$ -LIPSCHITZ and *locally definable* in  $R$ . Does  $f$  extend to a locally definable  $L$ -LIPSCHITZ function  $\mathbb{R}^m \rightarrow \mathbb{R}^n$ ? (Yes if  $f$  is bounded, or if in the conclusion  $L$  is replaced by  $L + \varepsilon$ .)
- 2 What about the  $C^m$  case of WHITNEY extension for definable functions in  $R$ ? [BIERSTONE & MILMAN, 2009]

Let  $R$  be an o-minimal structure on  $\mathbb{R}$ .

- ① Suppose  $f: A \rightarrow \mathbb{R}^n$  ( $A \subseteq \mathbb{R}^m$ ) is  $L$ -LIPSCHITZ and *locally definable* in  $R$ . Does  $f$  extend to a locally definable  $L$ -LIPSCHITZ function  $\mathbb{R}^m \rightarrow \mathbb{R}^n$ ? (Yes if  $f$  is bounded, or if in the conclusion  $L$  is replaced by  $L + \varepsilon$ .)
- ② What about the  $C^m$  case of WHITNEY extension for definable functions in  $R$ ? [BIERSTONE & MILMAN, 2009]
- ③ The  $p$ -adic case; e.g.: Let  $f, g_1, \dots, g_m: X \rightarrow \mathbb{Q}_p$  ( $X \subseteq \mathbb{Q}_p^m$ ) be semialgebraic such that there are continuous  $y_i$  with

$$f = g_1 y_1 + \dots + g_m y_m.$$

Are there also semialgebraic continuous  $y_i$  with this property?

(FEFFERMAN-KOLLÁR: yes if  $f, g_i$  are polynomials.)