

# Definability-theoretic dividing lines

Philipp Hieronymi

University of Illinois at Urbana-Champaign

Konstanz 2018

Throughout this talk, we will consider expansions of

- $(\mathbb{R}, <, +)$ ,
- $(\mathbb{R}, <, +, x \mapsto \lambda x \text{ for each } \lambda \in \mathbb{R})$ .
- $\overline{\mathbb{R}} := (\mathbb{R}, <, +, \cdot)$ .

**Motivating goal.** Classify such expansions!

Throughout this talk, we will consider expansions of

- $(\mathbb{R}, <, +)$ ,
- $(\mathbb{R}, <, +, x \mapsto \lambda x \text{ for each } \lambda \in \mathbb{R})$ .
- $\overline{\mathbb{R}} := (\mathbb{R}, <, +, \cdot)$ .

**Motivating goal.** Classify such expansions!

**A disclaimer.** Two perspectives on first-order expansions of  $(\mathbb{R}, <, +)$ :

- ① as a concrete collection of (definable) subsets of  $\mathbb{R}^n$ ,
- ② in terms of its theory, up to bi-interpretability.

For this talk, we take perspective 1.

Throughout this talk, we will consider expansions of

- $(\mathbb{R}, <, +)$ ,
- $(\mathbb{R}, <, +, x \mapsto \lambda x \text{ for each } \lambda \in \mathbb{R})$ .
- $\overline{\mathbb{R}} := (\mathbb{R}, <, +, \cdot)$ .

**Motivating goal.** Classify such expansions!

**A disclaimer.** Two perspectives on first-order expansions of  $(\mathbb{R}, <, +)$ :

- 1 as a concrete collection of (definable) subsets of  $\mathbb{R}^n$ ,
- 2 in terms of its theory, up to bi-interpretability.

For this talk, we take perspective 1.

**More realistic goal.** Classify such expansions according to the geometric/topological complexity/tameness of its definable set?

This is an instance of a larger program:

*'A lot of model theory is concerned with discovering and charting the "tame" regions of mathematics, where wild phenomena like space filling curves and Gödel incompleteness are absent, or at least under control. As Hrushovski put it recently: Model Theory = Geography of Tame Mathematics.'* - Lou van den Dries

This is an instance of a larger program:

*'A lot of model theory is concerned with discovering and charting the "tame" regions of mathematics, where wild phenomena like space filling curves and Gödel incompleteness are absent, or at least under control. As Hrushovski put it recently: Model Theory = Geography of Tame Mathematics.'* - Lou van den Dries

So what is tameness precisely?

*'The notion of tameness carries with it some judgement by the user as to what constitutes acceptable behavior of the definable sets, as well as what constitutes acceptable knowledge of the behavior.'* - Chris Miller

There are large number of expansions of  $(\mathbb{R}, <, +)$  known to be tame:

- $(\mathbb{R}, <, +)$ ,  $(\mathbb{R}, <, +, \cdot)$ ,  $\mathbb{R}_{an}$ , (o-minimal)
- $(\mathbb{R}, <, +, \mathbb{Z})$ , (locally o-minimal)
- $(\mathbb{R}, <, +, \mathbb{Q})$ , (o-minimal open core)
- $(\mathbb{R}, <, +, \cdot, 2^{\mathbb{Z}})$ , (d-minimal)
- $(\mathbb{R}, <, +, \cdot, 2^{\mathbb{Z}}3^{\mathbb{Z}}, 2^{\mathbb{Z}})$ , (d-minimal open core)
- $(\mathbb{R}, <, +, \mathbb{Q}, \mathbb{Z})$ , (locally o-minimal open core)

There are large number of expansions of  $(\mathbb{R}, <, +)$  known to be tame:

- $(\mathbb{R}, <, +)$ ,  $(\mathbb{R}, <, +, \cdot)$ ,  $\mathbb{R}_{an}$ , (o-minimal)
- $(\mathbb{R}, <, +, \mathbb{Z})$ , (locally o-minimal)
- $(\mathbb{R}, <, +, \mathbb{Q})$ , (o-minimal open core)
- $(\mathbb{R}, <, +, \cdot, 2^{\mathbb{Z}})$ , (d-minimal)
- $(\mathbb{R}, <, +, \cdot, 2^{\mathbb{Z}}3^{\mathbb{Z}}, 2^{\mathbb{Z}})$ , (d-minimal open core)
- $(\mathbb{R}, <, +, \mathbb{Q}, \mathbb{Z})$ , (locally o-minimal open core)

Compare to this, the expansion of  $(\mathbb{R}, <, +)$  by predicates for every compact subset of  $\mathbb{R}^n$  - call it  $\mathbb{R}_{cph}$ . It defines every projective subset of  $[0, 1]^k$  for every  $k \in \mathbb{N}$ .



There are large number of expansions of  $(\mathbb{R}, <, +)$  known to be tame:

- $(\mathbb{R}, <, +)$ ,  $(\mathbb{R}, <, +, \cdot)$ ,  $\mathbb{R}_{an}$ , (o-minimal)
- $(\mathbb{R}, <, +, \mathbb{Z})$ , (locally o-minimal)
- $(\mathbb{R}, <, +, \mathbb{Q})$ , (o-minimal open core)
- $(\mathbb{R}, <, +, \cdot, 2^{\mathbb{Z}})$ , (d-minimal)
- $(\mathbb{R}, <, +, \cdot, 2^{\mathbb{Z}}3^{\mathbb{Z}}, 2^{\mathbb{Z}})$ , (d-minimal open core)
- $(\mathbb{R}, <, +, \mathbb{Q}, \mathbb{Z})$ , (locally o-minimal open core)

Compare to this, the expansion of  $(\mathbb{R}, <, +)$  by predicates for every compact subset of  $\mathbb{R}^n$  - call it  $\mathbb{R}_{cph}$ . It defines every projective subset of  $[0, 1]^k$  for every  $k \in \mathbb{N}$ .

For future reference:  $(\mathbb{R}, <, +, \cdot, \mathbb{Z})$  defines all compact sets.

There are large number of expansions of  $(\mathbb{R}, <, +)$  known to be tame:

- $(\mathbb{R}, <, +)$ ,  $(\mathbb{R}, <, +, \cdot)$ ,  $\mathbb{R}_{an}$ , (o-minimal)
- $(\mathbb{R}, <, +, \mathbb{Z})$ , (locally o-minimal)
- $(\mathbb{R}, <, +, \mathbb{Q})$ , (o-minimal open core)
- $(\mathbb{R}, <, +, \cdot, 2^{\mathbb{Z}})$ , (d-minimal)
- $(\mathbb{R}, <, +, \cdot, 2^{\mathbb{Z}}3^{\mathbb{Z}}, 2^{\mathbb{Z}})$ , (d-minimal open core)
- $(\mathbb{R}, <, +, \mathbb{Q}, \mathbb{Z})$ , (locally o-minimal open core)

Compare to this, the expansion of  $(\mathbb{R}, <, +)$  by predicates for every compact subset of  $\mathbb{R}^n$  - call it  $\mathbb{R}_{cph}$ . It defines every projective subset of  $[0, 1]^k$  for every  $k \in \mathbb{N}$ .

For future reference:  $(\mathbb{R}, <, +, \cdot, \mathbb{Z})$  defines all compact sets.

**Observation:** There is a large class of structures whose definable sets can be understood by model-theoretic (tame topology) methods. But there are also structures whose definable sets can only be studied in terms of descriptive set theory.

What do we expect from tameness notion?

# What do we expect from tameness notion?

- 1 'Natural' examples.

What do we expect from tameness notion?

- 1 'Natural' examples.
- 2 Implies many (if not all) properties of these natural examples.

What do we expect from tameness notion?

- 1 'Natural' examples.
- 2 Implies many (if not all) properties of these natural examples.
- 3 appears naturally when classifying structures.

What do we expect from tameness notion?

- ① 'Natural' examples.
- ② Implies many (if not all) properties of these natural examples.
- ③ appears naturally when classifying structures.
- ④ Its failure has strong consequences.

What do we expect from tameness notion?

- ① 'Natural' examples.
- ② Implies many (if not all) properties of these natural examples.
- ③ appears naturally when classifying structures.
- ④ Its failure has strong consequences.
- ⑤ ...



What do we expect from tameness notion?

- 1 'Natural' examples.
- 2 Implies many (if not all) properties of these natural examples.
- 3 appears naturally when classifying structures.
- 4 Its failure has strong consequences.
- 5 ...

Condition 4 is very debatable, but goes back to Shelah and has a long history in model theory. His idea was to identify important tame/wild dividing lines in the class of first-order theories and he proposed stability as maybe the first such dividing line.

What do we expect from tameness notion?

- 1 'Natural' examples.
- 2 Implies many (if not all) properties of these natural examples.
- 3 appears naturally when classifying structures.
- 4 Its failure has strong consequences.
- 5 ...

Condition 4 is very debatable, but goes back to Shelah and has a long history in model theory. His idea was to identify important tame/wild dividing lines in the class of first-order theories and he proposed stability as maybe the first such dividing line.

In the rest of the talk, I will discuss Condition 3 and 4.

A common theme in model theory is the classification of all structures on the tame side of a given dividing line.

A common theme in model theory is the classification of all structures on the tame side of a given dividing line.

Classical examples: Zilber's (false) trichotomy conjecture on strongly minimal theory and the Peterzil-Starchenko trichotomy theorem for o-minimal structures.

A common theme in model theory is the classification of all structures on the tame side of a given dividing line.

Classical examples: Zilber's (false) trichotomy conjecture on strongly minimal theory and the Peterzil-Starchenko trichotomy theorem for o-minimal structures.

In our setting: Classify (non-elementary) classes of expansions of the real field by objects of the same type according to common tameness conditions they satisfy.

A common theme in model theory is the classification of all structures on the tame side of a given dividing line.

Classical examples: Zilber's (false) trichotomy conjecture on strongly minimal theory and the Peterzil-Starchenko trichotomy theorem for o-minimal structures.

In our setting: Classify (non-elementary) classes of expansions of the real field by objects of the same type according to common tameness conditions they satisfy.

Tameness conditions that appear in such classifications, should be consider significant.

## Classification of expansions of $(\mathbb{R}, <, +)$ by cyclic multiplicative subgroups of $\mathbb{C}^*$ .

Let  $S := (ae^{i\varphi})^{\mathbb{Z}} \subseteq \mathbb{R}^2$  be infinite, where  $a \in \mathbb{R}_{>0}$  and  $\varphi \in \mathbb{R}$ . Then exactly one of the following holds:

- 1 the open core of  $(\overline{\mathbb{R}}, S)$  is o-minimal,
- 2  $(\overline{\mathbb{R}}, S)$  is d-minimal,
- 3  $(\overline{\mathbb{R}}, S)$  defines  $\mathbb{Z}$ .

## Classification of expansions of $(\mathbb{R}, <, +)$ by cyclic multiplicative subgroups of $\mathbb{C}^*$ .

Let  $S := (ae^{i\varphi})^{\mathbb{Z}} \subseteq \mathbb{R}^2$  be infinite, where  $a \in \mathbb{R}_{>0}$  and  $\varphi \in \mathbb{R}$ . Then exactly one of the following holds:

- 1 the open core of  $(\overline{\mathbb{R}}, S)$  is o-minimal,
- 2  $(\overline{\mathbb{R}}, S)$  is d-minimal,
- 3  $(\overline{\mathbb{R}}, S)$  defines  $\mathbb{Z}$ .

Caulfield (2016): fails for finitely generated subgroups. There is a subgroup  $S$  generated by two elements such that  $(\overline{\mathbb{R}}, S)$  has d-minimal open core, but is not d-minimal.



## Classification of expansions of $(\mathbb{R}, <, +)$ by discrete multiplicative subgroups of $\mathrm{GL}_n(\mathbb{C})$ . (H.-Walsberg 2018)

Let  $\Gamma$  be an infinite discrete subgroup of  $\mathrm{GL}_n(\mathbb{C})$ . Then either

- $(\overline{\mathbb{R}}, \Gamma)$  defines  $\mathbb{Z}$  or
- $(\overline{\mathbb{R}}, \Gamma)$  is d-minimal.

If  $\Gamma$  is not virtually abelian, then  $(\overline{\mathbb{R}}, \Gamma)$  defines  $\mathbb{Z}$ .

## Classification of expansions of $(\mathbb{R}, <, +)$ by discrete multiplicative subgroups of $\mathrm{GL}_n(\mathbb{C})$ . (H.-Walsberg 2018)

Let  $\Gamma$  be an infinite discrete subgroup of  $\mathrm{GL}_n(\mathbb{C})$ . Then either

- $(\overline{\mathbb{R}}, \Gamma)$  defines  $\mathbb{Z}$  or
- $(\overline{\mathbb{R}}, \Gamma)$  is d-minimal.

If  $\Gamma$  is not virtually abelian, then  $(\overline{\mathbb{R}}, \Gamma)$  defines  $\mathbb{Z}$ .

**Conjecture.** Let  $G$  be a finitely generated subgroup of  $\mathrm{GL}_n(\mathbb{C})$ . Then one of the following holds:

- 1  $(\overline{\mathbb{R}}, G)$  defines  $\mathbb{Z}$ ,
- 2  $(\overline{\mathbb{R}}, G)$  is d-minimal,
- 3  $(\overline{\mathbb{R}}, G)$  has o-minimal open core,
- 4  $(\overline{\mathbb{R}}, G)$  has d-minimal open core.

**Conjecture (Miller).** Let  $K$  be a subfield of  $\mathbb{R}$ . Then one of the following holds:

- $(\overline{\mathbb{R}}, K)$  defines  $\mathbb{Z}$ ,
- $(\overline{\mathbb{R}}, K)$  has o-minimal open core.

**Conjecture (Miller).** Let  $K$  be a subfield of  $\mathbb{R}$ . Then one of the following holds:

- $(\overline{\mathbb{R}}, K)$  defines  $\mathbb{Z}$ ,
- $(\overline{\mathbb{R}}, K)$  has o-minimal open core.

Wide open. Alexi Block Gorman will report on some further evidence that the conjecture could actually be true.

**Tameness?** For  $r \in \mathbb{N}_{\geq 2}$ , consider a ternary predicate  $V_r(x, u, k)$  that holds if and only if  $u$  is an integer power of  $r$ ,  $k \in \{0, \dots, r-1\}$ , and the digit of some base  $r$  representation of  $x$  in the position corresponding to  $u$  is  $k$ .

Set  $\mathcal{T}_r := (\mathbb{R}, <, +, V_r)$ .

**Tameness?** For  $r \in \mathbb{N}_{\geq 2}$ , consider a ternary predicate  $V_r(x, u, k)$  that holds if and only if  $u$  is an integer power of  $r$ ,  $k \in \{0, \dots, r-1\}$ , and the digit of some base  $r$  representation of  $x$  in the position corresponding to  $u$  is  $k$ .

Set  $\mathcal{T}_r := (\mathbb{R}, <, +, V_r)$ .

**Observation 1:**  $\mathcal{T}_r$  is bi-interpretable with  $(\mathbb{N}, \mathcal{P}(\mathbb{N}), +1, \in)$  (ie. monadic second order theory of one successor).

**Tameness?** For  $r \in \mathbb{N}_{\geq 2}$ , consider a ternary predicate  $V_r(x, u, k)$  that holds if and only if  $u$  is an integer power of  $r$ ,  $k \in \{0, \dots, r-1\}$ , and the digit of some base  $r$  representation of  $x$  in the position corresponding to  $u$  is  $k$ .

Set  $\mathcal{T}_r := (\mathbb{R}, <, +, V_r)$ .

**Observation 1:**  $\mathcal{T}_r$  is bi-interpretable with  $(\mathbb{N}, \mathcal{P}(\mathbb{N}), +1, \in)$  (ie. monadic second order theory of one successor).

**Observation 2:** To understand definable sets in  $\mathcal{T}_r$  you need to use automata-theoretic methods.

**Tameness?** For  $r \in \mathbb{N}_{\geq 2}$ , consider a ternary predicate  $V_r(x, u, k)$  that holds if and only if  $u$  is an integer power of  $r$ ,  $k \in \{0, \dots, r-1\}$ , and the digit of some base  $r$  representation of  $x$  in the position corresponding to  $u$  is  $k$ .

Set  $\mathcal{T}_r := (\mathbb{R}, <, +, V_r)$ .


**Observation 1:**  $\mathcal{T}_r$  is bi-interpretable with  $(\mathbb{N}, \mathcal{P}(\mathbb{N}), +1, \in)$  (ie. monadic second order theory of one successor).

**Observation 2:** To understand definable sets in  $\mathcal{T}_r$  you need to use automata-theoretic methods.

**Observation 3:** The middle-thirds Cantor set is  $\emptyset$ -definable in  $\mathcal{T}_3$  (it is the set of those numbers in  $[0, 1]$  admitting a base 3 expansion that omits the digit 1.) and so is...




... the Sierpinski carpet



It is the set of pairs of real numbers  $(x, y) \in [0, 1]^2$  such that for every positive integer  $n$  the  $n$ -th digit of two ternary expansions of  $x$  and of  $y$  are not both equal to 1.

... the Sierpinski carpet



It is the set of pairs of real numbers  $(x, y) \in [0, 1]^2$  such that for every positive integer  $n$  the  $n$ -th digit of two ternary expansions of  $x$  and of  $y$  are not both equal to 1.

Same for the Menger sponge and other famous fractals. See also Adamczewski, Bell, *'An analogue of Cobham's Theorem for Fractals'*, TAMS 2011

Reminder:  $\mathcal{T}_r := (\mathbb{R}, <, +, V_r)$ .

Let  $D_r$  be the set of numbers in  $[0, 1)$  admitting a finite base  $r$  expansion.

Reminder:  $\mathcal{T}_r := (\mathbb{R}, <, +, V_r)$ .

Let  $D_r$  be the set of numbers in  $[0, 1)$  admitting a finite base  $r$  expansion.

**Observation:**  $D_r$  is definable in  $\mathcal{T}_r$  and dense in  $[0, 1]$ .

Reminder:  $\mathcal{T}_r := (\mathbb{R}, <, +, V_r)$ .

Let  $D_r$  be the set of numbers in  $[0, 1)$  admitting a finite base  $r$  expansion.

**Observation:**  $D_r$  is definable in  $\mathcal{T}_r$  and dense in  $[0, 1]$ .

Define  $\tau_r: D_r \rightarrow r^{-\mathbb{N}_{>0}}$  so that  $\tau_r(d)$  is the least  $u \in r^{-\mathbb{N}_{>0}}$  appearing with nonzero coefficient in the finite base  $r$  expansion of  $d$ .

Reminder:  $\mathcal{T}_r := (\mathbb{R}, <, +, V_r)$ .

Let  $D_r$  be the set of numbers in  $[0, 1)$  admitting a finite base  $r$  expansion.

**Observation:**  $D_r$  is definable in  $\mathcal{T}_r$  and dense in  $[0, 1]$ .

Define  $\tau_r: D_r \rightarrow r^{-\mathbb{N}_{>0}}$  so that  $\tau_r(d)$  is the least  $u \in r^{-\mathbb{N}_{>0}}$  appearing with nonzero coefficient in the finite base  $r$  expansion of  $d$ .

For  $d, e \in D_r$ , let

$$d \prec_r e \iff \tau_r(d) > \tau_r(e) \text{ or } (\tau_r(d) = \tau_r(e) \text{ and } d < e).$$

Reminder:  $\mathcal{T}_r := (\mathbb{R}, <, +, V_r)$ .

Let  $D_r$  be the set of numbers in  $[0, 1)$  admitting a finite base  $r$  expansion.

**Observation:**  $D_r$  is definable in  $\mathcal{T}_r$  and dense in  $[0, 1]$ .

Define  $\tau_r: D_r \rightarrow r^{-\mathbb{N}_{>0}}$  so that  $\tau_r(d)$  is the least  $u \in r^{-\mathbb{N}_{>0}}$  appearing with nonzero coefficient in the finite base  $r$  expansion of  $d$ .

For  $d, e \in D_r$ , let

$$d \prec_r e \iff \tau_r(d) > \tau_r(e) \text{ or } (\tau_r(d) = \tau_r(e) \text{ and } d < e).$$

**Observation:**  $(D_r, \prec)$  is definable in  $\mathcal{T}_r$  and has order type  $\omega$ .

**Definition.** An infinite definable subset of  $\mathbb{R}^n$  is  $\omega$ -**orderable** if it admits a definable ordering with order type  $\omega$ . We say an expansion  $\mathcal{R}$  of  $(\mathbb{R}, <, +)$  **defines a dense  $\omega$ -orderable set** if it defines an  $\omega$ -orderable subset of  $\mathbb{R}$  that is dense in some open interval.



**Definition.** An infinite definable subset of  $\mathbb{R}^n$  is  $\omega$ -**orderable** if it admits a definable ordering with order type  $\omega$ . We say an expansion  $\mathcal{R}$  of  $(\mathbb{R}, <, +)$  **defines a dense  $\omega$ -orderable set** if it defines an  $\omega$ -orderable subset of  $\mathbb{R}$  that is dense in some open interval.

### Observations.

- If  $D$  is  $\omega$ -orderable, so is  $D^k$ .

**Definition.** An infinite definable subset of  $\mathbb{R}^n$  is  $\omega$ -**orderable** if it admits a definable ordering with order type  $\omega$ . We say an expansion  $\mathcal{R}$  of  $(\mathbb{R}, <, +)$  **defines a dense  $\omega$ -orderable set** if it defines an  $\omega$ -orderable subset of  $\mathbb{R}$  that is dense in some open interval.

### Observations.

- If  $D$  is  $\omega$ -orderable, so is  $D^k$ .
- If  $D$  is  $\omega$ -orderable and  $f : D \rightarrow \mathbb{R}$  is definable, then  $f(D)$  is  $\omega$ -orderable (if  $f(D)$  is infinite).

**Definition.** An infinite definable subset of  $\mathbb{R}^n$  is  $\omega$ -**orderable** if it admits a definable ordering with order type  $\omega$ . We say an expansion  $\mathcal{R}$  of  $(\mathbb{R}, <, +)$  **defines a dense  $\omega$ -orderable set** if it defines an  $\omega$ -orderable subset of  $\mathbb{R}$  that is dense in some open interval.

### Observations.

- If  $D$  is  $\omega$ -orderable, so is  $D^k$ .
- If  $D$  is  $\omega$ -orderable and  $f : D \rightarrow \mathbb{R}$  is definable, then  $f(D)$  is  $\omega$ -orderable (if  $f(D)$  is infinite).
- If  $D \subseteq \mathbb{R}$  is infinite, closed and discrete, then  $D$  is  $\omega$ -orderable.

**Definition.** An infinite definable subset of  $\mathbb{R}^n$  is  $\omega$ -**orderable** if it admits a definable ordering with order type  $\omega$ . We say an expansion  $\mathcal{R}$  of  $(\mathbb{R}, <, +)$  **defines a dense  $\omega$ -orderable set** if it defines an  $\omega$ -orderable subset of  $\mathbb{R}$  that is dense in some open interval.

### Observations.

- If  $D$  is  $\omega$ -orderable, so is  $D^k$ .
- If  $D$  is  $\omega$ -orderable and  $f : D \rightarrow \mathbb{R}$  is definable, then  $f(D)$  is  $\omega$ -orderable (if  $f(D)$  is infinite).
- If  $D \subseteq \mathbb{R}$  is infinite, closed and discrete, then  $D$  is  $\omega$ -orderable.
- If  $D \subseteq \mathbb{R}$  is closed and discrete set and  $f : D^n \rightarrow \mathbb{R}$  is such that  $f(D^n)$  is somewhere dense, then  $f(D^n)$  is a dense  $\omega$ -orderable set.

**Definition.** An infinite definable subset of  $\mathbb{R}^n$  is  $\omega$ -**orderable** if it admits a definable ordering with order type  $\omega$ . We say an expansion  $\mathcal{R}$  of  $(\mathbb{R}, <, +)$  **defines a dense  $\omega$ -orderable set** if it defines an  $\omega$ -orderable subset of  $\mathbb{R}$  that is dense in some open interval.

### Observations.

- If  $D$  is  $\omega$ -orderable, so is  $D^k$ .
- If  $D$  is  $\omega$ -orderable and  $f : D \rightarrow \mathbb{R}$  is definable, then  $f(D)$  is  $\omega$ -orderable (if  $f(D)$  is infinite).
- If  $D \subseteq \mathbb{R}$  is infinite, closed and discrete, then  $D$  is  $\omega$ -orderable.
- If  $D \subseteq \mathbb{R}$  is closed and discrete set and  $f : D^n \rightarrow \mathbb{R}$  is such that  $f(D^n)$  is somewhere dense, then  $f(D^n)$  is a dense  $\omega$ -orderable set.

**Example.**  $(\mathbb{R}, <, +, \mathbb{Z}, x \mapsto \lambda x)$  defines  $n \in \mathbb{Z} \mapsto \lambda n - \lfloor \lambda n \rfloor$ , and thus a dense  $\omega$ -orderable set.

**Trivial Trichotomy.** An expansion  $\mathcal{R}$  of  $(\mathbb{R}, <, +)$  satisfies exactly one of the following three conditions:

- (A)  $\mathcal{R}$  does not define a dense  $\omega$ -orderable set.
- (B)  $\mathcal{R}$  defines a dense  $\omega$ -orderable, but avoids a compact set.
- (C)  $\mathcal{R}$  defines every compact set.

**Trivial Trichotomy.** An expansion  $\mathcal{R}$  of  $(\mathbb{R}, <, +)$  satisfies exactly one of the following three conditions:

- (A)  $\mathcal{R}$  does not define a dense  $\omega$ -orderable set.
- (B)  $\mathcal{R}$  defines a dense  $\omega$ -orderable, but avoids a compact set.
- (C)  $\mathcal{R}$  defines every compact set.

**Connections to Neostability.** A type B expansions interprets  $(\mathbb{N}, \mathcal{P}(\mathbb{N}), +1, \in)$  and therefore violates all known Shelah-style combinatorial tameness properties such as NIP or NTP2 or  $n$ -dependent.

**Trivial Trichotomy.** An expansion  $\mathcal{R}$  of  $(\mathbb{R}, <, +)$  satisfies exactly one of the following three conditions:

- (A)  $\mathcal{R}$  does not define a dense  $\omega$ -orderable set.
- (B)  $\mathcal{R}$  defines a dense  $\omega$ -orderable, but avoids a compact set.
- (C)  $\mathcal{R}$  defines every compact set.

**Connections to Neostability.** A type B expansions interprets  $(\mathbb{N}, \mathcal{P}(\mathbb{N}), +1, \in)$  and therefore violates all known Shelah-style combinatorial tameness properties such as NIP or NTP2 or  $n$ -dependent.

**Connections to Decidability.** All type C expansions have undecidable theories. There are expansions of type B with decidable theories. There are expansions of type A with undecidable theories.



**Trivial Trichotomy.** An expansion  $\mathcal{R}$  of  $(\mathbb{R}, <, +)$  satisfies exactly one of the following three conditions:

- (A)  $\mathcal{R}$  does not define a dense  $\omega$ -orderable set.
- (B)  $\mathcal{R}$  defines a dense  $\omega$ -orderable, but avoids a compact set.
- (C)  $\mathcal{R}$  defines every compact set.

**Connections to Neostability.** A type B expansions interprets  $(\mathbb{N}, \mathcal{P}(\mathbb{N}), +1, \in)$  and therefore violates all known Shelah-style combinatorial tameness properties such as NIP or NTP2 or  $n$ -dependent.

**Connections to Decidability.** All type C expansions have undecidable theories. There are expansions of type B with decidable theories. There are expansions of type A with undecidable theories.

*'What about decidability of the theory? Just as biological taxonomy does not tell us whether a species is tasty, the classification here does not deal with decidability.'* - Saharon Shelah

Type A:  $\mathcal{R}$  does not define a dense  $\omega$ -orderable set.

*Motto:* In Type A the definable sets are geometrically/topologically tame.

Type A:  $\mathcal{R}$  does not define a dense  $\omega$ -orderable set.

*Motto:* In Type A the definable sets are geometrically/topologically tame. This means (weak) analogues of results known for o-minimal expansions hold in this setting.

- Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a definable continuous function. Then there is a definable open dense  $U \subseteq \mathbb{R}$  such that  $f$  is strictly increasing, strictly decreasing, or constant on each connected component of  $U$ .

Type A:  $\mathcal{R}$  does not define a dense  $\omega$ -orderable set.

*Motto:* In Type A the definable sets are geometrically/topologically tame. This means (weak) analogues of results known for o-minimal expansions hold in this setting.

- Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a definable continuous function. Then there is a definable open dense  $U \subseteq \mathbb{R}$  such that  $f$  is strictly increasing, strictly decreasing, or constant on each connected component of  $U$ .
- Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a definable continuous function and  $k \in \mathbb{N}$ . Then there is a definable open dense  $U \subseteq \mathbb{R}$  such that the restriction of  $f$  to  $U$  is  $C^k$ .

Type A:  $\mathcal{R}$  does not define a dense  $\omega$ -orderable set.

*Motto:* In Type A the definable sets are geometrically/topologically tame. This means (weak) analogues of results known for o-minimal expansions hold in this setting.

- Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a definable continuous function. Then there is a definable open dense  $U \subseteq \mathbb{R}$  such that  $f$  is strictly increasing, strictly decreasing, or constant on each connected component of  $U$ .
- Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a definable continuous function and  $k \in \mathbb{N}$ . Then there is a definable open dense  $U \subseteq \mathbb{R}$  such that the restriction of  $f$  to  $U$  is  $C^k$ .
- Every nowhere dense definable subset of  $\mathbb{R}^n$  is Lebesgue null.

Type A:  $\mathcal{R}$  does not define a dense  $\omega$ -orderable set.

*Motto:* In Type A the definable sets are geometrically/topologically tame. This means (weak) analogues of results known for o-minimal expansions hold in this setting.

- Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a definable continuous function. Then there is a definable open dense  $U \subseteq \mathbb{R}$  such that  $f$  is strictly increasing, strictly decreasing, or constant on each connected component of  $U$ .
- Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a definable continuous function and  $k \in \mathbb{N}$ . Then there is a definable open dense  $U \subseteq \mathbb{R}$  such that the restriction of  $f$  to  $U$  is  $C^k$ .
- Every nowhere dense definable subset of  $\mathbb{R}^n$  is Lebesgue null.
- There is no continuous definable bijection from  $\mathbb{R}^m$  to  $\mathbb{R}^n$  when  $m < n$ .

Type A:  $\mathcal{R}$  does not define a dense  $\omega$ -orderable set.

The previous results all depend on the following

### **Strong Baire Category Theorem.**

Let  $\{X_{r,s} : r, s > 0\}$  be a definable family of closed subsets of  $\mathbb{R}^n$  such that  $X_{r,s} \subseteq X_{r',s'}$  when  $r \leq r'$  and  $s \geq s'$ . If  $\bigcup_{r,s} X_{r,s}$  is somewhere dense, then there are  $r', s' > 0$  such that  $X_{r',s'}$  has interior.

Type A:  $\mathcal{R}$  does not define a dense  $\omega$ -orderable set.

The previous results all depend on the following

### **Strong Baire Category Theorem.**

Let  $\{X_{r,s} : r, s > 0\}$  be a definable family of closed subsets of  $\mathbb{R}^n$  such that  $X_{r,s} \subseteq X_{r',s'}$  when  $r \leq r'$  and  $s \geq s'$ . If  $\bigcup_{r,s} X_{r,s}$  is somewhere dense, then there are  $r', s' > 0$  such that  $X_{r',s'}$  has interior.

**Special case.** Let  $\{X_r : r > 0\}$  be a definable family of *finite* subsets of  $\mathbb{R}^n$  such that  $X_r \subseteq X_{r'}$  when  $r \leq r'$ . Then  $X_\infty := \bigcup_r X_r$  is nowhere dense.



Type A:  $\mathcal{R}$  does not define a dense  $\omega$ -orderable set.

The previous results all depend on the following

### Strong Baire Category Theorem.

Let  $\{X_{r,s} : r, s > 0\}$  be a definable family of closed subsets of  $\mathbb{R}^n$  such that  $X_{r,s} \subseteq X_{r',s'}$  when  $r \leq r'$  and  $s \geq s'$ . If  $\bigcup_{r,s} X_{r,s}$  is somewhere dense, then there are  $r', s' > 0$  such that  $X_{r',s'}$  has interior.

**Special case.** Let  $\{X_r : r > 0\}$  be a definable family of *finite* subsets of  $\mathbb{R}^n$  such that  $X_r \subseteq X_{r'}$  when  $r \leq r'$ . Then  $X_\infty := \bigcup_r X_r$  is nowhere dense.

**Proof.** For  $x \in X_\infty$ , let  $\delta(x) := \inf\{r > 0 : x \in X_r\}$ .

Type A:  $\mathcal{R}$  does not define a dense  $\omega$ -orderable set.

The previous results all depend on the following

### Strong Baire Category Theorem.

Let  $\{X_{r,s} : r, s > 0\}$  be a definable family of closed subsets of  $\mathbb{R}^n$  such that  $X_{r,s} \subseteq X_{r',s'}$  when  $r \leq r'$  and  $s \geq s'$ . If  $\bigcup_{r,s} X_{r,s}$  is somewhere dense, then there are  $r', s' > 0$  such that  $X_{r',s'}$  has interior.

**Special case.** Let  $\{X_r : r > 0\}$  be a definable family of *finite* subsets of  $\mathbb{R}^n$  such that  $X_r \subseteq X_{r'}$  when  $r \leq r'$ . Then  $X_\infty := \bigcup_r X_r$  is nowhere dense.

**Proof.** For  $x \in X_\infty$ , let  $\delta(x) := \inf\{r > 0 : x \in X_r\}$ . Then for  $x, y \in X_\infty$  set

$$x \prec y \text{ iff } (\delta(x), x) <_{\text{lex}} (\delta(y), y).$$

Type A:  $\mathcal{R}$  does not define a dense  $\omega$ -orderable set.

The previous results all depend on the following

### Strong Baire Category Theorem.

Let  $\{X_{r,s} : r, s > 0\}$  be a definable family of closed subsets of  $\mathbb{R}^n$  such that  $X_{r,s} \subseteq X_{r',s'}$  when  $r \leq r'$  and  $s \geq s'$ . If  $\bigcup_{r,s} X_{r,s}$  is somewhere dense, then there are  $r', s' > 0$  such that  $X_{r',s'}$  has interior.

**Special case.** Let  $\{X_r : r > 0\}$  be a definable family of *finite* subsets of  $\mathbb{R}^n$  such that  $X_r \subseteq X_{r'}$  when  $r \leq r'$ . Then  $X_\infty := \bigcup_r X_r$  is nowhere dense.

**Proof.** For  $x \in X_\infty$ , let  $\delta(x) := \inf\{r > 0 : x \in X_r\}$ . Then for  $x, y \in X_\infty$  set

$$x \prec y \text{ iff } (\delta(x), x) <_{\text{lex}} (\delta(y), y).$$

The order  $\prec$  has order type  $\omega$ . Thus  $X_\infty$  is nowhere dense.

Type B:  $\mathcal{R}$  defines a dense  $\omega$ -orderable, but avoids a compact set.

Type B:  $\mathcal{R}$  defines a dense  $\omega$ -orderable, but avoids a compact set.

**Fact.**  $\mathcal{R}$  interprets  $(\mathbb{N}, \mathcal{P}(\mathbb{N}), +1, \in)$ . To understand all definable sets in specific Type B expansions automata-theoretic methods are necessary. However, there are a few things we can say in general.

Type B:  $\mathcal{R}$  defines a dense  $\omega$ -orderable, but avoids a compact set.

**Fact.**  $\mathcal{R}$  interprets  $(\mathbb{N}, \mathcal{P}(\mathbb{N}), +1, \in)$ . To understand all definable sets in specific Type B expansions automata-theoretic methods are necessary. However, there are a few things we can say in general.

**Elgot-Rabin '66.** Let  $f : \mathbb{N} \rightarrow \mathbb{N}$  be such that  $f^{-1}(n)$  is infinite for every  $n \in \mathbb{N}$ . Then the theory of  $(\mathbb{N}, \mathcal{P}(\mathbb{N}), +1, \in, f)$  is undecidable.

Type B:  $\mathcal{R}$  defines a dense  $\omega$ -orderable, but avoids a compact set.

**Fact.**  $\mathcal{R}$  interprets  $(\mathbb{N}, \mathcal{P}(\mathbb{N}), +1, \in)$ . To understand all definable sets in specific Type B expansions automata-theoretic methods are necessary. However, there are a few things we can say in general.

**Elgot-Rabin '66.** Let  $f : \mathbb{N} \rightarrow \mathbb{N}$  be such that  $f^{-1}(n)$  is infinite for every  $n \in \mathbb{N}$ . Then the theory of  $(\mathbb{N}, \mathcal{P}(\mathbb{N}), +1, \in, f)$  is undecidable.

**H.-Tychonievich 2014.** If  $\mathcal{R}$  defines an order  $(D, \prec)$  and a function  $g : \mathbb{R}^3 \times D \rightarrow D$  such that

- ①  $(D, \prec)$  has order type  $\omega$  and  $D$  is dense in  $[0, 1]$ ,
- ② for every  $a, b \in [0, 1]$  and  $e, d \in D$  with  $a < b$  and  $e \preceq d$ ,

$\{c \in \mathbb{R} : g(c, a, b, d) = e\} \cap (a, b)$  has nonempty interior,

then  $\mathcal{R}$  is type C. Example

The previous theorem puts strong restriction on what kind of function are definable in type B expansions.

**Corollary.** Suppose  $\mathcal{R}$  defines a dense  $\omega$ -orderable set and a function  $f : [0, 1] \rightarrow \mathbb{R}$  that is  $C^2$ , but not affine. Then  $\mathcal{R}$  is type C.



The previous theorem puts strong restriction on what kind of function are definable in type B expansions.

**Corollary.** Suppose  $\mathcal{R}$  defines a dense  $\omega$ -orderable set and a function  $f : [0, 1] \rightarrow \mathbb{R}$  that is  $C^2$ , but not affine. Then  $\mathcal{R}$  is type C.

**Corollary.** Suppose  $\mathcal{R}$  defines a dense  $\omega$ -orderable set and a family  $\{f_x : x \in \mathbb{R}^I\}$  of linear functions  $[0, 1] \rightarrow \mathbb{R}$  which has infinitely many distinct elements. Then  $\mathcal{R}$  is type C.

The previous theorem puts strong restriction on what kind of function are definable in type B expansions.

**Corollary.** Suppose  $\mathcal{R}$  defines a dense  $\omega$ -orderable set and a function  $f : [0, 1] \rightarrow \mathbb{R}$  that is  $C^2$ , but not affine. Then  $\mathcal{R}$  is type C.

**Corollary.** Suppose  $\mathcal{R}$  defines a dense  $\omega$ -orderable set and a family  $\{f_x : x \in \mathbb{R}^I\}$  of linear functions  $[0, 1] \rightarrow \mathbb{R}$  which has infinitely many distinct elements. Then  $\mathcal{R}$  is type C.

**Corollary.** Suppose  $\mathcal{R}$  defines a dense  $\omega$ -orderable set and the function  $x \mapsto \lambda x$  for uncountably many  $\lambda$ . Then  $\mathcal{R}$  is type C.

The previous theorem puts strong restriction on what kind of function are definable in type B expansions.

**Corollary.** Suppose  $\mathcal{R}$  defines a dense  $\omega$ -orderable set and a function  $f : [0, 1] \rightarrow \mathbb{R}$  that is  $C^2$ , but not affine. Then  $\mathcal{R}$  is type C.

**Corollary.** Suppose  $\mathcal{R}$  defines a dense  $\omega$ -orderable set and a family  $\{f_x : x \in \mathbb{R}^I\}$  of linear functions  $[0, 1] \rightarrow \mathbb{R}$  which has infinitely many distinct elements. Then  $\mathcal{R}$  is type C.

**Corollary.** Suppose  $\mathcal{R}$  defines a dense  $\omega$ -orderable set and the function  $x \mapsto \lambda x$  for uncountably many  $\lambda$ . Then  $\mathcal{R}$  is type C.

Let  $\mathcal{M} := (\mathbb{R}, <, +, (x \mapsto \lambda x)_{x \in \mathbb{R}})$ .

The previous theorem puts strong restriction on what kind of function are definable in type B expansions.

**Corollary.** Suppose  $\mathcal{R}$  defines a dense  $\omega$ -orderable set and a function  $f : [0, 1] \rightarrow \mathbb{R}$  that is  $C^2$ , but not affine. Then  $\mathcal{R}$  is type C.

**Corollary.** Suppose  $\mathcal{R}$  defines a dense  $\omega$ -orderable set and a family  $\{f_x : x \in \mathbb{R}^I\}$  of linear functions  $[0, 1] \rightarrow \mathbb{R}$  which has infinitely many distinct elements. Then  $\mathcal{R}$  is type C.

**Corollary.** Suppose  $\mathcal{R}$  defines a dense  $\omega$ -orderable set and the function  $x \mapsto \lambda x$  for uncountably many  $\lambda$ . Then  $\mathcal{R}$  is type C.

Let  $\mathcal{M} := (\mathbb{R}, <, +, (x \mapsto \lambda x)_{x \in \mathbb{R}})$ .

**Corollary.** Expansions of  $\mathcal{M}$  are either type A or type C.

## Future direction.

**Connections to neostability.**  $\text{NTP2}, \text{NIP} \Rightarrow \text{Type A}$ . Does NIP (NTP2) imply stronger geometric tameness than Type A? (cp. Dolich and Goodrick's work on strong theories)

## Future direction.

**Connections to neostability.**  $\text{NTP}_2, \text{NIP} \Rightarrow \text{Type A}$ . Does NIP ( $\text{NTP}_2$ ) imply stronger geometric tameness than Type A? (cp. Dolich and Goodrick's work on strong theories)

**Connections to descriptive set theory.** We say an expansion is Borel if every definable set is Borel. Type B expansions can be Borel. Some Type A expansions are not Borel. What can we say about expansions that are Borel? (cp. work on Borel structures)

## Future direction.

**Connections to neostability.**  $\text{NTP2}, \text{NIP} \Rightarrow \text{Type A}$ . Does NIP (NTP2) imply stronger geometric tameness than Type A? (cp. Dolich and Goodrick's work on strong theories)

**Connections to descriptive set theory.** We say an expansion is Borel if every definable set is Borel. Type B expansions can be Borel. Some Type A expansions are not Borel. What can we say about expansions that are Borel? (cp. work on Borel structures)

**Tame geometry.** Let  $\mathcal{R}$  be a type A expansions of  $\mathcal{M}$  by closed sets. Does every definable subset of  $\mathbb{R}$  has interior or is nowhere dense?

Thank you!