

# Around and about real closed valued fields

Deirdre Haskell

presenting joint work with Clifton Ealy and Jana Maříková

supported by an NSERC discovery grant

McMaster University

Summer school in tame geometry  
Konstanz, Germany, July 18–23, 2016

# ordered valued field

Let  $\mathcal{R}$  be a real closed field, consider the field of Laurent series  $\mathcal{R}((t))$

$$\mathcal{R}((t)) = \left\{ \sum_{i=N}^{\infty} a_i t^i : a_i \in \mathcal{R}, N \in \mathbb{Z} \right\}.$$

$\mathcal{R}((t))$  can be ordered in many different ways; let's consider  $t > 0$ ,  $t < r$  for any positive  $r \in \mathcal{R}$ . Then

$$t > t^2 > t^3 > \dots > 0,$$

and in general,

$$\sum_{i=N}^{\infty} a_i t^i < \sum_{i=N}^{\infty} b_i t^i \iff a_N < b_N.$$

A picture

A picture  
Another picture

# valued field

A valuation is a function  $v : K \rightarrow \Gamma \cup \{\infty\}$ , where  $K$  is a field,  $\Gamma$  is an ordered abelian group, such that

$$\begin{aligned}v(xy) &= v(x) + v(y) \\v(x + y) &\geq \min\{v(x), v(y)\} \\v(x) = \infty &\leftrightarrow x = 0\end{aligned}$$

# valued field

A valuation is a function  $v : K \rightarrow \Gamma \cup \{\infty\}$ , where  $K$  is a field,  $\Gamma$  is an ordered abelian group, such that

$$\begin{aligned}v(xy) &= v(x) + v(y) \\v(x + y) &\geq \min\{v(x), v(y)\} \\v(x) = \infty &\leftrightarrow x = 0\end{aligned}$$

Valuation ring  $\mathcal{O}_K = \{x \in K : v(x) \geq 0\}$

Maximal ideal  $\mathfrak{m}_K = \{x \in K : v(x) > 0\}$

Residue field  $k = \mathcal{O}_K / \mathfrak{m}_K$ .

In the case of  $K = \mathcal{R}((t))$ ,

$$v\left(\sum_{i=N}^{\infty} a_i t^i\right) = N,$$

$\mathcal{O}_K = \mathcal{R}[[t]]$  and the residue field is  $\mathcal{R}$ .

In the case of  $K = \mathcal{R}((t))$ ,

$$v\left(\sum_{i=N}^{\infty} a_i t^i\right) = N,$$

$\mathcal{O}_K = \mathcal{R}[[t]]$  and the residue field is  $\mathcal{R}$ .

With the given ordering,  $\mathcal{O}_K$  is a *convex* subring: if  $x, y \in \mathcal{O}_K$  and  $x < z < y$  then  $z \in \mathcal{O}_K$ .



In the case of  $K = \mathcal{R}((t))$ ,

$$v\left(\sum_{i=N}^{\infty} a_i t^i\right) = N,$$

$\mathcal{O}_K = \mathcal{R}[[t]]$  and the residue field is  $\mathcal{R}$ .

With the given ordering,  $\mathcal{O}_K$  is a *convex* subring: if  $x, y \in \mathcal{O}_K$  and  $x < z < y$  then  $z \in \mathcal{O}_K$ .

More generally, a valuation  $v$  on a field is *convex* with respect to an ordering  $<$  on the field if for all  $0 < x < y$ ,  $v(x) \geq v(y)$ .

Let  $\mathcal{R}$  be an ordered field with a convex valuation.

1) If  $\mathcal{R}$  is real closed then  $\Gamma_{\mathcal{R}}$  is divisible and  $k_{\mathcal{R}}$  is real closed.

# closure properties

Let  $\mathcal{R}$  be an ordered field with a convex valuation.

- 1) If  $\mathcal{R}$  is real closed then  $\Gamma_{\mathcal{R}}$  is divisible and  $k_{\mathcal{R}}$  is real closed.
- 2) If  $\Gamma_{\mathcal{R}}$  is divisible,  $k_{\mathcal{R}}$  is real closed and  $\mathcal{R}$  is henselian then  $\mathcal{R}$  is real closed.

Analogous to algebraically closed valued fields.

# AKE theorem

## Ax-Kochen, Ersov, late 60's

Let  $K$  be a henselian valued field of characteristic 0. Then  $\text{Th}(K)$  in the language of valued fields is determined up to elementary equivalence by  $\text{Th}(\Gamma)$  in the language of ordered groups and  $\text{Th}(k)$  in the language of fields.

# AKE theorem

## Ax-Kochen, Ersov, late 60's

Let  $K$  be a henselian valued field of characteristic 0. Then  $\text{Th}(K)$  in the language of valued fields is determined up to elementary equivalence by  $\text{Th}(\Gamma)$  in the language of ordered groups and  $\text{Th}(k)$  in the language of fields.

Motto: a valued field is controlled by its value group and residue field.

Motivation: to what extent is this still true when further structure is added?

More generally: pursue analogies between algebraically closed valued fields and real closed valued fields.

# model theory of pure real closed valued fields

## Cherlin-Dickmann 1983

The theory RCVF of real closed fields with a proper convex valuation has quantifier elimination in the language of rings with predicates for the ordering and the valuation ring.

# model theory of pure real closed valued fields

## Cherlin-Dickmann 1983

The theory RCVF of real closed fields with a proper convex valuation has quantifier elimination in the language of rings with predicates for the ordering and the valuation ring.

## Mellor 2006

The theory RCVF has elimination of imaginaries in the sorted language  $\mathcal{G}$  with sorts for the finitely generated  $\mathcal{O}_K$ -submodules and their torsors.

Again, analogous to algebraically closed valued fields.

## some imaginaries in valued fields

The value group  $K^\times / (\mathcal{O}_K \setminus \mathfrak{m}_K)$

$aEb$  if and only if  $a/b \in \mathcal{O}_K \setminus \mathfrak{m}_K$  if and only if  $v(a) = v(b)$



## some imaginaries in valued fields

The value group  $K^\times/(\mathcal{O}_K \setminus \mathfrak{m}_K)$

$aEb$  if and only if  $a/b \in \mathcal{O}_K \setminus \mathfrak{m}_K$  if and only if  $v(a) = v(b)$

The residue field  $\mathcal{O}_K/\mathfrak{m}$

$aEb$  if and only if  $a - b \in \mathfrak{m}$  if and only if  $v(a - b) > 0$

## some imaginaries in valued fields

The value group  $K^\times / (\mathcal{O}_K \setminus \mathfrak{m}_K)$

$aEb$  if and only if  $a/b \in \mathcal{O}_K \setminus \mathfrak{m}_K$  if and only if  $v(a) = v(b)$

The residue field  $\mathcal{O}_K/\mathfrak{m}$

$aEb$  if and only if  $a - b \in \mathfrak{m}$  if and only if  $v(a - b) > 0$

The RV sorts  $K^\times / (1 + \mathfrak{m})$

$aEb$  if and only if  $a(1 + \mathfrak{m}) = b(1 + \mathfrak{m})$  if and only if  $v(a - b) > v(b)$

# a pure valued fields theorem

H.-Hrushovski-Macpherson 2008

Let  $L, M, C$  be substructures of  $\mathcal{U} \models \text{ACVF}$ . Assume that  $C$  is maximal and is a substructure of both  $L$  and  $M$ .

# a pure valued fields theorem

## H.-Hrushovski-Macpherson 2008

Let  $L, M, C$  be substructures of  $\mathcal{U} \models \text{ACVF}$ . Assume that  $C$  is maximal and is a substructure of both  $L$  and  $M$ .

Also assume that  $\Gamma_L \cap \Gamma_M = \Gamma_C$  and that  $k_L$  is linearly independent from  $k_M$  over  $k_C$ .

# a pure valued fields theorem

## H.-Hrushovski-Macpherson 2008

Let  $L, M, C$  be substructures of  $\mathcal{U} \models \text{ACVF}$ . Assume that  $C$  is maximal and is a substructure of both  $L$  and  $M$ .

Also assume that  $\Gamma_L \cap \Gamma_M = \Gamma_C$  and that  $k_L$  is linearly independent from  $k_M$  over  $k_C$ .

Suppose there is an isomorphism  $\sigma : L \rightarrow L'$  fixing  $C, \Gamma_L$  and  $k_L$ .

# a pure valued fields theorem

## H.-Hrushovski-Macpherson 2008

Let  $L, M, C$  be substructures of  $\mathcal{U} \models \text{ACVF}$ . Assume that  $C$  is maximal and is a substructure of both  $L$  and  $M$ .

Also assume that  $\Gamma_L \cap \Gamma_M = \Gamma_C$  and that  $k_L$  is linearly independent from  $k_M$  over  $k_C$ .

Suppose there is an isomorphism  $\sigma : L \rightarrow L'$  fixing  $C$ ,  $\Gamma_L$  and  $k_L$ .

Then  $\sigma$  can be extended to an isomorphism of the field generated by  $L$  and  $M$  over  $C$ , which is the identity on  $M$ .

# a pure valued fields theorem

## H.-Hrushovski-Macpherson 2008

Let  $L, M, C$  be substructures of  $\mathcal{U} \models \text{ACVF}$ . Assume that  $C$  is maximal and is a substructure of both  $L$  and  $M$ .

Also assume that  $\Gamma_L \cap \Gamma_M = \Gamma_C$  and that  $k_L$  is linearly independent from  $k_M$  over  $k_C$ .

Suppose there is an isomorphism  $\sigma : L \rightarrow L'$  fixing  $C$ ,  $\Gamma_L$  and  $k_L$ .

Then  $\sigma$  can be extended to an isomorphism of the field generated by  $L$  and  $M$  over  $C$ , which is the identity on  $M$ .

Equivalently,

$$\text{tp}(M/C\Gamma_L k_L) \vdash \text{tp}(M/L).$$

# an ordered valued fields theorem

Theorem (Ealy-H.-Maříková 2016)

The analogous statement with  $\mathcal{U} \models \text{RCVF}$ .



Define  $\sigma : C[L, M] \rightarrow C[L', M]$  by  $\sigma(\sum_{i=1}^n \ell_i m_i) = \sum_{i=1}^n \sigma(\ell_i) m_i$ .

Define  $\sigma : C[L, M] \rightarrow C[L', M]$  by  $\sigma(\sum_{i=1}^n \ell_i m_i) = \sum_{i=1}^n \sigma(\ell_i) m_i$ .

1) Show  $\sigma$  preserves the valuation (HHM, also Johnson).

Because  $C$  is maximal, WMA that the  $m_i$  are a *separated* basis for the finite-dimensional vector subspace of  $M$  that they generate over  $C$  such that in addition  $v(\sum \ell_i m_i) = \min_i \{v(\ell_i) + v(m_i)\}$ .

# outline of proof

Define  $\sigma : C[L, M] \rightarrow C[L', M]$  by  $\sigma(\sum_{i=1}^n \ell_i m_i) = \sum_{i=1}^n \sigma(\ell_i) m_i$ .

1) Show  $\sigma$  preserves the valuation (HHM, also Johnson).

Because  $C$  is maximal, WMA that the  $m_i$  are a *separated* basis for the finite-dimensional vector subspace of  $M$  that they generate over  $C$  such that in addition  $v(\sum \ell_i m_i) = \min_i \{v(\ell_i) + v(m_i)\}$ .

As  $\sigma$  fixes  $\Gamma_M$  and is an isomorphism on  $\Gamma_L$ , the result follows.

2) Show that  $\sigma$  also preserves the ordering.

2) Show that  $\sigma$  also preserves the ordering.

Suppose not. Then there is some  $a = \sum_{i=1}^n \ell_i m_i$  and  $m_{n+1} \in M$  with

$$0 < \sigma(a) < m_{n+1} < a$$

and WMA  $\{m_i\}$  is a separated basis, and that  $a$  is a shortest counterexample.

2) Show that  $\sigma$  also preserves the ordering.

Suppose not. Then there is some  $a = \sum_{i=1}^n \ell_i m_i$  and  $m_{n+1} \in M$  with

$$0 < \sigma(a) < m_{n+1} < a$$

and WMA  $\{m_i\}$  is a separated basis, and that  $a$  is a shortest counterexample. We know that  $v(\sigma(a)) = v(a)$ , hence also  $v(\sigma(m_{n+1})) = v(a)$ .

## outline of proof

2) Show that  $\sigma$  also preserves the ordering.

Suppose not. Then there is some  $a = \sum_{i=1}^n \ell_i m_i$  and  $m_{n+1} \in M$  with

$$0 < \sigma(a) < m_{n+1} < a$$

and WMA  $\{m_i\}$  is a separated basis, and that  $a$  is a shortest counterexample.

We know that  $v(\sigma(a)) = v(a)$ , hence also  $v(\sigma(m_{n+1})) = v(a)$ .

For any  $x = \sum c_i m_i$ , show that

- it is not the case that  $\sigma(a) < x < a$ ;

## outline of proof

2) Show that  $\sigma$  also preserves the ordering.

Suppose not. Then there is some  $a = \sum_{i=1}^n \ell_i m_i$  and  $m_{n+1} \in M$  with

$$0 < \sigma(a) < m_{n+1} < a$$

and WMA  $\{m_i\}$  is a separated basis, and that  $a$  is a shortest counterexample.

We know that  $v(\sigma(a)) = v(a)$ , hence also  $v(\sigma(m_{n+1})) = v(a)$ .

For any  $x = \sum c_i m_i$ , show that

- it is not the case that  $\sigma(a) < x < a$ ;
- it is not the case that  $v(a - x) > v(a - \sigma(a))$ ,

hence there is a closest (in the valuation sense) such element  $x$  to  $a$ .



## outline of proof

2) Show that  $\sigma$  also preserves the ordering.

Suppose not. Then there is some  $a = \sum_{i=1}^n \ell_i m_i$  and  $m_{n+1} \in M$  with

$$0 < \sigma(a) < m_{n+1} < a$$

and WMA  $\{m_i\}$  is a separated basis, and that  $a$  is a shortest counterexample.

We know that  $v(\sigma(a)) = v(a)$ , hence also  $v(\sigma(m_{n+1})) = v(a)$ .

For any  $x = \sum c_i m_i$ , show that

- it is not the case that  $\sigma(a) < x < a$ ;
- it is not the case that  $v(a - x) > v(a - \sigma(a))$ ,

hence there is a closest (in the valuation sense) such element  $x$  to  $a$ .

Now construct a pseudo-Cauchy sequence to find a closer such element  $x$  to  $a$ , giving a contradiction.

## a stronger theorem

### Theorem (Ealy-Haskell-Maříková 2016)

Assume  $L, M, C$  are elementary substructures of  $\mathcal{U} \models \text{RCVF}$ , with  $C$  a common substructure of  $L$  and  $M$ ,  $C$  maximal,  $\Gamma_L \subseteq \Gamma_M$ .

Assume that  $\text{kInt}_{C\Gamma_L}^L$  is algebraically independent from  $\text{kInt}_{C\Gamma_L}^M$  over  $C\Gamma_L$ .

## the $k$ -internal sorts

Given parameter set  $A$ ,  $kInt_A$  is the collection of sets, definable over  $A$ , that are *internal* to the residue field.

That is,  $E \subset \text{dcl}(k \cup a)$ , where  $a$  is a finite tuple from  $A$

Ex:  $E = a(1 + \mathfrak{m})$ .

## the $k$ -internal sorts

Given parameter set  $A$ ,  $kInt_A$  is the collection of sets, definable over  $A$ , that are *internal* to the residue field.

That is,  $E \subset \text{dcl}(k \cup a)$ , where  $a$  is a finite tuple from  $A$

Ex:  $E = a(1 + \mathfrak{m})$ .

Given fields  $C \subseteq L$ ,  $S \subseteq \Gamma(L)$

$$\text{kInt}_{CS}^L = \text{kInt}_{CS} \cap \text{acl}(Ck_L\{\text{RV}_\gamma(L)\}_{\gamma \in S})$$

## a stronger theorem

### Theorem (Ealy-Haskell-Maříková 2016)

Assume  $L, M, C$  are elementary substructures of  $\mathcal{U} \models \text{RCVF}$ , with  $C$  a common substructure of  $L$  and  $M$ ,  $C$  maximal,  $\Gamma_L \subseteq \Gamma_M$ .

Assume that  $\text{kInt}_{C\Gamma_L}^L$  is algebraically independent from  $\text{kInt}_{C\Gamma_L}^M$  over  $C\Gamma_L$ .

Let  $\sigma : L \rightarrow L'$  be an ordered valued field isomorphism fixing  $\text{kInt}_{C\Gamma_L}^M$ .

Then  $\sigma$  extends to an ordered valued field isomorphism from  $C(L, M)$  to  $C(L', M)$  fixing  $M$ .

## some remarks on the proof

Define  $\sigma$  as before.

## some remarks on the proof

Define  $\sigma$  as before.

Show that  $\sigma$  preserves the valuation: perturb the valuation to  $v'$  with  $\Gamma_{v'}(L) \cap \Gamma_{v'}(M) = \Gamma_{v'}(C)$ . Note that  $v'$  is no longer convex with respect to the ordering, so apply pure valued field version of previous theorem and deduce that  $\sigma$  preserves  $v'$  and hence (by construction) also preserves  $v$ .

## some remarks on the proof

Define  $\sigma$  as before.

Show that  $\sigma$  preserves the valuation: perturb the valuation to  $v'$  with  $\Gamma_{v'}(L) \cap \Gamma_{v'}(M) = \Gamma_{v'}(C)$ . Note that  $v'$  is no longer convex with respect to the ordering, so apply pure valued field version of previous theorem and deduce that  $\sigma$  preserves  $v'$  and hence (by construction) also preserves  $v$ .

Show that  $\sigma$  preserves the ordering: if not, then there is a change in an order relation on balls defined over  $M$ , which is a formula in the type of  $L$  over  $\text{kInt}_{\text{CT}_L}^M$ .



## perturbing the valuation

Choose  $a_1, \dots, a_r$  from  $L$  and  $e_1, \dots, e_r$  from  $M$  such that  $v(a_i) = v(e_i)$  and  $\{a_i\}$  is a  $\mathbb{Q}$ -basis for  $\Gamma_L$  over  $\Gamma_C$ .

Choose  $b_1, \dots, b_s$  from  $L$  such that  $\{\text{res}(b_i)\}$  is a transcendence basis for  $k_L$  over  $k_C$ .

The assumption that  $\text{kInt}_{C\Gamma_L}^L$  is algebraically independent from  $\text{kInt}_{C\Gamma_L}^M$  over  $C\Gamma_L$  is equivalent to saying the elements

$$\text{res}(a_1/e_1), \dots, \text{res}(a_r/e_r), \text{res}(b_1), \dots, \text{res}(b_s)$$

are algebraically independent over  $k_M$ .

## perturbing the valuation

Choose  $a_1, \dots, a_r$  from  $L$  and  $e_1, \dots, e_r$  from  $M$  such that  $v(a_i) = v(e_i)$  and  $\{a_i\}$  is a  $\mathbb{Q}$ -basis for  $\Gamma_L$  over  $\Gamma_C$ .

Choose  $b_1, \dots, b_s$  from  $L$  such that  $\{\text{res}(b_i)\}$  is a transcendence basis for  $k_L$  over  $k_C$ .

The assumption that  $\text{kInt}_{\Gamma_L}^L$  is algebraically independent from  $\text{kInt}_{\Gamma_L}^M$  over  $\Gamma_L$  is equivalent to saying the elements

$$\text{res}(a_1/e_1), \dots, \text{res}(a_r/e_r), \text{res}(b_1), \dots, \text{res}(b_s)$$

are algebraically independent over  $k_M$ .

For each  $0 \leq j \leq r-1$  choose a place

$$p^{(j)} : \text{dcl}(k_M, \text{res}(b_1), \dots, \text{res}(b_s), \text{res}(a_1/e_1), \dots, \text{res}(a_{j+1}/e_{j+1})) \rightarrow \\ \text{dcl}(k_M, \text{res}(b_1), \dots, \text{res}(b_s), \text{res}(a_1/e_1), \dots, \text{res}(a_j/e_j))$$

Let  $p_{v'} : \text{dcl}(C(L, M)) \rightarrow \text{dcl}(k_M, k_L)$  be the composition. Let  $v'$  be a valuation associated to the place  $p_{v'}$ .

Then  $\Gamma_{v'}(L) \cap \Gamma_{v'}(M) = \Gamma_{v'}(C)$  and  $k_{v'}(L), k_{v'}(M)$  are linearly disjoint over  $k_{v'}(C)$ .

## $\sigma$ preserves the ordering

Suppose not. Let  $a = \sum_{i=1}^n \ell_i m_i > 0$  be a minimal counterexample. As before, WMA that the  $m_i$  form a separated basis for the space that they generate over  $C$  with respect to  $v'$ . From the construction of  $v'$ , in fact, the basis is also separated over  $v$  and over  $L$ .

Hence we may assume that  $v(\ell_i m_i) = 0$ , so  $v(m_i) \in \Gamma_L$ .

## $\sigma$ preserves the ordering

Suppose not. Let  $a = \sum_{i=1}^n \ell_i m_i > 0$  be a minimal counterexample. As before, WMA that the  $m_i$  form a separated basis for the space that they generate over  $C$  with respect to  $v'$ . From the construction of  $v'$ , in fact, the basis is also separated over  $v$  and over  $L$ .

Hence we may assume that  $v(\ell_i m_i) = 0$ , so  $v(m_i) \in \Gamma_L$ .

Thus  $a > 0$  implies that  $a + \sum_{i=1}^n \ell_i d_i > 0$  for any  $d_i$  with  $v(d_i) > v(m_i) = -v(\ell_i)$ .

In other words, the formula

$$x_1 B_{v(m_1)}^{op}(m_1) + \cdots + x_n B_{v(m_n)}^{op}(m_n) > 0$$

is a formula in the type of  $L$  over  $\text{kInt}_{\Gamma_L}^M$ . Since we assumed  $\sigma$  preserves this type,  $\sigma(a) > 0$ .

## in the geometric sorts

As in previous theorems, let  $C, M$  be substructures of  $\mathcal{U} \models \text{RCVF}$ ,  $A = \text{dcl}(Ce)$ , where  $e$  is a tuple of imaginaries in  $\mathcal{G}$ .

- Suppose  $\Gamma_A \cap \Gamma_M = \Gamma_C$  and  $k_A$  and  $k_M$  are linearly independent over  $k_C$ . Then

$$\text{tp}(A/Ck_M\Gamma_M) \vdash \text{tp}(A/M).$$

- Suppose  $\text{kInt}_{C\Gamma_A}^M$  is independent from  $\text{kInt}_{C\Gamma_A}^A$ . Then

$$\text{tp}(A/C\Gamma_A \text{kInt}_{C\Gamma_A}^M) \vdash \text{tp}(A/M).$$

## in the geometric sorts

As in previous theorems, let  $C, M$  be substructures of  $\mathcal{U} \models \text{RCVF}$ ,  $A = \text{dcl}(Ce)$ , where  $e$  is a tuple of imaginaries in  $\mathcal{G}$ .

- Suppose  $\Gamma_A \cap \Gamma_M = \Gamma_C$  and  $k_A$  and  $k_M$  are linearly independent over  $k_C$ . Then

$$\text{tp}(A/Ck_M\Gamma_M) \vdash \text{tp}(A/M).$$

- Suppose  $\text{kInt}_{C\Gamma_A}^M$  is independent from  $\text{kInt}_{C\Gamma_A}^A$ . Then

$$\text{tp}(A/C\Gamma_A \text{kInt}_{C\Gamma_A}^M) \vdash \text{tp}(A/M).$$

Proof: Find a resolution of  $A$  in the field sort with same value group and residue field. Then apply previous theorems.

Extend to a general  $T$ -convex theory.

How do functions behave on the interaction of  $L$  and  $M$ ?

Resolutions still exist for the geometric sorts (provided the underlying o-minimal theory is power bounded). Are new sorts required to eliminate imaginaries?