

LOCALLY DEFINABLE SUBGROUPS OF SEMIALGEBRAIC GROUPS

ELÍAS BARO, PANTELIS E. ELEFThERIOU, AND YA'ACOV PETERZIL

ABSTRACT. We prove the following instance of a conjecture stated in [7].
Let G be an abelian semialgebraic group and X a semialgebraic subset of G . Then the group \mathcal{U} generated by X contains a generic semialgebraic set. In particular, \mathcal{U} is divisible.

1. INTRODUCTION

Let \mathcal{M} be an arbitrary κ -saturated o-minimal structure (for κ sufficiently large). A *locally definable group* is a group $\langle \mathcal{U}, \cdot \rangle$ whose universe is a directed union $\mathcal{U} = \bigcup_{k \in \mathbb{N}} X_k$ of definable subsets of M^n for some fixed n , and for every $i, j \in \mathbb{N}$, the restriction of group multiplication to $X_i \times X_j$ is a definable function (by saturation, its image is contained in some X_k). The dimension of \mathcal{U} is by definition $\dim(\mathcal{U}) = \max\{\dim(X_k) : k \in \mathbb{N}\}$.

A map $\phi : \mathcal{U} \rightarrow \mathcal{H}$ between locally definable groups is called *locally definable* if for every definable $X \subseteq \mathcal{U}$ and $Y \subseteq \mathcal{H}$, $\text{graph}(\phi) \cap (X \times Y)$ is a definable set. Equivalently, the restriction of ϕ to any definable set is a definable map.

For a locally definable group \mathcal{U} , we say that $\mathcal{V} \subseteq \mathcal{U}$ is a *compatible subset* of \mathcal{U} if for every definable $X \subseteq \mathcal{U}$, the intersection $X \cap \mathcal{V}$ is a definable set (note that in this case \mathcal{V} itself is a bounded union of definable sets). Note that if $\phi : \mathcal{U} \rightarrow \mathcal{V}$ is a locally definable homomorphism between locally definable groups, then $\ker(\phi)$ is a compatible locally definable normal subgroup of \mathcal{U} . Compatible subgroups are used in order to obtain locally definable quotients:

Fact 1.1. [5, Theorem 4.2] *If \mathcal{U} is a locally definable group and $\mathcal{H} \subseteq \mathcal{U}$ a locally definable normal subgroup then \mathcal{H} is a compatible subgroup of \mathcal{U} if and only if there exists a locally definable surjective homomorphism of locally definable groups $\phi : \mathcal{U} \rightarrow \mathcal{V}$ whose kernel is \mathcal{H} .*

We are mostly interested here in *definably generated* groups, namely locally definable groups \mathcal{U} which are generated as a group by a definable subset

Date: October 19, 2018.

2010 Mathematics Subject Classification. 03C64, 03C68, 22B99.

Key words and phrases. semialgebraic groups, locally definable groups, generic sets, lattices.

The first author was supported by the program MTM2017-82105-P. The second author was supported by an Independent Research Grant from the German Research Foundation (DFG) and a Zukunftskolleg Research Fellowship.

X . An important example of such groups is the universal cover of a definable group, see [9]. Another relevant class of such groups are the connected locally definable groups with a definable (*left*) *generic* set X , that is, a definable set such that $AX = \mathcal{U}$ for some countable subset $A \subseteq \mathcal{U}$ (see [6, Fact 3.2(2)]). In [7], the converse of the latter was conjectured in the abelian case:

Conjecture 1.2. *Let \mathcal{U} be an abelian, connected, definably generated group. Then \mathcal{U} contains a definable generic set.*

We will use the following notation:

Notation 1.3. For $X \subseteq \mathcal{U}$ we write $\Sigma_n X$ for the set $X \pm \cdots \pm X$. The set $X(m)$ denotes the addition of $X - X$ to itself m times.

It has been shown in recent papers that the above conjecture can be restated in several ways. For example, given \mathcal{U} abelian, connected, definably generated group we say that a locally definable normal subgroup $\Gamma < \mathcal{U}$ is a *lattice* if $\dim(\Gamma) = 0$ and \mathcal{U}/Γ is definable, that is, there exist a definable group G and a locally definable surjective homomorphism from \mathcal{U} onto G , whose kernel is Γ (note that Γ is necessarily compatible).

Fact 1.4 ([7, Proposition 3.5] and [6, Theorem 2.1]). *Let \mathcal{U} be an abelian, connected, definably generated group. Then there is k so that the following are equivalent:*

- (1) \mathcal{U} contains a definable generic set.
- (2) \mathcal{U} admits a lattice.
- (3) \mathcal{U} admits a lattice isomorphic to \mathbb{Z}^k .

Moreover, the above conclusions imply that \mathcal{U} is divisible.

Our aim in the present note is to study some cases of the above conjecture. In Section 2 we study the conjecture under the presence of an exact sequence, and we prove the following result:

Theorem. *Assume that we are given an exact sequence of abelian locally definable groups and maps.*

$$0 \longrightarrow \mathcal{H} \xrightarrow{i} \mathcal{G} \xrightarrow{\pi} \mathcal{V} \longrightarrow 0$$

Assume also that definably generated connected subgroups of \mathcal{V} and of \mathcal{H} contain a definable generic set. Then the same is true for \mathcal{G} .

This is a useful criterion that can be applied inductively in certain situations. For example, we will use it in Sections 3 and 4 to study semi-algebraically generated subgroups of semialgebraic groups. The following (Theorem 4.4) is one of the main results of the paper, which generalizes the results in [8].

Theorem. *Let G be an abelian semialgebraic group and let X be a semi-algebraic subset of G . Then the group generated by X contains a generic semialgebraic set.*

2. GROUP EXTENSIONS

In this section we study the existence of generic sets when dealing with abelian group extensions. As a corollary, we will prove that definable generated subgroups of abelian torsion-free definable groups contain generic sets.

Proposition 2.1. *Assume that we are given an exact sequence of abelian locally definable groups and maps,*

$$0 \longrightarrow \mathcal{H} \xrightarrow{i} \mathcal{G} \xrightarrow{\pi} \mathcal{V} \longrightarrow 0 ,$$

where \mathcal{V} is connected and admits a lattice. Let $Y \subseteq \mathcal{V}$ be a definable generic set and let $s : Y \rightarrow \mathcal{G}$ be a definable section. Then the intersection $\langle s(Y) \rangle \cap \mathcal{H}$ is definably generated.

Proof. By [6, Fact 3.2(2)], $\mathcal{V} = \langle Y \rangle$. In particular, π sends the group $\langle s(Y) \rangle$ onto \mathcal{V} .

Henceforth we will use that given a definable set $Z \subseteq \mathcal{V}$, we can assume that $Z \subseteq Y$. Indeed, by saturation there is n such that $Z \subseteq \Sigma_n Y$ and by definable choice there is a section $r : Z \rightarrow \Sigma_n s(Y) \subseteq \langle s(Y) \rangle$. Thus we can extend the section $s : Y \rightarrow \mathcal{G}$ to a section $\tilde{s} : Y \cup (Z \setminus Y) \rightarrow \mathcal{G}$ via r in such a way that $\langle s(Y) \rangle = \langle \tilde{s}(Y \cup Z) \rangle$. Therefore we can work with the generic set $Y \cup Z$ instead of Y , as required.

By Fact 1.4 and since Y is generic, the locally definable group \mathcal{V} admits a lattice $\Gamma \simeq \mathbb{Z}^k$. Since \mathcal{V}/Γ is definable and Y generic in \mathcal{V} , there is a finite set $A \subseteq \mathcal{V}$ such that $Y + A + \Gamma = \mathcal{V}$. Without loss we can assume that A contains a fixed set of generators $\gamma_1, \dots, \gamma_k$ of Γ . Therefore we can assume that $Y + \Gamma = \mathcal{V}$ and $\gamma_1, \dots, \gamma_k \in Y$ (extending the section s to $Y + A$). Moreover, we can assume that Y is symmetric (again extend the section s to set $-Y$).

Let $\Delta = \langle s(\gamma_1), \dots, s(\gamma_k) \rangle$ and note that $\pi|_{\Delta} : \Delta \rightarrow \Gamma$ is an isomorphism. Let Δ_0 be the symmetric finite set of all $\delta \in \Delta$ such that $\pi(\delta) \in \Sigma_3 Y$ (notice that $\Sigma_3 Y \cap \Gamma$ is finite). Consider the definable set

$$D := (\Delta_0 + \Sigma_3 s(Y)) \cap \mathcal{H} \subseteq \langle \Delta + s(Y) \rangle \cap \mathcal{H} = \langle s(Y) \rangle \cap \mathcal{H}.$$

To prove that $D \ni 0$ generates $\langle s(Y) \rangle \cap \mathcal{H}$ is enough to prove the following:

Claim. For all n and for every $\delta_1, \dots, \delta_{2^n} \in \Delta$ and $y_1, \dots, y_{2^n} \in Y$, if $\Sigma_{i=1}^{2^n} \pi(\delta_i) + y_i = 0$ then $\Sigma_{i=1}^{2^n} \delta_i + s(y_i) \in \langle D \rangle$.

Indeed, granted the claim suppose that $\Sigma_{i=1}^m s(y_i) \in \langle s(Y) \rangle \cap \mathcal{H}$. Recall that the generator γ_1 of Γ belongs to Y and $s(\gamma_1) \in \Delta$, so we can write

$$\Sigma_{i=1}^m s(y_i) = \Sigma_{i=1}^m (0 + s(y_i)) + \Sigma_{i=m+1}^{2^m} (s(\gamma_1) - s(\gamma_1)).$$

Since $\Sigma_{i=1}^m s(y_i) \in \mathcal{H}$ we have that $\Sigma_{i=1}^m y_i = 0$ and therefore we are in the hypothesis of the claim, so we deduce that $\Sigma_{i=1}^m s(y_i) \in \langle D \rangle$, as required.

Proof. By induction on n . The case $n = 0$ gives $\pi(\delta_1) + y_1 = 0$, hence $\pi(\delta_1) \in Y \subseteq \Sigma_3 Y$, so $\delta_1 \in \Delta_0$. Therefore $\delta_1 + s(y_1) \in D$.

Assume now that $\Sigma_{i=1}^{2^n} \pi(\delta_i) + y_i = 0$ in \mathcal{V} . We want to show that $\Sigma_{i=1}^{2^n} \delta_i + s(y_i)$ is in $\langle D \rangle$. We write the sum in pairs:

$$\Sigma_{i=1}^{2^n} (\delta_i + s(y_i)) = \Sigma_{k=1}^{2^{n-1}} (s(y_{2k-1}) + s(y_{2k}) + \delta_{2k-1} + \delta_{2k}).$$

Now, because $Y + \Gamma = \mathcal{V}$, for each $k = 1, \dots, 2^{n-1}$ there is $w_k \in Y$ and $\beta_k \in \Gamma$ such that $y_{2k-1} + y_{2k} = \beta_k + w_k$. Let $\alpha_k \in \Delta$ be such that $\pi(\alpha_k) = \beta_k$. Note that $\beta_k \in \Sigma_3 Y$, so that $\alpha_k \in \Delta_0$. Hence,

$$(s(y_{2k-1}) + s(y_{2k}) - \alpha_k - s(w_k)) \in D.$$

Thus the above sum also equals:

$$\begin{aligned} \Sigma_{i=1}^{2^n} (\delta_i + s(y_i)) &= \Sigma_{k=1}^{2^{n-1}} (s(y_{2k-1}) + s(y_{2k}) - \alpha_k - s(w_k)) + \\ &\quad \Sigma_{k=1}^{2^{n-1}} (\delta_{2k-1} + \delta_{2k} + \alpha_k + s(w_k)). \end{aligned}$$

We already showed that $\Sigma_{k=1}^{2^{n-1}} (s(y_{2k-1}) + s(y_{2k}) - \alpha_k - s(w_k)) \in \langle D \rangle$, so if we denote $\tilde{\delta}_k = \delta_{2k-1} + \delta_{2k} + \alpha_k \in \Delta$ then it remains to prove that

$$\Sigma_{k=1}^{2^{n-1}} (\tilde{\delta}_k + s(w_k)) \in \langle D \rangle.$$

Indeed, we have that

$$\Sigma_{k=1}^{2^{n-1}} (\pi(\tilde{\delta}_k) + w_k) = -\Sigma_{i=1}^{2^n} (\pi(\delta_i) + y_i) - \Sigma_{k=1}^{2^{n-1}} (y_{2k-1} + y_{2k} - \beta_k - w_k) = 0$$

and therefore by induction we deduce that $\Sigma_{k=1}^{2^{n-1}} (\tilde{\delta}_k + s(w_k)) \in \langle D \rangle$, as required. \square

Proposition 2.2. *With \mathcal{H} , \mathcal{G} and \mathcal{V} as above, assume that $X \subseteq \mathcal{G}$ is a definable set with $\langle \pi(X) \rangle = \mathcal{V}$. Then $\langle X \rangle \cap \mathcal{H}$ is definably generated.*

Proof. Since \mathcal{V} admits a lattice it contains a generic set Y . Without loss we can assume that $\Sigma_2 \pi(X) \subseteq Y$. By saturation $Y \subseteq \Sigma_n \pi(X)$ for some n and therefore by definable choice we can pick a section $s : Y \rightarrow \langle X \rangle$. Moreover, we can assume that $s(\Sigma_2 \pi(X)) \subseteq \Sigma_2 X$. By Proposition 2.1 we have that $\mathcal{H}_0 = \langle s(Y) \rangle \cap \mathcal{H}$ is definably generated. Let $E = \Sigma_4 X \cap \mathcal{H}$. Therefore to prove that $\langle X \rangle \cap \mathcal{H}$ is definably generated it suffices to show that $\langle X \rangle \cap \mathcal{H} = \langle E \rangle + \mathcal{H}_0$.

We shall prove that for every $x_1, \dots, x_{2^n} \in X$, if $\Sigma_{i=1}^{2^n} x_i \in \mathcal{H}$ then $\Sigma_{i=1}^{2^n} x_i \in \langle E \rangle + \mathcal{H}_0$. When $n = 1$ then $x_1 + x_2 \in \Sigma_2 X \cap \mathcal{H} \subseteq \Sigma_4 X \cap \mathcal{H} = E$ so we are done. For $n > 1$, write

$$\Sigma_{i=1}^{2^n} x_i = \Sigma_{k=1}^{2^{n-1}} (x_{2k-1} + x_{2k} - s(\pi(x_{2k-1} + x_{2k}))) + \Sigma_{k=1}^{2^{n-1}} s((\pi(x_{2k-1} + x_{2k}))).$$

Each

$$(x_{2k-1} + x_{2k} - s(\pi(x_{2k-1} + x_{2k}))) \in (\Sigma_2 X + s(\pi(\Sigma_2 X))) \cap \mathcal{H} \subseteq E$$

so the sum on the left is in $\langle E \rangle$. Finally, for the sum on the right we have that

$$\begin{aligned} \Sigma_{k=1}^{2^{n-1}} s((\pi(x_{2k-1} + x_{2k}))) &= \Sigma_{i=1}^{2^n} x_i \\ &\quad - \Sigma_{k=1}^{2^{n-1}} (x_{2k-1} + x_{2k} - s(\pi(x_{2k-1} + x_{2k}))) \in \mathcal{H} + \mathcal{H} = \mathcal{H} \end{aligned}$$

so that $\sum_{k=1}^{2^{n-1}} s((\pi(x_{2k-1} + x_{2k})) \in \langle s(Y) \rangle \cap \mathcal{H} = \mathcal{H}_0$, which finishes the proof. \square

Before the main corollary we need also:

Lemma 2.3. *Let \mathcal{H} be an abelian locally definable group. If \mathcal{H} is definably generated then its connected component is definably generated by a connected definable set. In particular, if every connected definably generated subgroup of \mathcal{H} contains a generic set then every definably generated subgroup of \mathcal{H} contains a generic set.*

Proof. Let $X \subseteq \mathcal{H}$ be a definable set which generates \mathcal{H} . Let X_1, \dots, X_k be its connected components. Fix an element a_i in each X_i , and let $\Gamma = \langle a_1, \dots, a_k \rangle$. Consider the connected set $\tilde{X} = \bigcup X_i - a_i$, and notice that $\langle X \rangle = \langle \tilde{X} \rangle + \Gamma$. Since $\langle \tilde{X} \rangle$ is a locally definable subgroup of \mathcal{H} of bounded index, it must be its connected component.

For the second part of the statement, let \mathcal{G} be a definably generated subgroup of \mathcal{H} . Then its connected component \mathcal{G}^0 is definably generated by the first part and therefore by hypothesis it contains a generic set Y , that is, there is a bounded $A \subseteq \mathcal{G}^0$ such that $A + X = \mathcal{G}^0$. Since \mathcal{G}^0 has bounded index in \mathcal{G} , there is a bounded $B \subseteq \mathcal{G}$ such that $B + \mathcal{G}^0 = \mathcal{G}$. In particular $A + B + X = \mathcal{G}$, as required. \square

Theorem 2.4. *Assume that we are given an exact sequence of abelian locally definable groups and maps.*

$$0 \longrightarrow \mathcal{H} \xrightarrow{i} \mathcal{G} \xrightarrow{\pi} \mathcal{V} \longrightarrow 0$$

Assume also that definably generated connected subgroups of \mathcal{V} and of \mathcal{H} contain a generic set. Then the same is true for \mathcal{G} .

Proof. Let $X \subseteq \mathcal{G}$ be a definable set which generates a connected subgroup of \mathcal{G} . Since $\pi(\langle X \rangle) = \langle \pi(X) \rangle$ is a definably generated connected group, we have the exact sequence of locally definable groups

$$0 \rightarrow \langle X \rangle \cap \mathcal{H} \rightarrow \langle X \rangle \rightarrow \langle \pi(X) \rangle \rightarrow 0$$

By hypothesis the connected group $\langle \pi(X) \rangle$ contains a generic set, that is, there exists a definable set $Z_1 \subseteq \langle X \rangle$ such that $\pi(Z_1)$ is generic in $\langle \pi(X) \rangle$. In particular the group $\langle \pi(X) \rangle$ admits a lattice (see Fact 1.4) and therefore by Proposition 2.2 the group $\langle X \rangle \cap \mathcal{H}$ is definably generated. Again by hypothesis and by Lemma 2.3 we have that $\langle X \rangle \cap \mathcal{H}$ contains a generic set Z_2 . Finally, it is not hard to see that $Z_1 + Z_2$ is generic in $\langle X \rangle$. \square

Next we apply the above result in different contexts. First, we can study definably generated subgroups of abelian torsion-free definable groups (see basic facts on torsion-free groups definable in o-minimal structures in Section 2.1 in [14]).

Corollary 2.5. *Let G be any definable, abelian torsion-free definable group. Then for any definable $X \subseteq G$, the group $\langle X \rangle$ contains a generic set.*

Proof. We prove it by induction on $\dim(G)$. Assume first that $\dim(G) = 1$. Then there is a definable order relation $<$ on G such that G with $<$ is a definable ordered group. By Lemma 2.3 it suffices to study a subgroup generated by a set of the form $(-b, b) := \{x \in G : -b < x < b\}$, that is,

$$\langle(-b, b)\rangle = \bigcup_{n \in \mathbb{N}} (-nb, nb)$$

But this follows straightforward since the zero-dimensional locally definable subgroup $\Gamma = \mathbb{Z}b$ of $\langle(-b, b)\rangle$ is a lattice because $\langle(-b, b)\rangle/\Gamma$ is isomorphic to the definable group $([0, b), \text{mod } b)$.

Now, assume that $\dim(G) > 1$. Then, by [16], there exists a subgroup H of G of dimension 1. In particular, we have the exact sequence

$$0 \rightarrow H \rightarrow G \rightarrow G/H \rightarrow 0$$

Since both H and G/H are abelian torsion-free definable groups, by induction we have that definably generated connected subgroups of G/H and of H contain a generic set. Thus by Theorem 2.4 and Lemma 2.3 we deduce that definably generated subgroups of G contain a generic set, as required. \square

Next, we study algebraic groups defined without parameters (but note that this result is only a particular case of Theorem 4.4).

Corollary 2.6. *Let \mathcal{R} be an expansion of a model of $T_{\text{an}, \text{exp}}$. Let $H \subseteq R^k$ be an abelian connected algebraic group defined without parameters. Let $X \subseteq H(R)^\circ$ be a definable connected and definably compact definable set in \mathcal{R} , possibly with parameters. The subgroup of H generated by X has a generic definable subset.*

Proof. Denote $G := H(\mathbb{R})^\circ$. By [10, Lemma 4.10] we have that

$$G(\mathbb{R}) = H_1(\mathbb{R}) \times H_2(\mathbb{R}) \times K$$

where $H_1(\mathbb{R})$ and $H_2(\mathbb{R})$ are semialgebraic subgroups of $G(\mathbb{R})$ semialgebraically isomorphic to $(\mathbb{R}, +)^s$ and $(\mathbb{R}^*, \cdot)^\ell$ respectively, and K is a closed and compact real analytic group. Since K is compact it is definable in $T_{\text{an}, \text{exp}}$ and therefore we can transfer the equality above to \mathcal{R} . Moreover, the subgroup K is analytically isomorphic to a torus $\mathbb{T}(\mathbb{R}) = (\mathbb{R}/\mathbb{Z})^p$, and $H_1(\mathbb{R})$ and $H_2(\mathbb{R})$ can be identified via the exponentiation. All in all, we have definable in \mathcal{R} an isomorphism

$$G = R^m \times \mathbb{T}.$$

Now, consider the definable universal covering $\pi : \tilde{G} \rightarrow G$, that is,

$$\tilde{G} = R^m \times \text{Fin}(R)^p,$$

where $\text{Fin}(R) := \{x \in R : |x| < n \text{ for some } n \in \mathbb{N}\}$.

By definable choice there is $S \subseteq \tilde{G}$ such that $\pi(S) = X$. Moreover, the group generated by S is in particular a definably generated subgroups of R^{m+p} and therefore by Corollary 2.5 we have that $\langle S \rangle$ contains a generic

set. In particular, the group generated by X contains a generic set, as required. \square

3. ABELIAN VARIETIES

For a positive $g \in \mathbb{N}$, by a *complex g -torus* we mean the quotient group \mathbb{C}^g/Λ where Λ is a lattice. It is a compact complex Lie group of dimension g . A torus \mathbb{C}^g/Λ is called an *abelian variety* if it is biholomorphic with a projective variety in $\mathbb{P}^k(\mathbb{C})$ for some k .

Let us denote by \mathbb{H}_g the set of $g \times g$ symmetric matrices with a positive definite imaginary part. Then the well-known *Riemann criterion* establishes that a complex g -torus E is an abelian variety if and only if it is bi-holomorphic with a torus $\mathbb{C}^g/(\tau\mathbb{Z}^g + D\mathbb{Z}^g)$ where $\tau \in \mathbb{H}_g$ and D is a diagonal matrix $D = \text{diag}(d_1, \dots, d_g)$ with positive integers $d_1|d_2|\dots|d_g$. We call D a *polarization type* of E .

Note that if E has polarization D , then it also has polarization kD for any positive integer k . Indeed, the map $\mathbb{C}^g/(\tau\mathbb{Z}^g + D\mathbb{Z}^g) \rightarrow \mathbb{C}^g/(k\tau\mathbb{Z}^g + kD\mathbb{Z}^g) : z \mapsto kz$ is a biholomorphism and therefore being biholomorphic with $\mathbb{C}^g/(\tau\mathbb{Z}^g + D\mathbb{Z}^g)$ is equivalent to being biholomorphic with $\mathbb{C}^g/(k\tau\mathbb{Z}^g + kD\mathbb{Z}^g)$. In particular, by taking $k = 6$ above, any abelian variety has a polarization type $D = \text{diag}(d_1, \dots, d_g)$ satisfying $d_1 \geq 4$, $2|d_1$ and $3|d_1$. If a polarization type satisfies the latter we call it a *Baily-polarization*.

Let us show that the family of abelian varieties with a fixed Baily-polarization is constructible.

First, let us recall the definition of theta functions. In some occasions, we will identify the symmetric $g \times g$ matrices with \mathbb{C}^n for $n := \frac{g(g+1)}{2}$, and therefore we will view \mathbb{H}_g as a subset of \mathbb{C}^n . The well-defined function

$$\begin{aligned} \vartheta : \mathbb{C}^g \times \mathbb{H}_g &\rightarrow \mathbb{C} \\ (z, \tau) &\mapsto \sum_{n \in \mathbb{Z}^g} \exp(\pi i({}^t n \tau n) + 2{}^t n z) \end{aligned}$$

is holomorphic and \mathbb{Z}^g -periodic in z and $(2\mathbb{Z})^n$ -periodic in τ . For any $a \in \mathbb{R}^g$ the associated *Riemann Theta function* is

$$\begin{aligned} \vartheta_a : \mathbb{C}^g \times \mathbb{H}_g &\rightarrow \mathbb{C} \\ (z, \tau) &\mapsto \vartheta_a(z, \tau) = \exp(\pi i({}^t a \tau a) + 2{}^t a z) \vartheta(z + \tau a, \tau). \end{aligned}$$

A consequence of the classical Lefschetz Theorem is the following. Fix a polarization $D = \text{diag}(d_1, \dots, d_g)$ with $d_1 \geq 3$ and fix a set of representatives $\{c_0, \dots, c_N\}$ of the cosets of \mathbb{Z}^g in the group $D^{-1}\mathbb{Z}^g$. Then

$$\varphi^D : \mathbb{C}^g \times \mathbb{H}_g \rightarrow \mathbb{P}^N(\mathbb{C})$$

$$\varphi^D(z, \tau) = (\vartheta_{c_0}(z, \tau) : \dots : \vartheta_{c_N}(z, \tau))$$

is a well-defined holomorphic map, and given $\tau \in \mathbb{H}_g$ we have that

$$\begin{aligned} \varphi_\tau^D : \mathbb{C}^g &\rightarrow \mathbb{P}^N(\mathbb{C}) \\ z &\mapsto \varphi_\tau^D(z) := \varphi^D(z, \tau) \end{aligned}$$

is an $(\tau\mathbb{Z}^g + D\mathbb{Z}^g)$ -periodic immersion and induces an analytic embedding of the abelian variety $\mathcal{E}_\tau^D = \mathbb{C}^g/(\tau\mathbb{Z}^g + D\mathbb{Z}^g)$ into $\mathbb{P}^N(\mathbb{C})$.

Moreover, if we denote

$$\Psi^D : \mathbb{H}_g \rightarrow \mathbb{P}^N(\mathbb{C}) : \tau \mapsto \varphi^D(0, \tau),$$

and

$$\Phi^D : \mathbb{C}^g \times \mathbb{H}_g \rightarrow \mathbb{P}^N(\mathbb{C}) \times \mathbb{P}^N(\mathbb{C}) : (z, \tau) \mapsto (\varphi^D(z, \tau), \Psi^D(\tau))$$

then we have the following properties for Baily-polarizations, see [3, Theorem 8.10.1. and Remark 8.10.4].

Fact 3.1. *Let D be a Baily-polarization.*

- 1) *The map Ψ^D is an immersion, the set $\Psi^D(\mathbb{H}_g)$ is a Zariski open subset of an algebraic subvariety of $\mathbb{P}^N(\mathbb{C})$. Moreover, if $\Psi^D(\tau) = \Psi^D(\tau')$ then*

$$\varphi_\tau^D(\mathcal{E}_\tau^D) = \varphi_{\tau'}^D(\mathcal{E}_{\tau'}^D).$$

- 2) *The image of Φ^D is a Zariski open subset of an algebraic subvariety of $\mathbb{P}^N(\mathbb{C}) \times \mathbb{P}^N(\mathbb{C})$.*

Remark 3.2. In other words, if we denote by $P^D := \Psi^D(\mathbb{H}_g)$ and for each $p := \Psi^D(\tau) \in P^D$ we denote $A_p^D := \varphi_\tau^D(\mathcal{E}_\tau^D)$, then the above result ensures that the family

$$\mathcal{A}^D := \{A_p^D : p \in P^D\}$$

is constructible.

Let \mathcal{M} be a first order structure. Let A , B and T be definable sets in a first order structure \mathcal{M} . We say that the family of functions $\{f_t : A \rightarrow B : t \in T\}$ is definable if the family $\{(a, b, t) : t \in T, b = f_t(a)\}$ is a definable set. If \mathcal{N} is an elementary extension of \mathcal{M} and X is a definable subset of M^n then we denote by $X(N)$ the realization of X in \mathcal{N} .

Let K be an algebraic closed field. Henceforth, when we write that $\mathcal{A} = \{A_t : t \in T\}$ is a constructible family of g -dimensional abelian varieties of $\mathbb{P}^N(K)$, we mean that each A_t is an irreducible projective subvariety of $\mathbb{P}^N(K)$ of dimension g and that there is a constructible family $\{F_t : t \in T\}$ of regular maps $F_t : A_t \times A_t \rightarrow A_t$ such that each F_t endows A_t with a group structure.

Proposition 3.3. *Let $\mathcal{A} = \{A_t : t \in T\}$ be a constructible family without parameters of g -dimensional abelian varieties of $\mathbb{P}^M(\mathbb{C})$. Then there exist finitely many polarizations D_1, \dots, D_k and $d \in \mathbb{N}$ such that for any $t \in T$ there exists an isomorphism of degree less than d between A_t and an abelian variety in \mathcal{A}^{D_j} for some $j = 1, \dots, k$.*

Proof. We already note above that for each $t \in T$ the abelian variety A_t has a Baily-polarization D and therefore it is bi-holomorphic isomorphic to an abelian variety in \mathcal{A}^D , so in particular they are bi-regularly isomorphic,

by Chow's Theorem. We claim that only finitely many Bayli-polarizations D_1, \dots, D_k are needed to cover all the abelian varieties in \mathcal{F} .

Indeed, assume that it is not true. For any fixed Bayli-polarization D and for any $d \in \mathbb{N}$ consider the formula $F_d^D(t)$ in the language of rings that says that $t \in T$ and for every $p \in P^D$ there is no bi-regular isomorphism from A_t to A_p^D of degree d . Now, consider

$$q(t) = \{F_d^D(t) : D \text{ a Bayli-polarization \& } d \in \mathbb{N}\}.$$

By our assumption, we get that q is a partial type over a countable set of parameters. On the other hand, the complex field \mathbb{C} in the language of rings is \aleph_1 -saturated. For, any ω -stable complete theory in a countable language has a saturated model of cardinality \aleph_1 (see [11, Thm. 6.5.4]). In particular, since ACF_0 is \aleph_1 -categorical, it follows that \mathbb{C} is \aleph_1 -saturated. Thus, there is $t_0 \in T$ that realises the type q . In other words, the abelian variety A_{t_0} is not isomorphic to A_p^D for any Bayli-polarization D and $p \in P^D$, which is a contradiction. It follows that there exist D_1, \dots, D_k Bayli-polarizations such that for any $t \in T$ we have that A_t is bi-regularly isomorphic to an abelian variety in \mathcal{A}^{D_j} for some $j \in \{1, \dots, k\}$. Moreover, we have that the degree of the isomorphisms is bounded. Otherwise, we would get a contradiction by considering the partial type

$$q(t) := \{F_d^{D_j}(t) : j = 1, \dots, k \text{ \& } d \in \mathbb{N}\},$$

as required. \square

Remark 3.4. Note that with the notation used in Proposition 3.3, for each $j \in \{1, \dots, k\}$ the set T_j of $t \in T$ such that there exists a bi-regular isomorphism of degree less than d between A_t and an abelian variety in \mathcal{A}^{D_j} is constructible without parameters. Indeed, fix $j \in \{1, \dots, k\}$. Given an abelian variety $A_p^{D_j} \subseteq \mathbb{P}^{N_j}(\mathbb{C})$ in \mathcal{A}^{D_j} and $t \in T$, the set $I_{t,p}$ of bi-regular isomorphisms of degree less than d from A_t to $A_p^{D_j}$ is clearly a constructible set. Moreover, the family $\{I_{t,p} : t \in T, p \in P^{D_j}\}$ is also constructible without parameters. In particular, the set $T_j = \{t \in T : \exists p \in P^{D_j}, I_{t,p} \neq \emptyset\}$ is constructible without parameters.

In the rest of the paper, we fix a real closed field R , and let $K = R(i)$ be its algebraic closure.

We will use the obvious identification of K^n with R^{2n} . We say that a subset of K^n is semialgebraic over $C \subseteq R$ if it is semialgebraic over C as a subset of R^{2n} . Note that if X is a constructible subset of K^n over $A \subseteq K$ then it is clearly semialgebraic over the real and imaginary parts of the elements in A . Finally, note that R^n can be identified with the real part of K^n , and that by elimination of quantifiers of the theory of real closed fields, a subset of R^n is semialgebraic in the usual sense if and only if it is semialgebraic as a subset of K^n .

Henceforth, given a semialgebraic subset T of K^n , when we write that $\mathcal{A} = \{A_t : t \in T\}$ is a semialgebraic family of g -dimensional abelian varieties of $\mathbb{P}^N(K)$, we mean that each A_t is an irreducible projective subvariety of $\mathbb{P}^N(K)$ of dimension g , that the family \mathcal{A} is semialgebraic in the obvious (complex) sense, that is, the set

$$\{(x, t) : t \in T, x \in A_t\}$$

is a semialgebraic subset of $\mathbb{P}^M(\mathbb{C}) \times T$, and that there is also a semialgebraic family $\{F_t : t \in T\}$ of regular maps $F_t : A_t \times A_t \rightarrow A_t$ such that each F_t endow A_t with a group structure. Note that it only makes sense to say that a semialgebraic family is defined over a real tuple.

Lemma 3.5. *Let $g, d \in \mathbb{N}$, let D be a polarization and $\mathcal{F} = \{A_t : t \in T\}$ be a semialgebraic family of subsets of $\mathbb{P}^M(\mathbb{C})$ such that each A_t is a g -dimensional abelian variety for which there exists a bi-regular isomorphism of degree less than d between A_t and an abelian variety in \mathcal{A}^D . Then there exists a semialgebraic family $\{g_t : t \in T\}$ of bi-regular isomorphisms g_t from A_t to an abelian variety in \mathcal{A}^D .*

Proof. As in Remark 3.4, given $t \in T$ and $p \in P^D$, denote by $I_{t,p}$ the non-empty constructible set of bi-regular isomorphisms of degree less than d from A_t to \mathcal{A}_p^D , and note that $\{I_{t,p} : t \in T, p \in P^D\}$ is a semialgebraic family. Thus, by definable Skolem functions of the theory of real closed fields, there is a semialgebraic map

$$p : T \rightarrow P^D$$

and for each $t \in T$ a bi-regular isomorphism

$$g_t : A_t \rightarrow A_{p(t)}^D$$

such that the family $\{g_t : t \in T\}$ is semialgebraic, as required. \square

Given a polarization D and $\tau \in \mathbb{H}_g$, we denote by $E_\tau^D \subseteq \mathbb{C}^g$ the fundamental parallelogram of \mathcal{E}_τ^D . The following fact follows from [15, Thm 8.10] (see also the comments above it), and recall that P^D denotes $\Psi^D(\mathbb{H}_g) \subseteq \mathbb{P}^N(\mathbb{C})$.

Fact 3.6. *Let D be a Baily-polarization. Then there is a set $S \subseteq \mathbb{H}_g$ such that*

$$\Psi^D|_S : S \rightarrow P^D$$

is a surjective map definable in $\mathbb{R}_{an,exp}$ and such that there is a family $\{h_\tau^D : \tau \in S\}$ definable in $\mathbb{R}_{an,exp}$ with

$$h_\tau^D : E_\tau^D \rightarrow \mathbb{P}^N(\mathbb{C})$$

an embedding of the abelian variety \mathcal{E}_τ^D into the projective space $\mathbb{P}^N(\mathbb{C})$.

Theorem 3.7. *Let $\mathcal{A} = \{A_t : t \in T\}$ be a semialgebraic family without parameters of g -dimensional abelian varieties of $\mathbb{P}^M(\mathbb{C})$. Let $d \in \mathbb{N}$ and D be a polarization such that for each $t \in T$ there exists a bi-regular isomorphism*

of degree less than d between A_t and an abelian variety in \mathcal{A}^D . Then there is definable in $\mathbb{R}_{an,exp}$ a family of analytic maps

$$\{h_t : t \in T\}$$

where $h_t : E_t \rightarrow A_t$ is such that E_t is a fundamental parallelogram in \mathbb{C}^g of an abelian variety \mathcal{E}_t and h_t induces an isomorphism between \mathcal{E}_t and A_t .

Proof. By Lemma 3.5 there is a semialgebraic map

$$\mathbf{p} : T \rightarrow P^D$$

and for each $t \in T$ a bi-regular isomorphism

$$g_t : A_t \rightarrow A_{\mathbf{p}(t)}^D$$

such that the family $\{g_t : t \in T\}$ is semialgebraic. On the other hand, pick S a definable set in $\mathbb{R}_{an,exp}$ as in Fact 3.6 and consider a definable section

$$\mathbf{s} : P^D \rightarrow S$$

of $\Psi^D|_S : S \rightarrow P^D$. Finally, define

$$h_t := h_{\mathbf{s}(\mathbf{p}(t))}^D \circ g_t^{-1}$$

where $h_\tau^D : E_\tau^D \rightarrow \mathbb{P}^N(\mathbb{C})$, $\tau \in S$, are the embeddings given by Fact 3.6. \square

We now prove that semialgebraic connected subsets of abelian varieties generate locally definable groups with a generic subset. We will need the following generalization of a result in [8]. Given a subset Y of \mathbb{R}^n we say that it is symmetric if $Y = -Y$. Recall that we write $Y(m)$ for the addition of $Y - Y$ to itself m times. If $0 \in Y$ then $Y \subseteq Y(m)$. Given a point $a \in \mathbb{R}^n$, we denote by $(-a, a)$ the open segment from $-a$ to a .

Definition 3.8. Let \mathcal{R} be an o-minimal expansion of a real closed field R . Let $\mathcal{G} = \{G_t : t \in T\}$ be a locally definable family of locally definable groups such that $G_t \subseteq R^\ell$ for some $\ell \in \mathbb{N}$. We say that \mathcal{G} has the *uniform generic property (UGP)* if for every definable family $\{Y_t : t \in T\}$ of closed, bounded, connected definable sets $Y_t \subseteq G_t$, there are $m, N, s \in \mathbb{N}$ such that for every $t \in T$ there exists a finite set $0 \in A_t \subseteq Y_t(s)$ of cardinality at most N such that $Y_t(m) + Y_t(m) \subseteq A_t + Y_t(m)$.

We say that a locally definable group G has the *UGP* if for any definable set T , the definable family $\{G : t \in T\}$ has the UGP.

Lemma 3.9. Let \mathcal{R} be an o-minimal expansion of a real closed field R . Let $\{\pi_t : \tilde{G}_t \rightarrow G_t : t \in T\}$ be a locally definable family of maps where $G_t \subseteq R^g$ is a definable group, $\tilde{G}_t \subseteq R^\ell$ is its universal covering and π_t is its covering map. Let $\{s_t : G_t \rightarrow \tilde{G}_t : t \in T\}$ be a locally definable family of sections of π_t which map bounded subsets to bounded subsets. If the family $\{\tilde{G}_t : t \in T\}$ has the UGP, then $\{G_t : t \in T\}$ also has the UGP.

Proof. Let $\{X_t : t \in T\}$ be a definable family of closed bounded connected subsets $X_t \subseteq G_t$. Denote $Z_t := s_t(X_t)$, so that $\{Z_t : t \in T\}$ is a definable family of bounded definable subsets of \tilde{G}_t such that $\pi_t(Z_t) = X_t$.

We can assume that the number of connected components of Z_t is k for all $t \in T$, say Z_t^1, \dots, Z_t^k . By definable choice, there are definable maps

$$a_i : T \rightarrow R^\ell$$

such that $a_i(t) \in Z_t^i$ for $i = 1, \dots, k$. Let Y_t be the topological closure of $\bigcup_{i=1}^k [Z_t^i - a_i(t)]$. Note that Y_t is connected, bounded, closed and definable in \mathcal{R} . Moreover, if we denote $B_t = \{\pm \pi_t(a_1(t)), \dots, \pm \pi_t(a_k(t))\}$ then clearly

$$\pi_t(Y_t) \subseteq X_t + B_t.$$

Indeed, $\pi(Y_t)$ is contained in $\overline{\pi(\bigcup_{i=1}^k [Z_t^i - a_i(t)])}$, which in turn is contained in the closed subset $X_t + B_t$. On the other hand, note also that

$$X_t \subseteq \pi(Y_t) + B_t.$$

For, pick $x \in X_t$. Then, there is y in Z_t such that $\pi_t(y) = x$. In particular, $y \in Z_t^i$ for some i , and so $y - a_i(t) \in Y_t$. Then

$$x = \pi_t(y) \in \pi_t(Y_t) + \pi_t(a_i(t)) \subseteq \pi(Y_t) + B_t,$$

as required.

By hypothesis there are $m, N, s \in \mathbb{N}$ and a finite set $0 \in A_t \subseteq Y_t(s)$ of cardinality at most N such that

$$Y_t(m) + Y_t(m) \subseteq A_t + Y_t(m).$$

Thus,

$$\begin{aligned} X_t(m) + X_t(m) &\subseteq [\pi_t(Y_t)(m) + \pi_t(Y_t)(m)] + [B_t(m) + B_t(m)] \\ &\subseteq \pi_t(A_t) + \pi_t(Y_t(m)) + [B_t(m) + B_t(m)] \\ &\subseteq \pi_t(A_t) + [X_t(m) + B_t(m)] + [B_t(m) + B_t(m)] \\ &\subseteq [\pi_t(A_t) + B_t(3m)] + X_t(m). \end{aligned}$$

The cardinal of the finite set $\pi_t(A_t) + B_t(3m)$ is bounded by a certain number N_0 for all $t \in T$. Moreover, since $\pi_t(A_t) \subseteq X_t(s)$ and $B_t \subseteq X_t$, we have that $\pi_t(A_t) + B_t(3m) \subseteq X_t(s+3m)$. Therefore, if we define $C_t := \pi_t(A_t) + B_t(3m)$ and we set $s_0 := s + 3m$, we get that for the finite subset C_t of $X_t(s_0)$ of cardinality at most $\leq N_0$ it holds that

$$X_t(m) + X_t(m) \subseteq C_t + X_t(m),$$

as required. \square

Lemma 3.10. *Let \mathcal{R} be an o-minimal structure whose universe is the real field. Then R^n has the UGP.*

Proof. Let $\{Y_t : t \in T\}$ be a family definable without parameters of bounded closed connected definable subsets of \mathbb{R}^n containing 0. We will prove that there are $m \in \mathbb{N}$ and, for every $t \in T$, a finite subset $A_t \ni 0$ of $Y_t(m)$ of cardinality less than $2n + 1$ such that

$$Y_t(m) + Y_t(m) \subseteq \Sigma_{4m}A_t + Y_t(m).$$

Assume first that Y_t is also symmetric and contains 0. Let Z_t denote the convex hull of Y_t , so by [8, Lemma 3.3 and its proof] the family $\{Z_t : t \in T\}$ is definable. Moreover, there is $\ell \in \mathbb{N}$ such that $Y_t \subseteq Z_t \subseteq Y_t(\ell)$ for all $t \in T$. Otherwise, there is an elementary extension \mathcal{R}' of \mathcal{R} and $t_0 \in T(\mathcal{R}')$ such that $Z_{t_0}(\mathcal{R}')$ is not contained in $Y_{t_0}(\mathcal{R}')(\ell)$ for all $\ell \in \mathbb{N}$, a contradiction with [8, Lem. 3.4].

Claim. *There is $m \in \mathbb{N}$ such for every $t \in T$ we have that, up to an isometry of \mathbb{R}^n , there are at most $k \leq n$ points $a_1, \dots, a_k \in Z_t(m)$ such that if we denote $I_i = (-a_i, a_i)$ then:*

- (1) *the segments I_1, \dots, I_k are linearly independent,*
- (2) *$Z_t \subseteq I_1 + \dots + I_k \subseteq Z_t(m)$.*

Proof of Claim. We prove it by induction. If $n = 1$ then fix $t_0 \in T$. Since Z_{t_0} is an interval, we can use a translation so that $Z_{t_0} = [-a, a]$, for some $a \in R$. In particular, $Z_{t_0} \subseteq (-2a, 2a) \subseteq Z_{t_0}(2)$, as required.

We prove for $n + 1$ assuming that it is true for n . Fix $t_0 \in T$. Consider all line segments contained in Z_{t_0} and let J_0 be such segment of maximal length (it exists by o-minimality and because Z_{t_0} is closed). Since we work in a field we may assume that J_0 is parallel to the x_{n+1} -coordinate and furthermore that $0 \in J_0$ divides it exactly into two equal parts. We can therefore write $J_0 = (a_{k+1}, a_{k+1})$ with $a_{k+1} \in Z_{t_0}$. Let π denote the projection onto the first n coordinates, and note that

$$\{\pi(Z_t) : t \in T\}$$

is a family of connected bounded symmetric subsets of \mathbb{R}^n . By induction, there is $N \in \mathbb{N}$ and there are points $a_1, \dots, a_k \in \pi(Z_{t_0})(N)$, $k \leq n$, such that (1) and (2) are true for the intervals $I_i = (-a_i, a_i)$. So if we denote $I = I_1 + \dots + I_k$ then $I \subseteq \pi(Z_{t_0})(N)$. Then by Claim after the Question in [8], we have that

$$Z_{t_0} \subseteq I + J_0 \subseteq Z_{t_0}(2N).$$

Finally, if we set $m := 2\ell N$, then it follows that $Y_{t_0} \subseteq I + J_0 \subseteq Y_{t_0}(m)$, as required. \square

Now, note that for each $t \in T$ we have that $\dim(Z_t(m))$ is the number of intervals of the Claim, and therefore we can assume that it is a constant k . Now, by definable Skolem functions and since conditions (1) and (2) are first-order definable, there are definable functions

$$a_1, \dots, a_k : T \rightarrow Z_t$$

such that for $I_{i,t} := (-a_i(t), a_i(t))$ we have that (1) and (2) hold true. Thus, if we define $I_t := I_{1,t} + \cdots + I_{k,t}$ then the family $\{I_t : t \in T\}$ is definable and $Z_t \subseteq I_t \subseteq Z_t(m)$. It is easy to prove by induction on k that

$$(*) \quad I_t + I_t = A_t + I_t$$

where A_t are the vertices of the box I_t . Note that the cardinality of $A_t \subseteq Z_t(m)$ is $2k \leq 2n$. Without loss, we can add 0 to A_t . and hence the cardinality is $\leq 2n + 1$. Finally, since $Y_t \subseteq Z_t \subseteq I_t$ it is easy to prove by induction on j that:

$$Y_t(j) \subseteq \Sigma_{2j-1} A_t + I_t.$$

On the other hand, since $I_t \subseteq Z_t(m) \subseteq Y_t(m\ell)$ and $0 \in A_t$, it follows that

$$Y_t(m\ell) + Y_t(m\ell) = Y_t(2m\ell) \subseteq \Sigma_{4m\ell-1} A_t + Y_t(m\ell) \subseteq \Sigma_{4m\ell} A_t + Y_t(m\ell),$$

and we are done.

In the general case in which Y_t is not necessarily symmetric, consider $\tilde{Y}_t := Y_t(1) = Y_t - Y_t$, which is definable, closed, connected symmetric and contains 0. Note that $\tilde{Y}_t(j) = Y_t(2j)$ for all $j \in \mathbb{N}$. By the previous case there is $m \in \mathbb{N}$ such that

$$\tilde{Y}_t(m) + \tilde{Y}_t(m) \subseteq \Sigma_{4m} A_t + \tilde{Y}_t(m)$$

and therefore $Y_t(2m) + Y_t(2m) \subseteq \Sigma_{4m} A_t + Y_t(2m)$. Since $0 \in A_t$ then $Y_t(2m) + Y_t(2m) \subseteq \Sigma_{8m} A_t + Y_t(2m)$ and we are done. \square

Proposition 3.11. *Let R be a real closed field and $K = R(i)$ its algebraic closure. Let $A \subseteq \mathbb{P}^N(K)$ be an abelian variety defined over R and let X be a semialgebraic subset of A . Then the group generated by X contains a generic semialgebraic subset.*

Proof. First, note that we can assume that X is connected by Lemma 2.3. Moreover, since the group generated by X is closed, we can replace X by its closure, so we can assume that X is closed.

Let $c \in K^\ell$ be a tuple of coefficients defining algebraically the irreducible projective subvariety A of $\mathbb{P}^N(K)$ and the regular group operation $A \times A \rightarrow A$. We can replace the parameter c by a tuple u of free variables and therefore we obtain (without parameters) a constructible family $\mathcal{A} = \{A_u : u \in U\}$ of irreducible projective subvarieties of $\mathbb{P}^N(K)$ of dimension $g := \dim(A)$, and a constructible family $\mathcal{F} = \{F_u : u \in U\}$ of regular maps such that F_u endow A_u with a group structure. Note that $c \in U$, for which we get $A_c = A$ and $F_c = F$. Consider the realization $\mathcal{A}(\mathbb{C})$ of \mathcal{A} in \mathbb{C} . By Proposition 3.3 we can assume that there exists $d \in \mathbb{N}$ and a polarization D such that for each $u \in U(\mathbb{C})$ there exists a bi-regular isomorphism of degree less than d between $A_u(\mathbb{C})$ and an abelian variety in \mathcal{A}^D .

Next, consider the semialgebraic subset X of A which is semialgebraically defined over the tuple $r \in R^\ell$. Again replace r by a tuple v of free variables, define $A_{(u,v)} := A_u$ and consider the semialgebraic set $X_{(u,v)}$ of $A_{(u,v)}$ obtained when v plays the role of the real parameter r . Let T be the set of

$t := (u, v) \in U \times R^\ell$ for which X_t is connected. Note that T is a semialgebraic subset of $U \times R^\ell \subseteq U \times K^\ell$ defined without parameters and that $(c, r) \in T$.

Consider the semialgebraic family $\mathcal{A}_0 = \{A_t : t \in T\}$ of g -dimensional abelian varieties of $\mathbb{P}^N(K)$. Again, take the realization $\mathcal{A}_0(\mathbb{C})$ of \mathcal{A}_0 in the complex numbers, and note that by Theorem 3.7 there is a definable family in $\mathbb{R}_{an,exp}$ of analytic maps

$$\{h_t : t \in T(\mathbb{C})\}$$

where $h_t : E_t \rightarrow A_t$ is such that E_t is a fundamental parallelogram in \mathbb{C}^g of an abelian variety \mathcal{E}_t , and h_t induces an isomorphism f_t between \mathcal{E}_t and A_t .

Henceforth, we will work over the complex numbers and we omit the script \mathbb{C} , for example, we write T instead of $T(\mathbb{C})$. For each $t \in T$, denote by π_t the projection of \mathbb{C}^g over \mathcal{E}_t , so that $\pi_t|_{E_t} \circ f_t = h_t$. Note that $(\pi|_{E_t})^{-1} : \mathcal{E}_t \rightarrow E_t$ is a definable section of π_t . Therefore by Lemmas 3.9 and 3.10, the definable family $\{\mathcal{E}_t : t \in T\}$ has the UGP. In particular, since $\{f_t : t \in T\}$ is a definable family of isomorphisms between \mathcal{E}_t and A_t , it follows that $\{A_t : t \in T\}$ has also the UGP. Thus, there are $m, N, s \in \mathbb{N}$ such that for every $t \in T$ there exists a finite set $0 \in A_t \subseteq X_t(s)$ of cardinality at most N such that $X_t(m) + X_t(m) \subseteq A_t + X_t(m)$. In particular, since $X_{(c,r)} = X$ and $A_{(c,r)} = A$, we obtain that $X(m)$ is a generic subset of the group generated by X , as desired. \square

4. SEMIALGEBRAICALLY GENERATED SUBGROUPS OF SEMIALGEBRAIC GROUPS

The main purpose of this section is to show that groups generated by a semialgebraic subset of a semialgebraic group contain a generic semialgebraic subset.

Lemma 4.1. *Let H be an irreducible abelian linear algebraic group defined over R . Then the subgroup generated by a connected semialgebraic subset of H contains a semialgebraic generic set.*

Proof. By [12, Corollary 17.19], the group H is isomorphic to $K^m \times (K^*)^n$ for some $m, n \in \mathbb{N}$ (but note that this isomorphism is defined over K). Thus, by Theorem 2.4 and Corollary 2.5 we can assume that $H = (K^*)^n$. By an inductive argument and Theorem 2.4 again, we may assume $H = K^*$. Next, consider the connected semialgebraic subgroup $R_{>0}$ of K^* , and note that the quotient $K^*/R_{>0}$ is semialgebraically isomorphic to

$$\mathbb{S}^1 = \left\{ \begin{pmatrix} x & -y \\ y & x \end{pmatrix} : x, y \in R, x^2 + y^2 = 1 \right\}.$$

Thus, we have the exact sequence

$$1 \rightarrow R_{>0} \rightarrow K^* \rightarrow \mathbb{S}^1 \rightarrow 1.$$

By Lemma 2.3, Theorem 2.4 and Corollary 2.5, it suffices to prove that every subgroup generated by a connected semialgebraic subset of \mathbb{S}^1 contains a semialgebraic generic set.

Denote by X a connected semialgebraic subset of \mathbb{S}^1 , say definable over $d \in R^\ell$, and note that X is bounded. If we replace d by some free-variables t then we obtain a semialgebraic family $\mathcal{X} := \{X_t : t \in T\}$ defined without parameters of connected semialgebraic bounded subsets of \mathbb{S}^1 . Consider the realization $\mathcal{X}(\mathbb{R})$ of this family in the real field. Note that the universal covering map of $\mathbb{S}^1(\mathbb{R})$,

$$\mathbb{R} \rightarrow \mathbb{S}^1 : t \mapsto \begin{pmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{pmatrix},$$

is locally definable in $\mathbb{R}_{an,exp}$. By Lemma 3.10, the group \mathbb{R}^n has the UGP, and therefore $\mathbb{S}^1(\mathbb{R})$ also has property the UGP by Lemma 3.9. In particular, there are $m, N, s \in \mathbb{N}$ such that for every $t \in T$ there is a finite subset $0 \in A_t \subseteq X_t(s)$ of cardinality at most N such that $X_t(m) + X_t(m) \subseteq A_t + X_t(m)$. This is a first-order property and therefore for our parameter d we obtain that $X(m)$ is a generic subset of $\langle X \rangle$, as required. \square

Theorem 4.2. *Let H be an irreducible algebraic group definable over R . Then the subgroup generated by a connected semialgebraic subset of H contains a semialgebraic generic set.*

Proof. By Chevalley theorem's, there is a linear group L and abelian variety A , both defined over R , such that

$$0 \rightarrow L \rightarrow H \rightarrow A \rightarrow 0.$$

Thus, by Corollary 2.4, Proposition 3.11 and Lemma 4.1, we get the result. \square

Note that if in the above result, if X is a subset of $H(R)^o$, then the group generated by X is contained in $H(R)^o$ and the result gives a semialgebraic generic subset (in the usual “real” sense) of $H(R)^o$.

Proposition 4.3. *Let G be an abelian definable group and let $p : \tilde{G} \rightarrow G$ be its universal covering. Assume that definably generated connected subgroups of \tilde{G} contain a definable generic set. Then the same is true for G .*

Proof. Let $s : G \rightarrow \tilde{G}$ be a definable section of p . Consider the definable subset $Z = s(X)$ of \tilde{G} and let Z_1, \dots, Z_k be its connected components. For each $i = 1, \dots, k$ take $a_i \in Z_i$ and consider the connected definable set $Y = \bigcup_{i=1}^k (Z_i - a_i)$ of \tilde{G} . As in the proof of Lemma 2.3, note that $\langle Y \rangle$ is the connected component of $\langle Z \rangle$. Since $p(\langle Y \rangle)$ is a connected compatible subgroup of $p(\langle Z \rangle)$, and $p(\langle Z \rangle) = \langle X \rangle$ is connected, we deduce that $p(\langle Y \rangle) = \langle X \rangle$. On the other hand, by hypothesis there is a definable subset E of $\langle Y \rangle$ which is generic. Thus, $p(E)$ is a generic definable subset of $\langle X \rangle$, as desired. \square

Theorem 4.4. *Let G be an abelian semialgebraic group and let X be a semialgebraic subset of G . Then the group generated by X contains a generic semialgebraic subset.*

Proof. By Lemma 2.3, we can assume that X is connected. By [4, Thm 1.2] the quotient of G by its maximal normal semialgebraic torsion-free subgroup $\mathcal{N}(G)$ is semialgebraically compact. Then by Corollary 2.4 and Corollary 2.5 we can assume that G is semialgebraically compact. By [3, Thm 7.2] there is an irreducible algebraic group H defined over R , an open connected locally semialgebraic subgroup \mathcal{W} of the \mathfrak{o} -minimal universal covering group $\widetilde{H(R)^o}$ of $H(R)^o$ and a locally semialgebraic homomorphism $\theta : \mathcal{W} \rightarrow G$ that is that \mathfrak{o} -minimal universal covering homomorphism of G .

Denote by $p : \widetilde{H(R)^o} \rightarrow H(R)^o$ the universal covering map. We have the exact sequence

$$0 \rightarrow \ker(p) \rightarrow \widetilde{H(R)^o} \rightarrow H(R)^o \rightarrow 0.$$

Note that $\ker(p)$ is discrete and therefore its only semialgebraically generated connected subgroup is the trivial one. Thus, by Corollary 2.4 and Theorem 4.2, we deduce that semialgebraically generated connected subgroups of $\widetilde{H^0(R)}$ contain a semialgebraic generic set. In particular, the same is true for \mathcal{W} , and by Proposition 4.3, for G , as required. \square

REFERENCES

- [1] E. Baro, E. Jaligot, M. Otero, *Commutators in groups definable in \mathfrak{o} -minimal structures*, Proc. Amer. Math. Soc. 140(10) (2012), 3629-3643.
- [2] A. Berarducci, M.J. Edmundo, and M.Mamino, *Discrete subgroups of locally definable groups*, Selecta Mathematica, pages 1-17, 2012.
- [3] E. Barriga, *Definably Compact groups definable in real closed fields. II.* arXiv:1705.07370.
- [4] A. Conversano, *A reduction to the compact case for groups definable in \mathfrak{o} -minimal structures*, Journal of Symbolic Logic, 79 (2014), 45-53.
- [5] M. Edmundo, *Locally definable groups in \mathfrak{o} -minimal structures*, J. Algebra, 301 (1), 194-223, 2006.
- [6] P.E. Eleftheriou and Y. Peterzil, *Lattices in locally definable subgroups of $\langle R^n, + \rangle$* , Notre Dame J. Form. Log. 54 (2013), no. 3-4, 449-461.
- [7] P.E. Eleftheriou and Y. Peterzil, *Definable quotients of locally definable groups*, Selecta Math. (N.S.) 18 (2012), no. 4, 885-903.
- [8] P.E. Eleftheriou and Y. Peterzil, *Lattices in Locally Definable Subgroups of $\langle R^n, + \rangle$* , Notre Dame J. Formal Logic (54), Number 3-4 (2013), 449-461.
- [9] P.E. Eleftheriou and M. Edmundo, *The universal covering homomorphism in \mathfrak{o} -minimal expansions of groups*, MLQ Math. Log. Q. 53 (2007), no. 6, 571-582.
- [10] E. Hrushovski, A. Pillay, *Groups definable in local fields and pseudo-finite fields*, Israel J. Math. 85 (1994), no. 1-3, 203-262.
- [11] D.Marker, *Model Theory : An Introduction*, Graduate Texts in Mathematics 217.
- [12] J.S. Milne, *Algebraic groups*, <https://www.jmilne.org/math/CourseNotes/iAG200.pdf>
- [13] Y. Peterzil and A. Pillay, *Generic sets in definably compact groups*, Fund. Math. 193 (2007), 153-170.
- [14] Y. Peterzil and S. Starchenko, *On torsion-free groups in \mathfrak{o} -minimal structures*. Illinois Journal of Mathematics, vol. 49 (2008), no. 4, pp. 1299-1321.

- [15] Y. Peterzil and S. Starchenko, *Definability of restricted theta functions and families of abelian varieties*. Duke Math. J. 162 (2013), no. 4, 731-765.
- [16] Y. Peterzil and C. Steinhorn, *Definable compactness and definable subgroups of o-minimal groups*, J. London Math. Soc., 59 (1999), pp. 769-786.

DEPARTAMENTO DE ÁLGEBRA, GEOMETRÍA Y TOPOLOGÍA, FACULTAD DE MATEMÁTICAS,
UNIVERSIDAD COMPLUTENSE DE MADRID, MADRID, SPAIN
E-mail address: `eliasbaro@pdi.ucm.es`

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF KONSTANZ, BOX
216, 78457 KONSTANZ, GERMANY
E-mail address: `panteleimon.eleftheriou@uni-konstanz.de`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HAIFA, HAIFA, ISRAEL
E-mail address: `kobi@math.haifa.ac.il`